



HAL
open science

Analysis of the biogeochemical and biogeophysical effects of cropland management changes to prioritize actions in a perspective of climate change mitigation

Eric Ceschia, Dominique Carrer, Morgan Ferlicoq, Gaétan Pique, Rémy Fieuzal, Raphaël Garisoain, Elias Azzi

► **To cite this version:**

Eric Ceschia, Dominique Carrer, Morgan Ferlicoq, Gaétan Pique, Rémy Fieuzal, et al.. Analysis of the biogeochemical and biogeophysical effects of cropland management changes to prioritize actions in a perspective of climate change mitigation. Biophysical and biogeochemical effects of nature based solutions workshop, Cland institute, May 2022, Online webinar, France. pp.1-30. hal-04222030

HAL Id: hal-04222030

<https://hal.inrae.fr/hal-04222030v1>

Submitted on 28 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Analysis of the biogeochemical and biogeophysical effects of cropland management changes to prioritize actions in a perspective of climate change mitigation

Eric Ceschia (INRAE/CESBIO)

And

Morgan Ferlicoq (CESBIO), Gaétan Pique (CESBIO), Dominique Carrer (CNRM),
Raphael Garisoain (CESBIO/CNRM), Elias Azzi (KTH), Remy Fieuzal (CESBIO)

Introduction

- First studies comparing biogeochemical and biogeophysical effects were on forest ecosystems (e.g. *Betts et al. 2000* ; *Rottenberg & Yakir 2010* ; *O'Halloran et al. 2011*)
- For cropland, during many decades, studies were either focussed on :
 - 1) Soil C storage and reduction of Green House Gases (GHG) emissions for climate mitigation,
 - 2) Or the effects of management practices on biogeophysical effects (e.g. RF_{α}) caused by changes in cropland management (e.g. *Genesio et al., 2012* ; *Davin et al. 2014* ; *Luyssaert et al., 2014*).
- To compare biogeochemical effects with the RF_{α} caused by cropland management changes, the latter had to be converted in CO₂-eq but stabilised methodologies to do so were missing,
- In recent years, though, methodological advances allowing to convert albedo effects in CO₂-eq raised awareness of the potential significant effects of RF_{α} on climate mitigation (see *Bright et al. 2015*).
- As a consequence, recent studies showed that for some management changes RF_{α} had impacts of the same order of magnitude than biogeochemical effects.

Introduction

In this presentation we will :

- First analyse the causes of surface albedo dynamics on croplands in order to identify land management changes that could contribute to climate change mitigation through both CDR and SRM approaches,
- Then we will compare short term and long terms biogeophysical and biogeochemical effects of some management changes at larger scale to analyse their direct and undirect effects on the net radiative forcing.

Various spatial and temporal scales of study

Gaillac (81)



Biogeophysical effects of cropping systems





Lamasquère (31)



Biogeophysical & biogeochemical effects on RFnet




Causes of fast surface albedo changes




European ICOS sites

 In situ measurements/Southwest France



MODIS

Surface albedo dynamics



RF α

Consequences of cropland management changes on biogeochemical & biogeophysical (mainly RF α) components of RFnet



Satellite data and /or modelling at European scale

What do local scale studies teach us ?

Methodology for in situ measurements



Dynamics of surface albedo :

① Daily weighted average albedo

Half-hourly measured albedo (CNR1) and weighted by incident solar radiation

② Radiative forcing equation. We choose a bare soil albedo (measured on each site) as a reference for croplands.

$$RF_{\alpha} \text{ (W.m}^{-2}\text{)} = - SW_{in} \times T_A \times \Delta \text{albedo}$$

$$T_A = \frac{SW_{IN}}{R_{TOA}}$$

$$\alpha_{\text{daily}} - \alpha_{\text{bare soil}}$$


③ Annual radiative forcing was calculated over a cropping year by using the dynamics of each terms of the previous equation.

if α increase, $FR_{\alpha} < 0$ (Eq. C sink)

if α decrease, $FR_{\alpha} > 0$ (Eq. C source)

④ Conversion in CO₂-eq based on AF method (*Betts et al. 2000*)

Albedo vs. cropland status (land cover)



The rapeseed suffered from November **drought and frost** that increased surface albedo because of leaf damage + snow.


$$LC : \alpha_{\text{residues}} \approx \alpha_{\text{crop}} \approx \alpha_{\text{s.regrowth}} > \alpha_{\text{bare soil}}$$

$$\alpha_{\text{bare soil}} = f(\text{soil humidity})$$

→ Dry soil albedo > wet soil albedo

$$\text{Crop type: } \alpha_{\text{rapeseed}} > \alpha_{\text{WW}} > \alpha_{\text{maize}} > \alpha_{\text{sunflower}}$$

How do crop development affects surface albedo ?




In general, surface albedo increases with the green plant area index (PAI) but the response is crop dependant;

- For winter wheat and rapeseed, PAI reaches its maximum at PAI_{max} ,
- For maize, the albedo response to PAI is less pronounced,
- For sunflower maximum albedo occurred before PAI_{max} .


How do crop development affects surface albedo ?

Crop phenology effect on surface albedo



Albedo dynamics differ according to crop species


Several years of measurements/crop species



RF α induced by cropland albedo dynamic in reference to bare soil


Illustrates the combined effect of albedo dynamics with those of Rg and TA

In situ measurements/Southwest France



- Soil coverage may contribute to a “cooling” albedo effect,
- Same observations at other European flux sites


European ICOS sites



Comparison of biogeophysical effects between cropping systems

Gaillac (France)

July 2016



- The two subplots are adjacent :
 - (Up) Agroecology practices since 5 years
 - (Down) in transition from conventional to agroecology practices


Depth	Agroecology		Transition	
	Corg	OM	Corg	OM
0 to 10	8.6 ± 0.4	14.9 ± 0.8	8.2 ± 1.0	14.3 ± 1.7
10 to 30	7.4 ± 0.4	12.8 ± 0.7	8.0 ± 0.9	13.9 ± 1.6
30 to 60	5.3 ± 0.4	9.1 ± 0.7	5.4 ± 0.5	9.4 ± 0.9
60 to 90	5.0 ± 0.3	8.7 ± 0.4	4.9 ± 0.3	8.4 ± 0.5

- Cover crop growing duration were about 6 to 9 months (common in our area).
- At the “agroecology “ site α were **always equal or higher** in spite of a higher top soil OM content because the soil was permanently covered by vegetation or crop residues.
- Punctually, we observed an increase in **LW radiation that overwhelmed the albedo effect** at the “agroecology “ site during summer at the beginning of CC development (not showed here).

Comparative in situ analysis of all RFnet components – bare soil vs cover crop



ICOS Lamasquère site



Measured variables :


- CO₂, N₂O, water & energy fluxes
- Soil temperature & humidity at 0-5 cm
- Soil heat fluxes
- Solar incident/reflected radiation (short & longwave)

Objectives :

- Difference in surface albedo and RF induced by cover crop (CC)
- Effect of CC on :
 - Surface IR radiations & soil temperature
 - Sensible heat fluxes (hot eddys at the surface)
 - Latent heat fluxes (evapotranspiration)
 - C and GHG budgets


Comparative in situ analysis – Radiative effects of cover crops

1. Shortwave (albedo) effect (RF_{α})




→ $\Delta\alpha$ causes a cooling effect

2. Longwave effects



→ Longwave effect $\approx RF_{\alpha}$
in term of intensity (not necessarily in term of cooling)


3. Soil temperature



→ Mean difference of 2.5°C
→ Likely **slowdown** in OM mineralisation (and consequences on soil CO₂/N₂O fluxes)


Comparative in situ analysis – Non Radiative effects of cover crops

Effects on latent and sensible heat fluxes




- ↑ evapotranspiration & ↓ sensible heat fluxes causes local surface climate cooling (Boucher et al., 2004) → Natural air conditioner !! ;-)
- But this effect is difficult to express in term of radiative forcing (Pielke et al., 2002), especially at local scale

Summarizing cover crop biophysical effects



Global effect on climate of CC is difficult to estimate (requires coupled surface-atmosphere modelling exercises) but local/regional effect on perceived temperature at the surface could be significant (Georgescu et al., 2011).

Effect of cover crops on the components of the GHG Budget + RFA



Calculations of GHG emissions from field operations are based on Ceschia et al. (2010)

- The **differences in C & GHG budgets** were mainly **caused by the C storage effect** (but short term effect → very depleted soil in OM) in spite of a low CC biomass production (2.2 t DM/ha) compared to mean regional figures (4 t DM/ha),
- Increase in N₂O emissions and GHG emissions from field operations were negligible,
- Albedo RF in CO₂-eq was calculated considering that CC would be maintained over the next 100 yrs
- Very low RFA because **CC was grown in late fall** with low TA and Rg (and destroyed in early December) → this effect would have been close to 10 times larger if cover crop had been grown till spring (common in our area ; see Ferlicog & Ceschia, 2015),

But is it appropriate to compare RFA in CO₂-eq with the C/GHG budget components? → It will be discussed at the end of the presentation.


What do studies at larger spatial and temporal scales teach us ?

Carbon storage effect of cover crops (vs bare soil) in time


Several studies tend to show that :

- the carbon storage effect of the CC could be limited in time : new equilibrium reached after 45-50 year,


Meta-analysis based on in-situ data
(Poeplau & Don, 2015)



STICS simulations in France
(Tribouillois et al., 2018)



DayCent simulations over Europe
(Lugato et al., 2020) : **red line**




GHG budget of cover crops (vs bare soil) in time


Several studies tend to show that :

- the carbon storage effect of the cover crops could be limited in time : new equilibrium reached after 45-50 year,
- N₂O emissions may decrease on the short term but then increase 15-50 years after cover crop introduction → [Adapt N fertilisation after cover crop destruction](#) → [integrated soil fertility management](#) (Guardia et al. 2019 ; MERCI Meth.)


In-situ data in Spain
(Guardia et al. 2019)



STICS simulations in France
(Tribouillois et al., 2018)



DayCent simulations over Europe
(Lugato et al., 2020) : orange line




GHG budget of cover crops (vs bare soil) in time


Several studies tend to show that :

- the carbon storage effect of the cover crops could be limited in time : new equilibrium reached after 45-50 year,
- N₂O emissions may decrease on the short term but then increase 15-50 years after cover crop introduction → [Adapt N fertilisation after cover crop destruction](#) → [integrated soil fertility management](#) (Guardia et al. 2019 ; MERCI Meth.)


In-situ data in Spain
(Guardia et al. 2019)



STICS simulations in France
(Tribouillois et al., 2018)




DayCent simulations over Europe
(Lugato et al., 2020) : red + orange




Analysis of the cover crop albedo effect (vs bare soil) over Europe


Carrer et al. (2018) in ERL




Desagregated vegetation index, bare soil albedo & vegetation albedo (snow free) derived from MODIS data at 5*5 km (Kalman filter ; Carrer et al., 2014) → albedo of C3-C4 crop rotation



Analysis of where and when cover crops are introduced



Daily albedo increase with cover crops




Daily global radiation & atmospheric transmittance (ERA-INTERIM)

RFCC
Radiative Forcing of Cover Crop


$$RF\alpha = - R_g \times TA \times \Delta\alpha$$

Radiative forcing ($W.m^{-2}$)



Analysis of the cover crop albedo effect (vs bare soil) over Europe



(Carrer et al. 2018)



- Conversion in CO₂-eq with the constant airborne fraction method, e.g. see *Betts et al. (2000)* (and with GWP method by *Myhre et al. 2013*)
- ← - 3 month duration cover crop scenario → the cumulative RFA over EU-28 is 3.2 (2.9) MtCO₂-eq.year⁻¹.
- Same but accounting for rain limitation → the cumulative RFA over EU-28 was 2.3 (2.1) MtCO₂-eq.year⁻¹
- 6 month duration cover crop scenario + rain limitation → the cumulative RFA over EU-28 was 4.3 (4.0) MtCO₂-eq.year⁻¹ i.e. a compensation of up to 1.0 (0.9)% of the EU-28 agricultural GHG emissions.

Analysis of the cover crop albedo effect (vs bare soil) over Europe

- In general the introduction of CC increase surface albedo compared to the bare soil (snow effect not accounted for) but for some soil types (e.g. calcisoils) with high albedo introducing CC could be counter productive.




➔ Remote sensing data are useful to identify where/when cover crops should be introduced (or not) in order to increase the current surface albedo (even better when high resolution products available)


Analysis of the cover crop albedo effect (vs bare soil) over Europe

However...

Pique et al.
(submitted to ERL)



Adapted from Tribouillois et al (2018) and considering Corg max similar to Romanian soils




Modelled bare soil albedo decrease takes into account the progressive incorporation of organic matters in the soil (in the whole soil profile while in reality OM accumulates first in the top soil)



Analysis of the cover crop albedo effect (vs bare soil) over Europe

However...

Gaétan Pique's PhD (paper in prep)



Adapted from Tribouillois et al (2018) and considering Corg max similar to Romanian soils





Modelled bare soil albedo decrease takes into account the progressive incorporation of organic matters in the soil (in the whole soil profile while in reality OM accumulates first in the top soil)




Same method as in Carrer et al. (2018) but over 50 years (current climate) + soil darkening (50, 80, 100% of C storage potential) + maximising soil coverage with CC (as in Pellerin et al. 2019)

Once cover crop are adopted, soil should be covered permanently to avoid this drawback (as in Gaillac). This can be achieved by different means...




t = 0 t = 50

RF α induced by soil coverage with crop residues vs ploughing



- Generalising this practice to the whole Europe could decrease air temperature during summer heat waves by $\approx 2^{\circ}\text{C}$,
- However most of the albedo cooling effect is lost : why ?
- The mulch effect reduces evaporation \rightarrow higher surface temperature,
- This change in surface energy partitioning increases sensible heat flux and thermal IR radiations (interact strongly with GHGs in the atmosphere).



\rightarrow Better cover the soil with CC. But in areas where CC cannot be grown during the fallow period (e.g. too dry, too cold), or in the interval between a crop and a cover crop, maintaining crop residues at the soil surface is to be encouraged (avoids soil darkening effect on albedo).

Discussions

- Other ecosystem services, trade offs and drawbacks of CC...see *Justes et al. (2012)*, *Kaye & Kemada (2017)*, *Pellerin et al. (2019)*, *Runk et al. (2020)*.
- They are still many things to investigate :
 - What is the potential increase in albedo cooling effect through the choice of CC species in the rotations and through varietal selection ?
 - Whats is the true effect of snow + CC ? More realistic approaches accounting for stand architecture/species, plant and snow height are needed ?
 - Whats are the CC effects on soil temperature/humidity → consequences for soil mineralisation, CO₂ and N₂O emissions ?
 - Consequences of CC on soil water retention & water resources for the following cash crop ?
 - What is the durability of the C stored in the soil by CC (climate change) ?
- Appart from CC and no till, what are the biogeophysical effects of other cropland management changes ?
 - For biochar application, see *Genesio et al. (2012)*,
 - What about agroforestry ?...

Discussions

What is/will be the net climatic effect of cover crops ?

Difficult to answer now !!!

Because :

- Mitigations based on soil C storage or reduced GHGs emissions (CDR) have a global diffused effect on temperature, since GHGs are well mixed in the atmosphere. On the contrary, biogeophysical effects trigger predominantly local variation in temperature + difficult to predict non-local effects due to teleconnection in the climate system (e.g. mediated by clouds, advection of heat, etc.) → the SRM effect caused by surface $\Delta\alpha$ (e.g. with cover crop), should not be considered as CO₂ accountable quotes equal to those generated by GHG reduction, but rather as an indication of the intensity and location of the albedo effect,
- Current Earth System Models do not have a sufficiently fine spatial resolution and detailed management schemes to represent local practices in a realistic way → makes the overall biogeochemical + biogeophysical effects of CC difficult to quantify for now. Most (if not all) IPCC models only have 2 crop PFTs (C3 & C4) for cropland... and none of those models account for CC...



Where the levers tested in the 2018 IPCC special report to define the pathways allowing to stay below 1.5 °C global warming by the end of the century the best ones ?

Conclusions

- We have analysed the causes of fast albedo changes for cropland at a range of cropland sites over Europe and identified solutions for climate change mitigation through nature based SRM approaches,
- In several studies, cover crops appear as the perfect solution for climate change mitigation as synergies between C storage effects, radiative effects (short and longwave), changes in energy partitioning (e.g. sensible/latent) are observed + many other ecosystem services at an acceptable cost for the farmer (+ CAP subsidies and C market),
- Also additional N₂O emissions caused by CC could be limited/neutralised through ISFM + GHG emissions associated to seeding/destruction are low compared to the C storage effect,
- However once C farming practices (e.g. Cover Crops) are introduced → permanent soil cover to avoid the soil darkening effect,
- Yet, the net mitigation effect (+ retroaction) of CC is unknown → must be addressed through coupled surface-atmosphere modelling exercises at global scale (including all biogeophysical and biogeochemical effects). At this point, it is not possible to do such exercises as Earth System models do not account for CC.

Key messages

- So yes, we should consider biogeochemical and biogeophysical effects to prioritize changes in cropland management in order to implement more efficient climate change mitigation strategies but difficult to compare directly those effects,
- It is urgent to reduce the gap between agronomists/soil scientists... and Earth System modellers to obtain a more realistic quantification of the true climatic effect of cropland management changes.
- One starting point to achieve this could be to assimilate higher resolutions satellite products in the ES models.

Many thanks for your attention !!!



H2020
ClieNFarm project