



HAL
open science

Experimental heat transfer measurement applied to CO₂ hydrate formation kinetic study

Véronique Osswald, Pascal Clain, Didier Dalmazzone, Anthony Delahaye,
Laurence Fournaison

► **To cite this version:**

Véronique Osswald, Pascal Clain, Didier Dalmazzone, Anthony Delahaye, Laurence Fournaison. Experimental heat transfer measurement applied to CO₂ hydrate formation kinetic study. ECGH2022 European Conference on Gas Hydrate, Jun 2022, Lyon, France. hal-04222135

HAL Id: hal-04222135

<https://hal.inrae.fr/hal-04222135v1>

Submitted on 29 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0
International License

Experimental heat transfer measurement applied to CO₂ hydrate formation kinetic study

Véronique OSSWALD¹, Pascal CLAIN^{1,2}, Didier DALMAZZONE³,
Anthony DELAHAYE¹, Laurence FOURNAISON¹

¹ Université Paris-Saclay, INRAE, FRISE, Antony, 92761, France, veronique.osswald@inrae.fr

² Léonard de Vinci Pôle Universitaire, Research Center, Paris la Défense, 92916, France

³ ENSTA-Paris, UCP, Palaiseau 91120, France

CO₂ hydrate slurry is a promising phase change material for secondary refrigeration cold distribution issues, due to its high latent heat (about 500 kJ.kg⁻¹ of water, higher than that of ice - 333 kJ.kg⁻¹) and wide melting temperature range suitable for air conditioning application. While the thermodynamic properties of CO₂ hydrates are now well established, knowledge of crystallization kinetics phenomena is still a challenge. Trying to understand and control the formation of gas hydrates is a key factor since their discovery in pipeline plugs by Hammerschmidt, but unlike that case, the goal here is to promote the rate of CO₂ hydrate formation. Most laboratory-scale hydrate reactors are equipped with pressure and temperature loggers, and mass balance on CO₂ allows formation kinetic studies, but with assumptions on CO₂ concentration in liquid phase in the liquid phase and the hydration number.

The present work investigates the kinetics of CO₂ hydrate crystallization for different types of stirrers and stirring speed conditions in a jacketed stirred batch reactor. Heat balance obtain experimentally using a specially developed sensor installed on the cooling jacket. The mass fraction of crystallized hydrate determine directly from the heat balance on the cooling jacket. First, tests on water heating and cooling steps and ice crystallization allowed the method validation before to apply it to CO₂ hydrate crystallization. Experimental results described with an empirical model and compared to other mass balance based kinetic determinations. Finally, the development of a simplified thermal model improved this kinetic determination by estimating heat flows inside the reactor and with the environment.

- Candidature to the best PhD poster presentation
- Candidature to the best Young Researcher oral presentation (PhD student or young researcher – PhD defended < 5 years)
- I want to deposit (end of June, I will be contacted by reviewers) a full document for reviewing and publication