



HAL
open science

Poplar rust never sleeps: evolutionary characterization of the first avirulence gene in the poplar rust fungus *Melampsora larici-populina*

Clémentine Louet, Méline Saubin, Sébastien Duplessis, Fabien Halkett, Pascal
Frey

► To cite this version:

Clémentine Louet, Méline Saubin, Sébastien Duplessis, Fabien Halkett, Pascal Frey. Poplar rust never sleeps: evolutionary characterization of the first avirulence gene in the poplar rust fungus *Melampsora larici-populina*. 12th International Congress of Plant Pathology, Aug 2023, Lyon, France. 2023. hal-04230809

HAL Id: hal-04230809

<https://hal.inrae.fr/hal-04230809v1>

Submitted on 6 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Reference: A44360PF

Category: 18.02 - Understanding emergence of pathogens in commercial and public forest ecosystems

Author contact: Frey Pascal (pascal.frey@inrae.fr)

Passage preference: Oral communication

POPLAR RUST NEVER SLEEPS: EVOLUTIONARY CHARACTERIZATION OF THE FIRST AVIRULENCE GENE IN THE POPLAR RUST FUNGUS MELAMPSORA LARICI-POPULINA

LOUET C. (1), SAUBIN M. (1), DUPLESSIS S. (1), HALKETT F. (1), **FREY P. (1)**

(1) INRAE, Nancy, FRANCE

Text

Poplar rust, caused by the pathogenic fungus *Melampsora larici-populina* (Basidiomycota, Pucciniales), is the main phytosanitary constraint for commercial poplar cultivation in Europe and other parts of the world. In the last 50 years, many rust-resistant cultivars were bred and released, but all the qualitative resistance genes released were overcome by pathogen evolution within a short period. In 1994, breakdown of the *RMIp7* resistance gene was detected in Belgium and Northern France. New virulent *M. larici-populina* individuals spread all over Western Europe in less than five years, causing very destructive epidemics, and lead to a complete replacement of the pathogen's populations. Through a genome-wide association study (GWAS), we identified a locus in the genome of *M. larici-populina*, that corresponds to the candidate avirulence gene *AvrMlp7*, whose mutation is responsible for *RMIp7* resistance breakdown. To further characterize this effector, we used a population genetics approach on a set of almost 300 individuals collected throughout a 28-year period encompassing the resistance breakdown event. Genotyping at the avirulence locus highlighted two different mutations: a non-synonymous mutation and a complete deletion of this locus. The temporal survey at the candidate locus revealed that both mutations pre-existed long before the breakdown, but at a heterozygous state. Using a reverse ecology approach, we identified the first candidate avirulence gene in this rust fungus.