



**HAL**  
open science

## Genetic variability of grapevine vegetative development parameters as described with LiDAR data and associated quantitative trait loci

Elsa Chedid, Komlan Avia, Vincent Dumas, Lionel Ley, Nicolas Reibel, Gisèle Butterlin, Maxime Soma, Raul Lopez-Lozano, Frédéric Baret, Didier Merdinoglu, et al.

### ► To cite this version:

Elsa Chedid, Komlan Avia, Vincent Dumas, Lionel Ley, Nicolas Reibel, et al.. Genetic variability of grapevine vegetative development parameters as described with LiDAR data and associated quantitative trait loci. XIII. International Symposium on Grapevine Breeding and Genetics, Institute for Grapevine Breeding Geilweilerhof, Jul 2022, Landau in der Pfalz, Germany. pp.108. hal-04231650

**HAL Id: hal-04231650**

**<https://hal.inrae.fr/hal-04231650>**

Submitted on 10 Oct 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## **P26 – Genetic variability of grapevine vegetative development parameters as described with LiDAR data and associated quantitative trait loci**

Chedid, Elsa<sup>1</sup>; Avia, Komlan<sup>1</sup>; Dumas, Vincent<sup>1</sup>; Ley, Lionel<sup>2</sup>; Reibel, Nicolas<sup>2</sup>; Butterlin, Gisèle<sup>1</sup>; Soma, Maxime<sup>4</sup>; Lopez-Lozano, Raul<sup>3</sup>; Baret, Frédéric<sup>3</sup>; Merdinoglu, Didier<sup>1</sup>; Duchêne, Éric<sup>1\*</sup>

<sup>1</sup>SVQV, INRAE-University of Strasbourg, Colmar, France

<sup>2</sup>UEAV, INRAE, Colmar, France

<sup>3</sup>EMMAH, INRAE-Avignon University, Avignon, France

<sup>4</sup>TETIS, INRAE, Montpellier, France

\*eric.duchene@inrae.fr

### **Abstract**

Estimating growth traits of grapevine plants is common practice to describe the effects of cultural practices, training systems, environmental conditions, or genotypic effects, both for rootstocks and scions varieties. Among these traits, the simple pruning weight is one of the most popular, while being time-consuming when hundreds of genotypes are to be characterized. To face the challenge of adaptation to climate change and the global demand for grapevine varieties resistant to disease, there is an increasing need for high throughput phenotyping methods. We studied in the vineyard the variability of plant vigor in the progeny of the cross between two genotypes carrying resistance genes for powdery and downy mildew. The pruning weight was measured for more than 200 genotypes and, in parallel, we used a newly developed high throughput LiDAR acquisition system to characterize the plants both in summer and in winter. We will present the LiDAR system and the comparison between LiDAR data, pruning weights and leaf area. Using “genotyping by sequencing” technology to describe the genetic heritage of the offspring, we performed quantitative trait loci (QTL) detection and identified several genomic regions associated with the genetic variability both for pruning weights; leaf area and LiDAR data.

**Keywords:** grapevine, pruning wood, leaf area, LiDAR, QTL