



HAL
open science

Genetic analysis of grapevine (*Vitis vinifera*) wood composition

Hadrien Chepca, Patricia Claudel

► **To cite this version:**

Hadrien Chepca, Patricia Claudel. Genetic analysis of grapevine (*Vitis vinifera*) wood composition. XIII. International Symposium on Grapevine Breeding and Genetics, Jul 2022, Landau in der Pfalz, Germany. Genetic analysis of grapevine (*Vitis vinifera*) wood composition. hal-04231659

HAL Id: hal-04231659

<https://hal.inrae.fr/hal-04231659v1>

Submitted on 8 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

P46 – Genetic analysis of grapevine (*Vitis vinifera*) wood composition

Chepca, Hadrien*; Avia, Komlan; Claudel, Patricia; Duchêne, Éric; Maia-Grondard, Alessandra; Hugueney, Philippe

¹Université de Strasbourg, INRAE, SVQV UMR-A 1131, Colmar, France

*hadrien.chepca@inrae.fr

Abstract

Grapevine woody stems represent a valuable source of polyphenols with a broad spectrum of health benefits. In contrast to the large number of studies about genetic determinants of grape berry composition, few works have investigated the genetic bases of grapevine wood composition.

In this work, we have used both targeted and non-targeted metabolomic approaches to perform a global metabolic quantitative trait loci (mQTL) analysis with a progeny from a cross between 'Riesling' (Ri) and 'Gewürztraminer' (Gw).

Analysis of woody canes extracts using high performance liquid chromatography coupled to mass spectrometry has revealed genetic variability in the Ri x Gw progeny for metabolites from different families such as stilbenoids and flavonoids. High-density genetic maps based on single nucleotide polymorphism (SNPs) allowed the detection of significant mQTLs for a large number of metabolites, including major polyphenols.

This poster presents information about the detected QTLs impacting the accumulation of major grapevine wood polyphenols. Furthermore, characterization of major determinants of grapevine wood composition may provide bases for a better understanding of sensitivity to wood diseases such grapevine trunk diseases.

Keywords: Grapevine, wood, metabolomics, genetics, QTLs