



HAL
open science

Inflammatory profile of AAV6 in a pig and human model

A Pereira, Jb Ducloyer, L Libeau, J Demilly, C Morival, M Guilbaud, E
Toublanc, F Roux, P Hulin, D Augereau, et al.

► To cite this version:

A Pereira, Jb Ducloyer, L Libeau, J Demilly, C Morival, et al.. Inflammatory profile of AAV6 in a pig and human model. ESGCT, Oct 2023, Bruxelles, Belgium. 2023. hal-04234863

HAL Id: hal-04234863

<https://hal.inrae.fr/hal-04234863>

Submitted on 10 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Inflammatory profile of AAV6 in a pig and human model

Pereira A¹, Ducloyer JB¹, Libeau L¹, Demilly J¹, Morival C¹, Guilbaud M¹, Toublanc E¹, Roux F^{2,3}, Hulin P⁴, Augereau D⁵, Jounier C⁵, Larcher T⁵, Mellet A¹, Weber M¹, Adjali O¹, Cronin T¹

¹Nantes University, CHU de Nantes, TaRGeT - Translational Research in Gene Therapy, INSERM UMR 1089, F-44200 Nantes, France

²Emergency Unit, Oniris, Nantes-Atlantic College of Veterinary Medicine, Food Science and Engineering, 44307 Nantes, France

³Nutrition, PathoPhysiology and Pharmacology (NP3) Unit, Oniris, Nantes-Atlantic College of Veterinary Medicine, Food Science and Engineering, 44307 Nantes, France

⁴Nantes Université, CHU Nantes, CNRS, Inserm, BioCore, US16, SFR Bonamy, F-44000 Nantes, France


⁵INRAE, Oniris, PANTher, APEX, Nantes, France.

Introduction / Background


Adverse effects related to inflammatory reactions compromise the safety and long-term efficacy of retinal gene therapy. AAV6 has previously shown promise for transduction of immune cells. We examined retinal immune cell activation following subretinal injection of AAV6, comparing it to the pan-retinal vector, AAV8 in Yucatan minipigs. We further compared the effects of these vectors on a culture of microglial cells derived from human iPSCs. An overt inflammatory response was found in 2 of 3 pigs injected with a moderate to high dose of AAV6-CAG.EGFP while pigs injected with AAV8-CAG.EGFP or PBS did not show clinical effects. A proliferation of invading phagocytic mononuclear cells were found in AAV-injected eyes. Cytokine secretion following AAV6 and AAV8 treatment of microglial cultures was subsequently evaluated.

Materials & Methods

Subretinal bilateral injection of rAAV (4,6e11 total vg/eye) was performed in the eyes of minipigs (6-month-old, 20kg):



A subconjunctival injection of 0.5ml of corticosteroid (methylprednisolone, 40mg/ml) was performed at the end of the procedure. Optical coherence tomography (OCT) and IHC on flat-mount retinas was used to assess outcome.



The *in vitro* study involved the use of microglial cells derived from human iPSCs tested with AAV at MOI of 1E5. RNA from cells were analysed by RT-qPCR and cytokine secretion into culture medium was tested by an electrochemiluminescence-based multiplex ELISA, MSD (Mesoscale Discover assay).

Conclusion

The results highlight the risk for retinal inflammation and the need to further study the serotype-specific response of microglial cells in rAAV-based gene therapy studies. The intracellular response of microglia to AAV transduction may result in a specific cytokine signalling cascade. While we find AAV6 expressing EGFP to be toxic in the pig retina, a pronounced cytokine secretion was not found *in vitro*.

Results

Retinal structure was assessed by OCT of eyes before injection and 1 month after injection. Iba1+ cells were counted on confocal slices of flat-mount retinas from the pig eyes in 6 random squares of 500µm side length in the injected zone (IZ), peripheral non-injected zone (NIZ) and macular NIZ.




Fig 1 Effects of AAV6 compared to AAV8 in a pig retina: Colour fundus photographs of retinas injected with AAV6-CAG.EGFP (A), AAV8-CAG.EGFP (B) show transgene spread while the structure for AAV6 (C) and AAV8 (D) are shown with a proliferation of fluorescent phagocytic cells (E). IBA1+ cells counted on retinal flat-mounts were more numerous in IZ of AAV6-injected pig (61.3 +/- 35.0) than AAV8-injected pig (mean 23.5 cells/square +/-sd 6.3) (G). SRI: subretinal injection




Fig 2 Validation of inducible human microglial (iMGL) culture model: cytokine secretion upon LPS induction show the expected profile.





Fig 3 Effect of AAV-CAG.EGFP treatment on iMGL culture model: cytokine secretion upon AAV treatment.



Note that...
No significant GFP expression in iMGL cells was observed for any AAV serotype tested!

Fig 4 Effect of AAV6-CAG.EGFP treatment on iMGL gene expression: cellular expression measured 24 hours after transfection and shown normalised to untransfected cells.