



HAL
open science

Using a combined stable isotope -speciation approach to understand the impact of long-term spreading of organic effluents on agricultural soils

A Pappoe, Z Fekiacova, E Doelsch

► To cite this version:

A Pappoe, Z Fekiacova, E Doelsch. Using a combined stable isotope -speciation approach to understand the impact of long-term spreading of organic effluents on agricultural soils. ICOTTE & ICHMET 2023, Sep 2023, Wuppertal, Germany. hal-04235040

HAL Id: hal-04235040

<https://hal.inrae.fr/hal-04235040>

Submitted on 10 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Using a combined stable isotope – speciation approach to understand the impact of long-term spreading of organic effluents on agricultural soils

A. Pappoe^{1,2}, Z. Fekiacova¹, E. Doelsch^{1,2}

¹ CEREGE, CNRS, Aix Marseille Univ, IRD, INRAE, Aix-en-Provence, France

² CIRAD, UPR Recyclage et risque, F-34398 Montpellier, France

Email : pappoe@cerege.fr

Recycling of organic waste (OW) as fertilizers on farmlands is a common practice as it provides a cost effective and sustainable way of managing OW. However, it represents a major source of contaminants such as copper (Cu) and zinc (Zn) that may pose potentially negative environmental impacts (Lavado et al., 2005).

To limit the environmental impact of OW spreading on farmlands, agronomical doses are calculated based on the nitrogen, phosphorus and potassium requirements of crops. Consequently, lower quantities of Cu and Zn are introduced in soil following OW application (López-Rayó et al., 2016). However, the long-term impacts of OW spreading at the set agronomical doses on the accumulation and fate of Cu and Zn in soils are scant. This study was designed to fill this gap by using a combination of Cu and Zn speciation characterization and isotopic compositions in OW and soils. We studied four soils with contrasting physicochemical properties (luvisols, nitisols, calcisols and arenosols) and the OW used in four long-term field experiments.

We observed, in agreement with published literature, that in OW, Cu and Zn speciation is affected by the OW treatment (anaerobic digestion and composting) and by the physicochemical conditions during storage. In the studied OW samples, Cu and Zn occurred mainly or entirely as sulfides in raw and anaerobically digested pig slurries whereas only oxidized species were present in composted household wastes (Zn-phosphate, Cu bound to organic matter).

In two field experiments selected for this study, little or no change in Cu and Zn speciation was observed between control and amended soil due to OW application at agronomical doses. Thus, the interest in analyzing the isotopic signatures to investigate whether Cu and Zn originating from OW application can be traced in such cases.

Keywords: copper, zinc, organic waste, agricultural recycling, spectroscopy, isotopes

References

- Lavado, R. S., Rodríguez, M. B., & Taboada, M. A. (2005). Treatment with biosolids affects soil availability and plant uptake of potentially toxic elements. *Agriculture, Ecosystems and Environment*, 109(3–4), 360–364. <https://doi.org/10.1016/j.agee.2005.03.010>
- López-Rayó, S., Laursen, K. H., Lekfeldt, J. D. S., Delle Grazie, F., & Magid, J. (2016). Long-term amendment of urban and animal wastes equivalent to more than 100 years of application had minimal effect on plant uptake of potentially toxic elements. *Agriculture, Ecosystems and Environment*, 231, 44–53. <https://doi.org/10.1016/j.agee.2016.06.019>