



**HAL**  
open science

# Taking into account soils and climate change in assessing the production potential of a legume crop of interest: pea

Annabelle Larmure, Jean-Marc Brayer, Mathieu Chanis, Éric Lucot,  
Jean-Claude Monnet, Thierry Castel, Marjorie Ubertosì

## ► To cite this version:

Annabelle Larmure, Jean-Marc Brayer, Mathieu Chanis, Éric Lucot, Jean-Claude Monnet, et al.. Taking into account soils and climate change in assessing the production potential of a legume crop of interest: pea. ARFAGRI Seminario Agroecología, Nov 2022, Rosario, Argentina. , 2022. hal-04250249

**HAL Id: hal-04250249**

**<https://hal.inrae.fr/hal-04250249>**

Submitted on 19 Oct 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Taking into account soils and climate change in assessing the production potential of a legume crop of interest: pea



Seminario Agroecología  
Rosario, 22-23 nov. 2022

Larmure A.<sup>1</sup>, Brayer J.<sup>1</sup>, Chanis M.<sup>1</sup>, Lucot E.<sup>2</sup>,  
Monnet J-C.<sup>2</sup>, Castel T.<sup>3</sup>, Ubertosi M.<sup>1</sup>

<sup>1</sup>Agroécologie, Institut Agro Dijon, INRAE, Univ. Bourgogne, France.

<sup>2</sup>UMR 6249 Chrono-Environnement CNRS/Université Franche-Comté, France

<sup>3</sup>UMR 6282 Biogéosciences CNRS/ Université Bourgogne Franche-Comté - Institut Agro Dijon, France




## Context

Climate change and the increasing pressure on resources are challenging the mode of agricultural production. To meet the challenge of more autonomous, efficient and sustainable protein production, it is necessary to acquire knowledge on the evolution of the protein crops production potential.

Pea (*Pisum sativum* L) is an annual seed legume cultivated for its high protein content and for its environmental services (e.g. symbiotic fixation of atmospheric N<sub>2</sub>)

## Processing of "SOIL" semantic and cartographic data necessary to simulate pea production potential at an agricultural scale in the Burgundy Franche-Comté county (France)

1 The majority Soil Type Unit (STU) (excluding forest) is assigned to each SMU



Map of soil mapping units (SMU) in Burgundy Franche-Comté: soil available water capacity (AWC) of the majority STU

AWC at pF 1,5: Available water capacity (mm) (Bruand *et al.* 2004)

Bd: Bulk density (Keller *et al.* 2010)


Soil\_d: Soil depth (cm)

CaCO<sub>3</sub>: Total calcium carbonate content (%)

Arg: Clay content (%)

Corg: Organic carbon content (%)


2 The majority Soil Map Unit - SMU (excluding forest and urban areas) is assigned to each 8 km x 8 km grid cell



Majority land use map by 8 km x 8 km grid cell in Burgundy Franche-Comté: (data Corine Land Cover 2018)


- Discontinuous urban fabric
- Arable land outside irrigation perimeter
- Vineyards
- Grasslands
- Cropping systems and complex plots
- Mainly agricultural areas interrupted by important natural areas
- Hardwood forests
- Coniferous forests
- Mixed forests
- Natural grasslands and pastures
- Rivers

## Majority SMU in each grid cell of 8 km x 8 km grid: simulation scale




Flowering date in each 8 km x 8 km grid cell, simulated as a function of temperature since sowing and photoperiod.

Winter pea variety (Isard) - Near future (2006-2049)



Daily climate data, 8 km x 8 km grid cell simulated for the 1980-2005 and 2006-2100 periods (RCP 4.5 and 8.5).

- Temperature (Tmin, Tmax)
- Solar radiation (Rs)
- Precipitation (P)
- Potential evapotranspiration (ETP)



**Simulation results:**  
PRODUCTION Yield, seed protein content  
ABIOTIC STRESSES encountered by the crop

AZODYN-Pea simulates the functioning of a pea plot at a daily time step, integrating the effect of different abiotic stresses (water deficit, frost, high temperatures, soil compaction) and of some varietal characteristics.