



HAL
open science

Species Distribution Model using Remote-Sensed Dynamic Habitat Index

Mairi Souza Oliveira, Clémentine Préau, Maxime Lenormand, Samuel
Alleaume, Sandra Luque

► **To cite this version:**

Mairi Souza Oliveira, Clémentine Préau, Maxime Lenormand, Samuel Alleaume, Sandra Luque. Species Distribution Model using Remote-Sensed Dynamic Habitat Index. IUFRO Forest Environment DIV8 Conference 2023, Oct 2023, Evora, Portugal. hal-04256967

HAL Id: hal-04256967

<https://hal.inrae.fr/hal-04256967v1>

Submitted on 24 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Species Distribution Model using Remote-Sensed Dynamic Habitat Index

Mairi Souza Oliveira, Clémentine Préau, Samuel Alleaume
Maxime Lenormand & Sandra Luque

UMR TETIS, INRAE, France

IUFRO 2023 | Evora, Portugal


Tuesday, 24 October 2023



Motivation



Presence data



Predicted suitable habitat

**Predict (and explain) species suitable habitat
using remote sensing data**

Species Distribution Models


Presence data



+




Suitability map




Threshold

Suitable habitat



Environmental variables




Package 'biomod2'
Thuiller et al. (2023)



Environmental Predictors


Land Use (LU)



- ▶ Data preprocessing ('expert knowledge')
- ▶ Categorized data
- ▶ Morphological information

└ Distance to the closest...
└ Area...

Remote Sensing (RS)





- ▶ 'Raw' data
- ▶ Continuous radiometric indices of vegetation
- ▶ Temporal information


└ Annual statistics...
└ Seasonality...

Research question

Can we retrieve a similar information from both data sources?





Case Study Site






- ▶ 4 species of birds
- ▶ 5 amphibians
- ▶ 2 mammals

Results - Garbage in, garbage out?







Results - Overlaps





Bufo bufo

Predicted niche parts

-  RS Off-ovlap
-  LULC Off-ovlap
-  Overlap
-  Occurrences


Results - Response curves



Distance to the forest

Results - R ratio

$$R = \text{Green} / \text{Blue}$$




Bufo bufo

Predicted niche parts

- RS Off-overlap
- LULC Off-overlap
- Overlap
- Occurrences

Results - R ratio



Take home message

- ▶ It is not possible to determine which approach is better than the other in this study.
- ▶ The RS approach predicts a larger habitat than the LU approach.
- ▶ Strong influence of the (biased) observed data.

Biogeography



<https://biorgeo.github.io/bioregion/>