



**HAL**  
open science

## Proteomics as an emerging tool in equine meat research: an overview

Antonella Della Malva, Mohammed Gagaoua, Agostino Sevi, Marzia Albenzio

### ► To cite this version:

Antonella Della Malva, Mohammed Gagaoua, Agostino Sevi, Marzia Albenzio. Proteomics as an emerging tool in equine meat research: an overview. *Meat Technology*, 2023, 64 (2), pp.401-406. 10.18485/meattech.2023.64.2.77 . hal-04261526

**HAL Id: hal-04261526**

**<https://hal.inrae.fr/hal-04261526>**

Submitted on 27 Oct 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



# Proteomics as an emerging tool in equine meat research: an overview

Antonella della Malva<sup>a\*</sup>, Mohammed Gagaoua<sup>b\*</sup>, Agostino Sevi<sup>a</sup> and Marzia Albenzio<sup>a</sup>

<sup>a</sup> University of Foggia, Department of Agriculture, Food, Natural Resources and Engineering (DAFNE), Via Napoli 25, 71121 Foggia, Italy

<sup>b</sup> PEGASE, INRAE, Institut Agro, 35590 Saint-Gilles, France

## ARTICLE INFO

### Keywords:

Horse  
Donkey  
Meat quality  
Biochemical pathways  
Biomarkers  
Muscle Proteome

## ABSTRACT

Proteomics tools in the field of equine meat research have been very recently applied to explore the changes in the post-mortem muscle proteome and to discover biomarkers to monitor the variations in its different meat quality traits. The current advances achieved by proteomics in equine meat research are reviewed. Different proteomics techniques (sodium dodecyl sulphate-polyacrylamide gel electrophoresis; two-dimensional polyacrylamide gel electrophoresis; fluorescent two-dimensional difference gel electrophoresis; targeted proteomics; tandem-mass tag labeled proteomics; data-independent analysis proteomics) have been applied in the study of the equine muscle/meat. The studies revealed the biochemical pathways involved in the development of several donkey and horse (foal) meat quality variation. The current knowledge would be useful to develop high-quality products.

## 1. Introduction

In the last years, the increase in consumer health consciousness led to increased demand and consumption of alternative meat sources worldwide, with consumers choosing products with high nutritional value. Equine meat (from donkeys and horses) is widely recognized as a health-beneficial food due to its greater content of vitamins, minerals, conjugated fatty acids, and low fat and cholesterol content (Marino *et al.*, 2022). Despite its excellent nutritional value, preferences and perceptions of equine meat differ significantly due to the social, historical, ethical, and psychological characteristics of consumers (Lopez-Pedrouso *et al.*, 2023). In fact, recent trends show an increase in production and consumer demand for equine meat (FAOSTAT, 2021). Improving consumer confidence in equine meat consump-

tion including its nutritional features could offer an opportunity for farming systems to enlarge the livestock species, to differentiate the market, and to promote environmental sustainability. In terms of sensory properties, equine meat is considered a tough and dark red meat, which drives its consumer acceptability, and consequently, the purchasing decision at the point of sale. Proteomics-based techniques have been employed to decipher the proteome changes and the underlying molecular mechanisms related to different meat organoleptic traits in different species (Di Luca *et al.*, 2011; della Malva *et al.*, 2017; Lopez-Pedrouso *et al.*, 2020; Gagaoua *et al.*, 2020; Gagaoua *et al.*, 2021; Lamri *et al.*, 2023). In the particular case of equids, proteomics has been very recently applied, and to the best of our knowledge, in less than 10 studies (Table 1). Such studies have as their ultimate goal the development of high-quality

\*Corresponding authors: Antonella della Malva, [antonella.dellamalva@unifg.it](mailto:antonella.dellamalva@unifg.it); Mohammed Gagaoua, [mohammed.gagaoua@inrae.fr](mailto:mohammed.gagaoua@inrae.fr)

Paper received July 3<sup>rd</sup> 2023. Paper accepted July 18<sup>th</sup> 2023.

Published by Institute of Meat Hygiene and Technology — Belgrade, Serbia

This is an open access article under CC BY licence (<http://creativecommons.org/licenses/by/4.0>)

ity equine products and, more specifically, the goal of better understanding the biochemical pathways behind the variability of equine meat quality. Therefore, this paper aims to briefly introduce the basis of proteomic approaches in the assessment of equine meat quality variation by highlighting the applications of this powerful tool in both donkey and horse meat research with a focus on i) post-mortem changes and underlying mechanisms and ii) meat quality traits.

## 2. Brief overview of proteomics in meat research

In the field of meat research, proteomics is an adapted tool for in-depth characterization and to explore the biochemical processes taking place during the conversion of muscle into meat or during the post-mortem time (D'Alessandro & Zolla, 2013; Purslow et al., 2021; Gagaoua et al., 2022). Several strategies and methodologies can be used. Gel-based approaches (sodium dodecyl sulphate-polyacrylamide gel electrophoresis, SDS-PAGE; two-dimensional polyacrylamide gel electrophoresis, 2D-PAGE; fluorescent two-dimensional difference gel electrophoresis, 2D-DIGE) coupled with mass spectrometry (MS) are widely applied and have, so far, been the main methods used to investigate muscle proteome changes (For review: Gagaoua et al., 2022). However, in recent years, gel-free quantitative techniques (label-free or label-based approaches) have gradually taken over, leading to high accuracy and sensitivity in the quantification of proteins, including intact proteoforms and post-translational protein modifications (Li et al., 2021; Lamri et al., 2023). Furthermore, the tremendous progress in powerful bioinformatics tools and software have recently enabled the expansion and identification of new features related to several meat quality traits (Kiyimba et al., 2022). Generally, the muscle undergoes several dynamic modifications during the animal's life due to intrinsic and extrinsic factors. Several studies on beef evidenced that those factors are responsible for huge variations in the muscle proteome, which consequently impact the final quality traits of muscle foods (Sierra et al., 2021; Di Luca et al., 2022; Gagaoua et al., 2022). Proteomics technologies were successfully applied to decipher several of these changes and mechanisms, including in the field of biomarkers discovery to monitor meat tenderness, color, and water holding capacity (Gagaoua et al., 2022; Gagaoua and Picard, 2022).

## 3. Proteomics applications in horse meat research

Proteomics tools in horsemeat research have been applied with the objectives of monitoring and exploring the tenderization rate and the quality traits of different muscles during aging and also for achieving a better understanding of the muscle proteome differences among different breeds and diets (Table 1). A 2D-PAGE approach combined with liquid chromatography-tandem mass spectrometry (LC/MS-MS) was used by della Malva et al. (2019) to investigate the post-mortem changes in the myofibrillar muscle proteome and the tenderization rate during the aging time of different muscles (*Longissimus lumborum*, *Semimembranosus* and *Semitendinosus*) from Italian Heavy Draft Horses. The authors identified 22 myofibrillar and sarcoplasmic protein biomarkers to follow up on the influence of aging time on proteolysis. The 22 proteins belong, based on Gene Ontology analysis, to six biological cluster pathways, these being muscle contraction (GO: 0006936), NADH regeneration (GO: 0006735), regulation of ATP-dependent activity (GO: 0043462), muscle structure development (GO: 0061061), response to estradiol (GO: 0032355), and organophosphate biosynthetic process (GO: 0090407). In addition, della Malva et al. (2019) revealed differences in the rates of tenderization among muscles during aging, showing a greater accumulation of MYL1 and MYL2 fragments in *Semitendinosus* muscle with an extending aging time (14 days). When investigating the changes in the sarcoplasmic proteome in comparison with the meat organoleptic characteristics, another study by della Malva et al. (2022) found 24 muscle-specific protein patterns during aging, from which TPM1 and TPM2 cytoskeletal proteins were potential biomarkers of intense proteolysis for *Longissimus* muscle. The authors also indicated mitochondrial and glycolytic proteins (SOD, PGM1, and MB) as putative biomarkers to monitor the meat quality characteristics of horse *Semitendinosus* muscle.

Beldarrain et al. (2022) applied the OFFGEL proteomics-based approach on the Hispano-Breton *Longissimus thoracis et lumborum* (LTL) horse muscle to unveil changes in the myofibrillar proteome during three weeks of aging time. The authors indicated that aging-induced significant abundance changes of muscle structure proteins (MYL1, MYBPC1, TNNT3, and TNNI2) as key players of horse meat tenderization. To better understand the changes

found in the myofibrillar proteome of horse meat during aging, a 2D-DIGE approach was further applied by the same authors (Beldarrain *et al.*, 2023) to gain insights into the biochemistry of horse muscles and identify candidate protein biomarkers to monitor meat tenderness. Five putative protein biomarkers (TNNT3, MYBPC1, MYBPC2, ACTA1, and GAPDH) evidenced/expressed a great potential to monitor the changes in horse meat tenderization.

Targeted proteomics (SWATH-MS: Sequential Window Acquisition of all Theoretical Mass Spectra) has been recently applied by Lopez-Pedrouso *et al.* (2023) on two horse breeds (Burguete vs Jaca Navarra) finished with conventional concentrate and straw *or* silage and organic feed diets to identify biomarkers of multiple horsemeat quality traits (tenderness, color, and intramuscular fat). The authors built a database of 294 proteins, from which 23 proteins were candidate biomarkers of intramuscular fat content, while eight proteins, from the energy metabolism (ALDOA, CKM, TPI1, and PGMA2) and the muscle structure (ACTA1, MYBPH, MYL1, and TNNC1) pathways were identified as biomarkers to monitor the tenderization process. Regarding horse meat color determination, seven potential protein biomarkers related to energy metabolism (ALDOA, PKM, PFKM, and CKM), and oxidative stress (HSPA1A, SOD2, and PRDX2) were identified.

#### 4. Proteomics applications in donkey meat research

Proteomics investigations in donkey meat has been applied to decipher the biological mechanisms governing the meat tenderness and intramuscular fat variation and, consequently, to reveal the underlying pathways including those for the identification of protein biomarkers of the desirable meat quality (Table 1). In the frame of understanding the tenderization rate and the proteins changes occurring during the aging time of Martina Franca donkey meat, a gel-based 2DE proteomics approach coupled with LC/MS-MS was applied by della Malva *et al.* (2022). The authors proposed the first repertoire of 15 meat tenderness biomarkers for the donkey meat species. Using bioinformatics, the candidate biomarkers were allocated to three interconnected pathways: nine proteins were from muscle contraction, structure pathway (MYH1, MYH2, ACTA1, MYLPF, MYL6B, MYL1, TNNC2, TPM1, and TPM2); five from energy metabolism (ATP5PD, UQCRC1, COX5A, GAPDH, and CKM),

and one protein from the response to stress pathway (HSPB1), thus evidencing a key insight into the pathways and processes involved in the tenderness development of donkey meat.

Intramuscular fat content plays a pivotal role in the quality of muscle foods, thus affecting the flavor, juiciness, and tenderness of the end product. A tandem-mass tag (TMT) labeled proteomics study conducted by Tan *et al.* (2022) identified 30 differentially abundant proteins strictly related to the intramuscular fat deposition of the donkey *Longissimus thoracis* muscle. The functional enrichment analysis confirmed that the main biological pathways are involved in lipid metabolism and adipogenesis, thus confirming their role in the biological mechanisms that regulate meat quality variation.

Chai *et al.* (2022) used a data-independent analysis (DIA) proteomics approach to investigate the proteome differences of donkey muscles (*Semitendinosus* (ST), *Longissimus thoracis* (LT), and *Gluteus maximus* (GM)) related to meat quality parameters. The pairwise comparisons of the ST/LT and GM/LT allowed identification of, respectively, 111 and 127 differentially abundant proteins involved mainly in the MARK signaling pathway, fat digestion and absorption, and regulation of actin cytoskeleton. The focus given by these studies on the strong role of different biological pathways in the post-mortem processes linked with the donkey meat quality variation emphasizes the need for future research to explain the molecular basis of variations in donkey meat quality for developing high-quality products from this sustainable species.

#### 5. Future perspectives/Conclusion

The potential of proteomic tools to decipher and understand biological mechanisms and pathways underlying the equine meat quality variations has barely/recently been explored. The current few proteomics studies, developed on equids, allowed us to gain more insights about the biological mechanisms responsible for the variations in meat quality, in terms of tenderness, color, and intramuscular fat content. Several biological pathways have been discovered including proteins from the energy metabolism, muscle structure, oxidative stress, lipid metabolism, and adipogenesis, although further post-mortem muscle proteome studies and multi-omics approaches will be necessary to validate the biochemical pathways and the proposed candidate biomarkers, with the aim of monitoring equine meat quality

**Table 1.** Proteomics approaches in the selection of biomarkers related to meat quality in equids.

Proteomic approach	Muscle/cut (breed)	Considered variable(s)/ effects	Identified proteins/candidate biomarkers <sup>1</sup>	References
<b>Horse</b>				
SDS-PAGE, 2DE, HPLC/Q-TOF mass spectrometry, and Western Blotting	<i>Longissimus lumborum</i> (LL), <i>Semitendinosus</i> (ST) and <i>Semimembranosus</i> (SM) (Italian Heavy Draft Horse)	Aging – Muscle	MYL1, TNNT3, MYLPF, MYL3, TPM2, CKM, ENO2, AK1, PGK1, TPI1, GPX1	della Malva et al., 2019
SDS-PAGE, 2DE, and HPLC/Q-TOF mass spectrometry	<i>Longissimus lumborum</i> (LL), <i>Semitendinosus</i> (ST) and <i>Semimembranosus</i> (SM) (Italian Heavy Draft Horse)	Aging – Muscle	PGM1, CKM, TPM1, TPM2, ENO3, ALDOB, GPD1, GAPDH, TPI1, AK1, MB, SOD1	della Malva et al., 2022
OFFGEL, SDS-PAGE, LC-MS/MS	<i>Longissimus thoracis et lumborum</i> (LTL) (Hispano-Breton horse)	Aging	MYL1, TUBB4A, TNNT3, CRYAB, CKM, ENO3, ALDOA, GAPDH, LDHA, TNNT2, MYBC1, TPM2, ALDOA, CAPZA2, LDHA, MDH2, VDAC3, ATP5F1C, CA3, PGAM2, MYL3, ATP5PO, MYL1, MYLPF	Beldarrain et al., 2022
2-D DIGE, LC-MS/MS and Immunoblotting	<i>Longissimus thoracis et lumborum</i> (LTL) (Hispano-Breton horse)	Aging	ACTA1, MYBPC2, MYBPC1, PYGM, HSPA1A, DLAT, ALB, MYBPC2, SDHA, DES, TNNT3, ALDOA, CKM, LDHA, MYOZ1, ENO3, PHB, NDUFS3, HSPB1, ATP5PD, GAPDH,	Beldarrain et al., 2023
Shotgun data-dependent acquisition proteomic approach by micro-LC-MS/MS, Data-independent acquisition (DIA), SWATH-MS	<i>Longissimus thoracis and lumborum</i> (LTL) (Jaca Navarra and Burguete horses)	Breed – Feeding – Meat quality	<b>WBSF</b> <i>Burguete</i> : OBSCN, MYBPC1, MYL1, AHNAK, FLNC, NEB, SMTNL1, PDLIM5, CKM, MYOM1, LDB3, ACTN3, MSN, PAICS, ALDOA, ACTA1, HBA2, PGAM2, ARHGDI, NME2, MYBPH, WARS1, TTN, GSTO1, MYBPC2, MYOM2, PFN1, TPI1, MACROD1 <i>Jaca Navarra</i> : TNNC2, CKM, EEF1G, NDUFV2, ADSS1, FABP4); <b>Lightness (L*)</b> <i>Burguete</i> : IGL, PGK1, ALDOA, CKM, MACROD1, ORM1, ACYP2, GSTP1, PCMT1, SDHB, PKM, ALB, A1BG, ATP5F1B, UAB1, GSTO1, PDLIM5, CES1 <i>Jaca Navarra</i> : HIBADH, BIN1, CKM, HSPA1A, PCMT1, SOD2, EEF1A2, GAPDH, GOT2, MYBPH, GOT1); <b>Redness (a*)</b> <i>Burguete</i> : ECH1, VDAC2, PFKM, EEF1G, DES, SOD2, TRIM72, PGP, <i>Jaca Navarra</i> : HBA2, PPIA, GOT1, GLNA1, PRDX2, MYL1, CS, PHGDH); <b>Yellowness (b*)</b> <i>Burguete</i> : A1BG, TRIM72, VCL, ALDOA, ANXA7, ST13, NPEPPS, PSMA4, ORM1, CSRP3, MACROD1, DES <i>Jaca Navarra</i> : PCMT1, MYBPC1, ARHGDI, AHCY); <b>IMF</b> <i>Burguete</i> : ACTA1, MSN, HSPD1, MYH1, PDLIM3, NME2, ART3, ALDH2 <i>Jaca Navarra</i> : HSPA5, FBP1, TF, LDHB, ANXA1, STIP1, EEF1G, A1BG, GPD1, TNNC2	Lopez-Pedrouso et al., 2023



Proteomic approach	Muscle/cut (breed)	Considered variable(s)/ effects	Identified proteins/candidate biomarkers <sup>1</sup>	References
<i>Donkey</i>				
SDS-PAGE, 2DE, LC-MS/MS, and Western Blotting	<i>Longissimus thoracis et lumborum</i> (LTL) (Martina Franca donkey)	Aging	MYLPF, TPM2, TPM1, MYL1, ACTA1, GAPDH, TNNC2, MYH1, MYH2, UQCRC1, HSPB1, MYL6B, COX5A, ATP5PD, CKM	della Malva et al., 2022
Tandem-mass-tag, LC-MS/MS	<i>Longissimus dorsi</i> (LD) (Liaoxi donkey)	Meat quality	MAP4K4, PDLIM3, ADGRV1, ACTB, H2AJ, FLOT1, TPM1, UBR4, TRMT61A, PRMT3, C9, RPL32, MPC2, MBP, FHL3, LHPP, TAPT1, F11R, RPL27A, GLDN, PYCARD, COG3, CFAP251, SPAG8, GALNS, ROCK2, PRKCQ, K9KFH7, LUC7L2, ATP5MJ	Tan et al., 2022
Data-independent acquisition (DIA) and nano-LC-MS/MS	<i>Longissimus thoracis</i> (LT), <i>Gluteus maximus</i> (GM), and <i>Semitendinosus</i> (ST) (Dezhou donkey)	Muscle	111 proteins in the ST/LT and 127 proteins in the GM/LT muscles comparison were differentially expressed	Chai et al., 2022

<sup>1</sup>The full names of the proteins (gene names) are retrieved from Uniprot database (<https://www.uniprot.org>).

**Disclosure statement:** No potential conflict of interest was reported by the authors.

## References

- Beldarrain, L. R., Sentandreu, E., Aldai, N. & Sentandreu, M. A. (2022). Horse meat tenderization in relation to post-mortem evolution of the myofibrillar sub-proteome. *Meat Science*, 188, 108804.
- Beldarrain, L. R., Sentandreu, E., Aldai, N., Sentandreu, M. Á. & Miller, I. (2023). Application of 2-D DIGE to study the effect of ageing on horse meat myofibrillar sub-proteome. *Journal of Proteomics*, 272, 104770.
- Chai, W., Xu, J., Qu, H., Ma, Q., Zhu, M., Li, M... & Wang, C. (2022). Differential proteomic analysis to identify potential biomarkers associated with quality traits of Dezhou donkey meat using a data-independent acquisition (DIA) strategy. *LWT*, 166, 113792.
- D'Alessandro, A. & Zolla, L. (2013). Meat science: From proteomics to integrated omics towards system biology. *Journal of Proteomics*, 78, 558–577.
- della Malva, A., Marino, R., Santillo, A., Annicchiarico, G., Caroprese, M., Sevi, A. & Albenzio, M. (2017). Proteomic approach to investigate the impact of different dietary supplementation on lamb meat tenderness. *Meat Science*, 13, 74–81.
- della Malva, A., De Palo, P., Lorenzo, J. M., Maggolino, A., Albenzio, M. & Marino, R. (2019). Application of proteomic to investigate the post-mortem tenderization rate of different horse muscles. *Meat Science*, 157, 107885.
- della Malva, A., Maggolino, A., De Palo, P., Albenzio, M., Lorenzo, J. M., Sevi, A. & Marino, R. (2022a). Proteomic analysis to understand the relationship between the sarcoplasmic protein patterns and meat organoleptic characteristics in different horse muscles during aging. *Meat Science*, 184, 108686.
- della Malva, A., Gagaoua, M., Santillo, A., De Palo, P., Sevi, A. & Albenzio, M. (2022b). First insights about the underlying mechanisms of Martina Franca donkey meat tenderization during aging: A proteomic approach. *Meat Science*, 193, 108925.
- Di Luca, A., Mullen, A. M., Elia, G., Davey, G. & Hamill, R. M. (2011). Centrifugal drip is an accessible source for protein indicators of pork ageing and water-holding capacity. *Meat Science*, 88(2), 261–270.
- Di Luca, A., Ianni, A., Bennato, F., Henry, M., Meleady, P. & Martino, G. (2022). A label-free quantitative analysis for the search of proteomic differences between goat breeds. *Animals*, 12(23), 3336.
- FAOSTAT, (2021). Production quantities of Meat, horse. Retrieved October 14, 2021, from <https://www.fao.org/faostat/en/#data/QCL/visualize>.
- Gagaoua, M., Hughes, J., Terlouw, E. M. C., Warner, R. D., Purslow, P. P., Lorenzo, J. M. & Picard, B. (2020). Proteomic biomarkers of beef colour. *Trends in Food Science & Technology*, 101, 234–252.
- Gagaoua, M., Terlouw, E. C., Mullen, A. M., Franco, D., Warner, R. D., Lorenzo, J. M. ... & Picard, B. (2021). Molecular signatures of beef tenderness: Underlying mechanisms based on integromics of protein biomarkers from multi-platform proteomics studies. *Meat Science*, 172, 108311.
- Gagaoua, M., Schilling, W. M., Zhang, X. & Suman, S. P. (2022). Applications of proteomics in meat research. Reference module in food science. Elsevier.
- Gagaoua, M. & Picard, B., 2022. Proteomics to explain and predict meat quality. Chapter 14. In: Purslow, P. (Ed.), *New Aspects of Meat Quality*, second ed. Woodhead Publishing, pp. 393–431.

- Kiyimba, F., Gagaoua, M., Suman, S., Mafi, G. & Ramathan, R. (2022).** Bioinformatics: in-depth analyses of omics data in the field of muscle biology and meat biochemistry. Reference module in food science. Elsevier.
- Lamri, M., della Malva, A., Djenane, D., Lopez-Pedrouso, M., Franco, D., Albenzio, M., ... Gagaoua, M. (2023).** Towards the discovery of goat meat quality biomarkers using label-free proteomics. *Journal of Proteomics*, 278, 104868.
- Li, X., Zhang, D., Ren, C., Bai, Y., Ijaz, M., Hou, C. & Chen, L. (2021).** Effects of protein posttranslational modifications on meat quality: A review. *Comprehensive Reviews in Food Science and Food Safety*, 20(1), 289–331.
- Lopez-Pedrouso, M., Lorenzo, J. M., Gagaoua, M. & Franco, D. (2020).** Application of proteomic technologies to assess the quality of raw pork and pork products: An overview from farm-to-fork. *Biology*, 9(11), 393.
- López-Pedrouso, M., Lorenzo, J. M., Cittadini, A., Sarries, M. V., Gagaoua, M. & Franco, D. (2023).** A proteomic approach to identify biomarkers of foal meat quality: A focus on tenderness, color and intramuscular fat traits. *Food Chemistry*, 405, 134805.
- Marino R., della Malva A., Maggiolino A., De Palo P., d'Angelo F., Lorenzo J. M., Sevi A. & Albenzio M. (2022).** Nutritional profile of donkey and horse meat: Effect of muscle and aging time. *Animals*, 12(6), 746.
- Purslow, P. P., Gagaoua, M. & Warner, R. D. (2021).** Insights on meat quality from combining traditional studies and proteomics. *Meat Science*, 174, 108423.
- Sierra, V., Gonzalez-Blanco, L., Dineiro, Y., Díaz, F., García-Espina, M. J., CotoMontes, A., Gagaoua, M. & Olivan, M. (2021).** New insights on the impact of cattle handling on post-mortem myofibrillar muscle proteome and meat tenderization. *Foods*, 10(12), 3115.
- Tan, X., He, Y., Qin, Y., Yan, Z., Chen, J., Zhao, R., Zhou, S., Irwin, D.M., Li, B. & Zhang, S. (2023).** Comparative analysis of differentially abundant proteins between high and low intramuscular fat content groups in donkeys. *Frontiers in Veterinary Science*, 9, 951168