



HAL
open science

Identification of shoot architectural traits to promote winter oilseed rape vigour during the vegetative growth: a simulation approach

Laurène Perthame, Christophe Pradal, Alexandra Jullien, Frédéric Rees, Céline Richard-Molard, Grumel Arman

► To cite this version:

Laurène Perthame, Christophe Pradal, Alexandra Jullien, Frédéric Rees, Céline Richard-Molard, et al.. Identification of shoot architectural traits to promote winter oilseed rape vigour during the vegetative growth: a simulation approach. IRC 2023 - 16th International Rapeseed Congress, Sep 2023, Sydney, Australia. hal-04263001

HAL Id: hal-04263001

<https://hal.inrae.fr/hal-04263001>

Submitted on 27 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

Identification of shoot architectural traits to promote winter oilseed rape vigour during the vegetative growth: a simulation approach.

L. Perthame¹, C Pradal^{2,3}, A Jullien¹, F Rees¹, C Richard-Molard¹, A. Grumel¹

¹ *Université Paris-Saclay, INRAE, AgroParisTech, UMR EcoSys, 91120 Palaiseau, France*

² *AGAP, Univ. Montpellier, CIRAD, INRAE, SupAgro, Montpellier, France*

³ *INRIA, Univ. Montpellier, France*

Background:

Winter oilseed rape (WOSR) cultivation can often face implantation difficulties due to the lack of water at the beginning of growth and/or to pest damages, recently increased by climate change and environmental regulations. Farmers are therefore looking for more vigorous cultivars, which are able to better resist to such stresses thanks to their high carbon uptake efficiency, and ensure crop establishment and yield.

Objective:

Our work aimed to identify and classify important shoot architectural traits of WOSR during the vegetative growth to improve light capture, and favour the selection of vigorous WOSR cultivars.

Methods:

Our strategy was to build a model of leaf area establishment of WOSR at vegetative stage, to calibrate and evaluate it with experimental architectural data on isolated plants, and to connect it to a light interception model.

Our model describes the establishment of leaf area through time (expressed in degree.days), with a 3D representation of the leaves and their geometry. We use phyllochron (e.g. time between the emergence of two leaves) to parameterize the appearance of a new leaf over time. Leaves are separated by an angle of 137,5°. Each leaf is characterized by its length of the limb, from which infers its width, leaf area, petiole length and curvature angle through allometric relationships. Limb length, and thus leaf area, grows depending on thermal time, according to a speed growth parameter and a maximal leaf area, which varies with the total leaf area of the plant. This architectural model was connected the architectural model to a light interception model (Caribu; Chelle and Andrieu, 1998), in OpenAlea modelling platform (Pradal *et al.*, 2008).

Future results that will be showcase during the presentation are our model assemble (light interception model and leaf area establishment model) used to simulate a wide range of phenotypes and estimate which parameters (e.g. phyllochron, leaf length, leaf curvature angle, leaf speed growth) can maximize light capture, depending on plant density.

Conclusions:

We developed a new data-driven 3D architectural model of leaf area establishment for WOSR during the vegetative growth. Such model represents a powerful tool to design ideotypes that improve light capture and guide the selection of vigorous WOSR cultivars. In the near future, we intend to add metabolic processes (C acquisition and effects on leaf area expansion and biomass) and complete the model during the entire crop cycle.

References:

Chelle, M., Andrieu, B. (1998). The nested radiosity model for the distribution of light within plant canopies. *Ecological Modelling*, 111, 75–91. [https://doi.org/10.1016/S0304-3800\(98\)00100-8](https://doi.org/10.1016/S0304-3800(98)00100-8)

Pradal C., et al. (2008). OpenAlea: A visual programming and component-based software platform for plant modeling. *Functional Plant Biology*, 35 (10), pp.751-760. 10.1071/FP08084