



HAL
open science

Tracking bacilli swimmers in microalgae biofilms

Simon Manceau, Andrea Fanesi, Julia Mougin, Julien Deschamps, Filipa Lopes, Romain Briandet

► **To cite this version:**

Simon Manceau, Andrea Fanesi, Julia Mougin, Julien Deschamps, Filipa Lopes, et al.. Tracking bacilli swimmers in microalgae biofilms. *Bacell*, Nov 2023, Kobe, Japan. pp.1-2. hal-04263849

HAL Id: hal-04263849

<https://hal.inrae.fr/hal-04263849>

Submitted on 29 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Introduction

Some *Bacillus* biofilms have been shown to spontaneously contain a subpopulation of motile bacteria capable of navigating through the matrix in three dimensions (Houry *et al.*, 2012). These swimming bacteria vascularize the biofilms and facilitate the penetration of specific substances to the deeper layers. Microalgae are rich in molecules of interest, and are at the heart of the challenges of the 21st century. Their biofilm culture is attracting growing interest due to their high productivity potential. In this innovative study, the aim is to characterize *Bacillus* swimming in a *Cylindrotheca closterium* biofilm using swimming bacilli to further increase biomass productivity.

The swimmers criss-cross a biofilm of bacteria, but are they capable of vascularizing a biofilm of microalgae, given that these microorganisms are at least three times larger?

Using confocal microscopy (CLSM), an experimental method was developed to study the swimming of these bacilli. Although the microalgae biofilm becomes thicker with increasing matrix production (Fanesi *et al.*, 2019), *Bacillus* swimming is still observable. Precise characterization of the swimmers requires further optimization and trajectory analyses.

Materials and Methods

Culture of microalgae biofilms



Inoculation

50 μL .puits⁻¹
Each day, inoculated wells are sacrificed


Culture of swimming bacteria:

Bacillus thuringiensis Bt407-GFP
Growth medium : TSB
DO_{600nm} = 0,4




CLSM Acquisitions 30 minutes after inoculation:

Biofilm biovolume extraction




Autofluorescent chloroplasts of *C. closterium*




Surface tool of Imaris®




Biovolume over time




Bacteria tracking at the bottom of the wells



0s



15s



30s




Motility caracterisation:

- Position
- Swimming parameters (speed, length, duration...)
- Mean square displacement
- Straightness index


Results

Normalized trajectories of swimming bacteria over time




Heterogeneity of microalgal biofilms

Autofluorescent clusters (560 nm):



C. closterium (> 5 days)


Mesoscopic structure by Optical Coherence Tomography (B-scan):




C. closterium (> 3 weeks)

Swimming parameters over time


Control : Control Biofilm * : mean significantly different from control (p-value < 0,05, test de Mann-Whitney-Wilcoxon)



Control mean speed (16 $\mu\text{m}\cdot\text{s}^{-1}$) > Biofilm mean speed (13 $\mu\text{m}\cdot\text{s}^{-1}$)
There is a decrease in the number of bacteria over time.



There is no decrease in mean speed over time.



Track Length in control < Track Length in biofilm

Conclusion

This project, at the convergence of the fields of biofilms, microalgae and bacterial motility, has demonstrated the remarkable ability of bacilli to move within the matrix of a microalgal biofilm. However, the swimmers used have a reduced average speed of movement, more confined trajectories and a reduced number of swimmers observed at the bottom of the well. This is consistent with the increased biovolume of the *C. closterium* biofilm and its substantial thickening.

This groundbreaking study highlights the heterogeneity of this phenomenon in a complex biofilm. The impact of biofilm vascularization by bacteria swimming through the matrix has not yet been assessed. Exploring these vascularization phenomena could encourage the entry of nutrients or the release of molecules such as O₂ trapped in biofilms, and thus promote their growth.