



HAL
open science

Yeast-based heterologous production of the Colletochlorin family of fungal secondary metabolites

Aude Geistodt-Kiener, Jean Chrisologue Totozafy, Géraldine Le Goff, Justine Vergne, Kaori Sakai, Jamal Ouazzani, Grégory Mouille, Muriel Viaud, Richard O'Connell, Jean-Félix Dallery

► To cite this version:

Aude Geistodt-Kiener, Jean Chrisologue Totozafy, Géraldine Le Goff, Justine Vergne, Kaori Sakai, et al.. Yeast-based heterologous production of the Colletochlorin family of fungal secondary metabolites. *Metabolic Engineering*, 2023, 80, pp.216 - 231. 10.1016/j.ymben.2023.10.002 . hal-04263969v2

HAL Id: hal-04263969

<https://hal.inrae.fr/hal-04263969v2>

Submitted on 29 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Contents lists available at [ScienceDirect](https://www.sciencedirect.com)

Metabolic Engineering

journal homepage: www.elsevier.com/locate/meteng

Yeast-based heterologous production of the Colletochlorin family of fungal secondary metabolites

Aude Geistodt-Kiener^a, Jean Chrisologue Totozafy^b, Géraldine Le Goff^c, Justine Vergne^a, Kaori Sakai^a, Jamal Ouazzani^c, Grégory Mouille^b, Muriel Viaud^a, Richard J. O'Connell^a, Jean-Félix Dallery^{a,*}^a Université Paris-Saclay, INRAE, UR BIOGER, 91120, Palaiseau, France^b Université Paris-Saclay, INRAE, AgroParisTech, Institut Jean-Pierre Bourgin, 78000, Versailles, France^c Centre National de La Recherche Scientifique, Institut de Chimie des Substances Naturelles ICSN, 91190, Gif-sur-Yvette, France

ARTICLE INFO

Keywords:

Secondary metabolism
Heterologous expression
Polycistronic vector
Colletotrichum higginsianum
Saccharomyces cerevisiae

ABSTRACT

Transcriptomic studies have revealed that fungal pathogens of plants activate the expression of numerous biosynthetic gene clusters (BGC) exclusively when in presence of a living host plant. The identification and structural elucidation of the corresponding secondary metabolites remain challenging. The aim was to develop a polycistronic system for heterologous expression of fungal BGCs in *Saccharomyces cerevisiae*. Here we adapted a polycistronic vector for efficient, seamless and cost-effective cloning of biosynthetic genes using *in vivo* assembly (also called transformation-assisted recombination) directly in *Escherichia coli* followed by heterologous expression in *S. cerevisiae*. Two vectors were generated with different auto-inducible yeast promoters and selection markers. The effectiveness of these vectors was validated with fluorescent proteins. As a proof-of-principle, we applied our approach to the Colletochlorin family of molecules. These polyketide secondary metabolites were known from the phytopathogenic fungus *Colletotrichum higginsianum* but had never been linked to their biosynthetic genes. Considering the requirement for a halogenase, and by applying comparative genomics, we identified a BGC putatively involved in the biosynthesis of Colletochlorins in *C. higginsianum*. Following the expression of those genes in *S. cerevisiae*, we could identify the presence of the precursor Orsellinic acid, Colletochlorins and their non-chlorinated counterparts, the Colletorins. In conclusion, the polycistronic vectors described herein were adapted for the host *S. cerevisiae* and allowed to link the Colletochlorin compound family to their corresponding biosynthetic genes. This system will now enable the production and purification of infection-specific secondary metabolites of fungal phytopathogens. More widely, this system could be applied to any fungal BGC of interest.

1. Introduction

Fungi are rich sources of structurally diverse, small molecule natural products, as illustrated by the fact that more than 60% (20,304) of the 33,372 microbial natural products currently catalogued in the Natural Products Atlas were isolated from fungi (<https://www.npatlas.org>, May 2023) (van Santen et al., 2022). Well-known for their uses in medicine or agriculture, fungal natural products play important roles in the adaptation of fungi to their ecological niches, for example as toxins to compete with other microorganisms, as protection against environmental stresses, or in the case of pathogens, as effectors to facilitate the infection of plant or animal hosts (Collemare et al., 2019; Oberlie et al.,

2018).

Fungal genes involved in the biosynthesis of natural products are typically located side-by-side in the genome as so-called biosynthetic genes clusters (BGCs). These clusters usually contain genes encoding one or two key enzymes that generate the backbone of the molecule and a varying number of genes encoding accessory enzymes that decorate the initial molecule. Genes coding for transporters or transcription factors can also form part of the BGC (Keller, 2019). This colocalization in the genome facilitates the identification of BGCs, and several bioinformatic tools and databases (e.g. SMURF, antiSmash, MIBiG, CusProSe) have been developed to mine and annotate the ever-increasing number of sequenced fungal genomes released in public databases (Blin et al., 2021; Khaldi et al., 2010; Oliveira et al., 2023; Terlouw et al., 2023).

* Corresponding author. UR BIOGER, 22 place de l'Agronomie, 91120, Palaiseau, France.

E-mail address: jean-felix.dallery@inrae.fr (J.-F. Dallery).<https://doi.org/10.1016/j.ymben.2023.10.002>

Received 5 July 2023; Received in revised form 15 September 2023; Accepted 12 October 2023

Available online 19 October 2023

1096-7176/© 2023 The Authors. Published by Elsevier Inc. on behalf of International Metabolic Engineering Society. This is an open access article under the CC BY-NC license (<http://creativecommons.org/licenses/by-nc/4.0/>).

Abbreviations

BGC	biosynthetic gene cluster	NRPS	non-ribosomal peptide synthetase
CDS	coding sequence	OA	Orsellinic acid
DAPI	4',6-diamidino-2-phenylindole	OSMAC	one strain many compounds
DMAPP	dimethylallyl pyrophosphate	PCR	polymerase chain reaction
ELSD	evaporative light scattering detector	PDA	potato dextrose agar
GPP	geranyl pyrophosphate	PKS	polyketide synthase
HPLC	high-performance liquid chromatography	SM	secondary metabolite
IVA	<i>in vivo</i> assembly	TAR	transformation-assisted recombination
LC	liquid chromatography	TEV	<i>Tobacco Etch Virus</i>
MS	mass spectrometry	UV	ultraviolet
NLS	nuclear localization signal	WT	wild type
		YNB	yeast nitrogen broth
		YPD	yeast extract peptone dextrose

Analysis with these tools has shown that a single fungal genome can contain more than 80 non-redundant BGCs (Han et al., 2016; Inglis et al., 2013; Liang et al., 2018; Valero-Jiménez et al., 2020). However, for the vast majority of these predicted BGCs, the chemical products are currently unknown.

Colletotrichum higginsianum is a plant-pathogenic ascomycete fungus that causes disease on various cultivated *Brassicaceae* as well as the model plant *Arabidopsis thaliana* (O'Connell et al., 2004). Resequencing the genome of *C. higginsianum* revealed the presence of 77 non-redundant BGCs, of which only 14 (18%) can be linked to known chemical products based on their similarity to BGCs characterized in other fungi (Dallery et al., 2017; O'Connell et al., 2012). Transcriptional analysis showed that 19 *C. higginsianum* BGCs are induced at particular stages of plant infection. Among these, 14 are specifically expressed during the initial biotrophic phase of penetration and growth inside living plant cells, two are upregulated during later necrotrophic growth in dead tissues, while three are expressed at both infection stages (Dallery et al., 2017; O'Connell et al., 2012). These clusters are poorly expressed, or not at all, by axenic cultures of *C. higginsianum*, which has hindered isolation of the corresponding fungal metabolites in sufficient amounts for determination of their chemical structures and analysis of their biological activities.

Various strategies can be used to activate the expression of cryptic fungal BGCs in axenic cultures. For example, variation of the culture conditions (media composition, static/liquid cultures), as in the 'One strain many compounds' (OSMAC) technique (Bode et al., 2002; Hewage et al., 2014) and co-cultivation with other microbes (Yu et al., 2021). Other approaches include over-expressing global or cluster-specific transcriptional activators (Chiang et al., 2009; von Bargen et al., 2013), and opening up chromatin structure by the chemical or genetic manipulation of epigenetic regulator proteins, such as CCLA, KMT6 and LAEA (Lyu et al., 2020; Pimentel-Elardo et al., 2015). In *C. higginsianum*, we previously deleted the CCLA subunit of the COMPASS protein complex, which mediates mono-, di- and trimethylation of lysine 4 in histone H3. Cultures of the resulting mutant over-produced a number of terpenoid compounds belonging to the Higginsianin, Colletorin/Colletochlorin and Sclerosporide families (Dallery et al., 2019).

Although these genome-wide strategies have been successfully used to isolate new compounds from some fungi, they are untargeted and do not activate all silent BGCs. Heterologous expression provides a way to activate specific BGCs of interest and facilitates the large-scale production of metabolites in axenic cultures. In this approach, an entire BGC is cloned into a heterologous microbial host that can be easily cultivated (Ahmed et al., 2020; Bond and Tang, 2019; Chiang et al., 2013; Gomez-Escribano and Bibb, 2011; Pfeifer et al., 2001; Zhang et al., 2018). For heterologous production of fungal secondary metabolites, the most frequently used hosts have been *Escherichia coli* (Kealey et al., 1998; Pfeifer Blaine et al., 2003), the yeasts *Saccharomyces cerevisiae* and *Pichia pastoris* (Cochrane et al., 2016; Gao et al., 2013; Kealey et al.,

1998; Xue et al., 2017) and some filamentous fungi, including species of *Aspergillus*, *Trichoderma* and *Penicillium* (Heneghan et al., 2010; Nielsen et al., 2013; Pohl et al., 2020; Shenouda et al., 2022). *S. cerevisiae* has the advantage of being a GRAS (generally recognized as safe) organism that can be easily genetically manipulated and cultured at scale. Moreover, yeast can support the protein folding and post-translational modifications occurring in filamentous fungi and produces very few endogenous secondary metabolites, which facilitates the purification and isolation of heterologous compounds (Bond et al., 2016; Ishiuchi et al., 2012; Yu et al., 2013; Zhao et al., 2020).

To synchronously activate all the genes in a biosynthetic pathway during heterologous expression it is often necessary to laboriously change the native promoter and terminator of each gene (Harvey et al., 2018; Pahirulzaman et al., 2012). To avoid this, Hoefgen et al. (2018) designed an expression vector that allows the concerted expression of multiple genes as a single polycistron, where all the genes are placed under the control of a single promoter, allowing their co-expression. The self-splicing porcine teschovirus P2A DNA sequence is inserted at the 3' end of each gene of the polycistron, which induces bond skipping by the ribosome during translation, thereby releasing the individual proteins (Kim et al., 2011). The vector also incorporates a split fluorescent reporter gene so that correct translation of the polycistronic transcript can be monitored by microscopy. The vector was successfully used to transfer the Austinoid and Psilocybin biosynthetic pathways into *A. nidulans* (Hoefgen et al., 2018).

In the present study, our objective was to adapt the polycistronic vector of Hoefgen et al. (2018) for the heterologous expression of fungal BGCs in baker's yeast. To validate this expression system, we chose the Colletochlorins and their non-chlorinated counterparts, the Colletorins, a well-characterized family of prenylated polyketide compounds produced by *C. higginsianum* (Dallery et al., 2019). Using genome mining and comparative genomics, we identified a candidate BGC that could be involved in producing this family of molecules. Four genes, encoding a polyketide synthase, a non-ribosomal peptide synthase-like enzyme, a putative prenyltransferase and a halogenase, were expressed in *S. cerevisiae*. The resulting yeast culture supernatants were found to contain the expected products and intermediates of this biosynthetic pathway, namely Orsellinic acid, Colletorins B and D, Colletochlorin B and D and Colletorin D acid.

2. Materials and methods

2.1. Biological material and growth conditions

A summary of the *Saccharomyces cerevisiae* strains used in this study is given in Supplementary File 1. *In vivo* assembly and propagation of plasmids was performed using the *E. coli* TOP10 strain. Bacteria were maintained and propagated in LB broth, supplemented with antibiotics and agar as required. The yeast strains were cultivated at 28 °C in the

selective medium YNB (cat. No. Y0626, Sigma), supplemented with drop-out without leucine (cat. No. Y1376, Sigma), without uracil (cat. No. Y1501, Sigma) or without both leucine and uracil (cat. No. Y1771, Sigma), and supplemented with 2% glucose and 2% agar. For the heterologous production of metabolites, the yeast strains were cultivated in YPD liquid medium (yeast extract 10 g·L⁻¹, peptone 20 g·L⁻¹ and glucose 20 g·L⁻¹) for 72 h at 30 °C with agitation on a rotary shaker at 250 rpm. For quantification of Colletochlorins in *Colletotrichum higginsianum* strain IMI 349063A, the fungus was grown as described in Dallery et al. (2019).

2.2. Yeast expression vector construction

All vectors developed in the present study were built by *in vivo* assembly (IVA) of DNA fragments in *E. coli*. The PCR primers used to amplify the fragments were designed to provide a minimum of 22 bp of overlapping sequence between adjacent fragments for homologous recombination in the bacteria. The amount of fragment used for IVA varied according to the fragment size: 200–300 bp: 1.5 pmol, 300–600 bp: 1.0 pmol, 600–1000 bp: 0.5 pmol, 1000–3000 bp: 0.2 pmol, 3000–5000 bp: 0.1 pmol, >5000 bp: 0.05 pmol. Residual plasmid matrix in the PCR reactions was removed by digestion at 37 °C for 2 h with 10U of *DpnI* enzyme before proceeding further. All the cloning fragments were obtained by PCR amplification using Q5 polymerase according to the manufacturer's instructions (cat.no. M0491L, New England Biolabs). Diagnostic PCR was performed using GoTaq polymerase (cat.no. M7805, Promega).

The plasmid pHYX104 was constructed by amplifying by PCR the polycistronic fragment (*VenusN-P2A-TEV-P2A-VenusC*) from pV2A-T (Hoefgen et al., 2018) using primers P953 and P954, the *URA3* marker gene, 2 μ origin, *AmpR* and *ColE1* origin from pNAB-OGG (Schumacher, 2012) with primers P949 and P950, the *ScADH2* promoter with primers P951 and P952 and the *ScHIS5* terminator with primers P955 and P956. The four fragments were assembled by IVA in *E. coli*. The resulting plasmid pHYX104 was amplified by PCR with primers P963 and P964 to remove the *AmpR* gene and replace it with the modified *KanR*⁺ gene amplified with primers P965 and P966 from pV2A-T to take advantage of the *SwaI/PmeI* restriction sites. The resulting plasmid pHYX105 was further modified with an *EcoRV*-free version of the *URA3* gene obtained by gene synthesis to give rise to pHYX138.

The plasmid pHYX106 was constructed by amplifying the polycistronic fragment (*VenusN-P2A-TEV-P2A-VenusC*) from pV2A-T with primers P953 and P954, the *LEU2* marker gene from M4755 (Voth et al., 2003) with primers P971 and P972, the 2 μ origin, *KanR*⁺ and *ColE1* origin from pHYX105 with primers P967 and P950, the *ScPCK1* promoter with primers P975 and P976 and the *ScPRM9* terminator with primers P977 and P978. The five fragments were assembled by IVA in *E. coli*. The resulting pHYX106 was further modified by PCR with primers P1204 and P1205 to insert a single nucleotide mutation to remove the *EcoRV* restriction site present in the *LEU2* marker gene, giving rise to the plasmid pHYX137 (Supplementary File 2).

The plasmid pHYX163 was constructed by digesting the plasmid pHYX153 with *SwaI* and the pHYX154 with *PmeI* following the strategy described by Hoefgen et al. (2018). The digestion was performed overnight (nearly 16 h). After heat inactivation, the digested fragments (0.1 pmol of each fragment of interest) were assembled by IVA in *E. coli*. The resulting plasmid pHYX163 was digested by *PmeI* and assembled with the plasmid pHYX152 previously digested with *SwaI*, to obtain the pHYX164 plasmid. The plasmid pHYX172 was derived from plasmid pCfB2312, and was constructed by amplifying pCfB2312 with primers P1312 and P1313, and the prenyltransferase (codon-adapted for yeast) from pHYX154 with primers P1310 and P1311. The two fragments were gel-purified and assembled by IVA in *E. coli*. All primers used are listed in Supplementary File 3. The following plasmids were deposited with Addgene: pHYX137 (#202814), pHYX138 (#202815), pHYX143

(#202816) and pHYX173 (#202817).

2.3. Modifications of the yeast chassis strain

The yeast strain *S. cerevisiae* BJ5464-NpgA (Ma et al., 2009) was modified by inserting the *Botrytis cinerea* NADPH cytochrome P450 reductase gene *BcCPR1* (Bcin12g03180), a protein required for activation of cytochrome P450s, at the yeast locus XI-3, as described by Mikkelsen et al. (2012). The Colletochlorins BGC contained no cytochrome P450, so this modification was not necessary but was present in the chassis strain used. Competent yeast cells were prepared according to Knop et al. (1999). The CRISPR-Cas9 transformation was performed as described by Jessop-Fabre et al. (2016) using the plasmids pCfB2312, pCfB3045 and XI-3-bccpr1 yielding the strain BJNBC (Supplementary File 1).

2.4. Comparative analyses of the biosynthetic gene cluster 16

The protein sequences of the genes belonging to the BGC16 of *C. higginsianum* IMI 349063 (CH63R_05468 to CH63R_05483) as predicted in Dallery et al. (2017) were used to query the NCBI nr database using cblaster v1.3.9 (Gilchrist et al., 2021) with the `search` module and default parameters but limiting the investigations to fungi with `-eq "txid4751[orgn]"` for the Colletochlorin part of BGC16, or limiting the investigations to *Colletotrichum* spp. with the parameter `-eq "txid5455[orgn]"` for the entire BGC16. The BGCs identified were retrieved with the module `extract_clusters` and used as input for clinker software v0.0.23 using default parameters (Gilchrist and Chooi, 2021). Manual editing of the cartoons was performed to represent the contig ends where appropriate.

2.5. Cloning of reporter genes and biosynthetic genes

The gene predictions of CH63R_05468 to CH63R_05471 were manually inspected for correctness using RNA-Seq datasets previously published (O'Connell et al., 2012). The CDS of CH63R_05468 appeared to be composed of six exons instead of only five in the initial prediction. Each gene was then synthesized without stop codons and with optimization of the codons for *S. cerevisiae* and exclusion of common restriction enzyme sites using the GenSmart™ Codon Optimization tool (Genscript Biotech B.V., Netherlands). The optimized sequences were deposited at Genbank with the following accession numbers: OR389982, OR389983, OR389984 and OR389985. Each coding sequence was individually cloned into the *EcoRV* linearized pHYX137 and subsequently assembled together following the strategy described by Hoefgen et al. (2018) using *SwaI* and *PmeI* restriction enzymes followed by IVA in *E. coli*. The gene coding mScarlet-I was amplified from the plasmid Double UP mNeonGreen to mScarlet (Addgene #125134) by PCR using primers P1214 and P1215 and cloned into the *EcoRV* linearized pHYX137 by IVA in *E. coli*. The gene coding Tobacco Etch Virus (TEV) protease was synthesized (Genscript Biotech B.V., Netherlands), then amplified by PCR using primers P1177 and P1178 and cloned into the *EcoRV* linearized pHYX137 by IVA in *E. coli*. All coding sequences were verified by Sanger sequencing after their initial cloning and their presence was further verified by PCR after subsequent combination of vectors for multigene expression. A summary of the plasmids constructed in this study is presented in Supplementary File 4.

2.6. Yeast transformation

Established protocols were used for the transformation of plasmids into yeast strain BJNBC and the preparation of frozen competent yeast cells (Knop et al., 1999).

2.7. Microscopy

For confocal microscope observations, yeast strains were cultivated in YPD media at 28 °C for 2 days. For staining nuclei, samples were fixed with 4% formaldehyde in PBS for 30 min, spun down and rinsed once in PBS, permeabilized with 0.2% Triton X-100 in PBS for 5 min and rinsed three times in PBS. Samples were then incubated in 15 µg·mL⁻¹ of DAPI (4',6-diamidino-2-phenylindole) in PBS for 30 min, washed for 5 min and mounted on microscope slides prior to observation. Samples were imaged by sequential scanning using a Leica TCS SPE laser scanning microscope (Leica Microsystems) equipped with an APO 40× (1.15 NA) oil immersion objective. Venus, mScarlet-I and DAPI were excited using the 488 nm, 532 nm and 405 nm laser lines, respectively.

2.8. Protein extraction

For total protein extraction, yeast strains were cultivated in the appropriate YNB medium until OD reached 0.4. The cultures were then centrifuged (3000×g) for 10 min and the pellets resuspended in fresh YNB medium supplemented with 2% (w/v) glucose and 3% (v/v) ethanol. Yeast cells were pelleted by centrifugation at 3000×g for 5 min at 4 °C and immediately resuspended in lysis buffer (8 M urea, 5% [w/v] SDS, 40 mM Tris-HCl pH 6.8, 0.1 mM EDTA, 0.4 mg·mL⁻¹ bromophenol blue) supplemented with 1% (v/v) β-mercaptoethanol, 1× protease inhibitor cocktail (cat.no. P8215, Sigma-Aldrich), 5 µg·mL⁻¹ leupeptin (cat.no. L2884, Sigma-Aldrich) and 1 mM PMSF (cat.no. P7626, Sigma-Aldrich). PMSF was renewed every 7 min until the samples were frozen at -80 °C or loaded on a gel. Immediately after resuspension, glass beads were added up to the meniscus and the mixture incubated at 70 °C for 10 min. Cells were disrupted using a vortex mixer for 1 min and debris were pelleted by centrifugation at 18,000×g for 5 min at 4 °C. After the supernatants were recovered on ice, 75 µL of lysis buffer was added to the pellets, boiled at 100 °C for 5 min, centrifuged again and the supernatants were finally combined with those from the first centrifugation.

2.9. Immunoblot assay

Proteins were separated by SDS-PAGE on 4–15% gradient Mini-Protein TGX Stain-free gels (cat.no. 4568083, Bio-Rad), transferred onto PVDF membranes (cat.no. 1704273, Bio-Rad) and subsequently blocked with 5% (w/v) BSA in TBST buffer. The membranes were incubated for 1 h at RT with a mouse anti-2A primary antibody (cat.no. MABS2005, Merck) diluted 1:2000 in 1% (w/v) BSA in TBST. The membranes were then rinsed 15 min in TBST then 3 × 5 min in TBST before incubation for 1 h at ambient temperature with HRP-coupled goat anti-mouse secondary antibody diluted 1:5000 (cat.no. ab6728, Abcam). The membranes were then rinsed with TBST as above before chemiluminescence detection using the Clarity Western ECL substrate kit (cat.no. 1705060, Bio-Rad). Gels and blots were recorded with a ChemiDoc Imaging System (Bio-Rad).

2.10. General chemistry procedures

The cultivation of yeast strains harboring each plasmid and isolation of the chemical compounds were as described previously by Harvey et al. (2018). After preculturing the yeast strains in selective medium supplemented with 2% (w/v) glucose (2 days at 28 °C with shaking at 250 rpm), the preculture (10 mL) was inoculated into 1 L of YPD medium in a 2 L Erlenmeyer flask (total 4 × 1 L) and incubated for 72 h at 30 °C with shaking at 250 rpm. The YPD culture was then centrifuged aseptically (5000×g, 5 min) and the supernatant was incubated overnight with sterile XAD-16N resin (Dow Chemicals) for solid phase extraction (Dallery et al., 2019). The resin was collected by filtration and extracted for 2 h in ethyl acetate (100 mL) followed by 2 h in methanol (100 mL). Lyophilized cell pellets were resuspended in acetone (3 × 30

mL), sonicated for 3 × 15 min, centrifuged at 5000×g, 5 min between each acetone addition, followed by extraction with methanol (3 × 30 mL). Ethyl acetate extracts were dried over anhydrous sodium sulphate. Similar extracts were pooled, evaporated under reduced pressure and resuspended in HPLC grade methanol. The crude extracts were then analysed on an Alliance 2695 HPLC instrument equipped with a 2998 photodiode array, a 2420 evaporative light scattering and an Acquity QDa mass detector (Waters Corporation). The HPLC column used was a 3.5 µm C-18 column (Sunfire 150 × 4.6 mm) operating a linear gradient from H₂O to CH₃CN, both containing 0.1% formic acid, for 50 min at 0.7 mL·min⁻¹. Thin layer chromatography plates (Si gel 60 F 254) were purchased from Merck. Purified standards of Orsellinic acid, Colletorin D, Colletorin D acid, Colletochlorin D, Colletochlorin B and Colletochlorin A were dissolved at 1 mg·mL⁻¹ in methanol. All standards were purified as previously described (Dallery et al., 2019) except Orsellinic acid that was purchased from ThermoFisher Scientific (cat.no. 453290010).

2.11. Untargeted analysis of different Colletochlorin derivatives

Untargeted analysis was performed using a UHPLC system (Ultimate 3000 Thermo) coupled to a quadrupole time of flight mass spectrometer (Q-ToF Impact II Bruker Daltonics).

Separation was performed on an EC 100/2 Nucleoshell Phenyl-Hexyl column (2 × 100 mm, 2.7 µm; Macherey-Nagel) at 40 °C, with a flow rate of 0.4 mL·min⁻¹, for 1 µL injected. The mobile phases used for the chromatographic separation were: (A) 0.1% formic acid in H₂O; and (B) 0.1% formic acid in acetonitrile. Elution was as follows: 5% phase B for 2 min, the gradient elution increased linearly to 50% phase B in 13 min, followed by a further linear increase to 100% phase B in 10 min, then 100% phase B for 3 min and the final gradient linear elution decreased to 5% phase B for 7 min.

Data-dependent acquisition methods were used for mass spectrometer data in negative ESI mode using the following parameters: capillary voltage, 4.5 kV; nebulizer gas flow, 2.1 bar; dry gas flow, 6 L·min⁻¹; drying gas in the heated electrospray source temperature, 200 °C. Samples were analysed at 8 Hz with a mass range of 100–1500 *m/z*. Stepping acquisition parameters were created to improve the fragmentation profile with a collision RF from 200 to 700 Vpp, a transfer time from 20 to 70 µsec, and collision energy from 20 to 40 eV. Each cycle included a MS fullscan and 5 MS/MS CID on the 5 main ions of the previous MS spectrum.

2.12. Data processing of untargeted metabolomic data

The .d data files (Bruker Daltonics) were converted to .mzXML format using the MSConvert software (ProteoWizard package 3.0; Chambers et al., 2012). MzXML data processing, mass detection, chromatogram building, deconvolution, samples alignment and data export were performed using MZmine-2.37 software (<http://mzmine.github.io/>) for negative data files. The ADAP chromatogram builder (Myers et al., 2017) method was used with a minimum group size of scan 9, a group intensity threshold of 1000, a minimum highest intensity of 1000 and *m/z* tolerance of 10 ppm. Deconvolution was performed with the ADAP wavelets algorithm using the following setting: S/N threshold 10, peak duration range = 0.01–2 min RT wavelet range 0.02–0.2 min, MS2 scan were paired using a *m/z* tolerance range of 0.05 Da and RT tolerance of 0.5 min. Then, isotopic peak grouper algorithm was used with a *m/z* tolerance of 10 ppm and RT tolerance of 0.1min. All the peaks were filtered using a feature list row filter keeping only peaks with MS2 scan. The alignment of samples was performed using the join aligner with an *m/z* tolerance of 10 ppm, a weight for *m/z* and RT at 1, a retention time tolerance of 0.2 min. Metabolites accumulation was normalized according to the weight of dried extract for the relative quantification. Molecular networks were generated with MetGem software (Olivon et al., 2018; <https://metgem.github.io>) using the .mgf and .csv files

obtained with MZmine2 analysis. The molecular network of ESI–datasets was generated using cosine score thresholds of 0.60.

2.13. Metabolite annotation of untargeted metabolomic data

Metabolite annotation was performed in three consecutive steps. First, the obtained RT and m/z data of each feature were compared with our library containing the 6 standards based on their RT and m/z . Second, the ESI– metabolomic data used for molecular network analyses were searched against the available MS² spectral libraries (Massbank NA, GNPS Public Spectral Library, NIST14 Tandem, NIH Natural Product and MS-Dial), with absolute m/z tolerance of 0.02, 4 minimum matched peaks and minimal cosine score of 0.60. Third, not-annotated metabolites that belong to molecular network clusters containing annotated metabolites from steps 1 and 2 were assigned to the same chemical family and annotation was carried out on the basis of MS/MS spectrum comparisons.

3. Results

3.1. Heterologous expression vectors and modification of the yeast recipient strain

In their study, Hoefgen et al. (2018) described the polycistronic plasmid pV2A-T designed to express multiple secondary metabolism (SM) genes under the control of a single promoter. In this polycistronic system each gene is separated by a TEV-P2A sequence. The P2A sequence encodes a self-cleaving peptide releasing the upstream protein with a 33 amino acids tail in C-term and the downstream protein with a proline in N-term. The TEV peptide is recognized and cut by the TEV-protease enzyme, reducing the C-term tail to 6 amino acids. Each polycistron contained the *VenusN* and *VenusC* genes on the first and last position of the coding sequence, respectively. Because both genes contain a nuclear localization signal (NLS), when the polycistronic transcript is translated, the *VenusN* and *VenusC* proteins accumulate in the yeast nucleus, where they self-assemble to produce a yellow fluorescent protein. The presence of this fluorescence in the nucleus is thus an indicator of the production of the polycistronic proteins. By digestion with the *EcoRV* restriction enzyme, biosynthetic genes from the cluster of interest can be introduced between the *VenusN* and *VenusC* genes. Polycistronic plasmids containing the desired genes can then be fused after digestion by *SwaI* or *PmeI* (Hoefgen et al., 2018).

Here, we generated the plasmids pHYX137 and pHYX138, both of which can be used in *S. cerevisiae* (Fig. 1; Supplementary File 2). Both vectors harbour auto-inducible promoters from yeast, namely *pPCK1* or *pADH2*, respectively, which are repressed in the presence of glucose and activated after the diauxic shift during ethanol-anaerobic fermentation

(Harvey et al., 2018). This allows to disconnect biomass accumulation from SM production, which is a valuable feature when the SM are toxic. Both promoters are poorly induced in selective medium, in contrast to rich medium (Lee and DaSilva, 2005).

In addition to the polycistronic gene, the plasmids pHYX137 and pHYX138 possess a yeast 2 μ origin of replication and the pUC19 origin of replication was also replaced by the *ColE1* origin. The nutritional selection genes, either *LEU2* or *URA3*, were included and silent mutations were introduced to remove the *EcoRV* restriction sites. This allows to linearize the pHYX137 or pHYX138 with *EcoRV* and to clone individually the coding sequences by *in vivo* assembly (IVA) in *E. coli*, which is a fast, cost-effective and simple method. Likewise, the coding sequences are successively assembled in a single polycistron by digesting the plasmids with either *SwaI* or *PmeI* and direct transformation of *E. coli* with the unpurified plasmid fragments for IVA. This IVA method uses the natural DNA repair capability of *E. coli* to assemble DNA fragments with short overlapping homologies (18–25 bp), avoiding the use of expensive commercial kits or ligases.

To enhance the production of heterologous polyketides in the yeast *S. cerevisiae*, the strain BJ5464-npga was previously modified by the deletion of two vacuolar proteases, PEP4 and PRB1 (Lee et al., 2009) and the introduction of the *npgA* gene (Bond et al., 2016) involved in activating the ACP domain of PKS enzymes. Here, we used the strain BJNBC, a derivative of BJ5464-npga that we generated in the frame of a project involving BGCs with numerous cytochrome P450 enzymes. The modification involved the integration of an NADPH-cytochrome reductase from a filamentous fungus. This NADPH-cytochrome reductase was successfully used to enhance the heterologous production of *B. cinerea* abscisic acid in *S. cerevisiae* (Otto et al., 2019).

3.2. Fluorescence-based validation of the yeast heterologous expression system

To verify the proper transcription of the polycistronic gene, and correct translation and separation of the individual proteins, a gene coding for the mScarlet-I red fluorescent protein was introduced between the coding sequences of the *VenusN* and *VenusC* genes. The *mScarlet-I* gene was first introduced into plasmid pHYX137, giving the plasmid pHYX143, which was then transferred into the BJNBC yeast strain. The transformed yeast was cultivated 24 h in YNB medium supplemented with 2% (w/v) glucose without leucine. Epi-fluorescence microscopy revealed that the yellow fluorescence of *Venus* was present in the yeast nucleus, where it colocalized with the blue fluorescent DNA stain DAPI, whereas the red fluorescence of mScarlet-I, which lacked an NLS, was distributed throughout both the cytoplasm and nucleus (Fig. 2). These observations confirm that proteins encoded by the polycistronic gene had been well-transcribed and separately translated




Fig. 1. Features of the pV2A-T plasmid described by Hoefgen et al. (2018) and of the two plasmids, pHYX137 and pHYX138, adapted for *S. cerevisiae* expression, described in this study. Complete maps are shown in Supplementary File 2.




Fig. 2. Confocal microscopy of the yeast strain BJNBC-003 expressing Venus-NLS and mScarlet-I fluorescent proteins and stained with DAPI to detect DNA. A-D, Bar = 50 μm . E-H, from left to right, bright-field image showing the yeast cells, red fluorescence corresponding to mScarlet-I, yellow fluorescence corresponding to Venus and, blue fluorescence corresponding to DAPI. The mScarlet-I signal is distributed throughout the cell whereas the Venus signal is co-localized with DNA in the nucleus. Bar = 5 μm .

in yeast, and that Venus was correctly assembled in the yeast nucleus from the two complementary non-fluorescent fragments.

3.3. Selection of a biosynthetic pathway to test the expression system

Previously, we isolated several members of the Colletochlorin family of secondary metabolites from *C. higginsianum* and we proposed a

hypothetical biosynthetic pathway based on the isolated molecules and the plausible expected enzymatic activities (Dallery et al., 2019). In order to identify the biosynthetic gene cluster (BGC) responsible for producing Colletochlorins, we looked for putative halogenases (InterPro signature IPR006905) in the *C. higginsianum* IMI 349063 genome. The only BGC with an IPR006905 signature was BGC16, comprising genes CH63R_05468 to CH63R_05483 in the original prediction (Dallery et al.,

2017). Recently, Tsukada et al. (2020) reported the heterologous production of Higginsianins as well as other decalin-containing diterpenoid pyrones by expressing 8 of the 16 genes in BGC16. None of the Higginsianins are chlorinated and only one PKS (*ChPKS11*) was required for the biosynthesis of Higginsianins despite the presence of a second PKS (*ChPKS10*) in the cluster, suggesting the BGC16 comprises two BGCs side-by-side or intertwined. To test this hypothesis, we examined the conservation of BGC16 in 57 genome-sequenced *Colletotrichum* spp. (NCBI taxid 5455) using the cblaster tool (Fig. 3). Interestingly, the BGC16 was found in six species belonging to four different species complexes, namely *C. higginsianum* and *C. tanacetii* (Destructivum complex), *C. musicola* and *C. sojae* (Orchidearum complex), *C. spaethianum* (Spaethianum complex) and *C. chlorophyti*. In *C. tanacetii* and *C. sojae*, the BGC16 homologous genes were found respectively on two and three different contigs with each part being located at contig ends, suggesting problems of genome assembly rather than locations on different chromosomes. Interestingly, in *C. chlorophyti* only the genes required for making Higginsianin-like molecules were retrieved. Homologues of the genes CH63R_05468 to CH63R_05472 were absent from the *C. chlorophyti* NTL11 genome and were hypothesized to be involved in the biosynthesis of Colletochlorins (Fig. 3).

Using the cblaster tool, we investigated the conservation of genes putatively responsible for Colletochlorins biosynthesis (CH63R_05468 to CH63R_05472) in other fungi. Clustered gene homologues were found mostly in Sordariomycetes belonging to the *Glomerellaceae*, *Nectriaceae*, *Hypocreaceae* and *Stachybotryaceae* families (Fig. 4). Only three Eurotiomycetes had homologous BGCs (*Aspergillus ellipticus* and two *Talaromyces* spp.). None of the four homologous clusters found in *Stachybotrys* species contained a halogenase-encoding gene. Consistently, *Stachybotrys bisbyi* cultures produced only non-chlorinated prenylated derivatives of Orsellinic acid, notably LL-Z1272 β , also called Ilicicolin B (Li et al., 2016). Among the retrieved homologues, we also found the BGC in *Acremonium egyptiacum* responsible for biosynthesis of Ascochlorin, another prenylated yet chlorinated derivative of Orsellinic acid (Fig. 4). Based on these findings and knowledge of the experimentally-determined biosynthetic route for LL-Z1272 β and Ascochlorin, as well as the predicted pathway for Colletochlorins, we selected genes CH63R_05468 (*ctc2*, halogenase), CH63R_05469 (*ctc1*,

also known as *ChPKS10*), CH63R_05470 (*ctc3*, also known as *ChNRPS-like04*) and CH63R_05471 (*ctc4*, prenyltransferase) for heterologous expression in *S. cerevisiae*.

3.4. Heterologous production of Orsellinic acid in yeast

After careful examination of the gene models and intron borders using available RNA-Seq data, the coding sequences (CDS) of *ctc1* to *ctc4* were codon-optimized for *S. cerevisiae* and *de novo*-synthesized. Each CDS was then cloned individually into the plasmid pHYX137. A polycistron containing the four *ctc* CDS under control of the *PCK1* promoter of *S. cerevisiae* was constructed by successive digestions with either *SwaI* or *PmeI* followed by *in vivo* assembly (Transformation-Assisted Recombination, TAR) directly in *E. coli*. This plasmid was named pHYX164 and was transferred into the adapted yeast strain BJNBC, giving BJNBC-008. All strains and plasmids used in this study are described in Supplementary Files 1 and 4.

The yeast strains BJNBC-001 (containing the empty polycistronic plasmid pHYX137) and BJNBC-008 (harboring the Colletochlorin gene cluster in pHYX164) were cultured for three days and then metabolites were extracted and analysed by HPLC. Only one molecule was detected in BJNBC-008 that was not present in BJNBC-001. This molecule had a retention time of 16.5 min, a molecular weight of 168 and a UV spectrum with maxima at 227, 259 and 298 nm (Fig. 5). These same characteristics were also shown by an Orsellinic acid (OA) standard, indicating that the molecule detected in culture extracts is OA (Fig. 5). The presence of OA, the first molecule in the proposed biosynthetic pathway, but none of the other expected molecules, suggests that the polyketide synthase (CTC1) was functional but not the prenyltransferase (CTC4; Fig. 6).

All the enzymes expressed from this polycistron retain a P2A tag at their carboxyl terminus. To verify that all the enzymes in the pathway were present in the transgenic yeast BJNBC-008, we used immunoblotting with antibodies raised against the P2A peptide (Hoefgen et al., 2018). Protein samples were collected at two time-points, the first (t0) corresponded to when the culture reached an OD₆₀₀ of 0.4 and the culture medium was changed to a new one containing 2% glucose and 3% ethanol. At that point, the polycistronic gene under control of the




Fig. 3. General overview of the biosynthetic gene cluster BGC16 (A) and its conservation and microsynteny in genome-sequenced *Colletotrichum* spp. (B). The BGC is actually composed of two BGCs side-by-side or intertwined, one for biosynthesis of Colletochlorins and the other for biosynthesis of Higginsianins. The Higginsianin genes (dpch) were characterized by Tsukada et al. (2020). The gene CH63R_05474 is a pseudogene whereas the gene CH63R_05472 encodes a predicted transcription factor. When present, regions composed of repeated transposable elements are shown as double-headed arrows with their length. Vertical red bars denote contig ends. Fungal BGCs are often misassembled during genome sequencing and split over several contigs due to difficult-to-assemble long stretches of repeats. The intensity of grey/black shading represents the percentage of amino acid identity.




Fig. 4. Conservation and microsynteny of the Colletochlorin (*ctc*) biosynthetic genes cluster in fungi. In *Colletotrichum higginsianum* IMI 349063A, this BGC was initially predicted to be a single BGC16 but actually comprises the *dpch* (Higginsianins) and *ctc* BGCs located side-by-side. Genes are color-coded according to their putative or experimentally-confirmed functions. The intensity of grey/black shading represents the percentage of amino acid identity. Experimentally-verified BGCs are shown with one of their molecular products. Hy, Hypocreaceae; is, incertae sedis; Mg, Magnaporthaceae; St, Stachybotryaceae.

PCK1 promoter was expected to be repressed, as *PCK1* is repressed in glucose-containing media. The second time-point (t24) corresponds to 24 h after t0. At t24, all the glucose was supposed to be consumed by the yeast and ethanol fermentation had started (Lee and DaSilva, 2005), thus activating the polycistronic gene expression.

Before induction (t0), a band at 29 kDa corresponding to the VenusN protein was detected in both BJNBC-001 and BJNBC-008. The detection of VenusN in non-induced conditions (t0) showed that the polycistron was expressed at a very low level in these conditions. After induction (t24 h), only the VenusN protein was found in BJNBC-001, whereas in




Fig. 5. Monitoring the production of Colletochlorin biosynthetic intermediates using HPLC-PDA-ELSD-MS. (A) ELSD chromatogram of crude extracts of strains BJNBC-001 (empty vector) and BJNBC-008 (*ctc1* to *ctc4* genes expressed from a polycistron) and the Orsellinic acid standard. Only Orsellinic acid could be detected among the known intermediates of the Colletochlorin family. (B) UV spectra of the differential peak identified in BJNBC-008 (asterisk) and of the Orsellinic acid standard.

the BJNBC-008 yeast, four bands were detected at 29, 61, 119 and 233 kDa, corresponding to VenusN, CTC2, CTC3 and CTC1, respectively (Fig. 7). However, the prenyltransferase CTC4 (expected Mr = 40 kDa) was not detectable. This apparent absence of the prenyltransferase could explain why only the first molecule in the pathway, Orsellinic acid, was obtained from BJNBC-008 cultures.

3.5. Heterologous production of the colletochlorin metabolites in yeast

To overcome the absence of the CTC4 protein in the BJNBC-008 protein extract, a new plasmid pHYX172 was made containing only the prenyltransferase gene under the control of the strong and constitutive *TEF1* promoter. Two new yeast strains were generated: BJNBC-015 containing plasmids pYHX164 (polycistron with Colletochlorin genes) and pYHX172 (*ctc4* alone), and as a control, the strain BJNBC-017 containing the pYHX137 and pYHX138 empty vectors. Metabolites were extracted from 3-day-old cultures and then analysed by LC-

QToF-MS.

The expected molecular ions corresponding to Colletochlorins and Colletochlorins were readily found in samples from BJNBC-015 and with retention times and masses similar to those for the purified standards Orsellinic acid (RT, 6.40; m/z 167.0344 [M-H]⁻), Colletochlorin D acid (RT, 13.51; m/z 235.0970 [M-H]⁻), Colletochlorin D (RT, 15.22; m/z 219.1021 [M-H]⁻), Colletochlorin B (RT, 20.02; m/z 321.1257 [M-H]⁻) and Colletochlorin D (RT, 17.06; m/z 235.0631 [M-H]⁻) (Fig. 8).

Next, we performed a non-targeted analysis, comparing the BJNBC015 extract with the control BJNBC-017 extract. This confirmed the detection of all the standard molecules previously described in the BJNBC-015 extract. In addition, another molecule (RT, 18.71; m/z 287.1666 [M-H]⁻) was detected. The molecular mass of this molecule corresponds to that of Colletochlorin B, the non-chlorinated form of Colletochlorin B. In order to validate this hypothesis, we carried out a comparative analysis of the fragmentation pattern of Colletochlorin B and Colletochlorin B (Fig. 9; Supplementary File 5). Taken together, the




Fig. 6. Proposed biosynthetic pathways of the Colletochlorins. (A) Domain structure of CTC1 and CTC3 proteins. (B) Hypothetical scenario 1 where the CTC4 prenyltransferase accepts only DMAPP (dimethylallylpyrophosphate) as isoprene donor. (C) Hypothetical scenario 2 where the CTC4 prenyltransferase accepts both DMAPP and GPP (geranylpyrophosphate) as isoprene donors. Domains: ACP, acyl-carrier protein; AT, acyl transferase; KS, ketosynthase; PT, product template; SAT, starter-unit acyltransferase; TE, thioesterase; A, adenylation; PCP, peptidyl-carrier protein; R, reduction.

results show that Orsellinic acid, Colletorin D acid, Colletorin B and D, and Colletochlorin B and D were detected in the yeast harbouring genes *ctc1* to *ctc4* of *C. higginsianum* BGC16, and validated that the proposed *ctc* gene cluster does indeed encode the Colletochlorin biosynthetic pathway. The *C. higginsianum* *ctc* BGC was deposited at the MIBiG repository with accession number BGC0002812.

For relative quantification of Colletochlorins and their biosynthetic intermediates, crude extracts of the BJNBC-015 culture supernatants and cell pellets were analysed. Orsellinic acid and Colletorin D acid were largely found in the supernatant and nearly absent in cell pellets. The chlorinated end products, Colletochlorin B and D, were predominantly found in the cell pellets (Fig. 10). In extracts of BJNBC-015, Colletochlorin B production was 8.2-fold higher than in extracts of *C. higginsianum*, the original producer of the Colletochlorins. Likewise, in yeast cell pellets the production of Colletorin D and Colletochlorin D was, respectively, 35 and 3.3-fold higher than in *C. higginsianum*. Moreover, Orsellinic acid and Colletorin D acid were not detected in extracts of *C. higginsianum* (Fig. 10). Taken together, these results showed the utility of the system to overproduce molecules of fungal

origin and to produce biosynthetic intermediates.

4. Discussion

Fungi are a huge and underestimated reservoir of bioactive natural products. While genetic and chemical manipulations of fungi are common strategies to activate biosynthetic pathways in laboratory conditions, they require extensive trial-and-error. In order to facilitate the discovery of new fungal secondary metabolites, we developed the system described in this study to provide an easily-applicable tool for the heterologous expression of entire secondary metabolite gene clusters in engineered *S. cerevisiae*. The expression system should facilitate the discovery of new fungal secondary metabolites, especially those produced by silent BGCs, or BGCs that are only expressed at low levels *in vitro* or uniquely during interactions with the host plant. Apart from the possible discovery of high-value natural products including new medicines or biopesticides, this will provide a better understanding of the ecological role of these molecules, including their contribution to the pathogenesis of plant pathogens.




Fig. 7. Immunodetection of the CTC proteins and VenusN. (A) Immunoblot of whole-cell protein extracts from the strains BJNBC-001 (empty vector) and BJNBC-008 (ctc genes) at t0 (optical density of 0.4; repressive medium replaced by inductive medium) and t24 after induction. The proteins were detected with an anti-2A antibody. Note that VenusN is present also in the empty vector. Equal loading was assessed using TGX Stain-Free gels. (B) Schematic representation of the expected proteins with their size.

The expression system was firstly validated in yeast by introducing the *mScarlet* reporter gene into the adapted polycistronic plasmid. We then introduced the genes coding the key and tailoring enzymes (CTC1 to CTC4) of *C. higginsianum* BGC16 and showed that the resulting yeast cultures heterologously produced Orsellinic acid, Colletorin D acid, Colletochlorin B and D and Colletorin B. The chemical structure of these metabolites validates the correct production and enzymatic activity of the PKS, prenyltransferase, NRPS-like and halogenase enzymes in the heterologous system. Moreover, it demonstrates that *ctc1* to *ctc4* encode all the enzymes necessary for the biosynthesis of Colletochlorins B and D, Colletorin B and D, Colletorin D acid and Orsellinic acid. Importantly, we demonstrated that these molecules were all produced in greater quantities when heterologously expressed in yeast compared to the natural producer of these molecules, *C. higginsianum*. Using our approach, we could produce two known biosynthetic intermediates that were not detected in the wild-type strain of *C. higginsianum*, namely Orsellinic acid and Colletorin D acid and one intermediate, Colletorin B, that, to the best of our knowledge, was never described from any *Colletotrichum* species producing this family of molecules.

Our system offers several important advantages for the heterologous expression of fungal secondary metabolites. Firstly, the host organism is *S. cerevisiae* which is easily cultured on a large scale, produces few unwanted small molecules that may cause background noise and lacks endogenous BGCs that could cause crosstalk with heterologous pathways or deplete substrate pools. Secondly, our vectors rely on strong auto-inducible and co-regulated yeast promoters, namely *pADH2* and *pPCK1*, each with a different selection marker. Being activated after the diauxic shift (Harvey et al., 2018), these promoters could allow the production of toxic metabolites by enabling yeast biomass to increase before potentially toxic metabolites start to be produced. Our expression system provides several advantages over classical cloning strategies (e.g. multiple plasmids with bidirectional promoters). Firstly, polycistron-based systems allow to synchronously express the enzymes from a single promoter. Secondly, the pHYX vectors described here take advantage of the split Venus reporter from the pV2A-T vector of Hoefgen et al. (2018), which allows real-time monitoring of polycistron expression. At a practical level, our system provides versatile cloning of various combinations of genes into a polycistron, facilitating the dissection of biosynthetic pathways. This is especially true when coupled with *in vitro* assembly in *E. coli*, which eliminates the need for cloning kits.

There are relatively few reports of the use of polycistronic

biosynthetic pathways in yeast hosts. One notable example was the pioneering study of Beekwilder et al. (2014) where a polycistronic 2A sequence was used for the production of β -carotene from three genes separated by the *Thosea asigna* virus 2A sequence. More recently, Mukherjee and Wang (2023) described a flexible polycistronic-based system using modular cloning (MoClo) with Type IIS restriction enzymes. However, these authors also reported that fluorescent proteins with a C-terminal 2A peptide showed significantly reduced fluorescence, raising the possibility that this viral peptide could interfere with protein functionality.

In the present study, we likewise encountered a loss of functionality of the prenyltransferase CTC4 when expressed from a 2A-based polycistron. This enzyme was a predicted UbiA-like membrane-bound prenyltransferase, possessing the typical NDXDXXXD motif and potentially seven transmembrane domains according to TMHMM (Chang et al., 2021; Krogh et al., 2001). One hypothesis explaining a non-functional prenyltransferase is the absence of insertion of the protein into the membrane or misfolding followed by rapid degradation of the protein. Although not detected by the signal peptide predictor SignalP (Petersen et al., 2011), this enzyme is predicted to be localized at the plasma membrane by WoLF PSORT (Horton, 2007). Generally, proteins destined to be transported into the membrane are synthesized with a targeting-sequence, usually at the N-terminus and possibly at the C-terminus (Schatz and Dobberstein, 1996). The porcine tescho virus P2A self-cleaving peptide (P2A) has the peculiarity to add 21 amino acids at the C-terminus of the upstream protein and a proline at the N-terminus of the downstream protein (Kim et al., 2011). The presence of these extra residues likely caused the loss of function observed for the prenyltransferase CTC4. An example of non-recognition of an N-terminal signal peptide due to the presence of the P2A proline was reported previously for an N-myristoylated protein (Hadpech et al., 2018).

Another hypothesis could be that the prenyltransferase is correctly translocated into the membrane but the catalytic site of the enzyme is non-active due to presence of the P2A tail, as was reported by Mattern et al. (2017) for an O-acetyltransferase. However, this explanation is less probable because in that case the prenyltransferase would have been detected by Western blot. Whatever the explanation, proper functioning of the prenyltransferase appears to have been prevented by the P2A peptide, because Colletochlorins were successfully produced when the prenyltransferase was expressed from a separate plasmid with a conventional construction involving a strong promoter and terminator, in




Fig. 8. Combined extracted ion chromatograms (EIC) of the culture supernatant of the strain BJNBC-015 expressing all four *ctc* genes in a polycistron and an additional copy of *ctc4* (prenyltransferase) with its own promoter. EIC of standards of the different biosynthetic intermediates in the Colletochlorin pathway are also represented.

addition to the polycistronic plasmid containing the other *ctc* genes. In similar studies involving prenyltransferases, or other enzymes that are targeted to specific subcellular locations or membranes, we recommend to check for the presence of predicted signal peptide and transmembrane domains in the enzymes of each pathway of interest and to clone genes encoding this type of enzyme in a separate plasmid. Alternatively, the TEV protease may be used to cut the P2A C-terminal tail to avoid interference with the enzyme activity.

Another possible limitation of expression from a polycistron is the position or the number of genes to be expressed. The positional effect was highlighted by Jiao et al. (2018) who found that the first gene in the polycistron was more highly expressed than the two following ones. Two other studies likewise found that gene expression level progressively decreases with distance from the N-terminus of the polycistron (Liu et al., 2017; Mukherjee and Wang, 2023). For BGCs containing numerous genes, distributing them between two polycistronic plasmids may allow a more homogeneous expression level. For this purpose, we

designed two polycistronic plasmids pHYX137 and pHYX138 with different markers of prototrophy, namely *LEU2* and *URA3*, respectively.

In this study, we have formally linked the biosynthesis of Colletochlorin family of molecules to four genes within the BGC16 of *C. higginsianum*. Among these genes, one encoded a halogenase responsible for the chlorination of those secondary metabolites. Halogenases are valuable enzymes that can be used for the functionalization of drugs or other biologically active small-molecules (Crowe et al., 2021). Eight other genes in the BGC16 were previously assigned to Higginsianins biosynthesis (Tsukada et al., 2020), while the gene CH63R_05474 is a methyltransferase relict that underwent pseudogenization, CH63R_05472 is a putative transcription factor and CH63R_05481 has no characterized function. Two biosynthetic pathways for the production of Colletochlorins are possible (Fig. 6). Various lines of evidence suggest that the second pathway is the most probable. Li et al. (2016) described the cluster involved in LL-Z1272 β (Ilicicolin B) synthesis, which contains three genes coding for a PKS *StbA*, a




Fig. 9. LC-MS/MS fragmentation pattern of, A, a Colletochlorin B standard and, B, a molecule annotated as Colletorin B from the BJNBC-015 strain. Detailed fragmentation patterns are presented in Supplementary File 5.

prenyltransferase *StbC* and an NRPS-like *StbB*. These authors showed that the PKS is involved in the formation of Orsellinic acid, which is then converted into Grifolic acid by the prenyltransferase. The NRPS-like *StbB* is only able to convert the prenylated form of Orsellinic acid (i.e. Grifolic acid) into LL-Z1272 β and does not accept Orsellinic acid as a substrate. In the literature, NRPS-like enzymes that require prenylated substrates have been rarely described. Comparison of the Adenylation domains of the NRPS-like accepting Orsellinic acid (ATEG_03630) or only prenylated-Orsellinic acid (*StbB*) as substrate showed differences in their protein sequence. At position 334, ATEG_03630 possesses a leucine and *StbB* a glycine, while at position 358, essential for ATEG_03630 substrate specificity (Wang and Zhao, 2014), ATEG_03630 has a histidine and *StbB* a phenylalanine. The *C. higginsianum* NRPS-like enzyme CTC3 has the same amino acids involved in substrate specificity as *StbB*, suggesting that it may have a similar substrate specificity towards prenylated Orsellinic acid. Finally, the proposed ability of prenyltransferase CTC4 to accept both DMAPP and GPP moieties as substrates is known to occur in other aromatic prenyltransferases (Chen et al., 2017; Cheng and Li, 2014; Kalén et al., 1990; Suzuki et al., 1994; Swiezewska et al., 1993). Further experiments are now needed to confirm the

Colletochlorin biosynthetic pathway, notably by purifying the prenyltransferase and NRPS-like enzymes for assessing their substrate specificity.

The Colletochlorins were previously isolated from a *C. higginsianum* mutant with a partially deficient COMPASS complex (Dallery et al., 2019) and several of them were shown previously to be biologically active. For example, Colletorin B and Colletochlorin B displayed moderate herbicidal, antifungal and antibacterial activities towards *Chlorella fusca*, *Ustilago violacea*, *Fusarium oxysporum*, and *Bacillus megaterium* (Hussain et al., 2015), while Colletochlorin B had a significant antibacterial effect against *Bacillus subtilis* (minimum inhibitory concentration, 2 $\mu\text{g}\cdot\text{mL}^{-1}$) (Kemkuignou et al., 2022).

5. Conclusions

Our findings demonstrate the utility of this synthetic biology tool for the metabolic engineering of yeast to produce fungal metabolites from BGCs of interest in bulk liquid cultures. This is a prerequisite for subsequent structural characterization and bioactivity profiling of SM products from BGCs that are otherwise silent in their native organisms




Fig. 10. Relative abundance of Colletochlorins and their biosynthetic intermediates by LC-QToF. Crude extracts of the culture supernatants and cell pellets were analysed for *S. cerevisiae* cultures whereas crude extracts of the mixed supernatant and mycelia was analysed for the organism of origin of the ctc BGC, *C. higginsianum* (Ch63A). Extracts from an empty-vector strain (no biosynthetic genes) were included as controls for the yeast. The values from two independent experiments are presented and are expressed as relative abundance per mg of crude extract, resuspended in methanol.

when cultured in laboratory conditions.

Funding

This work was supported by the ‘Département de Santé des Plantes et Environnement’ (SPE) of INRAE (grant ‘Appel à projets scientifiques SPE 2021’ to JFD and MV). AGK was supported by a doctoral grant from Saclay Plant Sciences-SPS. This work was partly funded by Agence Nationale de la Recherche (reference n°ANR-16-CE20-0023) to MV and has benefited from a French State grant (Saclay Plant Sciences, reference n°ANR-17-EUR-0007, EUR SPS-GSR) under a France 2030 program (reference n°ANR-11-IDEX-0003). The Funders had no role in study design, the collection, analysis and interpretation of data, or writing of the manuscript.

CRedit authorship contribution statement

Aude Geistodt-Kiener: Methodology, Investigation, Formal analysis, Visualization, Writing – original draft, Writing – review & editing. **Jean Chrisologue Totozafy:** Investigation, Formal analysis, Visualization, Writing – original draft. **Géraldine Le Goff:** Investigation, Formal analysis. **Justine Vergne:** Investigation, Visualization. **Kaori Sakai:** Investigation, Visualization. **Jamal Ouazzani:** Resources, Supervision, Funding acquisition. **Grégory Mouille:** Resources, Supervision, Funding acquisition. **Muriel Viaud:** Conceptualization, Writing – original draft, Writing – review & editing, Supervision, Funding acquisition. **Richard J. O’Connell:** Conceptualization, Writing – original draft, Writing – review & editing, Supervision. **Jean-Félix Dallery:** Conceptualization, Methodology, Investigation, Formal analysis, Visualization, Writing – original draft, Writing – review & editing, Supervision, Funding acquisition, Project administration.

Declaration of competing interest

The authors declare no conflict of interest.

Data availability

Data will be made available on request.

Acknowledgments

This work has benefited from the support of IJPB’s ‘‘Plant Observatory – Chemistry and Metabolism’’ platform. The authors would like to thank Axel A. Brakhage and Maria Stroe (Hans Knoll Institute, Jena, Germany), Nancy DaSilva (UC Irvine, California) and Verena Siewers (Chalmers Univ. Of Technology, Gothenburg, Sweden) for kindly providing the pV2A-T plasmid, the BJ5464-NpgA strain and the XI-3-pCfb2904-bccpr1 plasmid, respectively. The EasyClone-MarkerFree Vector Set was a gift from Irina Borodina (Addgene kit #1000000098).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ymben.2023.10.002>.

References

- Ahmed, Y., Rebets, Y., Estévez, M.R., Zapp, J., Myronovskiy, M., Luzhetskyy, A., 2020. Engineering of *Streptomyces lividans* for heterologous expression of secondary metabolite gene clusters. *Microb. Cell Factories* 19, 5.
- Beekwilder, J., van Rossum, H.M., Koopman, F., Sonntag, F., Buchhaupt, M., Schrader, J., Hall, R.D., Bosch, D., Pronk, J.T., van Maris, A.J.A., Daran, J.-M., 2014. Polycistronic expression of a β -carotene biosynthetic pathway in *Saccharomyces cerevisiae* coupled to β -ionone production. *J. Biotechnol.* 192, 383–392.
- Blin, K., Shaw, S., Kloosterman, A.M., Charlop-Powers, Z., van Wezel, G.P., Medema, M.H., Weber, T., 2021. antiSMASH 6.0: improving cluster detection and comparison capabilities. *Nucleic Acids Res.* 49, W29–w35.

- Bode, H.B., Bethe, B., Höfs, R., Zecek, A., 2002. Big effects from small changes: possible ways to explore nature's chemical diversity. *Chembiochem* 3, 619–627.
- Bond, C., Tang, Y., Li, L., 2016. Saccharomyces cerevisiae as a tool for mining, studying and engineering fungal polyketide synthases. *Fungal Genet. Biol.* 89, 52–61.
- Bond, C.M., Tang, Y., 2019. Engineering *Saccharomyces cerevisiae* for production of simvastatin. *Metab. Eng.* 51, 1–8.
- Chambers, M.C., Maclean, B., Burke, R., Amodei, D., Ruderman, D.L., Neumann, S., Gatto, L., Fischer, B., Pratt, B., Egertson, J., Hoff, K., Kessner, D., Tasman, N., Shulman, N., Frewen, B., Baker, T.A., Brusniak, M.-Y., Paulse, C., Creasy, D., Flashner, L., Kani, K., Moulding, C., Seymour, S.L., Nuwaysir, L.M., Lefebvre, B., Kuhlmann, F., Roark, J., Rainer, P., Detlev, S., Hemenway, T., Huhmer, A., Langridge, J., Connolly, B., Chadick, T., Holly, K., Eckels, J., Deutsch, E.W., Moritz, R.L., Katz, J.E., Agus, D.B., MacCoss, M., Tabb, D.L., Mallick, P., 2012. A cross-platform toolkit for mass spectrometry and proteomics. *Nat. Biotechnol.* 30, 918–920.
- Chang, H.-Y., Cheng, T.-H., Wang, A.H.J., 2021. Structure, catalysis, and inhibition mechanism of prenyltransferase. *IUBMB Life* 73, 40–63.
- Chen, R., Gao, B., Liu, X., Ruan, F., Zhang, Y., Lou, J., Feng, K., Wunsch, C., Li, S.-M., Dai, J., Sun, F., 2017. Molecular insights into the enzyme promiscuity of an aromatic prenyltransferase. *Nat. Chem. Biol.* 13, 226–234.
- Cheng, W., Li, W., 2014. Structural insights into ubiquinone biosynthesis in membranes. *Science* 343, 878–881.
- Chiang, Y.-M., Oakley, C.E., Ahuja, M., Entwistle, R., Schultz, A., Chang, S.-L., Sung, C.T., Wang, C.C.C., Oakley, B.R., 2013. An efficient system for heterologous expression of secondary metabolite genes in *Aspergillus nidulans*. *J. Am. Chem. Soc.* 135, 7720–7731.
- Chiang, Y.M., Szewczyk, E., Davidson, A.D., Keller, N., Oakley, B.R., Wang, C.C., 2009. A gene cluster containing two fungal polyketide synthases encodes the biosynthetic pathway for a polyketide, asperfuranone, in *Aspergillus nidulans*. *J. Am. Chem. Soc.* 131, 2965–2970.
- Cochrane, R.V.K., Sanichar, R., Lambkin, G.R., Reiz, B., Xu, W., Tang, Y., Vederas, J.C., 2016. Production of new cladospirin analogues by reconstitution of the polyketide synthases responsible for the biosynthesis of this antimalarial agent. *Angew. Chem. Int. Ed.* 55, 664–668.
- Collemare, J., O'Connell, R., Lebrun, M.-H., 2019. Nonproteinaceous effectors: the terra incognita of plant–fungal interactions. *New Phytol.* 223, 590–596.
- Crowe, C., Molyneux, S., Sharma, S.V., Zhang, Y., Kotsi, D.S., Connaris, H., Goss, R.J.M., 2021. Halogenases: a palette of emerging opportunities for synthetic biology–synthetic chemistry and C–H functionalisation. *Chem. Soc. Rev.* 50, 9443–9481.
- Dallery, J.-F., Lapalu, N., Zampounis, A., Pigné, S., Luyten, I., Amselem, J., Wittenberg, A.H.J., Zhou, S., de Queiroz, M.V., Robin, G.P., Auger, A., Hainaut, M., Henrissat, B., Kim, K.-T., Lee, Y.-H., Lespinet, O., Schwartz, D.C., Thon, M.R., O'Connell, R.J., 2017. Gapless genome assembly of *Colletotrichum higginsianum* reveals chromosome structure and association of transposable elements with secondary metabolite gene clusters. *BMC Genom.* 18, 667.
- Dallery, J.F., Le Goff, G., Adelin, E., Iorga, B.I., Pigne, S., O'Connell, R.J., Ouazzani, J., 2019. Deleting a chromatin remodeling gene increases the diversity of secondary metabolites produced by *Colletotrichum higginsianum*. *J. Nat. Prod.* 82, 813–822.
- Gao, L., Cai, M., Shen, W., Xiao, S., Zhou, X., Zhang, Y., 2013. Engineered fungal polyketide biosynthesis in *Pichia pastoris*: a potential excellent host for polyketide production. *Microb. Cell Factories* 12, 77.
- Gilchrist, C.L.M., Booth, T.J., van Wersch, B., van Grieken, L., Medema, M.H., Chooi, Y.-H., 2021. cblaster: a remote search tool for rapid identification and visualization of homologous gene clusters. *Bioinf. Adv.* 1.
- Gilchrist, C.L.M., Chooi, Y.-H., 2021. Clinker & clustermap.js: automatic generation of gene cluster comparison figures. *Bioinformatics* 37, 2473–2475.
- Gomez-Escribano, J.P., Bibb, M.J., 2011. Engineering *Streptomyces coelicolor* for heterologous expression of secondary metabolite gene clusters. *Microb. Biotechnol.* 4, 207–215.
- Hadpech, S., Jinathep, W., Saoin, S., Thongkum, W., Chupradit, K., Yasamut, U., Moonmuang, S., Tayapiwatana, C., 2018. Impairment of a membrane-targeting protein translated from a downstream gene of a “self-cleaving” T2A peptide conjunction. *Protein Expr. Purif.* 150, 17–25.
- Han, X., Chakraborti, A., Zhu, J., Liang, Z.-X., Li, J., 2016. Sequencing and functional annotation of the whole genome of the filamentous fungus *Aspergillus westerdijkiae*. *BMC Genom.* 17, 633.
- Harvey, C.J.B., Tang, M., Schlecht, U., Horecka, J., Fischer, C.R., Lin, H.-C., Li, J., Naughton, B., Cherry, J., Miranda, M., Li, Y.F., Chu, A.M., Hennessy, J.R., Vandova, G.A., Inglis, D., Aiyar, R.S., Steinmetz, L.M., Davis, R.W., Medema, M.H., Sattely, E., Khosla, C., St Onge, R.P., Tang, Y., Hillenmeyer, M.E., 2018. HEX: a heterologous expression platform for the discovery of fungal natural products. *Sci. Adv.* 4, eaar5459.
- Henehan, M.N., Yakasai, A.A., Halo, L.M., Song, Z., Bailey, A.M., Simpson, T.J., Cox, R. J., Lazarus, C.M., 2010. First heterologous reconstruction of a complete functional fungal biosynthetic multigene cluster. *Chembiochem* 11, 1508–1512.
- Hewage, R.T., Aree, T., Mahidol, C., Ruchirawat, S., Kittakoop, P., 2014. One strain-many compounds (OSMAC) method for production of polyketides, azaphilones, and an isochromanone using the endophytic fungus *Dothideomycete* sp. *Phytochemistry* 108, 87–94.
- Hoefgen, S., Lin, J., Fricke, J., Stroe, M.C., Mattern, D.J., Kufs, J.E., Hortschansky, P., Brakhage, A.A., Hoffmeister, D., Valiante, V., 2018. Facile assembly and fluorescence-based screening method for heterologous expression of biosynthetic pathways in fungi. *Metab. Eng.* 48, 44–51.
- Horton, P., 2007. WoLF PSORT: protein localization predictor. *Nucleic Acids Res.* 35, W585–W587.
- Hussain, H., Drogies, K.-H., Al-Harrasi, A., Hassan, Z., Shah, A., Rana, U.A., Green, I.R., Draeger, S., Schulz, B., Krohn, K., 2015. Antimicrobial constituents from endophytic fungus *Fusarium* sp. *Asian Pac. J. Trop. Dis.* 5, 186–189.
- Inglis, D.O., Binkley, J., Skrzypek, M.S., Arnaud, M.B., Cerqueira, G.C., Shah, P., Wymore, F., Wortman, J.R., Sherlock, G., 2013. Comprehensive annotation of secondary metabolite biosynthetic genes and gene clusters of *Aspergillus nidulans*, *A. fumigatus*, *A. niger* and *A. oryzae*. *BMC Microbiol.* 13, 23.
- Ishuchi, K.I., Nakazawa, T., Ookuma, T., Sugimoto, S., Sato, M., Tsunematsu, Y., Ishikawa, N., Noguchi, H., Hotta, K., Moriya, H., Watanabe, K., 2012. Establishing a new methodology for genome mining and biosynthesis of polyketides and peptides through yeast molecular genetics. *Chembiochem* 13, 846–854.
- Jessop-Fabre, M.M., Jakóciunas, T., Stovicek, V., Dai, Z., Jensen, M.K., Keasling, J.D., Borodina, I., 2016. EasyClone-MarkerFree: a vector toolkit for marker-less integration of genes into *Saccharomyces cerevisiae* via CRISPR-Cas9. *Biotechnol. J.* 11, 1110–1117.
- Jiao, X., Sun, W., Zhang, Y., Liu, X., Zhang, Q., Wang, Q., Zhang, S., Zhao, Z.K., 2018. Exchanging the order of carotenogenic genes linked by porcine teschovirus-1 2A peptide enable to optimize carotenoid metabolic pathway in *Saccharomyces cerevisiae*. *RSC Adv.* 8, 34967–34972.
- Kalén, A., Appelkvist, E.L., Chojnacki, T., Dallner, G., 1990. Nonaprenyl-4-hydroxybenzoate transferase, an enzyme involved in ubiquinone biosynthesis, in the endoplasmic reticulum-Golgi system of rat liver. *J. Biol. Chem.* 265, 1158–1164.
- Kealey, J.T., Liu, L., Santi, D.V., Betlach, M.C., Barr, P.J., 1998. Production of a polyketide natural product in nonpolyketide-producing prokaryotic and eukaryotic hosts. *Proc. Natl. Acad. Sci. U.S.A.* 95, 505–509.
- Keller, N.P., 2019. Fungal secondary metabolism: regulation, function and drug discovery. *Nat. Rev. Microbiol.* 167–180.
- Kemkuignou, B.M., Moussa, A.Y., Decock, C., Stadler, M., 2022. Terpenoids and meroterpenoids from cultures of two grass-associated species of *amyloporus* (basidiomycota). *J. Nat. Prod.* 85, 846–856.
- Khalidi, N., Seifuddin, F.T., Turner, G., Haft, D., Nierman, W.C., Wolfe, K.H., Fedorova, N. D., 2010. SMURF: genomic mapping of fungal secondary metabolite clusters. *Fungal Genet. Biol.* 47, 736–741.
- Kim, J.H., Lee, S.-R., Li, L.-H., Park, H.-J., Park, J.-H., Lee, K.Y., Kim, M.-K., Shin, B.A., Choi, S.-Y., 2011. High cleavage efficiency of a 2A peptide derived from porcine teschovirus-1 in human cell lines, zebrafish and mice. *PLoS One* 6, e18556.
- Knop, M., Siegers, K., Pereira, G., Zachariae, W., Winsor, B., Nasmyth, K., Schiebel, E., 1999. Epitope tagging of yeast genes using a PCR-based strategy: more tags and improved practical routines. *Yeast* 15, 963–972.
- Krogh, A., Larsson, B., von Heijne, G., Sonnhammer, E.L.L., 2001. Predicting transmembrane protein topology with a hidden markov model: application to complete genomes. *J. Mol. Biol.* 305, 567–580.
- Lee, K.K.M., Silva, N.A.D., Kealey, J.T., 2009. Determination of the extent of phosphopantetheinylation of polyketide synthases expressed in *Escherichia coli* and *Saccharomyces cerevisiae*. *Anal. Biochem.* 394, 75–80.
- Lee, M.K., DaSilva, N.A., 2005. Evaluation of the *Saccharomyces cerevisiae* ADH2 promoter for protein synthesis. *Yeast* 22, 431–440.
- Li, C., Matsuda, Y., Gao, H., Hu, D., Yao, X.S., Abe, I., 2016. Biosynthesis of LL-z1272β: discovery of a new member of NRPS-like enzymes for aryl-aldehyde formation. *Chembiochem* 17, 904–907.
- Liang, X., Wang, B., Dong, Q., Li, L., Rollins, J.A., Zhang, R., Sun, G., 2018. Pathogenic adaptations of *Colletotrichum fungi* revealed by genome wide gene family evolutionary analyses. *PLoS One* 13, e0196303.
- Liu, Z., Chen, O., Wall, J.B.J., Zheng, M., Zhou, Y., Wang, L., Ruth Vaseghi, H., Qian, L., Liu, J., 2017. Systematic comparison of 2A peptides for cloning multi-genes in a polycistronic vector. *Sci. Rep.* 7, 2193–2193.
- Lyu, H.-N., Liu, H.-W., Keller, N.P., Yin, W.-B., 2020. Harnessing diverse transcriptional regulators for natural product discovery in fungi. *Nat. Prod. Rep.* 37, 6–16.
- Ma, S.M., Li, J.W.-H., Choi, J.W., Zhou, H., Lee, K.K.M., Moorthe, V.A., Xie, X., Kealey, J. T., Da Silva, N.A., Vederas, J.C., Tang, Y., 2009. Complete reconstitution of a highly reducing iterative polyketide synthase. *Science* 326, 589–592.
- Mattern, D.J., Valiante, V., Horn, F., Petzke, L., Brakhage, A.A., 2017. Rewiring of the austinoid biosynthetic pathway in filamentous fungi. *ACS Chem. Biol.* 12, 2927–2933.
- Mikkelsen, M.D., Buron, L.D., Salomonsen, B., Olsen, C.E., Hansen, B.G., Mortensen, U. H., Halkier, B.A., 2012. Microbial production of indolylglucosinolate through engineering of a multi-gene pathway in a versatile yeast expression platform. *Metab. Eng.* 14, 104–111.
- Mukherjee, M., Wang, Z.Q., 2023. A well-characterized polycistronic-like gene expression system in yeast. *Biotechnol. Bioeng.* 120, 260–271.
- Myers, O.D., Sumner, S.J., Li, S., Barnes, S., Du, X., 2017. One step forward for reducing false positive and false negative compound identifications from mass spectrometry metabolomics data: new algorithms for constructing extracted ion chromatograms and detecting chromatographic peaks. *Anal. Chem.* 89, 8696–8703.
- Nielsen, M.T., Nielsen, J.B., Anyaogu, D.C., Holm, D.K., Nielsen, K.F., Larsen, T.O., Mortensen, U.H., 2013. Heterologous reconstitution of the intact geodin gene cluster in *Aspergillus nidulans* through a simple and versatile PCR based approach. *PLoS One* 8, e72871.
- O'Connell, R., Herbert, C., Sreenivasaprasad, S., Khatib, M., Esquerre-Tugaye, M.T., Dumas, B., 2004. A novel *Arabidopsis*-*Colletotrichum* pathosystem for the molecular dissection of plant-fungal interactions. *Mol. Plant Microbe Interact.* 17, 272–282.
- O'Connell, R.J., Thon, M.R., Hacquard, S., Amyotte, S.G., Kleemann, J., Torres, M.F., Damm, U., Buiaite, E.A., Epstein, L., Alkan, N., Altmüller, J., Alvarado-Balderrama, L., Bauser, C.A., Becker, C., Birren, B.W., Chen, Z., Choi, J., Crouch, J. A., Duvick, J.P., Farman, M.A., Gan, P., Heiman, D., Henrissat, B., Howard, R.J., Kabbage, M., Koch, C., Kracher, B., Kubo, Y., Law, A.D., Lebrun, M.H., Lee, Y.H.,

- Miyara, I., Moore, N., Neumann, U., Nordstrom, K., Panaccione, D.G., Panstruga, R., Place, M., Proctor, R.H., Prusky, D., Rech, G., Reinhardt, R., Rollins, J.A., Rounsley, S., Schardl, C.L., Schwartz, D.C., Shenoy, N., Shirasu, K., Sikkakolli, U.R., Stuber, K., Sukno, S.A., Sweigard, J.A., Takano, Y., Takahara, H., Trail, F., van der Does, H.C., Voll, L.M., Will, I., Young, S., Zeng, Q., Zhang, J., Zhou, S., Dickman, M. B., Schulze-Lefert, P., Ver Loren van Themaat, E., Ma, L.J., Vaillancourt, L.J., 2012. Lifestyle transitions in plant pathogenic *Colletotrichum* fungi deciphered by genome and transcriptome analyses. *Nat. Genet.* 44, 1060–1065.
- Oberlie, N.R., McMillan, S.D., Brown, D.W., McQuade, K.L., 2018. Investigating the role of trehalose metabolism in resistance to abiotic stress in the filamentous fungus *Fusarium verticillioides*. *Faseb. J.* 32, 665.3–665.3.
- Oliveira, L., Chevrollier, N., Dallery, J.-F., O'Connell, R.J., Lebrun, M.-H., Viaud, M., Lepinet, O., 2023. CusProSe: a customizable protein annotation software with an application to the prediction of fungal secondary metabolism genes. *Sci. Rep.* 13, 1417.
- Olivon, F., Elie, N., Grelier, G., Roussi, F., Litaudon, M., Touboul, D., 2018. MetGem software for the generation of molecular networks based on the t-SNE algorithm. *Anal. Chem.* 90, 13900–13908.
- Otto, M., Teixeira, P.G., Vizcaino, M.I., David, F., Siewers, V., 2019. Integration of a multi-step heterologous pathway in *Saccharomyces cerevisiae* for the production of abscisic acid. *Microb. Cell Factories* 18, 205.
- Pahirulzaman, A.K., Williams, K., Lazarus, C.M., 2012. Chapter twelve - a toolkit for heterologous expression of metabolic pathways in *Aspergillus oryzae*. In: Hopwood, D.A. (Ed.), *Methods in Enzymology*, vol. 517. Academic Press, pp. 241–260.
- Petersen, T.N., Brunak, S., von Heijne, G., Nielsen, H., 2011. SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nat. Methods* 8, 785–786.
- Pfeifer, B.A., Admiraal, S.J., Gramajo, H., Cane, D.E., Khosla, C., 2001. Biosynthesis of complex polyketides in a metabolically engineered strain of *E. coli*. *Science* 291, 1790–1792.
- Pfeifer Blaine, A., Wang Clay, C.C., Walsh Christopher, T., Khosla, C., 2003. Biosynthesis of yersiniabactin, a complex polyketide-nonribosomal peptide, using *Escherichia coli* as a heterologous host. *Appl. Environ. Microbiol.* 69, 6698–6702.
- Pimentel-Elardo, S.M., Sørensen, D., Ho, L., Ziko, M., Bueler, S.A., Lu, S., Tao, J., Moser, A., Lee, R., Agard, D., Fairn, G., Rubinstein, J.L., Shoichet, B.K., Nodwell, J. R., 2015. Activity-independent discovery of secondary metabolites using chemical elicitation and cheminformatic inference. *ACS Chem. Biol.* 10, 2616–2623.
- Pohl, C., Polli, F., Schütze, T., Viggiano, A., Mózsik, L., Jung, S., de Vries, M., Bovenberg, R.A.L., Meyer, V., Driessen, A.J.M., 2020. A *Penicillium rubens* platform strain for secondary metabolite production. *Sci. Rep.* 10, 7630.
- Schatz, G., Dobberstein, B., 1996. Common principles of protein translocation across membranes. *Science* 271, 1519–1526.
- Schumacher, J., 2012. Tools for *Botrytis cinerea*: new expression vectors make the gray mold fungus more accessible to cell biology approaches. *Fungal Genet. Biol.* 49, 483–497.
- Shenouda, M.L., Ambilika, M., Skellam, E., Cox, R.J., 2022. Heterologous expression of secondary metabolite genes in *Trichoderma reesei* for waste valorization. *J. Fungi* 8, 355.
- Suzuki, K., Ueda, M., Yuasa, M., Nakagawa, T., Kawamukai, M., Matsuda, H., 1994. Evidence that *Escherichia coli* ubiA product is a functional homolog of yeast COQ2, and the regulation of ubiA gene expression. *Biosci. Biotechnol. Biochem.* 58, 1814–1819.
- Swiezewska, E., Dallner, G., Andersson, B., Ernster, L., 1993. Biosynthesis of ubiquinone and plastoquinone in the endoplasmic reticulum-Golgi membranes of spinach leaves. *J. Biol. Chem.* 268, 1494–1499.
- Terlouw, B.R., Blin, K., Navarro-Muñoz, J.C., Avalon, N.E., Chevrette, M.G., Egbert, S., Lee, S., Meijer, D., Recchia, M.J.J., Reitz, Z.L., van Santen, J.A., Selem-Mojica, N., Tørring, T., Zaroubi, L., Alanjary, M., Aleti, G., Aguilar, C., Al-Salih, S.A.A., Augustijn, H.E., Avelar-Rivas, J.A., Avitia-Domínguez, L.A., Barona-Gómez, F., Bernaldo-Agüero, J., Bielinski, V.A., Biermann, F., Booth, T.J., Carrion Bravo, V.J., Castelo-Branco, R., Chagas, F.O., Cruz-Morales, P., Du, C., Duncan, K.R., Gavriilidou, A., Gayraud, D., Gutiérrez-García, K., Haslinger, K., Helfrich, E.J.N., van der Hooft, J.J.J., Jati, A.P., Kalkreuter, E., Kalyvas, N., Kang, K.B., Kautsar, S., Kim, W., Kunjapur, A.M., Li, Y.X., Lin, G.M., Loureiro, C., Louwen, J.J.R., Louwen, N. L.L., Lund, G., Parra, J., Philmus, B., Pourmohsenin, B., Pronk, L.J.U., Rego, A., Rex, D.A.B., Robinson, S., Rosas-Becerra, L.R., Roxborough, E.T., Schorn, M.A., Scobie, D.J., Singh, K.S., Sokolova, N., Tang, X., Udway, D., Vigneshwari, A., Vind, K., Vromans, S., Waschulin, V., Williams, S.E., Winter, J.M., Witte, T.E., Xie, H., Yang, D., Yu, J., Zdouc, M., Zhong, Z., Collemare, J., Linington, R.G., Weber, T., Medema, M.H., 2023. MIBiG 3.0: a community-driven effort to annotate experimentally validated biosynthetic gene clusters. *Nucleic Acids Res.* 51, D603–d610.
- Tsukada, K., Shinki, S., Kaneko, A., Murakami, K., Irie, K., Murai, M., Miyoshi, H., Dan, S., Kawaji, K., Hayashi, H., Kodama, E.N., Hori, A., Salim, E., Kuraishi, T., Hirata, N., Kanda, Y., Asai, T., 2020. Synthetic biology based construction of biological activity-related library of fungal decalin-containing diterpenoid pyrones. *Nat. Comms* 11, 1830.
- Valero-Jiménez, C.A., Steentjes, M.B.F., Slot, J.C., Shi-Kunne, X., Scholten, O.E., van Kan, J.A.L., 2020. Dynamics in secondary metabolite gene clusters in otherwise highly syntenic and stable genomes in the fungal genus *Botrytis*. *Genome Biol. Evol.* 12, 2491–2507.
- van Santen, J.A., Poynton, E.F., Iskakova, D., McMann, E., Alsup, Tyler A., Clark, T.N., Fergusson, C.H., Fewer, D.P., Hughes, A.H., McCadden, C.A., Parra, J., Soldatou, S., Rudolf, J.D., Janssen, E.M.L., Duncan, K.R., Linington, R.G., 2022. The Natural Products Atlas 2.0: a database of microbially-derived natural products. *Nucleic Acids Res.* 50, D1317–D1323.
- von Barga, K.W., Niehaus, E.-M., Bergander, K., Brun, R., Tudzynski, B., Humpf, H.-U., 2013. Structure elucidation and antimicrobial activity of apicidin F: an apicidin-like compound produced by *Fusarium fujikuroi*. *J. Nat. Prod.* 76, 2136–2140.
- Voth, W.P., Wei Jiang, Y., Stillman, D.J., 2003. New 'marker swap' plasmids for converting selectable markers on budding yeast gene disruptions and plasmids. *Yeast* 20, 985–993.
- Wang, M., Zhao, H., 2014. Characterization and engineering of the adenylation domain of a NRPS-like protein: a potential biocatalyst for aldehyde generation. *ACS Catal.* 4, 1219–1225.
- Xue, Y., Kong, C., Shen, W., Bai, C., Ren, Y., Zhou, X., Zhang, Y., Cai, M., 2017. Methylophilic yeast *Pichia pastoris* as a chassis organism for polyketide synthesis via the full citrinin biosynthetic pathway. *J. Biotechnol.* 242, 64–72.
- Yu, D., Xu, F., Zi, J., Wang, S., Gage, D., Zeng, J., Zhan, J., 2013. Engineered production of fungal anticancer cyclooligomer depsipeptides in *Saccharomyces cerevisiae*. *Metab. Eng.* 18, 60–68.
- Yu, G., Sun, Y., Han, H., Yan, X., Wang, Y., Ge, X., Qiao, B., Tan, L., 2021. Coculture, an efficient biotechnology for mining the biosynthesis potential of macrofungi via interspecies interactions. *Front. Microbiol.* 12.
- Zhang, J.J., Moore, B.S., Tang, X., 2018. Engineering *Salinispora tropica* for heterologous expression of natural product biosynthetic gene clusters. *Appl. Microbiol. Biotechnol.* 102, 8437–8446.
- Zhao, M., Zhao, Y., Yao, M., Iqbal, H., Hu, Q., Liu, H., Qiao, B., Li, C., Skovbjerg, C.A.S., Nielsen, J.C., Nielsen, J., Frandsen, R.J.N., Yuan, Y., Boeke, J.D., 2020. Pathway engineering in yeast for synthesizing the complex polyketide bikaverin. *Nat. Comms* 11, 6197.