

Monoclonal Antibodies against Bovine β -Casein: Production and Epitope Characterization

D. Senocq, Didier Dupont, O. Rolet-Répécaud, D. Levieux

▶ To cite this version:

D. Senocq, Didier Dupont, O. Rolet-Répécaud, D. Levieux. Monoclonal Antibodies against Bovine β -Casein: Production and Epitope Characterization. Food and Agricultural Immunology, 2010, 13 (4), pp.213-224. 10.1080/09540100120094483. hal-04301391

HAL Id: hal-04301391 https://hal.inrae.fr/hal-04301391

Submitted on 23 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1	Production and epitopic characterisation of monoclonal
2	antibodies against bovine β-casein
3	
4	
5	
6	BY DANIEL SENOCQ, DIDIER DUPONT, ODILE ROLET-RÉPÉCAUD AND DIDIER LEVIEUX*
7	
8	INRA- Station de Recherches en Technologie et Analyses laitières BP 89 F-39801 Poligny Cedex France
9	* INRA- Station de Recherches sur la Viande 63122 Saint Genès-Champanelle France
10	
11	
12	
13	RUN TITLE: Monoclonal antibodies against bovine β -casein
14	
15	
16	
17	Author to whom correspondence should be addressed: DIDIER DUPONT, INRA SRTAL B.P. 89 F-39801 Poligny Cedex
18	France, fax +33 3 84 37 37 81; e-mail dupont@poligny.inra.fr
19	
20	
21	

1

2 SUMMARY. Twenty one murine monoclonal antibodies (MAbs) were produced against bovine β -casein (β -CN). Eight 3 epitopic groups of MAbs recognising six discrete determinants (4-8, 14-24, 33-48, 49-91, 177-183, 184-209) were 4 defined from ELISA reactivity with peptides and related homologues from 3 species (ovine, human and equine). 5 Thirteen distinct epitopes were discriminated by cross-examination of the MAbs reactivity in BIAcore. Determinants 14-24, 49-91 and 184-209 were found to encompass at lest 4, 2 and 4 epitopes respectively. No MAb was bovine 6 7 specific, but one at least discriminated ovine/caprine casein from bovine. Terminal sequences 1-28 and 177-209 included ~80% of the discriminated epitopes, and the most conserved ones. This panel of MAbs can provide specific 8 9 probes for β-CN proteolysis fragments.

1 The caseins make up the major protein fraction of bovine milk (~80%). They represent the products of 4 linked 2 genes, the caseins α_{S1} , a_{S2} , β and κ . Bovine casein β (β -CN) accounts for ~40% of the casein content. Its primary 3 structure is known from chemical and gene sequencing (Swaisgood, 1992). It consists of 209 residues (~24000 Da), with phosphorylated serine residues clustered at N-termini (5 mole P/mole). It occurs in milk as a mixture of several 4 5 variants (Ng-Kwai-Hang & Grosclaude, 1992). Eight variants (A¹, A², A³, B, C, D, E, F) differing by 2-5 single-point substitutions have been fully characterised, and variants A¹, A², and B account for the large majority of Western cattle 6 7 heterogeneity. None was found to arise from post-translational modification, although substitutions in variants C and D 8 cause defect of phosphorylation. Homologue β -casein was found in milks of all species. Its sequence in human, sheep, 9 rat, mouse, rabbit and other species indicates a rapid divergence arising from point mutations, without segmental 10 rearrangements (Stewart *et al.*, 1987). Bovine β -CN differs from the other closely related caseins by (i) a higher rate of 11 hydrophobic residues, (ii) a low phosphorylation rate and (iii) a high proline content (17%). Its secondary structure 12 reflects these characteristics (Swaisgood, 1992). Experimental and theoretical data agree on a low rate of periodic 13 structure (30%), especially α -helix (10%), but many proline-induced β -turns (34%)(Kumosinski *et al.*, 1993). Owing to 14 the impossibility to crystallise the molecule, secondary and tertiary structure (Kumosinski et al., 1993) could be only 15 predicted. All data suggest an organisation opposite to that of typical globular proteins: conserved domains with strictly 16 defined structure occur at both termini, and are linked by an open and flexible intermediate region.

17 The products of β -CN cleavage in milk reflect both its structure and the specificity of the native enzymes. The main 18 native protease, plasmin (EC 3.4.21.7), cleaves apart the hydrophilic and heat resistant domain β-CN (f1-28), the 19 intermediate region β -CN (f29-105/-107) and the hydrophobic C-termini β -CN (f106/108-209). These major fragments 20 occur in large amount in fresh milk (typically 10-20% of β-CN in weight) (Andrews, 1983; Andrews & Alichanidis, 1983). 21 Plasmin is associated with casein micelles, and is incorporated in dairy foods, e.g. cheeses, where it contributes 22 significantly to the induced proteolysis. The contribution of plasmin to β -CN proteolysis is thought to significantly 23 increase the release of peptides and hydrophobic amino acids in the sapid fraction of cheeses. The objective 24 monitoring of plasmin action could contribute to modelling desirable pathways of flavour development. Biochemical 25 fractionation of peptides is complex, and is not specific for β -CN (Ferranti *et al.*, 1997b). In contrast, immunochemical 26 detection has a great potential for descriptive analysis of proteolysis of specific casein fractions (Addeo et al., 1995). 27 Provided that specific probes are produced, the degradation of β -CN segments can be monitored and quantified 28 (Pizzano et al., 2000). Specific antibodies are currently raised against peptides (Pizzano et al., 1999), although the 29 reactivity of anti-peptide antibodies with the intact parent protein is ambiguous. Moreover, these reagents come in 30 limited supply, and may not be reproduced. In contrast, hybridoma technology (Köhler & Milstein, 1975) allows the 31 continuous production of monospecific antibodies, which corresponding epitopes might be investigated in detail 32 (Kobayashi *et al.*, 1991). β -CN presents many β -turn, and a high interspecies divergence which should contribute to its

1 immunogenicity. Four previous studies reported however only four specific MAbs (Nagaune et al., 1988; Kuzmanoff et 2 al., 1991; Oudshorn et al., 1994; Anguita et al., 1995)., and thus supported the current opinion that caseins are poorly 3 immunogenic. Since the MAbs raised to a protein usually vary in specificity and reactivity, a large collection is 4 preferable to select probes with pre-requisite characteristics. The objective of this study was to produce a large panel 5 of characterised MAbs against β-CN, as probes for its plasmin-induced fragments. Twenty-one murine MAbs were compared for isotypes, ELISA reactivity with proteins and peptides, and reactivity with β-CN in BIAcore. Thirteen non-6 7 redundant MAbs bound six discrete determinants distributed along β -CN sequence, with a marked bias in favour of the 8 termini.

1

MATERIALS AND METHODS

2 Reagents

Unless otherwise stated, chemicals of analytical grade were purchased from Prolabo (Merck eurolab, D-64293
 Darmstadt, Germany). BIAcore 3000 was used with registered chemicals (BIAcore AB, S-75450 Uppsala, Sweden).

Inactive plasminogen was isolated from fresh bovine blood according to the procedure of Deutsch & Mertz (1970),
and frozen at -20°C. Immediately prior to use, aliquots (70 g/l) were treated for 10 min at 37°C with urokinase (EC
3.4.21.73) (Choay, Sanofi Winthrop) at an enzyme:substrate ratio of 100 Ul/mg, and thus activated to plasmin (EC
3.4.21.7).

9

10 β -casein isolation

11 Individual milk was obtained from the genetically typed () herd of the Coopérative agricole Graindorge (14140

12 Livarot, France). β -CN from a cow typed as homozygous for β -CN A² was analysed in electrospray ionisation mass-

13 spectroscopy (ESI-MS) according to Léonil et al. (1995). It was found in agreement with the variant reported by

14 Jimenez-Flores et al. (1987), and its primary sequence (A² substituted for leucine at position 93) was used.

15 Milk fat was removed after 15 min centrifugation (1800 g, 4°C). Whole casein was obtained from defatted milk by 16 isoelectric precipitation at pH 4.6 according to the procedure of Wei & Whitney (1984).

17 Crude β -CN was obtained in gram amounts by the batch-wise ion exchange procedure of Leaver & Law (1992), 18 dialysed against water, and lyophilised. The purity determined on urea-polyacrylamide gel electrophoresis (PAGE) 19 (Andrews, 1983) was ~ 90%. Crude β -CN was purified to homogeneity by the fast protein liquid chromatography 20 (FPLC) ion exchange procedure of Andrews et al. (1985), scaled to fit a HR16/10 column (Pharmacia Biotech. AB, S-21 75450 Uppsala, Sweden). Briefly, 600 mg of crude β-CN were loaded onto 20 ml of Q Sepharose FF (Pharmacia), and 22 eluted at 1 ml/min with a linear gradient of NaCl over 0.09-0.32 M in 25 mM-Tris-HCl-4.5 M-urea-0.08 mM-DTT buffer. 23 pH 7.0. Fractions automatically collected over 5 min were checked in urea-PAGE for homogeneity, pooled, dialysed 24 against water, and lyophilised.

25

26 Other antigens

Individual mare's milk was obtained from the farm of the Lycée Agricole (21140 Semur-en-Auxois, France). Mare's whole casein was prepared like bovine's. β -CN from cow buffalo's milk (*Bubalus bubalis*) was a gift from D. Lebars (INRA, 78352 Jouy-en-Josas, France). Whole casein from cow's milk was purchased from Merck, that from ewe, goat, and human milk from Sigma (Sigma-Aldrich, 38297 St.Quentin-Fallavier, France), and bovine serum albumin (BSA) from Pierce (Interchim, 03103 Montluçon, France). Other bovine proteins were isolated from bovine bulk milk according to the following chromatographic procedures: α -Lactalbumin (α -La) and β -Lactoglobulin (β -Lg) according to

Jeanson *et al.* (1999), casein fractions α_s and κ according to Collin *et al.* (1991), immunoglobulins G (IgG) according to Levieux (1974).

3

4 Mapping of β -case antigenicity

 β -casein hydrolysis. Plasmin was used to perform limited cleavage of β-CN, specific for the bonds lys-X and arg-X. Aliquots of purified β-CN (10 g/l) in 75 mM-ammonium acetate were adjusted with NaCl at 0, 2·5 and 5 g/l, and at each pH 5·5, 6·0 and 6·5, then plasmin was added in a weight ratio of 1:100, and the solution was incubated at 37°C for 0·5 and 8 hours. The reaction was stopped by boiling for 10 min at 100°C. Prior to chromatographic separation, hydrolysates were diluted with an equal volume of trifluoroacetic acid (TFA) 1·05 ml/l 8 M-urea buffer and filtered through 0·45 μm.

11 *Reverse phase-high performance liquid chromatography (RP-HPLC) fractionation.* β -CN hydrolysates were 12 fractionated on a C₁₈ Zorbax column (4.6x150 mm, 300 Å, 3.5 µm, Agilent Technologies, Interchim), as described by 13 Beuvier *et al.* (1997). Briefly, 5 nmoles of hydrolysed β -CN were loaded on the column and eluted at 0.8 ml/min by a 14 linear gradient of acetonitrile over ~2-22 % volume, with TFA at 1ml/l as a modifier. Peptides were detected by u.v. 15 absorbance (214 nm), and identified by on-line ESI-MS analysis as described by Léonil *et al.* (1995). Resolved 16 homogenous peaks were collected from further injections of 20 nmoles of hydrolysed β -CN and vacuum-dried at 30°C. 17 Peptides were designated according to the recommended nomenclature (Eigel *et al.*,1984).

18

19 Production of monoclonal antibodies

Monoclonal antibodies were produced essentially as described by Jeanson *et al.* (1999). Immunisation with 20 μ g of purified β -CN was carried out by the footpad route (Mirza *et al.*, 1987). For each experiment, popliteal lymph nods from two immunised mice were pooled. Lymphocytes were fused in a ratio of 5:1 to Sp2/O-Ag-14 myeloma according to two procedures. Essentially, in mass fusion (Köhler & Milstein, 1975) polyethylene glycol (PEG 1000, Merck) (450g/I water) was drop-wise added onto pellets of mixed cells, whereas in filter fusion (Buttin *et al.*, 1978), it was allowed to permeate through monolayers of mixed cells adsorbed on membrane filters. Supernatant media of log phase hybridoma culture were screened in enzyme linked immunosorbent assay (ELISA) for Ig binding to β -CN.

27

28 ELISA

29 The reactivity of the Ig secreted in culture supernatant was assayed in the antigen coated on plate (ACP) mode.

For screening assay, β -CN was coated at 0.5 mg/l 0.1 M-carbonate buffer, pH 9.6 (CB). For cross-reactivity assays, bovine protein fractions (β -CN, BSA, α -La, β -Lg, α_{s} -CN, κ -CN, IgG) (0.5 mg/l CB) or whole caseins (cow, cow-buffalo, human, ewe, goat and mare) (1 mg/l CB) were coated simultaneously. For antigenicity mapping, peptides initially

1 dissolved in 1 ml 0.05 M-phosphate buffer-0.15 M-NaCl, pH 7.2 (PBS), were coated at the optimised dilution that 2 produced similar detection signal as β-CN (0.5 mg/l CB). Aliguots of 100 µl antigens were incubated, for two hours at 3 37°C, in wells of a flat-bottomed 96-well microtitration plate (Nunc Maxisorp, Polylabo 67023 Strasbourg, France). The 4 remaining binding sites were blocked by incubating 250 µl gelatine (Merck) (10 g/l PBS-T buffer, i.e. Tween 20 (Merck) 0.55 g/l PBS) for one hour at 37°C in the wells. Culture supernatants (100 µl) were incubated in the wells, for one hour 5 6 at 37°C, at proper dilution in PBS with a final concentration of 0.55 g/l Tween 20. They were diluted ½ for screening. 7 For the following assays, dilution was optimised whenever possible as to produce a ~2.0 units absorbance increase in 8 screening conditions. Bound Ig were detected by incubating 100 µl of goat anti mouse immunoglobulin alkaline 9 phosphatase conjugate (0.33 ml/l PBS-T) for one hour at 37°C in the wells (Jackson Immunoresearch, Interchim). 10 Wells were rinsed between each incubation step for 15 s with four changes of 250 µl PBS-T. Following the last rinsing, 11 100 µl p-nitrophenyl phosphate (1 g/l 1 M-diethanolamine-HCl-1 mM-MgCl₂-0.1 mM-Zinc acetate) (Sigma) were 12 incubated in the wells. After one hour, the absorbance at 405 nm was read against a blank, and corrected for 13 background signal in the absence of antigen. Threshold for alien casein recognition was set at 4% of bovine signal. All 14 assays, except screening, were performed in duplicate

Isotyping. Light and heavy Ig chains were specifically detected in screening conditions using secondary antibodies
 picked from the Kit 37502 from Pierce (Interchim).

17

18 Biosensor assay using BIAcore 3000

IgG MAbs were assayed for reactivity with β-CN using BIAcore 3000 (Fägerstam *et al.*, 1992). Kinetics
 measurement were performed from single runs. Additive assays were performed without labelling antibodies using on line detection (Daiss & Scalice, 1994).

22 Capture assay. The carboxy-methylated dextran surface (CM5) was conditioned for immobilisation of IgG MAbs as 23 follows: 35 µl of a 0.05 M-N-hydroxy-succinimide-0.2 M-N-ethyl-N'-(3-dimethyl-aminopropyl) solution, then 35 µl of 24 rabbit anti-mouse Fc y (30 mg/l 10 mM-sodium acetate buffer, pH 4.8), and 20 µl of 1 M-ethanolamine hydrochloride, 25 pH 8.5 solution were injected in sequence at 5 ul/min. Undiluted supernatants were injected at 10 ul/min for 3 min. 26 Unoccupied binding sites of the matrix were blocked with IgG1, IgG2 and IgG3 from mixed mice ascitic fluids non-27 reactive with β-CN. Purified β-CN (10 mg/l HBS buffer i.e. polysorbate 20 50 μl/l 10 mM-HEPES-3 mM-EDTA-0.15 M-28 NaCl, pH 7.4) was injected at 10 µl/min for 3 min. The total mass immobilised on the sensor surface was recorded on-29 line throughout all steps as a proportional increase of the surface plasmonic resonance. For comparison of β -CN 30 capture, resonance was read at 3 min after injection of IgG and β -CN. The mass ratios of bound antigen and antibody 31 were corrected to molar ratio using the following Mw values: 146 kDa for IgG1 and G2, 170 kDa for IgG3 and 24 kDa 32 for β -CN.

le

Pair-wise sandwich assay. MAbs were pair-wise assayed within the reactivity groups established for tryptic
 mapping (Table 2), with single MAb IV pooled with group V. Primary MAb was assayed for capture of β-CN, then
 secondary MAb was injected for binding to captured casein in the same conditions.

4

5 Sequence analysis

Primary sequence of β-CN from ewe, goat, human, cow-buffalo, cow, and zebra (*Equus grevyi*) (respectively n°
P11839, P33048, P05814, Q9T5I0, P02666, and Q28401 in SwissProt-trEMBL) were compared with free software
Multalin 5.4.1 (INRA, 31326 Toulouse, France) set on default parameters. Sequence for bovine β-CN was corrected
according to Jimenez-Flores *et al.* (1987). Percentage homology was calculated as aligned residues matching
bovine's. Scales for hydrophilicity (Hopp & Woods), solvent accessibility (Boger *et al.*) and antigenicity (Parker *et al.*)
were calculated from primary structure using free release ANTHEPROT V4.9 (IBCP-CNRS UPR 412, 69367 Lyon,
France) with default parameters and five residues window.

1

RESULTS

2 Production of hybridoma

3 Table 1 features the outcome of three independent fusion experiments carried each with a pair of mice. 21 4 hybridoma secreting anti β -CN Ig were produced. All carried κ light chains, whereas several heavy chain classes (G1, 5 G2a, G3, M and A) were found. One experiment (filter protocol) yielded IgG MAbs of sub-classes exclusively 2a and 3, 6 whereas other experiments yielded a marked majority of sub-classes 1.

- 7

8 displaying the highest levels of cross-reactivity were found directed at β -CN C-terminus (group V). They frequently 9 recognised multiple test antigens, which suggested non-specific binding rather than casual sequence homology.

In ACP-ELISA, low cross-reactivity was detected with β -Lg, lgG, α_s - and κ -CN, but not with α -La and BSA. MAbs

10 Systematic cross-reaction with β-Lg originated apparently from contamination of the test antigen solution with β-CN.

11 Mapping of β -casein antigenicity

12 The MAbs were assigned to five groups (I to V) according to their reactivity with 5 fragments of β-CN in ACP-ELISA 13 (Table 2). HPLC fractions containing peptides shorter than 25 residues were not detected. The epitopes recognised by 14 MAbs from groups I, III and V were unambiguously present in peptides β -CN (f1-25), β -CN (f49-97), and β -CN (f184-209) 15 respectively. MAbs designated as II and IV did not recognised these peptides, but recognised respectively B-CN (f33-16 97) and β-CN (f177-209). Tentative determinants appeared as 33-48 and 177-184 respectively. However, it is possible 17 that the residues 49-52 and 184-187 had been too involved in their epitopes.

18 Reactivity with alien sequence variants

19 In ACP-ELISA, no MAb discriminated the casein from goat and sheep, nor that from cow and cow-buffalo. Each MAb 20 detected casein from at least one non-bovine specie. They were discriminated into eight classes by their specificity 21 towards ovine, equine and human casein (Fig. 1). The number of MAbs cross-reacting with ovine (20), human (7), and 22 with both (6) decreased with decreasing sequence homology (respectively 91%, 53% and 51%). MAb I-1 reacted only 23 faintly with equine and human casein.

24 Epitope delineation

25 The sequences of cow, buffalo, ewe and human β -CN were compared for the substituted residues within the 26 plasmin-cleaved peptide antigens. The sequence for equine β -CN has not been reported, but a partial sequence for 27 Zebra is available, which is likely very close. Four determinants, 4-8, 14-24, 177-183 and 184-190 could be delineated 28 without ambiguity, and probable sites of binding were refined for other MAbs (Fig. 2). Residues critical to the epitope of 29 MAb IV were precisely identified at position 180-181. Similarly to MAb IV, MAb V-1 also recognised horse sequence, 30 and was thus attributed an adjacent epitope, i.e. 184-190. MAb I-2 was assigned to the sequence 4-8, common to 31 ruminants. MAbs I-4,6 and MAbs I-3 and I-7 were attributed to the conserved domain 14-24, common to human and 1 ruminants, and partially shared by equine. MAbs I-3 and I-7 recognised however an epitope not common to equine

2 casein, and thus included at least one unidentified mutated residue.

3 BIAcore epitope mapping

In BIAcore assay, capture of β-CN by any MAb from group I, II or III prevented totally the further binding of all MAbs
from the same group. Within the remaining MAbs IV and V, assayed pair-wise, non-interfering binding to β-CN was
observed for MAb IV, MAb V-2 and a set composed of MAbs V-1, V-4 and V-5, which mutually interfered (Table 3).

7 Capture of β -CN by immobilised IgG monoclonal antibodies

The MAbs assayed in biosensor were rated for their ability to capture β -CN in solution (Table 4). MAb I-1 achieved 9 91% of the optimal monovalent binding. In contrast, MAbs III-1 and III-3 did not bind β -CN, whereas they did in the ACP-10 ELISA screening. These MAbs were apparently directed against β -CN epitopes specifically exposed upon conformation 11 changes subsequent to coating. The contrasted capture ability discriminated markedly sub-groups of MAbs within the 12 groups I to V.

- 13
- 14

DISCUSSION

15 Previous constructions of hybridoma against β-CN relied on intra-peritoneal immunisation, and each highlighted 16 apparently a single dominant epitope, though clear data often lack. MAb 1C3 recognises β -CN (f193-202) (Nagaune et 17 al., 1988), whereas three MAbs CAS-OT# recognise redundantly β-CN (f105-193) (Oudshorn et al., 1994). In addition, 18 MAb AH4 recognises an unidentified bovine-specific epitope (Anguita et al., 1995), and five MAbs named 58-# 19 recognise an identical unknown epitope (Kuzmanoff et al., 1991). According to Mirza et al. (1987), footpad route 20 increases the positive clone rate (100% vs 8%), the specificity spectrum, and particularly emphasises conserved 21 epitopes. Using this protocol, we achieved only a low positive clone rate, possibly owing to low intrinsic 22 immunogenicity of β-CN, but we elicited an unprecedented panel of specificity. A single IgA was obtained. IgA may be 23 obtained in mice specifically upon oral immunisation route and/or collection of gut associated lymphocytes, and 24 seldom result from other protocols. Oral immunisation probably occurred, either previously to the primary injection, via 25 the diet, or simultaneously, by casual ingestion of the immunogenic mixture. Dietary β-CN digests enhances class-26 switch towards IgA in mice, possibly via cytokine(s) stimulation (Otani et al., 2000).

All MAbs recognised β -CN fragments, which indicated they recognised continuous epitopes. Typically 10% of MAbs raised to globular proteins recognise peptides (Pellequer *et al.*, 1991), which was observed for bovine β -Lg (~36 kDa) (Venien *et al.*, 1997). The striking difference for β -CN (100% *vs* 10%) accounts probably for the high proportion of aperiodic structure (70% *vs* 45%), and the lack of stable tertiary structure in β -CN. Owing to the generalised crossreactivity with peptides, all the continuous epitopes could be circumscribed.

Seven antigenic areas were defined in ELISA assays, including three individual epitopes, tentatively delineated as 4-8, 178-183 and 185-190, and four larger determinants, defined as 14-24, 33-48, 49-94 and 184-209. The

determinant 14-24 was also found to contain two epitopes at least, one common to all assayed species, and the other differing in equine sequence. The eight corresponding groups of MAbs were further split into 13 epitopic classes according to (i) the additivity assays and (ii) the contrasted rates of β -CN capture, in BIAcore assays. The determinants 14-24 and 49-94 were found to contain at least respectively 4 and 2 distinct but probably overlapping epitopes. The determinant 184-209 contained three distinct epitopes probably overlapping and one non-overlapping one (V-2).

6 The fusion process immortalises only a restricted population of the collected lymphocytes that constitute the individual immune response, typically 10⁵-10⁻⁶. Moreover, Otani et al. (1987) have demonstrated that individual 7 8 responses of rabbits immunised with β -CN varied dramatically. The specificity panel we achieved sums the responses 9 of six mice and thus features a cross section of β -CN immunogenic determinants. Otani et al. have investigated β -CN 10 immunogenicity by the analysis of individual rabbit polyclonal antibodies raised either to bovine β -CN (1984; 1987) ot to 11 β -CN digest (1990). They have identified discrete antigenic determinants at the sequences 1-25, 26-60, 61-93, 94-102, 12 103-109, 110-144 and 157-185. In addition, they have found the peptide β -CN (f186-209) immunogenic. Our 13 experiments failed to detect determinants in the sequence 94-144. This discrepancy may/might be originated in the 14 different host we used (mouse vs rabbit). The comparison of the sequence homology for the three species at 94-144 15 does/does not support this hypothesis. Alternatively, a bias should be considered in our collection. The screening step 16 relied exclusively on ACP-ELISA with coated β -CN, and we may have failed to detect some antibodies specific for a 17 native form. As much as 7% of specific MAbs non-reactive in ACP ELISA with β-Lg have been reported by Venien et 18 al. (1997). β -CN is dramatically less organised than β -Lg, as was illustrated by the lower rate of ELISA specific MAbs we 19 obtained (10% vs 25%). It is thus likely that a lesser rate, if any, of such MAbs would be elicited against B-CN.

The determinant 14-24, recognised by MAbs I-3,7, coincides with the phosphorylated cluster. This sequence has been found a potent determinant common to α_{s} - and β - casein in immunised mice, and in allergic humans, and its reactivity has been shown to depend upon proper state of phosphorylation (Bernard *et al.*, 2000). No cross-reactivity with α_{s} -CN was observed for MAbs I-3,7. The determinant 3-12 has not been found dependant upon phosphorylation (Pizzano *et al.*, 2000). It would be interesting to investigate the reactivity of MAbs I-1,7 with dephosporylated species, since phosphatases are acting in cheese (Ferranti *et al.*, 1997a).

 β -CN terminal sequences 1-25 and 184-209 represented each ~12% of the sequence in length. They carried each 5 out of the 13 distinct determinants (~38%) we identified, and particularly carried all the epitopes conserved in two or more non-bovine species. Stewart *et al.* (1987) reported that the N-terminal domain 1-50 was highly conserved in most species (including cow, human, rat, ewe), in particular at the phosphorylated cluster, but that the C-terminus was highly variable. It is now a widespread affirmation that protein termini are highly immunogenic. This opinion arises partly from the confusion of immunogenicity for the reactivity towards antipeptide antibodies. This characteristic implies a structural similarity with the peptide, and has thus been correlated with flexibility and/or surface exposure in globular

proteins (Pellequer *et al.*, 1991). These parameters are favourable in the N-terminus, and all three immunogenicity scales based on hydrophilicity, solvent accessibility, and statistical propensity predicted the determinant 14-24. On the contrary, these parameters are not favourable in the C-terminus, and all three scales failed to predict the epitope 193-209. The scales were generally not efficient for predicting β -CN determinants: the determinants 14-24 and 33-49 were predicted by all three scales, the epitope 177-184 only by solvent accessibility and statistical propensity, and the determinant 4-8 only by hydrophilicity scale. Conversely, all three predicted a consensual epitope in the area 100-110, that we did not observed, whereas Otani *et al.* did (1987).

8 The produced panel of specificity enables the specific detection of fragments of β -CN released by plasmin, i.e. β -CN 9 (f1-28), β -CN (f29-105/-107), and β -CN (f106/108-209). The immunodetection of these peptides should help in objective 10 assessment of plasmin contribution to proteolysis in cheese. In particular, phosphopeptides may be potential indexes 11 for cheese ageing certification. Pizzano et al. (2000) reported that the determinant 3-12 was rapidly degraded in 12 cheese, and thus useless as a marker. In contrast, the peptide β -CN (f16-22) accumulates in hard-type cheeses, and is 13 therefore a good candidate as an ageing index (Ferranti et al., 1997a). Five MAbs from group I are directed at four 14 distinct epitopes within 14-24, and may enable its immunometrical detection in the pH 4-6 soluble fraction of cheeses. 15 In other respects, considerable efforts have been devoted to the production of bovine specific reagents suitable to 16 detect minute adulteration of ovine or caprine milk in cheeses made thereof (Dupont ????). Antibodies directed at 17 caseins include only one MAb, raised against an unidentified epitope of β-CN (Anguita et al., 1995). MAb IV might 18 prove useful in this issue, since its epitope is a good marker for γ -caseins that accumulate upon degradation of β -CN in 19 cheese, and might therefore be detectable in extensively ripened cheeses (Pizzano et al., 1999).

- 20
- 21

We thank D. MOLLÉ (INRA Rennes) for performing ESI-MS, and members of SRTAL staff for contribution, S. POCHET in HPLC analysis, F. FAURIE in animals care, H. Lamprell for linguistic revision,0and Y. NOEL for commenting the manuscript.

25 D. SENOCQ thanks J. LEONIL, S. POCHET and J. GUZZO (ENSBANA) for scientific advice within his thesis board.

We acknowledge the support of the French Ministry for Education, Research, and Technology for this work (FRT 99 Y0098-13) and D. SENOCQ' s undergraduate PhD grant.

1 REFERENCES 2 ADDEO, F., GARBO, G., INTORCIA, N., PELLEGRINO, L., RESMINI, P. & CHIANESE, L. 1995 Gel electrophoresis and immunoblotting for the detection of 3 casein proteolysis in cheese. Journal of Dairy Research 62 297-309 4 ANDREWS, A. T., TAYLOR, M. D. & OWEN, A. J. 1985 Rapid analysis of bovine milk proteins by fast protein liquid chromatography. Journal of 5 Chromatography 348 177-185 6 ANDREWS, A., T. & ALICHANIDIS, E. 1983 Proteolysis of caseins and the proteose-peptone fraction of bovine milk. J. Dairy Res. 50 275-290 ANDREWS, A., T. 1983 Proteinases in normal bovine milk and their action on caseins. Journal of Dairy Research 50 45-55 7 ANGUITA, G., MARTIN, R., GARCIA, T., MORALES, P., HAZA, A. I., GONZALES I., SANZ, B. & HERNANDEZ, P. 1995 Indirect ELISA for detection of cow's milk 8 in ewe's and goat's milks using a monoclonal antibody against bovine β-casein. Journal of Dairy Research 62 655-659 9 BERNARD, H., MEISEL, H., CREMINON, C. & WAL, J.M. 2000. Post-translational phosphorylation affects the IgE binding capacity of caseins. FEBS 10 Letters 467 239-244 11 BEUVIER, E., BERTHAUD, K., CEGARRA, S., DASEN, A., POCHET, S., BUCHIN, S. & DUBOZ, G. 1997 Ripening and quality of swiss-type cheese made from 12 raw, pasteurised or microfiltered milk. International Dairy Journal 7 311-323 13 BUTTIN, G., LEGUERN, G., PHALENTE L., LIN, E. C. C., MEDRANO, L. & CAZENAVE, P. A.01978 Production of hybrids lines secreting antiidiotypic 14 antibodies by cell fusion on membrane filters. Current Topics in Microbiology and Immunology 81 27-36 15 COLLIN, J. C., KOKELAAR, A., ROLET-REPECAUD, O. & DELACROIX-BUCHET, A. 1991 [Determination of cow's milk caseins by electrophoresis and fast 16 protein liquid ion-exchange chromatography (FPLC®): comparative results] Lait 71 339-350 17 DAISS, J. L. & SCALICE, E. R. 1994 Epitope mapping on BIAcore: theoretical and practical considerations. METHODS : a companion to Methods in 18 Enzymology 6 143-156 19 DEUTSCH, D. G. & MERTZ, E. T. 1970 Plasminogen: purification from human plasma by affinity chromatography. Science 170 1095-1096 20 EIGEL, W. N., BUTLER, J. E., ERNSTROM, C. A.; FARRELL, H. M., HARWALKAR, V. R., JENNESS, R. & WHITNEY, R. MCL. 1984 Nomenclature for proteins of 21 cow's milk : fifth revision. Journal of Dairy Science 67 1599 22 FÄGERSTAM, L. G., FROSTELL-KARLSSON, Å., KARLSSON, R., PERSSON, B. & RÖNNBERG, I. 1992 Biospecific interaction analysis using surface plasmon 23 resonance detection aplied to kinetic, binding site and concentration analysis. Journal of Chromatography 597 397-410 24 FERRANTI, P., BARONE, F., CHIANESE, L., ADDEO, F., SCALONI, A., PELLEGRINO, L. & RESMINI P. 1997a Phosphopeptides from Grana Padano cheese: 25 nature, origin and changes during ripening. Journal of Dairy Research 64 601-615 26 FERRANTI, P., ITOLLI, E., BARONE, F., MALORNI, A., GARRO, G., LAEZZA, P., CHIANESE, L., MIGLIACCO, F., STINGO, V. & ADDEO, F. 1997 b Combined high 27 resolution chromatographic techniques (FPLC and HPLC) and mass spectrometry-based identification of peptides and proteins in Grana 28 Padano cheese. Lait 77 683-697 29 JEANSON, S., DUPONT, D., GRATTARD, N. & ROLET-REPECAUD, O. 1999 Characterisation of the heat treatment undergone by milk using two inhibition 30 ELISAs for quantification of native and heat denatured α-Lactalbumin. Journal of Agricultural and Food Chemistry 47 2249-2254 JIMENEZ-FLORES, R., KANG, Y. C. & RICHARDSON, T. 1987 Cloning and sequence analysis of bovine β-casein cDNA. Biochemical and Biophysical Research Communications 142 617-621 31 KOBAYASHI, T., AMETANI, A., YAMAUCHI, K. & KAMINOGAWA, S. 1991 Differences in defining residues relevant to antibody binding by ELISA and 32 proteolysis protection at the level of peptide antigenic determinant. Biochimica et Biophysica Acta 1077 11-18 33 KÖHLER, G. & MILSTEIN, C. 1975 Continuous cultures of fused cells secreting antibody of predefined specificity. Nature 256 495-497

34 KUMOSINSKI, T. F., BROWN, E. M. & FARELL, H. M. Jr. 1993 Three-dimensional molecular modeling of bovine caseins: an energy-minimized β-casein

35 structure. *Journal of Dairy Science* **76** 931-945

36 KUZMANOFF, M. K., ANDRESEN, J. & BEATIE, C. W. 1991 Isolation and characterisation of monoclonal antibodies monospecific for bovine α-casein and

37 β -CN. Journal of Dairy Science **74** 803-810

- 1 LEAVER, J. & LAW, A. J. R. 1992 Preparative scale purification of bovine caseins on a cation-exchange resin. Journal of Dairy Research 59 557-561
- LÉONIL, J., MOLLÉ, D., GAUCHERON, F., ARPINO, P., GUÉNOT, P. & MAUBOIS, J. L. 1995 Analysis of major bovine milk proteins by on-line high
 performance liquid chromatography and electrospray ionization-mass spectrometry. *Lait* **75** 193-210
- LEVIEUX, D. 1974 [Bovine Immunoglobulins and Brucellosis. Purification of immunoglobulins and preparation of their respective specific antisera]
 Annales de Recherches Vétérinaires 5 329-342
- MIRZA, I.H., WILKIN, T.J., CANTARINI, M. & MOORE, K. 1987 A comparison of spleen and lymph node cells as fusion partners for the raising of
 monoclonal antibodies after different routes of immunisation. *Journal of Immunological Methods* 105 235-243
- 8 NAGAUNE, S. I., KAMINOGAWA, S., ENOMOTO, A., KOBAYASHI, T., KURISAKI, J. I. & YAMAUCHI, K. 1988 Preparation of anti bovine β-casein monoclonal
 9 antibody and analysis of the interaction between the antibody and β-casein fragment. *Agricultural and Biological Chemistry* **52** 2577-2581
- OTANI, H., DONG, X. Y. & HOSONO, A. 1990 Antigen specificity of antibodies raised in rabbits injected with a chymotryptic casein digest with molecular
 weight less than 1000. Japanese Journal of Dairy and Food Science **39** A31-A37
- OTANI, H., HIGASHIYAMA, S. & TOKITA, F. 1984 Studies on the antigenic structure of bovine Casein. IV. Antigenic activities of peptides obtained by
 cyanogen bromide cleavage of 94-209 region. Milchwissenschaft 39 469-472
- 14 Otani, H., Kihara, Y. & Park M. 2000 The immunoenhancing property of a dietary casein phosphopeptide preparation in mice. Food and Agricultural
- 15 Immunology 12 165-173
- Otani, H., Mine, Y. & Hosono, A. 1987 Studies on the antigenic structure of bovine
 Casein. V. Antigenic activities of some peptides derived from
 residues 1-93. Milchwissenschaft 42 505-508
- Pellequer, J. L., Westhof, E. & Van Regenmortel, M. H. V. 1991 Predicting location of continuous eitopes in proteins from their primary structures.
 Methods in enzymology 203 176-201.
- Pizzano, R., Nicolai, M. A. & Addeo, F. 1999 Antipeptide antibodies in milk and cheese analysis. Recent Results and Developments in Agricultural &
 Food Chemistry 3 403-412
- Pizzano, R., Nicolai, M. A., Padovano, P., Ferranti, P., Barone, F. & Addeo, F. 2000 Immunometrical evaluation of bovine
 –casein and its 1-28
 phosphopeptide in cheese during ripening. Journal of Agricultural and Food Chemistry ASAP Article 10.1021, Web realease: September 23.
- 26 STEWART, A. F., BONSING, J., BEATTIE, C. W., SHAH, F., WILLIS, I. M., MACKINLAY, A. G. 1987 Complete nucleotide sequences of bovine α_{S2}- and β-
- 27 casein cDNAs : comparisons with related sequences in others species. Molecular Biology and Evolution 4 231-241
- SWAISGOOD, H. E. 1992 Chemistry of the caseins. In *Advanced Dairy Chemistry-1: Proteins*, 63-110 (Ed. P.F. Fox). London, UK & New York, USA:
 Elsevier Applied Science
- VENIEN, A., LEVIEUX, D., ASTIER, C., BRIAND, L., CHOBERT, J. M. & HAERTLE, T. 1997 Production and epitopic characterization of monoclonal
 antibodies against bovine β-Lactoglobulin. *Journal of Dairy Science* 80 1977-1987
- 32 WEI, T. M. & WHITNEY, R. M. 1984 Batch fractionation of bovine caseins with DEA-cellulose. Journal of Dairy Science 68 1630-1636

1	FIGURE LEGENDS
2	
3	Fig. 1. Distribution of 21 anti bovine β -casein MAbs according to their reactivity with ovine, human and equine whole
4	casein in ACP-ELISA. Group size when >1 is given in brackets.
5	
6	Fig. 2. Tentative epitopes of specific anti bovine β -casein MAbs from 7 reactivity classes as determined from their
7	reactivity with tryptic fragments, buffalo, ovine and human sequence variants in ACP-ELISA.
8	
9	
10	
11	

Table 1. Outcome of three independent fusion experiments of activated murine lymphocytes

	fusion	growing	positive	isotype					
		/lymphocytes	/hybrids	G1	G2a	G3	М	А	κ
1	mass	1·1.10 ⁻⁶	6.0%	4				1	5
2	filter	1·9 .10 ⁻⁶	6.6%		5	1	1		7
3	filter	3·3 .10 ⁻⁶	5.9%	9					9

1

molecular mass (Da)			peptide	rea	reactive MAb group §				
measure	d†	theoretical‡		Ι	II	III	IV	V	
3122·73±	0.1	3123-0	β-CN -4P(f1-25)	+					7
7344.96.±	1.0	7346.2	β-CN -1P(f33-97)		+	+			8
5302·17±	0.6	5301.2	β-CN (f49-97)			+			5
3722·6 ±	0.1	3722.5	β-CN (f177-209)				+		6
2910-9 ±	0.9	2910.5	β-cn (f184-209)				+	+	5

Table 2. Mapping of bovine β -case in antigenicity in ACP-ELISA towards 21 specific MAbs

2 + average molecular mass ± SD calculated for multiple multiprotonated ions in RP-HPLC analysis coupled to ESI-MS

3 detection of hydrolysed β-CN

4 ‡ average molecular mass calculated according to the sequence of probable match (Jimenez-Flores *et al.*, 1987)

5 § classification of reactive MAbs in ACP-ELISA

- 1 **Table 3.** Interfering (greyed) or non-interfering pair-wise binding to bovine β -casein of specific MAbs with close
- 2 epitopes. Binding is expressed as a molar % of maximal capture of the secondary MAb.



3 *nd* : non determined

1

Table 4. Capture of β -casein (molar % of Ig) by immobilised specific IgG MAbs in biosensor assay

MAb % Ig		MAb % lg		MAb % lg		MAb % lg	
I-1	91	II-2	17	111-4	31	V-1	78
I-7	39	II-3	11	III-2	21	V-5	41
I-2	35			III-5	21	V-2	31
I-4	28			III-1	0	V-4	6
I-5	23			III-3	0		
I-6	19						
I-3	9			IV	12		
MAb	% lg	MAb	% lg	MAb	% lg		
I-1	91	II-2	17	IV	12		

2

MAb	% lg	MAb	% lg	MAb % lg		
I-1	91	II-2	17	IV	12	
I-7	39	II-3	11	V-1	78	
I-2	35	111-4	31	V-5	41	
I-4	28	III-2	21	V-2	31	
I-5	23	III-5	21	V-4	6	
I-6	19	III-1	0			
I-3	9	III-3	0			

3

4

FIGURES