



HAL
open science

Intercropping on French farms: Reducing pesticide and N fertiliser use while maintaining gross margins

Elodie Yan, Nicolas Munier-Jolain, Philippe Martin, Marco Carozzi

► To cite this version:

Elodie Yan, Nicolas Munier-Jolain, Philippe Martin, Marco Carozzi. Intercropping on French farms: Reducing pesticide and N fertiliser use while maintaining gross margins. *European Journal of Agronomy*, 2024, 152, pp.art. 127036. 10.1016/j.eja.2023.127036 . hal-04301593

HAL Id: hal-04301593

<https://hal.inrae.fr/hal-04301593>

Submitted on 23 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Intercropping on French farms: reducing pesticide and N 2 fertiliser use while maintaining gross margins

3 Elodie Yan^{a,*}, Nicolas Munier-Jolain^b, Philippe Martin^a, Marco Carozzi^a

4 ^a Université Paris-Saclay, INRAE, AgroParisTech, UMR SADAPT, F-91120, Palaiseau, France

5 ^b Université Bourgogne-Franche-Comté, INRAE, Institut Agro Dijon, UMR Agroécologie, F-
6 21000, Dijon, France

7 * Corresponding author. *E-mail address*: elodie.yan@inrae.fr

8 **Abstract**

9 Experimental studies to date have demonstrated the agronomic and environmental benefits of
10 intercropping, making it a key diversification method for reducing chemical inputs in
11 agriculture. However, intercropping is still a niche practice in European cropping systems,
12 particularly for arable crops. Few studies have focused on farmers' perspectives or used farm
13 data to assess the on-farm performances of intercropping. Here, we present an analysis of data
14 collected from farms of the DEPHY network, a demonstration farm network that aims to show
15 that pesticide reduction is possible through changes in farming practices. We focused our study
16 on four main species in France: winter wheat (*Triticum aestivum* L.), winter barley (*Hordeum*
17 *vulgare* L.), pea (*Pisum sativum* L.) and rapeseed (*Brassica napus* L.). We carried out paired
18 comparison tests between sole crops and intercrops for each species to compare the use of
19 pesticides (herbicides, fungicides, insecticides and all pesticides combined), mineral nitrogen
20 fertiliser and the gross margin between the sole crop and the intercrop. We showed that pesticide
21 use was reduced on average by 50% in the case of wheat- and barley-based intercrops compared
22 with sole wheat and barley crops, respectively. Pesticide use for peas was reduced by 83% on
23 average. Nitrogen fertiliser use was also reduced by up to 50% for wheat. On the other hand, in
24 the case of rapeseed, which is mainly intercropped with unharvested companion plants, we
25 found no significant differences in pesticide or fertiliser use between the sole crops and
26 intercrops. This suggests that farmers must choose companion plants carefully depending on
27 their objectives and desired services. Finally, our results show that crop mixtures do not
28 negatively affect gross margins and can even improve them; intercropped peas in particular
29 showed an average 16-fold increase in gross margins.

30 **Keywords:** crop mixtures; crop diversification; arable crops; cropping systems; agricultural
31 practices; Treatment Frequency Index

32 **1 Introduction**

33 Intensive and specialised agriculture primarily uses chemical inputs that harm human health
34 (Tudi et al., 2021; WHO, 1990) and the environment, e.g., by eroding biodiversity and
35 degrading water and soil quality (Matson et al., 1997; Tilman et al., 2002). Meeting the global
36 demand for food and feed while reducing the negative impact of agriculture on the environment
37 is a major present-day challenge. Diversifying agricultural systems is a suitable strategy to
38 achieve these objectives as it enables the restoration of ecosystem services (Beillouin et al.,
39 2021; Lin, 2011; Raseduzzaman & Jensen, 2017) and reduces dependence on chemical inputs,
40 such as fertilisers and pesticides.

41 Intercropping, which consists of growing two or more crop species simultaneously on the same
42 field (Willey, 1979), is a promising diversification practice. Experimental studies have shown
43 that intercropping can improve weed control in arable farming by improving soil cover and
44 competition (Corre-Hellou et al., 2011; Hauggaard-Nielsen & Jensen, 2005). The
45 morphological and physiological differences between the cultivated species can create barriers
46 that limit the spread of diseases and pests (Boudreau, 2013; Finckh et al., 2000). Intercropping
47 further improves crop quality, particularly in the case of cereal–legume associations, by
48 increasing the amount of mineral nitrogen available in the soil and, therefore, the protein content
49 in cereal seeds (Bedoussac et al., 2015; Li et al., 2023). In some cases, intercropping can
50 stabilise or even increase yields (Lithourgidis et al., 2011; Malézieux et al., 2009). According
51 to experimental results, intercropping is a relevant lever for reducing the use of chemical inputs
52 while maintaining a certain quality and quantity of production (Bedoussac et al., 2015;
53 Lithourgidis et al., 2011). Intercropping could also substantially contribute to achieving the
54 goals of Ecophyto, the French National plan to reduce pesticide use by 50% by 2025. This plan
55 was developed under the framework of the European Directive 2009/128/EC and encourages
56 the demonstration of good farming practices to reduce dependence on pesticides.

57 Despite all its advantages, intercropping is still a niche practice in France as well as in Europe
58 (Mamine & Farès, 2020; Timaeus et al., 2022), particularly in non-organic farming (Verret et
59 al., 2020). According to Land Parcel Identification System data (IGN, 2023), intercropping
60 accounted for only 3% of French arable land in 2020. Several factors seem to limit the spread
61 of intercropping, including technical difficulties linked to sowing, harvesting and sorting crops
62 on farms (Mamine & Farès, 2020; Timaeus et al., 2022; Verret et al., 2020); economic issues
63 due to limited outlets when cooperatives do not collect grain mixtures because they are not

64 equipped to sort grains (Bedoussac et al., 2015; Mamine & Farès, 2020); and farmers' psycho-
65 social reluctance to adopt such an innovation (Bonke et al., 2021; Bonke & Musshoff, 2020).

66 Though addressing these obstacles to the broader adoption of intercropping is necessary, it is
67 also crucial to understand how farmers manage crop mixtures in practice and whether on-farm
68 intercropping actually delivers the advantages demonstrated by field or plot experiments.
69 Recent research based on farmer interviews, such as Enjalbert et al. (2019) and Verret et al.
70 (2020) in France and Timaeus et al. (2022) in Germany, has moved towards understanding
71 farmers' satisfaction with intercropping. These qualitative studies aimed to show the diversity
72 of intercrops effectively grown and highlight the obstacles to their adoption but included only
73 small samples of farmers.

74 To our knowledge, apart from field trials, no study has yet verified the performance of crop
75 mixtures on farms, looking at both inputs (pesticides and fertilisers) and production factors.
76 This paper presents a quantitative study of the effects of intercropping on pesticide use, nitrogen
77 fertilisation and gross margins of major crops in France based on data collected from farms
78 across the country. Our analysis aimed to show the effects of intercropping under actual farming
79 conditions and highlight this practice's agronomic and environmental benefits.

80 **2 Material and methods**

81 2.1 Data from a demonstration farm network: The DEPHY network

82 The DEPHY network is a French nationwide demonstration farm network created in 2010 as
83 part of the Ecophyto National Plan. The DEPHY network aims to demonstrate that reducing
84 the use of pesticides is possible through changes in agricultural practices. The network started
85 with 178 farmers at its inception, and by 2016, more than 3000 farmers had joined and
86 committed to reducing their pesticide use. The network covers seven major French production
87 sectors (i.e., arable field crop, crop–livestock mixed farming, vegetables [both outdoor and
88 protected], ornamental crops, tropical crops, viticulture and arboriculture). Participating
89 farmers are gathered into local groups of 10 to 15 and regularly share their experiences with
90 each other. One cropping system per farm is monitored and described year after year.
91 Information on crops grown and management details, such as fertilisation, pesticide use, tillage
92 and economic performance, is collected and entered into a database (the AGROSYST
93 Information System). The database used for this study included 27,711 farm–years between
94 2010 and 2023.

95 2.2 Crop species

96 We focused our study on four of the most cultivated crops in France that may also be grown as
97 intercrops: winter wheat (*Triticum aestivum* L.), winter barley (*Hordeum vulgare* L.), peas
98 (*Pisum sativum* L.) and rapeseed (*Brassica napus* L.). According to the French annual
99 agricultural statistics (Agreste, 2020), winter wheat (referred to as “wheat” hereafter) was the
100 most cultivated crop in France in 2020, covering approximately 34% of the country’s total
101 arable crop area. Winter barley (referred to as “barley” hereafter) was the third most cultivated
102 (behind maize) with approximately 10% of the arable crop area. Rapeseed was the most
103 cultivated oilseed crop and peas were the most cultivated protein crop, with 9% and 2% of the
104 arable crop area, respectively.

105 Because the DEPHY network aims at reducing pesticide use, farmers belonging to the network
106 are likely to implement multiple practices to this end. Therefore, reductions in pesticide use
107 cannot be easily attributed to a single practice – in our case, intercropping. To ensure that any
108 differences in pesticide use, nitrogen fertiliser use, and gross margin actually resulted from
109 intercropping, we compared the intercrops involving wheat, barley, peas or rapeseed with their
110 corresponding sole crop (i.e., sole wheat, barley, peas or rapeseed) grown in cropping systems
111 in which we noted similar characteristics.

112 2.3 Selection and characterisation of the cropping systems


113 We focused our analysis on two production sectors out of the seven available in the DEPHY
114 database: the arable field crop and crop–livestock mixed farming, as these production sectors
115 included wheat, barley, peas and rapeseed. Inspired by the work of Lechenet et al. (2016), we
116 selected cropping systems with wheat, barley, peas and rapeseed either intercropped or as sole
117 crops and, from the database, computed 36 agronomic variables that may affect the use of
118 pesticides. These variables concern diversity in the crop sequence (e.g., proportion and number
119 of different crops), length of the crop sequence, sowing periods, tillage and irrigation and are
120 detailed in Table S1.

121 Organic farms were excluded due to the restricted use of chemical pesticides, which would have
122 introduced bias in our analysis. Moreover, we excluded farms with incomplete information on
123 crop succession and sowing operations. Finally, our dataset was narrowed down to n=2252
124 cropping system–years (1067 for arable field crops, 1185 for crop–livestock mixed farming)
125 fully described by the 36 selected agronomic variables.

126 2.4 Paired comparison and statistical analysis

127 The intercrop–sole crop pairs were selected using the Euclidean distance between cropping
128 systems. First, we calculated a matrix of Euclidean distances (function “dist”, R package
129 “stats”, R-software version 4.1.3) between cropping systems based on the 36 agronomic
130 variables (Table S1). In the DEPHY database, pesticide use for wheat, barley and rapeseed is
131 generally lower in crop–livestock mixed farming than in arable field crop farming (Figure S1).
132 Therefore, we paired each cropping system that included an intercrop with the closest cropping
133 system that included the corresponding sole crop based on the minimal distance within the same
134 production sector (arable field crop or crop–livestock mixed farming). For example, each
135 wheat-based intercrop grown in a crop–livestock mixed farm was compared with the sole wheat
136 crop from the least-distant cropping system in another crop–livestock mixed farm.

137 Some intercrops had several corresponding sole crops with the same minimal distance.
138 Considering all the possible couples in our analysis would have biased the results as it would
139 have multiplied the observations from the same intercrop, thus over-estimating the effects of
140 the intercrop. To avoid such biases, we chose to perform the analysis on each possible
141 combination and evaluate the deviation between these combinations. For example, if an
142 intercrop had two possible corresponding sole crops (i.e., with the same minimal distance to the
143 intercrop), we created a dataset for each and performed the analysis (i.e., the paired comparisons
144 for pesticide use, nitrogen fertilisation and gross margin, detailed hereafter) on each dataset.
145 Then, we calculated the coefficient of variation between the results of each of the two analyses
146 to test the variability of the datasets. Figure 1 illustrates this methodology using an example
147 with three intercrops: intercrop A is paired with sole crop A1, and intercrops B and C both have
148 two possible sole crops to pair with (sole crops B1 and B2, and sole crops C1 and C2,
149 respectively). In this example, there are four possible combinations, leading to four datasets on
150 which the analyses would be performed.



151

152 Figure 1: Methodology used to test the variability of the datasets when one intercrop (IC)
 153 could be paired with several equidistant sole crops (SCs)

154 We obtained two possible combinations for wheat (one intercrop had two equidistant sole
 155 crops), one for barley (each intercrop only had one sole crop at the minimal distance), 512 for
 156 peas (five intercrops had two to four equidistant sole crops) and 225 for rapeseed (four
 157 intercrops had three to five equidistant sole crops). In each possible combination, we had 16
 158 pairs for wheat, 12 for barley, 30 for peas and 31 for rapeseed. Table 1 reports the number of
 159 intercrops and sole crops for each farming sector.

160 Table 1: Number of farm–years with sole crops and intercrops involving wheat, barley, peas or
 161 rapeseed, and number of intercrop–sole crop pairs used to perform the analysis.

	Wheat	Barley	Peas	Rapeseed
Number of cropping systems with sole crops	1813	936	122	860
- field crop production system	881	465	90	491

- crop–livestock production system	932	471	32	369
Number of cropping systems with intercrops	16	12	30	31
- field crop production system	4	4	5	21
- crop–livestock production system	12	8	25	10

162

163 Using those pairs, we compared pesticide use, fertiliser inputs and gross margins between
 164 intercrops and sole crops. We used the Treatment Frequency Index (TFI) as a metric for
 165 pesticide use. This index is used to monitor the frequency and intensity of pesticides applied at
 166 different spatial scales (e.g., crop, cropping system, groups of cropping systems). In our study,
 167 we considered the TFI at the crop scale following Pingault et al. (2009):

168

$$TFI = \sum_T \frac{Applied\ Dose_T}{Reference\ Dose_T} * \frac{Treated\ area_T}{Total\ area_T}$$

169 For each treatment T on the considered crop, the applied dose per hectare is normalised to the
 170 reference dose per hectare. The crop-scale TFI is then calculated as a weighted sum of all
 171 normalised doses for the considered crop, with weights being the ratio of the treated area over
 172 the total area of the considered crop. The reference dose we used to compute TFI was the
 173 registered dose for the plant protection product on the considered crop for the targeted
 174 pest/weed/disease, based on the marketing authorisation database (E-Phy, ANSES, 2023). Our
 175 study used TFI computed for four distinct categories:

- 176 • All types of pesticides (e.g., herbicides, fungicides, insecticides, seed coating,
 177 fumigants, molluscicides), excluding biopesticides and “low-risk” pesticides according
 178 to French legislation
- 179 • All herbicides, excluding biopesticides and “low-risk” pesticides according to French
 180 legislation
- 181 • All fungicides, excluding seed coating, biopesticides and “low-risk” pesticides
 182 according to French legislation
- 183 • All insecticides, excluding seed coating, biopesticides and “low-risk” pesticides
 184 according to French legislation

185 We used the total amount of mineral nitrogen applied to the crop during the whole growing
 186 period (in kg N ha⁻¹) as a metric for fertilisation inputs.

187 We used the gross margin (€ ha⁻¹) as a metric for crop profitability, defined as:


188

$$\text{Gross margin} = \text{Gross product} - \text{Operating costs}$$

189 where the gross product is the product of yield and selling price (provided by the database user)
190 and operating costs include, e.g., seed, seed coating, fertilisation, pesticide and irrigation costs.

191 In the database, crops that were self-used on the farm (e.g., for animal feed) were assigned a
192 gross product value corresponding to the value they would have had if they had been sold. In
193 order to compare gross margins between farms that may have different levels of self-use, the
194 farm gross margin was calculated taking into account both the self-used and the sold parts of
195 the production. We chose to analyse the gross margin rather than the gross product because we
196 assumed that intercrops can affect the gross product (by affecting yields) as well as the operating
197 costs, and the gross margin better reflects the balance between the effects of intercropping on
198 the gross product and those on the operating costs.

199 Finally, we compared the TFI, N fertilisation and gross margin between the sole crops and
200 intercrops using paired t-tests. This test compares two groups of observations with the
201 specification that an observation from one group is paired with another observation from the
202 other group. Figure 2 recapitulates the study methods, from cropping system selection to data
203 analysis.



204

205 Figure 2: Flowchart of the study methods, from data selection and pairing to statistical analysis.

206 TFI, treatment frequency index; N, nitrogen.

207 3 Results

208 In this section, we first present the results of the analyses conducted over all possible
 209 combinations between intercrops and sole crops for the studied crops. Then, we present the
 210 results of the paired comparisons of TFI, N fertilisation and gross margin for one combination.

211 In the following, an increase (or decrease) in an indicator refers to an increase (or decrease) in
 212 that indicator for the intercrop compared with the corresponding sole crop.

213 3.1 Variability of intercrop–sole crop combinations

214 To assess the variability of our results, we computed the coefficient of variation of the mean
 215 intercrop – sole crop difference over all the possible combinations (Figure 1) for each species
 216 and indicator (TFI, N fertilisation, gross margin). Between the two possible combinations for
 217 wheat crops, the coefficient of variation for TFI varied from 0% for insecticides to 1.98% for

218 fungicides. The coefficient of variation for N fertilisation of wheat was 0.09%, and that for the
219 gross margin was 0.98%. For the 512 combinations of pea crops, the coefficients of variation
220 were 0%, 1.15% and 0.42% for TFI, N fertilisation and gross margin, respectively. For the 225
221 combinations of rapeseed crops, the coefficients of variation were 0% for both the TFI and N
222 fertilisation and 4.34% for the gross margin. Barley had a unique combination, so calculating a
223 coefficient of variation was not relevant in this case.

224 For wheat and peas, the p-values were homogeneous for all indicators across all the different
225 combinations, which means that the significance of the results was stable across the
226 combinations ($p < 0.01$). For rapeseed, the results were not significant for the TFI and N
227 fertilisation ($p > 0.05$), but they were significant for the gross margin in 136 of the 225
228 combinations ($p < 0.05$). Overall, we found little variability in the results across the different
229 combinations. Therefore, in the following sections, we present the results for a random
230 combination of wheat, peas and rapeseed, and we present the results of the unique combination
231 of barley.


232 3.2 Pesticide use

233 Our results show a significant overall reduction in the TFI for intercrops compared with sole
234 crops for wheat, barley and peas (Figure 3). Considering all the pesticide categories, TFI was
235 reduced on intercropped wheat by -1.46 (or -49.5% , $p < 0.01$) on average compared with sole
236 wheat (Figure 6). This reduction was of a similar magnitude for intercropped barley compared
237 with sole barley crops (-1.80 or -49.7% , $p < 0.001$) and was the highest for intercropped peas
238 compared with sole pea crops (-3.34 or -83.2% , $p < 0.01$) (Figure 6). We also noticed that the
239 interquartile range for intercropped pea TFI was much smaller than that of sole peas, though
240 we did not observe such a difference between sole crops and intercrops for the other species
241 (Figure 3). Rapeseed sole crops and intercrops showed no significant difference in the TFI.

242 The herbicide TFI in intercrops was significantly reduced for wheat (-0.52 , $p < 0.01$), barley
243 (-0.57 , $p < 0.01$) and peas (-0.99 , $p < 0.001$) compared with sole crops (Figure 6). We noted an
244 increase of $+0.27$ in the herbicide TFI for rapeseed cultivated in intercrops (Figure 6). However,
245 this increase was not significant ($p > 0.05$).

246 A reduction in the fungicide TFI was observed for all intercrops, though this reduction was not
247 significant for wheat and rapeseed ($p > 0.05$). The fungicide TFI was reduced by -0.84 ($p < 0.01$)
248 for barley and -1.19 ($p < 0.001$) for peas, which resulted in an average fungicide TFI close to 0
249 for barley- and pea-based intercrops (Figure 6).

250 For insecticides, our results showed no significant difference in the TFI for wheat, barley and
 251 rapeseed ($p>0.05$), but we observed a significant average reduction of -0.485 ($p<0.01$) for peas
 252 (Figure 6).




253

254 Figure 3: Treatment Frequency Index for wheat, barley, pea and rapeseed sole crops and
255 intercrops. The boxplots are based on the TFI values for each individual in each group. The
256 significance codes (ns = $p>0.05$; * = $p\leq 0.05$; ** = $p\leq 0.01$; *** = $p\leq 0.001$) apply to the paired
257 t-test results. The red diamonds show the mean values.

258 3.3 Nitrogen fertilisation

259 Our results show a significant reduction of $-72.2 \text{ kg N ha}^{-1}$ (or -46.8%) applied to intercropped
260 wheat (Figure 6). Although not statistically significant ($p>0.05$), we observed an average
261 reduction of $-11.8 \text{ kg N ha}^{-1}$ (or -12.3%) for intercropped barley compared with sole barley
262 crops (Figure 6). Peas, on the other hand, showed a significant increase of $+45.8 \text{ kg N ha}^{-1}$
263 ($p<0.001$) (Figure 6) for intercrops compared with sole crops, which received an average of 6.8
264 kg N ha^{-1} (Figure 4). For rapeseed, the results did not show any significant difference ($p>0.05$)
265 between fertilisation on intercrops and sole crops. However, we observed lower variability in
266 the amount of mineral nitrogen applied to intercrops versus sole crops (Figure 4).




267

268 Figure 4: Amount of mineral nitrogen applied to sole crops and intercrops for wheat, barley,
 269 peas and rapeseed. The boxplots are based on the amount of mineral nitrogen applied to each
 270 individual in each group. The significance codes (ns = $p > 0.05$; * = $p \leq 0.05$; ** = $p \leq 0.01$; *** =
 271 $p \leq 0.001$) apply to the paired t-tests. The red diamonds show the mean values.

272 3.4 Gross margin


273 Our results show reductions in the gross margin for intercropped wheat (-184 € ha^{-1} on average,
274 or -17.1%) and barley (-97 € ha^{-1} on average, or -14.7%) and an increase in the gross margin
275 for intercropped rapeseed ($+327 \text{ € ha}^{-1}$ on average, or $+40.9\%$), but these differences were not
276 significant ($p>0.05$) (Figure 6). However, the gross margins for intercropped wheat and barley
277 had lower variability than the corresponding sole crops (Figure 5). For peas, we observed a
278 significant increase of 711 € ha^{-1} (or $+1514\%$, $p<0.001$) for intercrops compared with sole
279 crops (Figure 6).



280

281 Figure 5: Gross margin of sole and intercropped wheat, barley, peas and rapeseed. The boxplots
 282 are based on the gross margin of each individual in each group. The significance codes (ns =
 283 $p > 0.05$; * = $p \leq 0.05$; ** = $p \leq 0.01$; *** = $p \leq 0.001$) apply to the paired t-tests. The red diamonds
 284 show the mean values.

285



286

287 Figure 6: Average difference between the sole crop and intercrop groups for the Treatment
 288 Frequency Indexes (Δ TFI), nitrogen fertilisation (Δ Nfert) and gross margins (Δ GM), estimated
 289 by the paired t-tests. Points represent the average difference and lines represent the 95%
 290 confidence intervals. A negative value indicates a reduction in the intercrop group compared to
 291 the sole crop group. The significance codes (ns = $p > 0.05$; * = $p \leq 0.05$; ** = $p \leq 0.01$; *** =
 292 $p \leq 0.001$) apply to the paired t-tests.

293 **4 Discussion**

294 4.1 Limited number of situations

295 Our data show a limited number of non-organic farms growing wheat-, barley-, pea- or
296 rapeseed-based intercrops. The majority of the wheat, barley and pea intercrops studied were
297 grown in crop–livestock mixed farms. Verret et al. (2020) and Timaeus et al. (2022) observed
298 similar repartitions, which can be explained by the sorting difficulties that hinder the adoption
299 of intercrops in arable field crop farming (Mamine & Farès, 2020). On the contrary, most of the
300 rapeseed-based intercrops studied were grown in arable field crop farms. This difference can
301 be explained by the nature of the intercrops: rapeseed is mainly intercropped with companion
302 plants that are not harvested, so sorting difficulties are less critical.

303 4.2 Effects of intercropping on pesticide use

304 We showed that intercropping enabled TFI reductions of 50% for wheat- and barley-based
305 intercrops and 83% for pea-based intercrops. The overall TFI was, on average, 1.5 for
306 intercropped wheat, 1.8 for intercropped barley and 0.7 for intercropped peas (Figure 3), which
307 are below the national averages of 5.1 for wheat, 4.4 for barley and 4.6 for peas in France,
308 according to the 2017 “Pratiques culturales” (cropping practice) national survey (Agreste,
309 2017). Though our results do not show a significant overall reduction in the TFI for intercropped
310 rapeseed, the average TFI (4.7) was still lower than the national average of 5.7 (Agreste, 2017).
311 It is also worth noting that in the DEPHY data, the average TFIs for pure stands of wheat, barley
312 and peas were lower than the national average, whereas the TFI for pure rapeseed in arable field
313 crop farms in the DEPHY data was equal to the national average (Figure S1).

314 Our analysis showed a reduction in herbicide use for intercropped wheat, barley and peas. One
315 reason for this might be that the number of available herbicide molecules that are selective for
316 both/all mixed crops and can efficiently control weeds is often limited (more limited than for
317 sole crops). Moreover, in our data, wheat and barley were often mixed with legumes, such as
318 peas, thus creating an intercrop highly competitive with weeds (Corre-Hellou et al., 2011; Gu
319 et al., 2021; Hauggaard-Nielsen et al., 2008). Wheat, barley and peas were also involved in
320 more complex mixtures with at least another cereal and/or legume (e.g., wheat–triticale–oat–
321 pea–vetch; Table S2). Such intercrops have a high coverage potential that can help suppress
322 weeds (Pelzer et al., 2014), thus reducing the need for herbicides. Grown as a sole crop, peas
323 have low competitive ability against weeds (Lemerle et al., 1995). Our results confirmed that
324 intercropping peas can foster stronger competitiveness against weeds: of the 30 sole pea crops

325 studied, 25 received herbicide treatments, whereas only seven intercropped peas were treated
326 with herbicide. As for rapeseed, we did not observe a significant difference in herbicide use,
327 although it seems that in some cases, more treatments were applied to intercropped rapeseed
328 than sole rapeseed crops. In our dataset, rapeseed was mainly intercropped with legume
329 companion plants that are not harvested (e.g., faba beans, white clover and/or common vetch).
330 Companion plants can help reduce weeds in rapeseed crops but cannot totally suppress weeds
331 (Lorin et al., 2015; Verret et al., 2017), and some plants have better control over weeds than
332 others. According to Emery et al. (2021), spring faba beans and clover are better suited to
333 controlling weeds than winter faba beans and peas. Herbicides are then likely to be used if the
334 farmer is unsatisfied with the weed control provided by the companion plant (Verret et al.,
335 2020). Moreover, though most companion crops are frost-sensitive legumes, frost may not be
336 enough to destroy them, so chemical weeding may still be necessary.

337 Our results show a net reduction in fungicide use for wheat-, barley- and pea-based intercrops.
338 Although the reduction was not significant for wheat, we found that of the 16 sole wheat crops
339 studied, nine were treated with fungicides, whereas only four wheat intercrops received
340 fungicides. Notably, no wheat–legume intercrops received fungicides. Eleven of the 12 sole
341 barley crops studied received a fungicide treatment, whereas only four barley intercrops were
342 treated with fungicides. Only three pea intercrops were treated with fungicides (vs. 19 sole pea
343 crops). These observed reductions in fungicide treatments are consistent with the barrier effect
344 created by the species mixtures, which limit the spread of diseases (Boudreau, 2013; Finckh et
345 al., 2000). We did not observe any differences in fungicide treatments in rapeseed intercrops
346 versus sole crops. According to Cadoux & Sauzet (2016), no study has yet shown the effect of
347 companion plants on rapeseed diseases. Furthermore, the barrier effect observed in wheat-,
348 barley- and pea-based intercrops is less likely to work for rapeseed because the rapeseed
349 companion plants are destroyed (either by frost or weeding, usually during the winter) and
350 cannot mechanically protect the rapeseed until the end of its growing cycle.

351 We did not observe any decrease in the insecticide TFI, except for pea-based intercrops. The
352 insecticide TFI was already close to zero in the wheat and barley sole crops and could therefore
353 hardly be decreased by intercropping. Insecticide treatments for cereals, such as wheat and
354 barley, are primarily done through seed coating, which was not included in our insecticide TFI
355 metric. Our data (Table S2) show that of the 16 pure wheat stands, 13 were sown using coated
356 seeds, and the remaining three, which were not sown using coated seeds, received an insecticide
357 treatment. Among all the wheat-based intercrops, six were sown using coated seeds, whereas

358 none of the wheat–legume intercrops (n=9) used coated seeds. Barley seed coating was used
359 for both sole crops and intercrops (10 in each case). Regarding peas, we observed a reduction
360 in the insecticide TFI for intercrops, which suggests that the mixture may have helped establish
361 natural predators of pea pests (Puliga et al., 2023) and/or created a barrier effect (Ratnadass et
362 al., 2012). For rapeseed, there was no difference in the insecticide TFI between sole crops and
363 intercrops. The effects of companion plants on rapeseed pests in the literature are nuanced:
364 although companion plants can favour natural predators and create a barrier or even a repellent
365 effect, these effects may not necessarily be sufficient to reduce the use of insecticides (Cadoux
366 et al., 2015). Furthermore, depending on the chosen companion plants, the effects on insects
367 are not always beneficial. In an experiment by Emery et al. (2021), faba bean companion plants
368 did not reduce the damage caused by pollen beetles compared with sole rapeseed crops, and the
369 association with berseem clover even resulted in more significant damage. Lastly, the rapeseed–
370 companion plant association can be effective against adult cabbage stem flea beetles but not
371 always against larvae (Breitenmoser et al., 2020).

372 4.3 Effects of intercropping on mineral nitrogen fertilisation

373 Our results showed a 50% reduction in the amount of mineral nitrogen applied to intercropped
374 wheat compared with sole wheat. This result is in line with other research showing that cereal–
375 legume combinations enable better use of soil nitrogen (e.g. Jensen et al., 2020; Rodriguez et
376 al., 2020). However, this decrease was less marked for intercropped barley. Although
377 fertilisation needs can be reduced by cereal–legume intercrops, depending on the farmer’s
378 objectives, mineral fertilisation can be employed as an adaptation tool. If they aim to obtain
379 fodder, then even a small amount of nitrogen will result in higher productivity for a barley–pea
380 intercrop, as Cowden et al. (2020) reported. If the farmer needs to grow a larger proportion of
381 cereals than legumes, then applying mineral nitrogen will negatively affect biological nitrogen
382 fixation by the legumes, reducing their growth (Ghaley et al., 2005; Naudin et al., 2010). This
383 also explains the increase in nitrogen applied to intercropped peas in our results, whereas no
384 application was necessary for sole pea crops. Conversely, one of the main advantages of
385 combining rapeseed with companion plants is that the mineralisation of the companion plant
386 improves the nitrogen nutrition of the rapeseed (Cadoux et al., 2015; Lorin et al., 2016).
387 However, our analysis did not show any significant reduction in the amount of nitrogen applied
388 to the intercropped rapeseed, though it was reduced for 22 of 31 rapeseed intercrops. The need
389 for fertilisation can be highly dependent on various factors, such as the date of the destruction
390 of the companion plant (the use of herbicides can lead to later destruction than by frost, thus

391 reducing the period of mineralisation of the companion plant), the soil mineral nitrogen content
392 and the species chosen as companion plants (Lorin et al., 2016). However, for the 22 cases
393 where nitrogen fertilisation was reduced, the average reduction was 35 kg N ha⁻¹ (Figure 4).
394 This finding is consistent with the results of Lorin et al. (2016), which show that choosing the
395 best legumes for the circumstances can reduce nitrogen fertilisation by 20–40 kg ha⁻¹. This
396 reduction is also close to that of 30 kg N ha⁻¹ observed by Cadoux et al. (2015).

397 4.4 Effects of intercropping on gross margins

398 For wheat, there were only four cases out of 16 in which the gross margin of the intercrop was
399 better than that of the sole crop, thanks to improved gross production and lower production
400 costs for the intercrops (Table S2). This may be linked to the intercrops' greater resilience to
401 weeds and diseases (Li et al., 2023), possibly resulting in a reduction in herbicide and fungicide
402 use. Otherwise, in most cases, gross production was lower in intercrops than in sole crops.
403 However, lower production costs compensated for this reduction, so there was no significant
404 gross margin loss. Conversely, in the case of barley, a possible increase in intercropping
405 production costs was offset by higher gross production, which limited the reduction in the gross
406 margin or even improved it. For peas, we observed a significant increase in the gross margin of
407 intercrops compared with sole crops. In 19 of the 30 cases, the increase was linked to both
408 higher gross production and lower production costs. Pea production can be improved in
409 intercrops thanks to the staking function of cereals, which limits the lodging of the pea and
410 therefore allows for a better harvest (Kontturi et al., 2011; Verret et al., 2020). The reduction in
411 pesticide-related costs partly offset the increase in nitrogen-related costs.

412 We observed higher gross production for intercropped rapeseed in 18 of the 31 cases (Table
413 S2). In 17 cases, this resulted in a higher gross margin (even with additional production costs
414 in seven cases). We did not observe any significant differences in the use of pesticides or
415 nitrogen fertilisation between rapeseed intercrops and sole crops, but intercropping rapeseed
416 tended to improve GMs. Depending on the farmers' objectives, rather than reducing inputs,
417 intercropping rapeseed with companion plants could be a lever for improving production
418 without changing crop management practices. However, some farmers who grew rapeseed with
419 companion plants to promote biocontrol and limit pesticide use eventually abandoned the
420 practice as they did not consider it effective enough in this respect (Verret et al., 2020).

421 4.5 Limitations of the study

422 Our study compared sole crops and intercrops grown under conditions that were as similar as
423 possible. The DEPHY data did not allow us to locate the crops more precisely than at the
424 municipal level, so we could not combine our data with additional soil data. We also could not
425 construct intercrop–sole crop pairs based on climate, soil or seasonal pest pressure. As a result,
426 the paired intercrops and sole crops were in municipalities distant of 250 km on average, which
427 might have generated statistical noise but we assume that by comparing sole crops and
428 intercrops grown under similar farming practices, we limited bias in our analysis. Nevertheless,
429 our study showed an overall reduction in TFI for intercrops compared with sole crops for wheat,
430 barley and peas, suggesting that intercrops effectively reduced pesticide use in different soil
431 and climate contexts. Finally, farmers’ decisions regarding inputs are not conditioned solely by
432 the agronomic variables we used: social aspects, particularly those concerning both the farmer
433 and their farm, can influence these decisions (Darnhofer et al., 2012; Salembier et al., 2015),
434 but no information on these factors was reported in our database. Thus, a more comprehensive
435 study of farmers is needed to better understand their motivation for intercropping and the
436 associated crop management strategies.

437 **5 Conclusion and perspectives**

438 Our study showed that intercrops enabled the studied farms (conventional arable field crop and
439 crop–livestock mixed farming) to reduce pesticide use by 50% for wheat- and barley-based
440 intercrops and up to 83% for pea-based intercrops. The effect of intercropping on herbicide use
441 was particularly strong, with a decrease of more than 50% for wheat and barley and 86% for
442 peas. The effect of fungicides and insecticides was also striking for peas, with reductions of
443 93% and 100%, respectively. This effect was less clear for insecticides used on wheat and barley
444 as these two crops benefit more from seed coating than insecticide sprays. According to our
445 results, intercropping can also enable a reduction in mineral nitrogen fertilisation by up to 50%
446 for wheat without eliminating the need for fertilisation. Finally, by improving production and/or
447 reducing production costs, intercropping does not harm GMs and even improves margins on
448 intercropped peas compared with sole-cropped peas. Rapeseed intercropping consisted mostly
449 of growing rapeseed with sown but not harvested companion plants, and we could not
450 demonstrate any significant effects of this intercropping on pesticide use or nitrogen
451 fertilisation. Tools such as CAPS (Médiène et al., 2016) can be used to choose the best
452 companion plants for rapeseed. Finally, our results show that intercropping efficiently reduces

453 chemical inputs, but the effectiveness depends on the types of intercrops. We found that
454 intercropping is especially relevant for reducing the use of pesticides or mineral fertilisers and
455 should be integrated into a broader strategy of integrated pest management. According to the
456 DEPHY data, few farmers grow intercrops in France, especially in conventional arable field
457 crop farms. In order to better promote intercropping, it is necessary to highlight the farming
458 conditions under which intercrops are grown as well as the reasons leading farmers to adopt
459 this practice.

460 **CRedit authorship contribution statement**

461 Elodie Yan: Conceptualization, Methodology, Formal analysis, Writing – original draft,
462 Writing – review & editing. Nicolas Munier-Jolain: Conceptualization, Resources, Writing –
463 review & editing. Philippe Martin: Conceptualization, Writing – review & editing. Marco
464 Carozzi: Conceptualization, Writing – review & editing.

465 **Acknowledgements**

466 We want to thank the DEPHY farmers, DEPHY “hub coaches” and the AGROSYST Team for
467 granting us access to the DEPHY network database and assistance with the data. We are also
468 grateful to Emmanuelle Porcher for her valuable contribution and feedback on the
469 methodological approach. This study is part of the MoBiDiv project
470 (www.inrae.fr/mobidiv_eng/) and funded by the ANR program Agrobiosphere (ANR-20-
471 PCPA-0006).

472 **References**

473 Agreste. (2017). Enquête pratiques culturales en grandes cultures et prairies 2017—Principaux
474 résultats (Version modifiée)|Agreste, la statistique agricole.
475 <https://agreste.agriculture.gouv.fr/agreste-web/disaron/Chd2009/detail/>
476 Agreste. (2020). Cultures développées (hors fourrage, prairies, fruits, fleurs et vigne).
477 [https://agreste.agriculture.gouv.fr/agreste-](https://agreste.agriculture.gouv.fr/agreste-saiku/?plugin=true&query=query/open/SAANR_DEVELOPPE_2#query/open/SAANR_DEVELOPPE_2)
478 [saiku/?plugin=true&query=query/open/SAANR_DEVELOPPE_2#query/open/SAAN](https://agreste.agriculture.gouv.fr/agreste-saiku/?plugin=true&query=query/open/SAANR_DEVELOPPE_2#query/open/SAANR_DEVELOPPE_2)
479 [R_DEVELOPPE_2](https://agreste.agriculture.gouv.fr/agreste-saiku/?plugin=true&query=query/open/SAANR_DEVELOPPE_2#query/open/SAANR_DEVELOPPE_2)

480 ANSES. (2023). Le catalogue des produits phytopharmaceutiques et de leurs usages, des
481 matières fertilisantes et des supports de culture autorisés en France.
482 <https://ephy.anses.fr/>

483 Bedoussac, L., Journet, E.-P., Hauggaard-Nielsen, H., Naudin, C., Corre-Hellou, G., Jensen, E.
484 S., Prieur, L., & Justes, E. (2015). Ecological principles underlying the increase of
485 productivity achieved by cereal-grain legume intercroops in organic farming. A review.
486 *Agronomy for Sustainable Development*, 35(3), 911-935.
487 <https://doi.org/10.1007/s13593-014-0277-7>

488 Beillouin, D., Ben-Ari, T., Malézieux, E., Seufert, V., & Makowski, D. (2021). Positive but
489 variable effects of crop diversification on biodiversity and ecosystem services. *Global*
490 *Change Biology*, 27(19), 4697-4710. <https://doi.org/10.1111/gcb.15747>

491 Bonke, V., Michels, M., & Musshoff, O. (2021). Will Farmers Accept Lower Gross Margins
492 for the Sustainable Cultivation Method of Mixed Cropping? First Insights from
493 Germany. *Sustainability*, 13(4), Article 4. <https://doi.org/10.3390/su13041631>

494 Bonke, V., & Musshoff, O. (2020). Understanding German farmer's intention to adopt mixed
495 cropping using the theory of planned behavior. *Agronomy for Sustainable*
496 *Development*, 40(6), 48. <https://doi.org/10.1007/s13593-020-00653-0>

497 Boudreau, M. A. (2013). Diseases in Intercropping Systems. *Annual Review of*
498 *Phytopathology*, 51(1), 499-519. [https://doi.org/10.1146/annurev-phyto-082712-](https://doi.org/10.1146/annurev-phyto-082712-102246)
499 [102246](https://doi.org/10.1146/annurev-phyto-082712-102246)

500 Breitenmoser, S., Steinger, T., Hiltbold, I., Grosjean, Y., Nussbaum, V., Bussereau, F., Klötzli,
501 F., Widmer, N., & Baux, A. (2020). Effet des plantes associées au colza d'hiver sur les
502 dégâts d'altises. <https://doi.org/10.34776/AFS11-16>

503 Cadoux, S., & Sauzet, G. (2016). Colza associé à un couvert de légumineuses gélives (Terres
504 Inovia).

505 Cadoux, S., Sauzet, G., Valantin-Morison, M., Pontet, C., Champolivier, L., Robert, C., Lieven,
506 J., Flénet, F., Mangenot, O., Fauvin, P., & Landé, N. (2015). Intercropping frost-
507 sensitive legume crops with winter oilseed rape reduces weed competition, insect
508 damage, and improves nitrogen use efficiency. *OCL*, 22(3), Article 3.
509 <https://doi.org/10.1051/ocl/2015014>

510 Corre-Hellou, G., Dibet, A., Hauggaard-Nielsen, H., Crozat, Y., Gooding, M., Ambus, P.,
511 Dahlmann, C., von Fragstein, P., Pristeri, A., Monti, M., & Jensen, E. S. (2011). The
512 competitive ability of pea–barley intercrops against weeds and the interactions with crop
513 productivity and soil N availability. *Field Crops Research*, 122(3), 264-272.
514 <https://doi.org/10.1016/j.fcr.2011.04.004>

515 Cowden, R. J., Shah, A. N., Lehmann, L. M., Kiær, L. P., Henriksen, C. B., & Ghaley, B. B.
516 (2020). Nitrogen Fertilizer Effects on Pea–Barley Intercrop Productivity Compared to
517 Sole Crops in Denmark. *Sustainability*, 12(22), Article 22.
518 <https://doi.org/10.3390/su12229335>

519 Darnhofer, I., Gibbon, D., & Dedieu, B. (2012). Farming Systems Research : An approach to
520 inquiry. In I. Darnhofer, D. Gibbon, & B. Dedieu (Éds.), *Farming Systems Research*
521 *into the 21st Century: The New Dynamic* (p. 3-31). Springer Netherlands.
522 https://doi.org/10.1007/978-94-007-4503-2_1

523 Emery, S. E., Anderson, P., Carlsson, G., Friberg, H., Larsson, M. C., Wallenhammar, A.-C.,
524 & Lundin, O. (2021). The Potential of Intercropping for Multifunctional Crop
525 Protection in Oilseed Rape (*Brassica napus* L.). *Frontiers in Agronomy*, 3.
526 <https://www.frontiersin.org/articles/10.3389/fagro.2021.782686>

527 Enjalbert, J., Litrico, I., Fournier, E., Médiène, S., Gauffretau, A., Borg, J., Hellou, G.,
528 Goldringer, I., Hannachi, M., Journet, E.-P., Justes, E., Morel, J.-B., Naudin, C.,
529 Sanguin, H., Morison, M., Verret, V., & Bedoussac, L. (2019). Mélanges variétaux et

530 mélanges plurispécifiques – atouts et contraintes. 75, 49-71.
531 <https://doi.org/10.15454/ak5jpd>

532 Finckh, M. R., Gacek, E. S., Goyeau, H., Lannou, C., Merz, U., Mundt, C. C., Munk, L.,
533 Nadziak, J., Newton, A. C., Vallavieille-Pope, C. de, & Wolfe, M. S. (2000). Cereal
534 variety and species mixtures in practice, with emphasis on disease resistance.
535 *Agronomie*, 20(7), 813-837. <https://doi.org/10.1051/agro:2000177>

536 Ghaley, B. B., Hauggaard-Nielsen, H., Høgh-Jensen, H., & Jensen, E. S. (2005). Intercropping
537 of Wheat and Pea as Influenced by Nitrogen Fertilization. *Nutrient Cycling in*
538 *Agroecosystems*, 73(2), 201-212. <https://doi.org/10.1007/s10705-005-2475-9>

539 Gu, C., Bastiaans, L., Anten, N. P. R., Makowski, D., & van der Werf, W. (2021). Annual
540 intercropping suppresses weeds: A meta-analysis. *Agriculture, Ecosystems &*
541 *Environment*, 322, 107658. <https://doi.org/10.1016/j.agee.2021.107658>

542 Hauggaard-Nielsen, H., & Jensen, E. S. (2005). Facilitative root interactions in intercrops. *Plant*
543 *and Soil*, 274(1/2), 237-250.

544 Hauggaard-Nielsen, H., Jørnsgaard, B., Kinane, J., & Jensen, E. S. (2008). Grain legume–cereal
545 intercropping: The practical application of diversity, competition and facilitation in
546 arable and organic cropping systems. *Renewable Agriculture and Food Systems*, 23(1),
547 3-12. <https://doi.org/10.1017/S1742170507002025>

548 IGN. (2023). Institut National de l'Information Géographique et Forestière—IGN. Base de
549 Données. <https://geoservices.ign.fr/telechargement>

550 Jensen, E. S., Carlsson, G., & Hauggaard-Nielsen, H. (2020). Intercropping of grain legumes
551 and cereals improves the use of soil N resources and reduces the requirement for
552 synthetic fertilizer N: A global-scale analysis. *Agronomy for Sustainable Development*,
553 40(1), 5. <https://doi.org/10.1007/s13593-020-0607-x>

554 Kontturi, M., Laine, A., Niskanen, M., Hurme, T., Hyövelä, M., & Peltonen-Sainio, P. (2011).
555 Pea–oat intercrops to sustain lodging resistance and yield formation in northern
556 European conditions. *Acta Agriculturae Scandinavica, Section B — Soil & Plant*
557 *Science*, 61(7), 612-621. <https://doi.org/10.1080/09064710.2010.536780>

558 Lechenet, M., Makowski, D., Py, G., & Munier-Jolain, N. (2016). Profiling farming
559 management strategies with contrasting pesticide use in France. *Agricultural Systems*,
560 149, 40-53. <https://doi.org/10.1016/j.agsy.2016.08.005>

561 Lemerle, D., Verbeek, B., & Coombes, N. (1995). Losses in grain yield of winter crops from
562 *Lolium rigidum* competition depend on crop species, cultivar and season. *Weed*
563 *Research*, 35(6), 503-509. <https://doi.org/10.1111/j.1365-3180.1995.tb01648.x>

564 Li, C., Stomph, T.-J., Makowski, D., Li, H., Zhang, C., Zhang, F., & van der Werf, W. (2023).
565 The productive performance of intercropping. *Proceedings of the National Academy of*
566 *Sciences*, 120(2), e2201886120. <https://doi.org/10.1073/pnas.2201886120>

567 Lin, B. B. (2011). Resilience in Agriculture through Crop Diversification: Adaptive
568 Management for Environmental Change. *BioScience*, 61(3), 183-193.
569 <https://doi.org/10.1525/bio.2011.61.3.4>

570 Lithourgidis, A., Dordas, C., Damalas, C., & Vlachostergios, D. (2011). Annual intercrops : An
571 alternative pathway for sustainable agriculture. *Australian Journal of Crop Science*, 5,
572 396-410.

573 Lorin, M., Jeuffroy, M.-H., Butier, A., & Valantin-Morison, M. (2015). Undersowing winter
574 oilseed rape with frost-sensitive legume living mulches to improve weed control.
575 *European Journal of Agronomy*, 71, 96-105. <https://doi.org/10.1016/j.eja.2015.09.001>

576 Lorin, M., Jeuffroy, M.-H., Butier, A., & Valantin-Morison, M. (2016). Undersowing winter
577 oilseed rape with frost-sensitive legume living mulch : Consequences for cash crop

578 nitrogen nutrition. *Field Crops Research*, 193, 24-33.
579 <https://doi.org/10.1016/j.fcr.2016.03.002>

580 Malézieux, E., Crozat, Y., Dupraz, C., Laurans, M., Makowski, D., Ozier-Lafontaine, H.,
581 Rapidel, B., Tourdonnet, S., & Valantin-Morison, M. (2009). Mixing plant species in
582 cropping systems : Concepts, tools and models. A review. *Agronomy for Sustainable*
583 *Development*, 29(1), 43-62. <https://doi.org/10.1051/agro:2007057>

584 Mamine, F., & Farès, M. (2020). Barriers and Levers to Developing Wheat–Pea Intercropping
585 in Europe : A Review. *Sustainability*, 12(17), Article 17.
586 <https://doi.org/10.3390/su12176962>

587 Matson, P. A., Parton, W. J., Power, A. G., & Swift, M. J. (1997). Agricultural Intensification
588 and Ecosystem Properties. *Science*, 277(5325), 504-509.
589 <https://doi.org/10.1126/science.277.5325.504>

590 Médiène, S., Verret, V., Felix, J., & Valantin-Morison, M. (2016). A tool integrating and
591 sharing knowledge to select legume species for oilseed rape intercropping. Second
592 International Legumes Society Conference, Troia, Portugal.

593 Naudin, C., Corre-Hellou, G., Pineau, S., Crozat, Y., & Jeuffroy, M.-H. (2010). The effect of
594 various dynamics of N availability on winter pea–wheat intercrops : Crop growth, N
595 partitioning and symbiotic N₂ fixation. *Field Crops Research*, 119(1), 2-11.
596 <https://doi.org/10.1016/j.fcr.2010.06.002>

597 Pelzer, E., Bedoussac, L., Corre-Hellou, G., Jeuffroy, M.-H., Métivier, T., & Naudin, C. (2014).
598 Association de cultures annuelles combinant une légumineuse et une céréale : Retours
599 d’expériences d’agriculteurs et analyse. *Innovations Agronomiques*, 40, 73.

600 Pingault, N., Pleyber, É., Champeaux, C., Guichard, L., & Omon, B. (2009). Produits
601 phytosanitaires et protection intégrée des cultures : L’indicateur de fréquence de
602 traitement.

603 Puliga, G. A., Arlotti, D., & Dauber, J. (2023). The effects of wheat-pea mixed intercropping
604 on biocontrol potential of generalist predators in a long-term experimental trial. *Annals*
605 *of Applied Biology*, 182(1), 37-47. <https://doi.org/10.1111/aab.12792>

606 Raseduzzaman, Md., & Jensen, E. S. (2017). Does intercropping enhance yield stability in
607 arable crop production? A meta-analysis. *European Journal of Agronomy*, 91, 25-33.
608 <https://doi.org/10.1016/j.eja.2017.09.009>

609 Ratnadass, A., Fernandes, P., Avelino, J., & Habib, R. (2012). Plant species diversity for
610 sustainable management of crop pests and diseases in agroecosystems: A review.
611 *Agronomy for Sustainable Development*, 32(1), 273-303.
612 <https://doi.org/10.1007/s13593-011-0022-4>

613 Rodriguez, C., Carlsson, G., Englund, J.-E., Flöhr, A., Pelzer, E., Jeuffroy, M.-H., Makowski,
614 D., & Jensen, E. S. (2020). Grain legume-cereal intercropping enhances the use of soil-
615 derived and biologically fixed nitrogen in temperate agroecosystems. A meta-analysis.
616 *European Journal of Agronomy*, 118, 126077.
617 <https://doi.org/10.1016/j.eja.2020.126077>

618 Salembier, C., Elverdin, J. H., & Meynard, J.-M. (2015). Tracking on-farm innovations to
619 unearth alternatives to the dominant soybean-based system in the Argentinean Pampa.
620 *Agronomy for Sustainable Development*, 36(1), 1. [https://doi.org/10.1007/s13593-015-](https://doi.org/10.1007/s13593-015-0343-9)
621 [0343-9](https://doi.org/10.1007/s13593-015-0343-9)

622 Tilman, D., Cassman, K. G., Matson, P. A., Naylor, R., & Polasky, S. (2002). Agricultural
623 sustainability and intensive production practices. *Nature*, 418(6898), 671-677.
624 <https://doi.org/10.1038/nature01014>

625 Timaeus, J., Ruigrok, T., Siegmeyer, T., & Finckh, M. R. (2022). Adoption of Food Species
626 Mixtures from Farmers' Perspectives in Germany: Managing Complexity and

627 Harnessing Advantages. *Agriculture*, 12(5), Article 5.
628 <https://doi.org/10.3390/agriculture12050697>

629 Tudi, M., Daniel Ruan, H., Wang, L., Lyu, J., Sadler, R., Connell, D., Chu, C., & Phung, D. T.
630 (2021). Agriculture Development, Pesticide Application and Its Impact on the
631 Environment. *International Journal of Environmental Research and Public Health*,
632 18(3), Article 3. <https://doi.org/10.3390/ijerph18031112>

633 Verret, V., Gardarin, A., Makowski, D., Lorin, M., Cadoux, S., Butier, A., & Valantin-Morison,
634 M. (2017). Assessment of the benefits of frost-sensitive companion plants in winter
635 rapeseed. *European Journal of Agronomy*, 91, 93-103.
636 <https://doi.org/10.1016/j.eja.2017.09.006>

637 Verret, V., Pelzer, E., Bedoussac, L., & Jeuffroy, M.-H. (2020). Tracking on-farm innovative
638 practices to support crop mixture design : The case of annual mixtures including a
639 legume crop. *European Journal of Agronomy*, 115, 126018.
640 <https://doi.org/10.1016/j.eja.2020.126018>

641 WHO. (1990). Public health impact of pesticides used in agriculture. World Health
642 Organization. <https://apps.who.int/iris/handle/10665/39772>

643