



HAL
open science

Glucose-6-Phosphate Dehydrogenases: The Hidden Players of Plant Physiology

Zhengrong Jiang, Ming Wang, Michael Nicolas, Laurent Ogé, Maria-Dolores Pérez-Garcia, Laurent Crespel, Ganghua Li, Yanfeng Ding, José Le Gourrierc, Philippe Grappin, et al.

► **To cite this version:**

Zhengrong Jiang, Ming Wang, Michael Nicolas, Laurent Ogé, Maria-Dolores Pérez-Garcia, et al.. Glucose-6-Phosphate Dehydrogenases: The Hidden Players of Plant Physiology. International Journal of Molecular Sciences, 2022, 23 (24), pp.16128. 10.3390/ijms232416128 . hal-04309049

HAL Id: hal-04309049

<https://hal.inrae.fr/hal-04309049v1>

Submitted on 27 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Review

Glucose-6-Phosphate Dehydrogenases: The Hidden Players of Plant Physiology

Zhengrong Jiang^{1,2,†}, Ming Wang^{3,†}, Michael Nicolas⁴ , Laurent Ogé¹, Maria-Dolores Pérez-García¹ , Laurent Crespel¹, Ganghua Li², Yanfeng Ding² , José Le Gourrierc¹ , Philippe Grappin¹ and Soulaiman Sakr^{1,*}

¹ Institut Agro, University of Angers, INRAE, IRHS, SFR QUASAV, 49000 Angers, France

² College of Agronomy, Nanjing Agricultural University, Nanjing 210095, China

³ Dryland-Technology Key Laboratory of Shandong Province, College of Agronomy, Qingdao Agricultural University, Qingdao 266109, China

⁴ Plant Molecular Genetics Department, Centro Nacional de Biotecnología-CSIC, Campus Universidad Autónoma de Madrid, 28049 Madrid, Spain

* Correspondence: soulaiman.sakr@agrocampus-ouest.fr

† These authors contributed equally to this work.

Abstract: Glucose-6-phosphate dehydrogenase (G6PDH) catalyzes a metabolic hub between glycolysis and the pentose phosphate pathway (PPP), which is the oxidation of glucose-6-phosphate (G6P) to 6-phosphogluconolactone concomitantly with the production of nicotinamide adenine dinucleotide phosphate (NADPH), a reducing power. It is considered to be the rate-limiting step that governs carbon flow through the oxidative pentose phosphate pathway (OPPP). The OPPP is the main supplier of reductant (NADPH) for several “reducing” biosynthetic reactions. Although it is involved in multiple physiological processes, current knowledge on its exact role and regulation is still piecemeal. The present review provides a concise and comprehensive picture of the diversity of plant G6PDHs and their role in seed germination, nitrogen assimilation, plant branching, and plant response to abiotic stress. This work will help define future research directions to improve our knowledge of G6PDHs in plant physiology and to integrate this hidden player in plant performance.

Keywords: glucose-6-phosphate dehydrogenase; seed germination; apical dominance; sugar signaling; abiotic stress; ROS



Citation: Jiang, Z.; Wang, M.; Nicolas, M.; Ogé, L.; Pérez-García, M.-D.; Crespel, L.; Li, G.; Ding, Y.; Le Gourrierc, J.; Grappin, P.; et al. Glucose-6-Phosphate Dehydrogenases: The Hidden Players of Plant Physiology. *Int. J. Mol. Sci.* **2022**, *23*, 16128. <https://doi.org/10.3390/ijms232416128>

Academic Editor: Jozef Kovacik

Received: 23 November 2022

Accepted: 12 December 2022

Published: 17 December 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Plant performance intrinsically depends on balanced carbon flux through interconnected metabolic hubs. Glucose-6-phosphate dehydrogenases (G6PDHs) catalyze a metabolic node between glycolysis and the pentose phosphate pathway (PPP), i.e., the oxidation of glucose-6-phosphate (G6P)—A substrate generated by hexokinase during glycolysis—to 6-phosphogluconolactone, concomitant with the production of the reductant nicotinamide adenine dinucleotide phosphate (NADPH). This reaction is considered as the rate-limiting step of the oxidative pentose phosphate pathway (OPPP), and controls OPPP-dependent carbohydrate allocation [1–3]. The PPP is divided into two branches with two distinct functions. The OPPP is the irreversible oxidative branch of the PPP. It comprises three irreversible reactions that convert glucose 6 phosphate (G6P) into carbon dioxide (CO₂) and ribulose-5-phosphate (Ru5P, a pentose phosphate) while two molecules of NADPH are produced (Figure 1). Ru5P becomes the substrate of the reversible non-oxidative phase (NOPPP), which is the reversible branch of the PPP. The second branch is the non-oxidative pentose pathway (NOPP). It is composed of reversible transaldolase and transketolase reactions enabling the cell to control carbon flux between the PPP and glycolysis [3–6]. Including two NADPH-producing steps, the OPPP is the main provider of reductant (NADPH) for “reductive” biosynthetic reactions (e.g., assimilation of inorganic nitrogen,

fatty acid synthesis) and for ROS scavenging. The NOPPP rather provides precursors (ribose-5-phosphate; erythro-4-phosphate) for aromatic amino-acid, nucleotide, and co-factor synthesis [3,6]. The OPPP is an almost ubiquitous pathway, present in all eukaryotic cells and most bacteria, and seems to be more recent than the NOPPP in evolution [3,5].




Figure 1. Overview of the reactions of the pentose phosphate pathway (PPP) and its connection to glycolysis. The glycolytic pathway is colored in green. The oxidative part of the PPP is colored in orange, the non-oxidative part in purple; one-headed arrows designate physiologically irreversible reactions, two-headed arrows reversible ones; abbreviation meanings: G6PDH, glucose-6-phosphate dehydrogenase (EC 1.1.1.49); 6PGL, 6-phosphogluconolactonase (EC 3.1.1.31); 6PGDH, 6-phosphogluconate dehydrogenase (EC 1.1.1.44); RPI, ribose-5-phosphate isomerase (EC 5.3.1.6); RuPE, ribulose-5-phosphate 3-epimerase (EC 5.1.3.1); TKT, transketolase (EC 2.2.1.1); TAL, transaldolase (EC 2.2.1.2); PHI, hexose-6-phosphate isomerase (EC 5.3.1.9); SBPase, sedoheptulose-1,7-bisphosphatase (EC 3.1.3.37); other details in the text.

G6PDH was discovered in the early 1930s, when Otto Warburg et al. investigated the enzymatic oxidation of glucose-6-phosphate to 6-phosphogluconate (6PG) in yeast. It was initially called *Zwischenferment* (ZWF1) or intermediate enzyme [5]. Plant G6PDH-cDNA sequences were first isolated from potato [7–9] and have been identified in several monocot and dicot crops [4,10]. Although the available results are assigned to a variety of fundamental processes, including those related to plant growth and plant resistance to abiotic stress, most of them are still piecemeal. This situation hampers the building up of a comprehensive picture of its central role throughout plant life cycle and prevents us from obtaining further insights into the main regulatory network governing its involvement in plant growth and resistance to stresses. We propose a concise overview of the diversity

of plant G6PDHs and their mechanisms of regulation, and of their role in four main plant physiological processes: seed germination, nitrogen assimilation, plant branching, and plant response to abiotic stresses. This work will provide a solid basis for future lines of research aimed at improving our knowledge of G6PDHs in plant physiology and integrating this hidden player in plant resilience to climate change.

2. Classification and Regulation of G6PDH

In higher plants, G6PDHs reside in two cellular compartments, the cytosol and plastids [11,12]. Genome-wide analysis of Arabidopsis G6PDH indicates the existence of four plastidial (pla-G6PD) and two cytosolic (cy-G6PD) isoforms [13,14], also reported in several crops [15,16]. Plastidial G6PDH comprises three functional isoforms belonging to two distinct groups [P1 (G6PD1), P2 (G6PD2, G6PD3)] and a non-functional one (G6PD4) belonging to the P0 group. Cyttoplasmic G6PDH is divided into two groups [P5 (G6PDH5) and P6 (G6PDH6)], providing 60–80% of the total activity. Besides their respective location, these G6PDH isoforms differ by their amino-sequences and their mode of regulation (Figure 2).




Figure 2. Comparison of different G6PDH isoforms from 8 higher plants; (A), phylogenetic tree showing the relative positions of 39 different gene encoding isoforms from higher plants, inferring by the maximum likelihood method of complete protein sequences; 10 sequences are P1-G6PDH, 10 sequences are P2-G6PDH, 8 sequences are P0-G6PDH; and 11 sequences are Cy-G6PDH. Legend for plant species: At, *Arabidopsis thaliana*; Os, *Oryza sativa Japonica Group*; Zm, *Zea mays*; Rc, *Rosa chinensis*; Vit, *Vitis vinifera*; Pp, *Prunus persica*; Solye, *Solanum lycopersicum*; St, *Solanum tuberosum*. (B), List of G6PDH isoforms and their relative gene symbol from different higher plants, summarizing from the phylogenetic tree constructed in (A). The complete list of sequences using for tree construction is showed in Table S1.

2.1. P1-G6PDH

The chloroplastic G6PDH has been defined as a P1-G6PDH and is mainly post-transcriptionally inhibited by high redox status (a high content of NADPH) and accumulation of ribulose-5P [17,18]. During photosynthetic electron transport in the light, a redox chain (the ferredoxin/thioredoxin system) inactivates the P1-G6PDH to guarantee an efficient photosynthesis, which activates the Calvin cycle and several stromal target enzymes such as fructose-1,6-bisphosphatase, NADP-malate dehydrogenase, phosphoribulokinase, and others [7,19]. This inhibitory effect of P1-G6PDH is removed in the dark when the NADPH level dropped, enabling the activation of the OPPP pathway to produce reducing equivalents [20].

2.2. P2-G6PDH

The plastidial P2-G6PDH is highly related to the rate of the OPPP, and expressed in almost all plant organs including growing tissues, photosynthetic, and non-photosynthetic tissues [13,21]. The activity of P2-G6PDH is more likely to be regulated in a similar manner as P1-G6PD, but their sensitivity to NADPH differs [22,23]. P2-G6PDH exhibits a 5–10 fold higher inhibition by NADPH than Cy-G6PDH and P1-G6PDH [24]. In addition, P2-G6PDH is much less sensitive than P1-G6PDH to regulation by thioredoxin and glutathione (GSH). A relatively significant role of P2-G6PDH generally consists in providing reductant in heterotrophic tissues in the absence of photochemical generation of reductants.

2.3. The Enigmatic P0-G6PDH

P0-G6PDH is considered as an enigmatic isoform found in peroxisomes and without any enzymatic activity. P1-G6PDH is involved in cysteine-dependent interaction with P0-G6PDH [25]. Due to the presence of a peroxisome targeting sequence (PTS) at the C-terminus of P0-G6PDH, these heterodimers can enter peroxisomes [26]. It is generally believed that the oxidative portion of the PPP exists in peroxisomes [27,28]. Additionally, some irreversible reactions of PPP are catalyzed by 6-phosphogluconolactonase (6-PGL) and 6-phosphogluconate dehydrogenase (6PGDH), both localized in the Arabidopsis peroxisome; hence, it is considered an efficient NADPH production mechanism in this organelle [26,29,30].

2.4. Cytosolic G6PDH (Cy-G6PDH)

In higher plants, the two cytosolic isoforms G6PD5 and G6PD6 are differently expressed in various tissues, even though they were initially purified from roots [13,21,31,32]. The cy-G6PD isoform is lowly sensitive to energy changes, which are regulated by the NADPH/NADP⁺ ratio and inhibited by NADPH [31]. However, the activity of Cy-G6PDH is regulated by a sugar-sensing mechanism, and plays an important role in amino acid biosynthesis according to the carbon status of plants [33]. Moreover, Cy-G6PDH expression is tightly induced by abscisic acid (ABA; Hou et al., 2006) and sensitive to light [34], which mainly control the activity of P1-G6PDH [35].

3. G6PDH and Seed Germination

In the 1980s, a great deal of studies carried out on several species (e.g., *Gossypium*, *Pisum sativum*, *Arachis hypogaea*, *Prunus cerasus*, *Phaseolus mungo*, *Avena fatua*) reported a tight correlation between dormancy breaking treatments and increased G6PDH activity. These investigations were based on the activity of the two cytosolic enzymes of the OPPP (G6PDH and 6-PDGH) and one glycolysis-related enzyme (aldolase) in seed tissues during dormancy-breaking treatments (e.g., stratification, after-ripening). Aldolase catalyzes the conversion of fructose 1-6-diphosphate to glyceraldehyde 3-phosphate and dihydroxyacetone phosphate through the glycolysis pathway [36]. These authors hypothesized that activation of the OPPP in germinating seeds played a pivotal role in dormancy breaking and storage mobilization by controlling redox homeostasis and enzyme activities [37–43]. The OPPP was found highly induced during seed germination, much more so in the

endosperm than in the radicle [44]. The specificity of the spatial regulation of the OPPP pointed out the respective roles of these sub-compartments in the regulation of Arabidopsis seed germination.

The involvement of different isoforms of G6PDH—cytosolic as well as plastidial ones—in dormancy release was investigated. It was proposed that Cy-G6PDH controls germination by maintaining a steady-state level of ROS (Figure 3) required for breaking dormancy [45,46], yet a limited one to avoid excessive oxidative damage in the root apical meristems [47]. Genetic evidence supports that ROS provided by NADPH oxidase in germinating seeds under salt stress stimulate the two cytosolic G6PDHs (Cy-G6PDH5 and Cy-G6PDH6) and increase the cytosolic NADPH content, which in turn dampens ROS damage by activating the GSH–ascorbate cycle involved in H_2O_2 scavenging [48]. Using *cy-g6pdh* deficient mutants, researchers demonstrated that Cy-G6PD5 is required to release dormancy and reduce seed sensitivity to ABA—a hormone involved in germination inhibition—through the repression of the *Abscisic Acid Insensitive 5 (ABI5)* gene [49]. The authors described that ABA induces excessive accumulation of ROS in germinating seeds and seedlings of the *cy-g6pd5*-deficient mutant. ROS control dormancy through post-transcriptional regulation by selective proteins and via RNA oxidation [50,51], cell wall loosening for cell wall elongation, and endosperm weakening [52], the response to ethylene, or through the control of the ABA catabolism/GA biosynthesis [53,54]. Nevertheless, to avoid the deleterious effects of ROS, their level must be finely balanced by the reducing power or detoxifying enzymes that seem to be largely provided by OPPP-derived NADPH. Moreover, Cy-G6PD is described to be induced by ABA, hydrogen peroxide (H_2O_2), or nitric oxide (NO) [15,49,55] and is crucial for plant tolerance to stresses, more likely by maintaining the redox balance [33]. Controlling the redox status of the germinating seeds should be central to optimize plant fitness. At this developmental stage, cyt-G6PDH may modulate both the adaptive mechanisms of dormancy and the plant responses to biotic and abiotic stresses.




Figure 3. Cytosolic G6PDH in radicle of imbibed seed modulates ROS homeostasis and hormonal signaling in the control of seed germination. The NADPH provided by cytosolic G6PDH activity is required both for ROS production by NADPH oxidase and for ROS scavenging by activation of GSH ascorbate cycle; the working model based on genetic evidence proposed that G6PDH modulates ROS to a steady state level controlling ABA and GA activities and dormancy release; GSH, reduced glutathione; ROS, reactive oxygen species; ABA, abscisic acid; GA, gibberellic acid.

The contribution of G6PDH in reserve mobilization through the regulation of thioredoxin (Trx) at the end of germination has also been suggested. Many investigations have underlined the important role of NADPH-dependent Trx systems in reserve mobilization [56–61]. Cytosolic Trx-dependent reduction in storage proteins provides essential organic resources for the transition of germinating seeds to autotrophic seedlings. Moreover, seed plastidial γ -type Trxs (Trx γ) induce plastidial P1-G6PDH, a major source of reducing power in heterotrophic tissues [62]. A functional genetic approach documented that γ -type Trxs contribute to seed germination by regulating ROS levels through the activation of plastidial G6PDH [37,63].

Elucidating the molecular mechanisms whereby G6PDH controls the hormonal metabolism and hormone sensing, ROS detoxification, but also the carbon metabolism and carbon reallocation during the germinating-seed-to-autotrophic-seedling transition will be crucial in the near future (Figure 3). A recent analysis of the whole transcriptome changes following nitrate treatment during seed imbibition showed that genes involved in nitrate assimilation and transport as well as the plastidial P2-G6PDH were upregulated, highlighting a potential link between G6PDH and the well-known nitrate signaling effect on the ABA catabolism and dormancy release.

4. G6PDHs and Nitrogen Assimilation

Nitrogen (N) is one of the most limiting factors for plant growth and productivity. Consequently, plants have various mechanisms for maximum N efficiency [64]. Plant nitrogen nutrition occurs through organic (amino acids, urea) and inorganic (nitrate, ammonium) forms of nitrogen and is governed by a complex regulatory network [65,66]. The main route of nitrate assimilation involves nitrate reductase (NR) and nitrite reductase (NiR), resulting in its reduction into ammonium (NH_4), which is incorporated into amino acids through the joint action of glutamine synthetase (GS) and the glutamine oxoglutarate aminotransferase (GOGAT) cycle [67–69]. A link between the OPPP and inorganic nitrogen assimilation has mainly been reported in non-photosynthetic tissues (e.g., root systems), with a prevailing role of plastidial G6PDH [19], which can meet the high demand for reducing power upon nitrogen assimilation [3]. Early data showed that NiR activity in barley roots relies on elevated levels of NADPH, suggesting that G6PDH could play a key role during nitrate assimilation [70]. In accordance with this, synthesis of glutamate (a final product of the GOGAT cycle) by isolated pea root plastids fed with two GOGAT substrates (glutamine and 2-oxoglutarate) requires the OPPP substrates (Glc6P, ribose P), coordinated with the reducing activity of G6PDH [70,71]. Exogenous supply of inorganic nitrogen (nitrate or ammonium) to barley roots induced plastidial G6PDH at both the transcriptional and protein levels [21,72]. This nitrate-dependent upregulation of G6PDH was also reported in seedlings of NiR knockout mutants, assuming that it is directly triggered by the nitrate-related signaling pathway [73,74]. This fine tuning between nitrogen and the OPPP was transposed to the induction of the major root nitrate uptake transporters (NRT1.1 and NRT2.2) in *Arabidopsis* roots [75,76]. In this case, photosynthesis-derived sugar induces nitrate transporters in the roots, which are repressed by 6-amino-nicotinamide (6-AN), an inhibitor of both G6PDH and 6PGDH (6-phospho-gluco-dehydrogenase) [77,78]. This effect is independent of the hexokinase- and trehalose-6P signaling pathway [75], supporting that sugar-dependent upregulation of nitrate transporters involves an OPPP-dependent signaling mechanism. Taken together, the OPPP and nitrogen uptake/assimilation by heterotrophic organs (roots) are a tightly coordinated process [79], with a key role of root plastidial G6PDH [3]. Huge efforts are still required to disclose the molecular regulatory network mechanisms governing this coordination between nitrogen assimilation and G6PDH regulation. Early studies indicate that the promoter sequences of NiR and G6PDH present the same NIT2 motif, which is a N-metabolism regulating factor [80,81]. Based on this, a promising line of study would be to investigate this close relationship between the OPPP and nitrogen assimilation in other non-photosynthetic organs such as vegetative buds, which also require organic and inorganic nitrogen to sustain their outgrowing activ-

ity [65]. Such a study would obviously bring new insights into the common and specific mechanisms in different biological contexts.

5. G6PDHs and Plant Branching

Shoot branching is crucial for plant development and yield and is greatly dependent on the ability of axillary buds to grow out along the stem [82–84]. Bud outgrowth is very finely regulated by multiple endogenous (e.g., hormones, sugar) and exogenous (e.g., light, water stress) cues [85–92]. In this intricate regulation, sugars behave as signaling entities that promote bud outgrowth through several sugar signaling pathways corresponding to the trehalose 6P-, hexokinase-, glycolysis/tricarboxylic acid (TCA)-, and OPPP-dependent signaling pathways [83,93,94]. The involvement of the OPPP in plant branching was identified following the discovery that bud outgrowth relies on the bud H₂O₂ content: cytokinins (CKs) promote bud outgrowth by reducing H₂O₂ through the induction of the GSH/ascorbate cycle, and, conversely, H₂O₂ accumulation reduces bud ability to grow out [95,96]. The authors of [94] first evidenced the role of the OPPP in sugar branching by demonstrating that the promotive effect of sucrose on bud outgrowth is repressed by 6-AN—an inhibitor of G6PDH—in *Rosa* sp. Molecular experiments conducted on in vitro-cultured vegetative buds and on stably transformed *Rosa* calluses revealed that the OPPP-dependent signaling pathway is involved in both sugar-mediated transcriptional (promoter level) and posttranscriptional (3′ untranslated region) downregulation of Teosinte Branched 1/Branched1 (TB1/BRC1) [91,94,97]. BRC1 is the main inhibitor of shoot branching [98]. An OPPP-specific 300-bp region was identified in the *BRC1* promoter between 1900 and 1600 bp [94]; its 3′UTR contains six putative motifs of the Pumilio/FBF RNA-binding protein family (PUF) [97]. One future task will consist of deciphering the molecular mechanisms underlying the OPPP-mediated downregulation of *BRC1*.

6. G6PDHs and Sugar Signaling

Sugar perception and signaling enable plants to integrate various internal and external cues to achieve nutrient homeostasis, mediate developmental programs, and orchestrate their stress response [99,100]. To sense different sugars, plants have evolved a complex mechanistic system that includes hexose-, disaccharide-, and the OPPP–signaling pathways [101,102]. The OPPP–dependent signaling pathway has been reported in two biological contexts related to sink organs. It drives sugar-mediated stimulation of nitrogen and sulfur acquisition by the roots, downstream and independently of hexokinase signaling and the trehalose-6P signaling pathway [75,76]. It has also been described as a main signaling route of sugar-dependent stimulation of bud outgrowth [94]. However, OPPP-dependent sugar signaling might be more complex and operate through different pathways [75,79]. Three hypotheses are still plausible: (1) one of the carbon metabolites generated through the OPPP could act as a cue; (2) an enzyme of the OPPP—e.g., G6PDH—could exhibit a dual (catalytic and signaling) function like hexokinase (HXK) does [103]; and (3) NADPH resulting from G6PDH activity could be involved in redox regulation via the NADPH-dependent signaling pathway. This latter hypothesis was recently reported for the sugar-mediated regulation of *NRT2.1* expression in the roots, which is related to the redox status of the plant [104]. This regulatory mechanism might also be involved in the OPPP-mediated sugar stimulation of bud outgrowth. Recent data indicate that the ability of *Rosa* buds to grow out depends on their redox status: high levels of H₂O₂ in the buds strongly prevent outgrowth, while high activity of the GSH–ascorbate cycle, which is directly linked to the activity of G6PDH, decreases the H₂O₂ content and promotes bud outgrowth [95]. The relevance of the GSH–ascorbate cycle has been confirmed for CK-induced bud outgrowth under darkness [96]. In this case, CKs positively affect GSH synthesis to stimulate H₂O₂ scavenging, and this allows for bud outgrowth [96]. However, whether CKs regulate G6PDH inside the buds still remains to be unveiled, while we do know that G6PDH is upregulated by sugar [90]. It will be of great interest to deeply investigate the role of G6PDH in bud outgrowth regulation by sugars, hormones, and ROS, and to identify the

OPPP-dependent regulatory molecular network. All these findings lay the cornerstones for deciphering the exact role of G6PDH in shoot branching, for instance, by characterizing the branching phenotype of G6PDH-deficient Arabidopsis mutants.

7. G6PDH and Abiotic Stress

7.1. Identification of Link between G6PDH and Abiotic Stress

As sessile organisms, plants have to cope with various abiotic stresses such as salinity, drought, or temperature changes [105]. G6PDHs play a crucial function in modulating redox homeostasis when plants are exposed to abiotic stresses [3,106–108]. Some researchers found the role of a cytosolic G6PDH (*NbG6PDH-Cyto*) and two plastidial isoforms of G6PDH (*NbG6PDH-P1* and *NbG6PDH-P2*) in the stress physiology of *Nicotiana benthamiana* [109]. ROS production in link with hypersensitive-response (HR) cell death and NADPH oxidase (NOX, also known as respiratory burst oxidase, RBOH) activity decreased in *NbG6PDH-P2-silenced* plants. Silencing of the cytosolic NAD kinase NbNADK1, which phosphorylates NADH to NADPH, compromised HR cell death and ROS production. The concomitant silencing of NbG6PDH-P2 reduced HR cell death and ROS down to levels near those of NbG6PDH-P2-silenced plants. These authors suggested that NADPH produced by *NbG6PDH-P2* was responsible for HR cell death and ROS production mediated by RBOH. Increased G6PDH activity could also stimulate ROS generation by regulating NADPH oxidase activity in rice [110]. G6PDH may be involved in maintaining redox homeostasis by regulating the activity of superoxide dismutase (SOD), peroxidase (POD), and ascorbate peroxidase (APX) [111–113]. Treatment with Acibenzolar-S-methyl (an inducer of disease resistance in plants) increased G6PDH activity and the ascorbic acid (AsA) level, while the GSH and NADPH contents and the expression level of redox homeostasis-related genes such as SOD, APX, or dehydroascorbate reductase (DHAR) were reduced [114]. G6PDH activity markedly increased in soybean under drought stress [115]. Upon PEG6000 treatment, the activity of the antioxidant enzymatic machinery (superoxide dismutase (SOD), catalase (CAT), peroxidase (POD), glutathione reductase (GR), dehydroascorbate reductase (DHAR), monodehydroascorbate reductase (MDHAR)) increased in soybean, and GSH and ascorbic acid (AsA) reached elevated levels; once again, these results demonstrate that G6PDH plays a central role in redox homeostasis by maintaining the GSH and Asc levels.

7.2. Saline-Alkaline Stress and Aluminum Toxicity

Saline-alkaline stress is one of the most serious global issues in plant production [116–118]. The metabolic changes triggered by salt stress result in a high need for reductants supplied by the OPPP, consistently with increased total G6PDH activity [119]. *Arabidopsis thaliana* glycogen synthase kinase3 ($ASK\alpha$) regulates stress tolerance by activating G6PDH, which is essential for maintaining the cellular redox balance. Loss of stress-activated $ASK\alpha$ impairs G6PDH activity, increases ROS levels, and enhances sensitivity to salt stress, while $ASK\alpha$ -overexpressing plants exhibit high G6PDH activity, lower ROS levels and are more tolerant to salt stress [33]. In wheat (*Triticum aestivum* L.), G6PDH transcripts were rapidly induced at the early stage of NaCl treatment (almost a 2.2-fold increase), indicating that G6PDH is involved in the initial response of plants to salt stress [120]. In highland barley (*Hordeum vulgare* var. nudum Hook. f.), cy-G6PDH confers resistance to alkaline stress and ultimately improves fresh weight and photosynthetic activity through NADPH production and accumulation of reduced GSH [121]. In line with these results, some researchers cloned five G6PDH genes (*HvG6PDH1* to *HvG6PDH5*) from highland barley and characterized their respective roles in the response to abiotic stresses. The analysis of enzyme activities and gene expression showed that *HvG6PDH1* to *HvG6PDH4* were involved in the responses to salt and drought stresses [122]. Cytosolic *HvG6PDH2* is the major isoform against oxidative stress. *HvG6PDH1* to *HvG6PDH4* and their encoded enzymes responded to jasmonic acid (JA) and ABA treatments, implying that JA and ABA are probably key regulators of HvG6PDHs [122–124]. Some researchers characterized the root behavior of two Arabidopsis single null mutants (*g6pdh5* and *g6pdh6*), one double

mutant (*g6pdh5/6*), and two cy-G6PDH isoforms to salt stress exposure. The seed mutants displayed a reduced germination rate, reduced root elongation, and high accumulation of ROS under salt stress compared to the wild type [48]. Interestingly, the alteration of *G6PDH5* and *G6PDH6* expression affected the activities and transcript levels of various antioxidant enzymes, especially APX and GR. Exogenous application of ascorbic acid and GSH rescued the seed and root phenotypes of *g6pdh5/6*. In response to salt stress, some researchers characterized the nine members of the *G6PDH* gene family (*GmG6PDHs*) in soybean [125]. The activities and transcripts of *GmG6PDHs* were dramatically stimulated, with a notable role of *GmG6PDH2*—a cytosolic isoform. Enzymatic assays of recombinant *GmG6PDH2* proteins expressed in *Escherichia coli* (*E. coli*) showed that this enzyme has functional NADP⁺-dependent G6PDH activity, and *GmG6PDH2*-overexpressing plants exhibited a high degree of resistance to salt stress related to a close coordination of the redox state, the ascorbic acid pool and the GSH pool. The G6PDH activity was enhanced rapidly in the presence of 100mM NaCl in *Phaseolus vulgaris*, which is associated with a raise of G6PDH protein [126]. Application of a G6PDH inhibitor blocked the increase in G6PDH and nitrate reductase activity, as well as NO production. Therefore, G6PDH plays a pivotal role in nitrate-reductase-dependent NO production and in tolerance to salt stress.

Besides salt stress, exposure to high aluminum concentrations significantly induced total and cytosolic G6PDH activities in soybean roots, along with NO accumulation [127]. NADPH produced by NO-modulated cytosolic G6PDH is responsible for ROS accumulation mediated by NADPH oxidase under aluminum stress. Further investigations using pharmacological and transgenic approaches demonstrated that G6PDH positively regulates the activity of NADPH oxidase under aluminum treatment. These results suggest that G6PDH mediates Al-induced programmed cell death through NADPH oxidase-dependent ROS production [128].

7.3. Drought and Heat

Drought can significantly increase the enzymatic activities of cytosolic G6PD (Cyt-G6PD) and plastidial G6PD (P2-G6PD) possibly triggered by NO and H₂O₂ in soybean roots [55]. In winter wheat, *TaG6PDH* (*Triticum aestivum* G6PDH) expression was up-regulated under cold stress and exogenous ABA application, suggesting that *TaG6PDH* positively responds to cold stress and ABA [129]. Similarly, *FaG6PDH* positively regulated cold tolerance in strawberry [130], and in silico bioinformatics analysis of 19 *FaG6PDH* promoters revealed the presence of at least one stress-responsive cis-acting element [131]. An early stress response would involve the OPPP, which represents a true metabolic sensor during the response to various stresses. These results indicate that *G6PDH* might play a pivotal role in redox homeostasis, ROS signaling, and NO cascade signaling. In line with this statement, the activity of G6PDH and antioxidant enzymes (APX, CAT, POD, and SOD) in *Przewalskia tangutica* and tobacco (*Nicotiana tabacum* L.) calluses increased after 40 °C treatment. When G6PDH was partially inhibited by glucosamine pretreatment, the antioxidant enzyme activities, the H₂O₂ content and plasma membrane NADPH oxidase activity decreased, while H₂O₂ application increased the activity of G6PDH and antioxidant enzymes [132]. In *Phaseolus vulgaris*, the heat-sensitive genotype had a higher G6PDH activity than the control under normal temperature [133]. However, under elevated temperature treatment, G6PDH activity increased by approximately 78% in the heat-insensitive genotype but decreased by approximately 37% in the sensitive genotype. These results indicate that G6PDH confers plants heat stress tolerance by regulating H₂O₂ levels under heat stress. Tomato plants grown for 30 days and 45 days without irrigation exhibited 1.67- and 1.32-fold higher total G6PDH activity, respectively [10].

Although all these findings clearly underline the crucial role of G6PDH in plant tolerance to abiotic stress (Figure 4), many efforts should be deployed to provide a comprehensive picture of the molecular regulatory network governing G6PDH regulation in the context of plant responses to harmful conditions. One outstanding question will be to elucidate whether these different abiotic stresses mediate upregulation of G6PDH through

common or specific regulatory pathways and to decipher how G6PDH is regulated in response to multiple abiotic stresses.




Figure 4. Regulatory network of G6PDH in the context of plant responses to abiotic stress. The NADPH provided by cytosolic G6PDH activity is required both for ROS production by NADPH oxidase and for ROS scavenging by activation of GSH ascorbate cycle. The working model proposed that G6PDH plays a pivotal role in redox homeostasis, ROS signaling and NO cascade signaling. ASK α , *Arabidopsis thaliana* glycogen synthase kinase3 (GSK3)/SHAGGY-like kinase; ABA, abscisic acid; APX, ascorbate peroxidase; POD, peroxidase; SOD, superoxide dismutase; GSH, reduced glutathione; ROS, reactive oxygen species.

8. Conclusions

Glucose-6-phosphate dehydrogenases (G6PDHs) are cytosolic or plastidial enzymes that catalyze the oxidation of glucose-6-phosphate (G6P) to 6-phosphogluconolactone. Their activity diverts part of G6P from glycolysis to the oxidative pentose phosphate pathway (OPPP), so that it is key for determining the balance between these two metabolic pathways. As the OPPP is also a critical producer of the reductant component NADPH, G6PDHs play a determining role in ROS scavenging. Besides their role in physiological processes, G6PDHs play a role in plant life cycle and development, especially by influencing seeds (germination) and axillary bud dormancy (plant branching), nitrogen assimilation, responses to abiotic stresses, and by contributing to the recently identified sugar signaling pathway. Further studies are required to (1) understand the respective roles of the different G6PDH isoforms and their regulation, (2) decipher the molecular mechanisms allowing G6PDHs to influence the development and the responses of plants to their environment, and (3) unveil the role of ROS regulation in these processes. An additional interesting research question would be to better understand the interaction between G6PDHs and the molecular regulatory network of hormones and nutrients.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/ijms232416128/s1>.

Author Contributions: Supervision, S.S., G.L. and Y.D.; Writing original draft preparation, Z.J., M.W., G.L., P.G., M.N. and S.S.; software, L.O.; validation, L.C.; data curation, Z.J. and L.O.; review and editing, Z.J., M.W., M.N., L.O., M.-D.P.-G., J.L.G., L.C. and P.G.; funding acquisition, Z.J., G.L., Y.D., P.G. and S.S. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by the National Natural Science Foundation of China (31871573), the Provincial key R&D plan, Modern Agriculture in China (BE2021361), and the China Scholarships Council (no. 202106850042). This work was also supported by the ANR (Agence Nationale de la Recherche) project Labcom, called ESTIM (Evaluation de STIMulateurs de vitalité des plantes; Project ANR-15-LCV3-000).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Williams, J.F.; Blackmore, P.F.; Duke, C.C.; MacLeod, J.K. Fact, uncertainty and speculation concerning the biochemistry of d-erythrose-4-phosphate and its metabolic roles. *Int. J. Biochem.* **1980**, *12*, 339–344. [[CrossRef](#)] [[PubMed](#)]
2. Copeland, L.E.S.; Turner, J.F. The Regulation of Glycolysis and the Pentose Phosphate Pathway. In *Biochemistry of Metabolism*; Academic Press: Cambridge, MA, USA, 1987; pp. 107–128.
3. Esposito, S. Nitrogen assimilation, abiotic stress and glucose 6-phosphate dehydrogenase: The full circle of reductants. *Plants* **2016**, *5*, 24. [[CrossRef](#)] [[PubMed](#)]
4. Hauschild, R.D.; von Schaewen, A. Differential Regulation of Glucose-6-Phosphate Dehydrogenase Isoenzyme Activities in Potato. *Plant Physiol.* **2003**, *133*, 47–62. [[CrossRef](#)] [[PubMed](#)]
5. Masi, A.; Mach, R.L.; Mach-Aigner, A.R. The pentose phosphate pathway in industrially relevant fungi: Crucial insights for bioprocessing. *Appl. Microbiol. Biotechnol.* **2021**, *105*, 4017–4031. [[CrossRef](#)] [[PubMed](#)]
6. Rashida, Z.; Laxman, S. The pentose phosphate pathway and organization of Metabolic Networks Enabling Growth Programs. *Curr. Opin. Syst. Biol.* **2021**, *28*, 100390. [[CrossRef](#)]
7. Graeve, K.; von Schaewen, A.; Scheibe, R. Purification, characterization, and cDNA sequence of glucose-6-phosphate dehydrogenase from potato (*Solanum tuberosum* L.). *Plant J.* **1994**, *5*, 353–361. [[CrossRef](#)]
8. von Schaewen, A.; Langenkamper, G.; Graeve, K.; Wenderoth, I.; Scheibe, R. Molecular characterization of the plastidic glucose-6-phosphate dehydrogenase from potato in comparison to its cytosolic counterpart. *Plant Physiol.* **1995**, *109*, 1327–1335. [[CrossRef](#)]
9. Wendt, U.K.; Hauschild, R.; Lange, C.; Pietersma, M.; Wenderoth, I.; von Schaewen, A. Evidence for functional convergence of redox regulation in G6PDH isoforms of cyanobacteria and higher plants. *Plant Mol. Biol.* **1999**, *40*, 487–494. [[CrossRef](#)]
10. Landi, S.; Nurcato, R.; De Lillo, A.; Lentini, M.; Grillo, S.; Esposito, S. Glucose-6-phosphate dehydrogenase plays a central role in the response of tomato (*Solanum lycopersicum*) plants to short and long-term drought. *Plant Physiol. Biochem.* **2016**, *105*, 79–89. [[CrossRef](#)]
11. Heber, U.; Hallier, U.; Hudson, M.; Groeben, B.; Ernst, R.; Stang, H., II. Lokalisation von Enzymen des reduktiven und dem oxydativen Pentosephosphat-Zyklus in den Chloroplasten und Permeabilität der Chloroplasten-Membran gegenüber Metaboliten. *Z. Für. Nat. B* **1967**, *22*. [[CrossRef](#)]
12. Schnarrenberger, C.; Oeser, A.; Tolbert, N.E. Two isoenzymes each of glucose-6-phosphate dehydrogenase and 6-phosphogluconate dehydrogenase in spinach leaves. *Arch. Biochem. Biophys.* **1973**, *154*, 438–448. [[CrossRef](#)] [[PubMed](#)]
13. Wakao, S.; Benning, C. Genome-wide analysis of glucose-6-phosphate dehydrogenases in *Arabidopsis*. *Plant J.* **2005**, *41*, 243–256. [[CrossRef](#)] [[PubMed](#)]
14. Ruan, M.; He, W.; Sun, H.; Cui, C.; Wang, X.; Li, R.; Wang, X.; Bi, Y. Cytosolic glucose-6-phosphate dehydrogenases play a pivotal role in *Arabidopsis* seed development. *Plant Physiol. Biochem.* **2022**, *186*, 207–219. [[CrossRef](#)] [[PubMed](#)]
15. Wang, H.; Yang, L.; Li, Y.; Hou, J.; Huang, J.; Liang, W. Involvement of ABA-and H₂O₂-dependent cytosolic glucose-6-phosphate dehydrogenase in maintaining redox homeostasis in soybean roots under drought stress. *Plant Physiol. Biochem.* **2016**, *107*, 126–136. [[CrossRef](#)]
16. Landi, S.; Capasso, G.; Esposito, S. Different G6PDH isoforms show specific roles in acclimation to cold stress at various growth stages of barley (*Hordeum vulgare*) and *Arabidopsis thaliana*. *Plant Physiol. Biochem.* **2021**, *169*, 190–202. [[CrossRef](#)]
17. Lendzian, K.; Bassham, J.A. Regulation of glucose-6-phosphate dehydrogenase in spinach chloroplasts by ribulose 1,5-diphosphate and NADPH/NADP⁺ ratios. *Biochim. Biophys. Acta* **1975**, *396*, 260–275. [[CrossRef](#)]

18. Anderson, L.E.; Duggan, J.X. Light modulation of glucose-6-phosphate dehydrogenase: Partial characterization of the light inactivation system and its effects on the properties of the chloroplastic and cytoplasmic forms of the enzyme. *Plant Physiol.* **1976**, *58*, 135–139. [[CrossRef](#)]
19. Kruger, N.J.; von Schaewen, A. The oxidative pentose phosphate pathway: Structure and organisation. *Curr. Opin. Plant Biol.* **2003**, *6*, 236–246. [[CrossRef](#)]
20. Lenzian, K. Modulation of glucose-6-phosphate dehydrogenase by NADPH, NADP⁺ and dithiothreitol at variable NADPH/NADP⁺ ratios in an illuminated reconstituted spinach (*Spinacia oleracea* L.) chloroplast system. *Planta* **1980**, *148*, 1–6. [[CrossRef](#)]
21. Esposito, S.; Carfagna, S.; Massaro, G.; Vona, V.; Di Martino Rigano, V. Glucose-6-phosphate dehydrogenase in barley roots: Kinetic properties and localisation of the isoforms. *Planta* **2001**, *212*, 627–634. [[CrossRef](#)]
22. Ferrara, M.; Guerriero, G.; Cardi, M.; Esposito, S. Purification and biochemical characterisation of a glucose-6-phosphate dehydrogenase from the psychrophilic green alga *Koliella antarctica*. *Extremophiles* **2013**, *17*, 53–62. [[CrossRef](#)] [[PubMed](#)]
23. Née, G.; Aumont-Nicaise, M.; Zaffagnini, M.; Nessler, S.; Valerio-Lepiniec, M.; Issakidis-Bourguet, E. Redox regulation of chloroplastic G6PDH activity by thioredoxin occurs through structural changes modifying substrate accessibility and cofactor binding. *Biochem. J.* **2014**, *457*, 117–125. [[CrossRef](#)] [[PubMed](#)]
24. Esposito, S.; Guerriero, G.; Vona, V.; Di Martino Rigano, V.; Carfagna, S.; Rigano, C. Glutamate synthase activities and protein changes in relation to nitrogen nutrition in barley: The dependence on different plastidic glucose-6P dehydrogenase isoforms. *J. Exp. Bot.* **2005**, *56*, 55–64. [[CrossRef](#)] [[PubMed](#)]
25. Meyer, T.; Hölscher, C.; Schwöppe, C.; von Schaewen, A. Alternative targeting of Arabidopsis plastidic glucose-6-phosphate dehydrogenase G6PD1 involves cysteine-dependent interaction with G6PD4 in the cytosol. *Plant J.* **2011**, *66*, 745–758. [[CrossRef](#)]
26. Hölscher, C.; Meyer, T.; von Schaewen, A. Dual-targeting of Arabidopsis 6-phosphogluconolactonase 3 (PGL3) to chloroplasts and peroxisomes involves interaction with Trx m2 in the cytosol. *Mol. Plant* **2014**, *7*, 252–255. [[CrossRef](#)]
27. Yang, Y.T.; Fu, Z.W.; Su, Y.C.; Zhang, X.; Li, G.Y.; Guo, J.L.; Que, Y.X.; Xu, L.P. A cytosolic glucose-6-phosphate dehydrogenase gene, ScG6PDH, plays a positive role in response to various abiotic stresses in sugarcane. *Sci. Rep.* **2014**, *4*, 7090. [[CrossRef](#)]
28. Lansing, H.; Doering, L.; Fischer, K.; Baune, M.-C.; Schaewen, A.V. Analysis of potential redundancy among Arabidopsis 6-phosphogluconolactonase isoforms in peroxisomes. *J. Exp. Bot.* **2020**, *71*, 823–836. [[CrossRef](#)]
29. Hölscher, C.; Lutterbey, M.-C.; Lansing, H.; Meyer, T.; Fischer, K.; von Schaewen, A. Defects in peroxisomal 6-phosphogluconate dehydrogenase isoform PGD2 prevent gametophytic interaction in *Arabidopsis thaliana*. *Plant Physiol.* **2016**, *171*, 192–205. [[CrossRef](#)]
30. Kaur, N.; Hu, J. Defining the plant peroxisomal proteome: From Arabidopsis to rice. *Front. Plant Sci.* **2011**, *2*, 103. [[CrossRef](#)]
31. Castiglia, D.; Cardi, M.; Landi, S.; Cafasso, D.; Esposito, S. Expression and characterization of a cytosolic glucose 6 phosphate dehydrogenase isoform from barley (*Hordeum vulgare*) roots. *Protein Expr. Purif.* **2015**, *112*, 8–14. [[CrossRef](#)]
32. Wakao, S.; Andre, C.; Benning, C. Functional analyses of cytosolic glucose-6-phosphate dehydrogenases and their contribution to seed oil accumulation in *Arabidopsis*. *Plant Physiol.* **2008**, *146*, 277–288. [[CrossRef](#)] [[PubMed](#)]
33. Dal Santo, S.; Stampfl, H.; Krasensky, J.; Kempa, S.; Gibon, Y.; Petutschnig, E.; Rozhon, W.; Heuck, A.; Clausen, T.; Jonak, C. Stress-induced GSK3 regulates the redox stress response by phosphorylating glucose-6-phosphate dehydrogenase in *Arabidopsis*. *Plant Cell* **2012**, *24*, 3380–3392. [[CrossRef](#)] [[PubMed](#)]
34. Fickenscher, K.; Scheibe, R. Purification and properties of the cytoplasmic glucose-6-phosphate dehydrogenase from pea leaves. *Arch. Biochem. Biophys.* **1986**, *247*, 393–402. [[CrossRef](#)] [[PubMed](#)]
35. Wenderoth, I.; Scheibe, R.; von Schaewen, A. Identification of the cysteine residues involved in redox modification of plant plastidic glucose-6-phosphate dehydrogenase. *J. Biol. Chem.* **1997**, *272*, 26985–26990. [[CrossRef](#)] [[PubMed](#)]
36. Haschek, W.M.; Rousseaux, C.G.; Wallig, M.A.; Bolon, B.; Ochoa, R. *Haschek and Rousseaux's Handbook of Toxicologic Pathology*; Academic Press: Cambridge, MA, USA, 2013.
37. Swamy, P.; Sandhyarani, C. Contribution of the pentose phosphate pathway and glycolytic pathway to dormancy breakage and germination of peanut (*Arachis hypogaea* L.) seeds. *J. Exp. Bot.* **1986**, *37*, 80–88. [[CrossRef](#)]
38. Lacroix, L.; Jaswal, A. Metabolic changes in after-ripening seed of *Prunus cerasus*. *Plant Physiol.* **1967**, *42*, 479–480. [[CrossRef](#)] [[PubMed](#)]
39. Brown, A.; Wray, J. Correlated changes of some enzyme activities and cofactor and substrate contents of pea cotyledon tissue during germination. *Biochem. J.* **1968**, *108*, 437–444. [[CrossRef](#)]
40. Kovacs, M.I.; Simpson, G.M. Dormancy and enzyme levels in seeds of wild oats. *Phytochemistry* **1976**, *15*, 455–458. [[CrossRef](#)]
41. Gosling, P.G.; Ross, J.D. Pentose phosphate metabolism during dormancy breakage in *Corylus avellana* L. *Planta* **1980**, *148*, 362–366. [[CrossRef](#)]
42. Adkins, S.W.; Ross, J.D. Studies in Wild Oat Seed Dormancy: II. Activities of Pentose Phosphate Pathway Dehydrogenases. *Plant Physiol.* **1981**, *68*, 15–17. [[CrossRef](#)]
43. Côme, D.; Corbineau, F. Some aspects of metabolic regulation of seed germination and dormancy. In *Recent Advances in the Development and Germination of Seeds*; Springer: Berlin/Heidelberg, Germany, 1989; pp. 165–179.
44. Ponnaiah, M.; Gilard, F.; Gakière, B.; El-Maarouf-Bouteau, H.; Bailly, C. Regulatory actors and alternative routes for Arabidopsis seed germination are revealed using a pathway-based analysis of transcriptomic datasets. *Plant J.* **2019**, *99*, 163–175. [[CrossRef](#)] [[PubMed](#)]
45. Barba-Espín, G.; Hernández, J.A.; Diaz-Vivancos, P. Role of H₂O₂ in pea seed germination. *Plant Signal Behav.* **2012**, *7*, 193–195. [[CrossRef](#)] [[PubMed](#)]
46. Sharma, S.; Yadav, S.; Sibi, G. Seed germination and maturation under the influence of hydrogen peroxide—A review. *J. Crit. Rev.* **2020**, *7*, 6–10.

47. Leymarie, J.; Vitkauskaitė, G.; Hoang, H.H.; Gendreau, E.; Chazoule, V.; Meimoun, P.; Corbineau, F.; El-Maarouf-Bouteau, H.; Bailly, C. Role of reactive oxygen species in the regulation of *Arabidopsis* seed dormancy. *Plant Cell Physiol.* **2012**, *53*, 96–106. [[CrossRef](#)]
48. Yang, L.; Wang, X.; Chang, N.; Nan, W.; Wang, S.; Ruan, M.; Sun, L.; Li, S.; Bi, Y. Cytosolic Glucose-6-Phosphate Dehydrogenase Is Involved in Seed Germination and Root Growth Under Salinity in *Arabidopsis*. *Front. Plant Sci.* **2019**, *10*, 182. [[CrossRef](#)]
49. Yang, L.; Wang, S.; Sun, L.; Ruan, M.; Li, S.; He, R.; Zhang, W.; Liang, C.; Wang, X.; Bi, Y. Involvement of G6PD5 in ABA response during seed germination and root growth in *Arabidopsis*. *BMC Plant Biol.* **2019**, *19*, 44. [[CrossRef](#)]
50. Oracz, K.; Bouteau, H.E.M.; Farrant, J.M.; Cooper, K.; Belghazi, M.; Job, C.; Job, D.; Corbineau, F.; Bailly, C. ROS production and protein oxidation as a novel mechanism for seed dormancy alleviation. *Plant J.* **2007**, *50*, 452–465. [[CrossRef](#)]
51. El-Maarouf-Bouteau, H.; Meimoun, P.; Job, C.; Job, D.; Bailly, C. Role of protein and mRNA oxidation in seed dormancy and germination. *Front. Plant Sci.* **2013**, *4*, 77. [[CrossRef](#)]
52. Muller, K.; Linkies, A.; Vreeburg, R.A.; Fry, S.C.; Krieger-Liszka, A.; Leubner-Metzger, G. In vivo cell wall loosening by hydroxyl radicals during cress seed germination and elongation growth. *Plant Physiol.* **2009**, *150*, 1855–1865. [[CrossRef](#)]
53. Corbineau, F.; Xia, Q.; Bailly, C.; El-Maarouf-Bouteau, H. Ethylene, a key factor in the regulation of seed dormancy. *Front. Plant Sci.* **2014**, *5*, 539. [[CrossRef](#)]
54. Liu, Y.; Ye, N.; Liu, R.; Chen, M.; Zhang, J. H₂O₂ mediates the regulation of ABA catabolism and GA biosynthesis in *Arabidopsis* seed dormancy and germination. *J. Exp. Bot.* **2010**, *61*, 2979–2990. [[CrossRef](#)] [[PubMed](#)]
55. Wang, X.; Ruan, M.; Wan, Q.; He, W.; Yang, L.; Liu, X.; He, L.; Yan, L.; Bi, Y. Nitric oxide and hydrogen peroxide increase glucose-6-phosphate dehydrogenase activities and expression upon drought stress in soybean roots. *Plant Cell Rep.* **2020**, *39*, 63–73. [[CrossRef](#)] [[PubMed](#)]
56. Kobrehel, K.; Wong, J.H.; Balogh, A.; Kiss, F.; Yee, B.C.; Buchanan, B.B. Specific reduction of wheat storage proteins by thioredoxin h. *Plant Physiol.* **1992**, *99*, 919–924. [[CrossRef](#)] [[PubMed](#)]
57. Yano, H.; Wong, J.H.; Lee, Y.M.; Cho, M.-J.; Buchanan, B.B. A strategy for the identification of proteins targeted by thioredoxin. *Proc. Natl. Acad. Sci. USA* **2001**, *98*, 4794–4799. [[CrossRef](#)] [[PubMed](#)]
58. Marx, C.; Wong, J.H.; Buchanan, B.B. Thioredoxin and germinating barley: Targets and protein redox changes. *Planta* **2003**, *216*, 454–460. [[CrossRef](#)] [[PubMed](#)]
59. Alkhalfioui, F.; Renard, M.; Vensel, W.H.; Wong, J.; Tanaka, C.K.; Hurkman, W.J.; Buchanan, B.B.; Montrichard, F. Thioredoxin-linked proteins are reduced during germination of *Medicago truncatula* seeds. *Plant Physiol.* **2007**, *144*, 1559–1579. [[CrossRef](#)]
60. Montrichard, F.; Alkhalfioui, F.; Yano, H.; Vensel, W.H.; Hurkman, W.J.; Buchanan, B.B. Thioredoxin targets in plants: The first 30 years. *J. Proteom.* **2009**, *72*, 452–474. [[CrossRef](#)]
61. Lozano, R.M.; Wong, J.H.; Yee, B.C.; Peters, A.; Kobrehel, K.; Buchanan, B.B. New evidence for a role for thioredoxin h in germination and seedling development. *Planta* **1996**, *200*, 100–106. [[CrossRef](#)]
62. Née, G.; Zaffagnini, M.; Trost, P.; Issakidis-Bourguet, E. Redox regulation of chloroplastic glucose-6-phosphate dehydrogenase: A new role for f-type thioredoxin. *FEBS Lett.* **2009**, *583*, 2827–2832. [[CrossRef](#)]
63. Nee, G.; Chatel-Innocenti, G.; Meimoun, P.; Leymarie, J.; Montrichard, F.; Satour, P.; Bailly, C.; Issakidis-Bourguet, E. A New Role for Plastid Thioredoxins in Seed Physiology in Relation to Hormone Regulation. *Int. J. Mol. Sci.* **2021**, *22*, 10395. [[CrossRef](#)]
64. Fernandes, M.S.; Rossiello, R.O.P. Mineral nitrogen in plant physiology and plant nutrition. *Crit. Rev. Plant Sci.* **1995**, *14*, 111–148. [[CrossRef](#)]
65. Luo, L.; Zhang, Y.; Xu, G. How does nitrogen shape plant architecture? *J. Exp. Bot.* **2020**, *71*, 4415–4427. [[CrossRef](#)] [[PubMed](#)]
66. Zhang, Z.; Hu, B.; Chu, C. Towards understanding the hierarchical nitrogen signalling network in plants. *Curr. Opin. Plant Biol.* **2020**, *55*, 60–65. [[CrossRef](#)] [[PubMed](#)]
67. Coruzzi, G.M. Primary N-assimilation into amino acids in *Arabidopsis*. In *The Arabidopsis Book*; American Society of Plant Biologists: Portland, OR, USA, 2003; Volume 2.
68. Bussell, J.D.; Keech, O.; Fenske, R.; Smith, S.M. Requirement for the plastidial oxidative pentose phosphate pathway for nitrate assimilation in *Arabidopsis*. *Plant J.* **2013**, *75*, 578–591. [[CrossRef](#)] [[PubMed](#)]
69. Frungillo, L.; Skelly, M.J.; Loake, G.J.; Spoel, S.H.; Salgado, I. S-nitrosothiols regulate nitric oxide production and storage in plants through the nitrogen assimilation pathway. *Nat. Commun.* **2014**, *5*, 5401. [[CrossRef](#)]
70. Bowsher, C.; Boulton, E.; Rose, J.; Nayagam, S.; Emes, M. Reductant for glutamate synthase in generated by the oxidative pentose phosphate pathway in non-photosynthetic root plastids. *Annu. Rev. Plant Biol.* **1992**, *2*, 893–898. [[CrossRef](#)]
71. Esposito, S.; Massaro, G.; Vona, V.; Di Martino Rigano, V.; Carfagna, S. Glutamate synthesis in barley roots: The role of the plastidic glucose-6-phosphate dehydrogenase. *Planta* **2003**, *216*, 639–647. [[CrossRef](#)]
72. Redinbaugh, M.G.; Campbell, W.H. Nitrate regulation of the oxidative pentose phosphate pathway in maize (*Zea mays* L.) root plastids: Induction of 6-phosphogluconate dehydrogenase activity, protein and transcript levels. *Plant Sci.* **1998**, *134*, 129–140. [[CrossRef](#)]
73. Wang, R.; Okamoto, M.; Xing, X.; Crawford, N.M. Microarray analysis of the nitrate response in *Arabidopsis* roots and shoots reveals over 1,000 rapidly responding genes and new linkages to glucose, trehalose-6-phosphate, iron, and sulfate metabolism. *Plant Physiol.* **2003**, *132*, 556–567. [[CrossRef](#)]
74. Wang, R.; Tischner, R.; Gutiérrez, R.A.; Hoffman, M.; Xing, X.; Chen, M.; Coruzzi, G.; Crawford, N.M. Genomic analysis of the nitrate response using a nitrate reductase-null mutant of *Arabidopsis*. *Plant Physiol.* **2004**, *136*, 2512–2522. [[CrossRef](#)]

75. Lejay, L.; Wirth, J.; Pervent, M.; Cross, J.M.-F.; Tillard, P.; Gojon, A. Oxidative Pentose Phosphate Pathway-Dependent Sugar Sensing as a Mechanism for Regulation of Root Ion Transporters by Photosynthesis. *Plant Physiol.* **2008**, *146*, 2036–2053. [[CrossRef](#)] [[PubMed](#)]
76. De Jong, M.; George, G.; Ongaro, V.; Williamson, L.; Willetts, B.; Ljung, K.; McCulloch, H.; Leyser, O. Auxin and strigolactone signaling are required for modulation of *Arabidopsis* shoot branching by nitrogen supply. *Plant Physiol.* **2014**, *166*, 384–395. [[CrossRef](#)] [[PubMed](#)]
77. Kohler, E.; Barrach, H.; Neubert, D. Inhibition of NADP dependent oxidoreductases by the 6-aminonicotinamide analogue of NADP. *FEBS Lett.* **1970**, *6*, 225–228. [[CrossRef](#)] [[PubMed](#)]
78. Garlick, A.P.; Moore, C.; Kruger, N.J. Monitoring flux through the oxidative pentose phosphate pathway using [1-14C] gluconate. *Planta* **2002**, *216*, 265–272. [[CrossRef](#)]
79. Chaput, V.; Martin, A.; Lejay, L. Redox metabolism: The hidden player in carbon and nitrogen signaling? *J. Exp. Bot.* **2020**, *71*, 3816–3826. [[CrossRef](#)]
80. Neuhaus, H.; Emes, M. Nonphotosynthetic metabolism in plastids. *Annu. Rev. Plant Biol.* **2000**, *51*, 111. [[CrossRef](#)]
81. Kumar, V.; Mills, D.J.; Anderson, J.D.; Mattoo, A.K. An alternative agriculture system is defined by a distinct expression profile of select gene transcripts and proteins. *Proc. Natl. Acad. Sci. USA* **2004**, *101*, 10535–10540. [[CrossRef](#)]
82. Schneider, A.; Godin, C.; Boudon, F.; Demotes-Mainard, S.; Sakr, S.; Bertheloot, J. Light regulation of axillary bud outgrowth along plant axes: An overview of the roles of sugars and hormones. *Front. Plant Sci.* **2019**, *10*, 1296. [[CrossRef](#)]
83. Barbier, F.F.; Cao, D.; Fichtner, F.; Weiste, C.; Pérez-García, M.-D.; Caradeuc, M.; Le Gourrierc, J.; Sakr, S.; Beveridge, C.A. HEXOKINASE1 signalling promotes shoot branching and interacts with cytokinin and strigolactone pathways. *New Phytol.* **2021**, *231*, 1088–1104. [[CrossRef](#)]
84. Tao, Y.; An, L.; Xiao, F.; Li, G.; Ding, Y.; Paul, M.; Liu, Z.H. Integration of embryo–endosperm interaction into a holistic and dynamic picture of seed development using a rice mutant with notched-belly kernels. *Crop J.* **2021**, *10*, 729–742. [[CrossRef](#)]
85. Rabot, A.; Henry, C.; Ben Baaziz, K.; Mortreau, E.; Azri, W.; Lothier, J.; Hamama, L.; Boummaza, R.; Leduc, N.; Pelleschi-Travier, S. Insight into the role of sugars in bud burst under light in the rose. *Plant Cell Physiol.* **2012**, *53*, 1068–1082. [[CrossRef](#)]
86. Barbier, F.; Péron, T.; Lecerf, M.; Perez-Garcia, M.-D.; Barrière, Q.; Rolčík, J.; Boutet-Mercey, S.; Citerne, S.; Lemoine, R.; Porcheron, B.; et al. Sucrose is an early modulator of the key hormonal mechanisms controlling bud outgrowth in *Rosa hybrida*. *J. Exp. Bot.* **2015**, *66*, 2569–2582. [[CrossRef](#)] [[PubMed](#)]
87. Rabot, A.; Portemer, V.; Péron, T.; Mortreau, E.; Leduc, N.; Hamama, L.; Coutos-Thévenot, P.; Atanassova, R.; Sakr, S.; Le Gourrierc, J. Interplay of sugar, light and gibberellins in expression of *Rosa hybrida* vacuolar invertase 1 regulation. *Plant Cell Physiol.* **2014**, *55*, 1734–1748. [[CrossRef](#)] [[PubMed](#)]
88. Demotes-Mainard, S.; Huché-Thélier, L.; Morel, P.; Boumaza, R.; Guérin, V.; Sakr, S. Temporary water restriction or light intensity limitation promotes branching in rose bush. *Sci. Hortic.* **2013**, *150*, 432–440. [[CrossRef](#)]
89. Crespel, L.; Le Bras, C.; Amoroso, T.; Unda Ulloa, M.G.; Morel, P.; Sakr, S. Genotype × Light quality interaction on rose architecture. *Agronomy* **2020**, *10*, 913. [[CrossRef](#)]
90. Wang, M.; Le Gourrierc, J.; Jiao, F.; Demotes-Mainard, S.; Perez-Garcia, M.-D.; Ogé, L.; Hamama, L.; Crespel, L.; Bertheloot, J.; Chen, J. Convergence and divergence of sugar and cytokinin signaling in plant development. *Int. J. Mol. Sci.* **2021**, *22*, 1282. [[CrossRef](#)]
91. Wang, M.; Ogé, L.; Pérez-García, M.D.; Launay-Avon, A.; Clément, G.; Le Gourrierc, J.; Hamama, L.; Sakr, S. Antagonistic Effect of Sucrose Availability and Auxin on Rosa Axillary Bud Metabolism and Signaling, Based on the Transcriptomics and Metabolomics Analysis. *Front. Plant Sci.* **2022**, *13*, 830840. [[CrossRef](#)]
92. Djennane, S.; Hibrand-Saint Oyant, L.; Kawamura, K.; Lalanne, D.; Laffaire, M.; Thouroude, T.; Chalain, S.; Sakr, S.; Boumaza, R.; Foucher, F.; et al. Impacts of light and temperature on shoot branching gradient and expression of strigolactone synthesis and signalling genes in rose. *Plant Cell Environ.* **2014**, *37*, 742–757. [[CrossRef](#)]
93. Fichtner, F.; Barbier, F.F.; Annunziata, M.G.; Feil, R.; Olas, J.J.; Mueller-Roeber, B.; Stitt, M.; Beveridge, C.A.; Lunn, J.E. Regulation of shoot branching in arabidopsis by trehalose 6-phosphate. *New Phytol.* **2021**, *229*, 2135–2151. [[CrossRef](#)]
94. Wang, M.; Pérez-García, M.-D.; Davière, J.-M.; Barbier, F.; Ogé, L.; Gentilhomme, J.; Voisine, L.; Péron, T.; Launay-Avon, A.; Clément, G.; et al. Outgrowth of the axillary bud in rose is controlled by sugar metabolism and signalling. *J. Exp. Bot.* **2021**, *72*, 3044–3060. [[CrossRef](#)]
95. Porcher, A.; Guérin, V.; Montrichard, F.; Lebrec, A.; Lothier, J.; Vian, A. Ascorbate glutathione-dependent H₂O₂ scavenging is an important process in axillary bud outgrowth in rosebush. *Ann. Bot.* **2020**, *126*, 1049–1062. [[CrossRef](#)] [[PubMed](#)]
96. Porcher, A.; Guérin, V.; Leduc, N.; Lebrec, A.; Lothier, J.; Vian, A. Ascorbate–glutathione pathways mediated by cytokinin regulate H₂O₂ levels in light-controlled rose bud burst. *Plant Physiol.* **2021**, *186*, 910–928. [[CrossRef](#)] [[PubMed](#)]
97. Wang, M.; Ogé, L.; Voisine, L.; Pérez-García, M.-D.; Jeauffre, J.; Hibrand Saint-Oyant, L.; Grappin, P.; Hamama, L.; Sakr, S. Posttranscriptional Regulation of RhBRC1 (*Rosa hybrida* BRANCHED1) in Response to Sugars is Mediated via its Own 3′ Untranslated Region, with a Potential Role of RhPUF4 (Pumilio RNA-Binding Protein Family). *Int. J. Mol. Sci.* **2019**, *20*, 3808. [[CrossRef](#)] [[PubMed](#)]
98. Aguilar-Martínez, J.A.; Poza-Carrión, C.; Cubas, P. Arabidopsis BRANCHED1 acts as an integrator of branching signals within axillary buds. *Plant Cell* **2007**, *19*, 458–472. [[CrossRef](#)] [[PubMed](#)]
99. Li, L.; Sheen, J. Dynamic and diverse sugar signaling. *Curr. Opin. Plant Biol.* **2016**, *33*, 116–125. [[CrossRef](#)] [[PubMed](#)]

100. Jiang, Z.; Chen, Q.; Chen, L.; Liu, D.; Yang, H.; Xu, C.; Hong, J.; Li, J.; Ding, Y.; Sakr, S.; et al. Sink Strength Promoting Remobilization of Non-Structural Carbohydrates by Activating Sugar Signaling in Rice Stem during Grain Filling. *Int. J. Mol. Sci.* **2022**, *23*, 4864. [[CrossRef](#)]
101. Jiang, Z.; Chen, Q.; Chen, L.; Yang, H.; Zhu, M.; Ding, Y.; Li, W.; Liu, Z.; Jiang, Y.; Li, G. Efficiency of Sucrose to Starch Metabolism Is Related to the Initiation of Inferior Grain Filling in Large Panicle Rice. *Front. Plant Sci.* **2021**, *12*, 732867. [[CrossRef](#)]
102. Sakr, S.; Wang, M.; Dédaldéchamp, F.; Pérez-García, M.-D.; Ogé, L.; Hamama, L.; Atanassova, R. The Sugar-Signaling Hub: Overview of Regulators and Interaction with the Hormonal and Metabolic Network. *Int. J. Mol. Sci.* **2018**, *19*, 2506. [[CrossRef](#)]
103. Moore, B.; Zhou, L.; Rolland, F.; Hall, Q.; Cheng, W.H.; Liu, Y.X.; Hwang, I.; Jones, T.; Sheen, J. Role of the Arabidopsis glucose sensor HXK1 in nutrient, light, and hormonal signaling. *Science* **2003**, *300*, 332–336. [[CrossRef](#)]
104. Bellegarde, F.; Maghiaoui, A.; Boucherez, J.; Krouk, G.; Lejay, L.; Bach, L.; Gojon, A.; Martin, A. The Chromatin Factor HNI9 and ELONGATED HYPOCOTYL5 Maintain ROS Homeostasis under High Nitrogen Provision. *Plant Physiol.* **2019**, *180*, 582–592. [[CrossRef](#)]
105. Zhu, J.-K. Abiotic Stress Signaling and Responses in Plants. *Cell* **2016**, *167*, 313–324. [[CrossRef](#)] [[PubMed](#)]
106. Wendt, U.K.; Wenderoth, I.; Tegeler, A.; Von Schaewen, A. Molecular characterization of a novel glucose-6-phosphate dehydrogenase from potato (*Solanum tuberosum* L.). *Plant J.* **2000**, *23*, 723–733. [[CrossRef](#)] [[PubMed](#)]
107. De Freitas-Silva, L.; Rodriguez-Ruiz, M.; Houmani, H.; da Silva, L.C.; Palma, J.M.; Corpas, F.J. Glyphosate-induced oxidative stress in *Arabidopsis thaliana* affecting peroxisomal metabolism and triggers activity in the oxidative phase of the pentose phosphate pathway (OxPPP) involved in NADPH generation. *J. Plant Physiol.* **2017**, *218*, 196–205. [[CrossRef](#)] [[PubMed](#)]
108. Huang, H.; Liao, J.; Zheng, X.; Chen, Y.; Ren, H. Low-level free nitrous acid efficiently inhibits the conjugative transfer of antibiotic resistance by altering intracellular ions and disabling transfer apparatus. *Water Res.* **2019**, *158*, 383–391. [[CrossRef](#)] [[PubMed](#)]
109. Asai, S.; Yoshioka, M.; Nomura, H.; Tone, C.; Nakajima, K.; Nakane, E.; Doke, N.; Yoshioka, H. A plastidic glucose-6-phosphate dehydrogenase is responsible for hypersensitive response cell death and reactive oxygen species production. *J. Gen. Plant Pathol.* **2011**, *77*, 152–162. [[CrossRef](#)]
110. Kano, A.; Fukumoto, T.; Ohtani, K.; Yoshihara, A.; Ohara, T.; Tajima, S.; Izumori, K.; Tanaka, K.; Ohkouchi, T.; Ishida, Y.; et al. The rare sugar d-allose acts as a triggering molecule of rice defence via ROS generation. *J. Exp. Bot.* **2013**, *64*, 4939–4951. [[CrossRef](#)] [[PubMed](#)]
111. Zaka, R.; Vandecasteele, C.M.; Misset, M.T. Effects of low chronic doses of ionizing radiation on antioxidant enzymes and G6PDH activities in *Stipa capillata* (Poaceae). *J. Exp. Bot.* **2002**, *53*, 1979–1987. [[CrossRef](#)]
112. Lin, Y.; Lin, S.; Guo, H.; Zhang, Z.; Chen, X. Functional analysis of PsG6PDH, a cytosolic glucose-6-phosphate dehydrogenase gene from *Populus suaveolens*, and its contribution to cold tolerance improvement in tobacco plants. *Biotechnol. Lett.* **2013**, *35*, 1509–1518. [[CrossRef](#)]
113. Li, C.; Wei, M.; Ge, Y.; Zhao, J.; Chen, Y.; Hou, J.; Cheng, Y.; Chen, J.; Li, J. The role of glucose-6-phosphate dehydrogenase in reactive oxygen species metabolism in apple exocarp induced by acibenzolar-S-methyl. *Food Chem.* **2020**, *308*, 125663. [[CrossRef](#)]
114. Wei, M.; Ge, Y.; Li, C.; Han, X.; Qin, S.; Chen, Y.; Tang, Q.; Li, J. G6PDH regulated NADPH production and reactive oxygen species metabolism to enhance disease resistance against blue mold in apple fruit by acibenzolar-S-methyl. *Postharvest Biol. Technol.* **2019**, *148*, 228–235. [[CrossRef](#)]
115. Liu, J.; Wang, X.; Hu, Y.; Hu, W.; Bi, Y. Glucose-6-phosphate dehydrogenase plays a pivotal role in tolerance to drought stress in soybean roots. *Plant Cell Rep.* **2013**, *32*, 415–429. [[CrossRef](#)] [[PubMed](#)]
116. Wang, S.; Li, X.; Liu, W.; Li, P.; Kong, L.; Ren, W.; Wu, H.; Tu, Y. Degradation of pyrene by immobilized microorganisms in saline-alkaline soil. *J. Environ. Sci.* **2012**, *24*, 1662–1669. [[CrossRef](#)]
117. Deinlein, U.; Stephan, A.B.; Horie, T.; Luo, W.; Xu, G.; Schroeder, J.I. Plant salt-tolerance mechanisms. *Trends Plant Sci.* **2014**, *19*, 371–379. [[CrossRef](#)]
118. Chuamnakthong, S.; Nampei, M.; Ueda, A. Characterization of Na⁺ exclusion mechanism in rice under saline-alkaline stress conditions. *Plant Sci.* **2019**, *287*, 110171. [[CrossRef](#)] [[PubMed](#)]
119. Cardi, M.; Castiglia, D.; Ferrara, M.; Guerriero, G.; Chiurazzi, M.; Esposito, S. The effects of salt stress cause a diversion of basal metabolism in barley roots: Possible different roles for glucose-6-phosphate dehydrogenase isoforms. *Plant Physiol. Biochem.* **2015**, *86*, 44–54. [[CrossRef](#)] [[PubMed](#)]
120. Nemoto, Y.; Sasakuma, T. Specific expression of glucose-6-phosphate dehydrogenase (G6PDH) gene by salt stress in wheat (*Triticum aestivum* L.). *Plant Sci.* **2000**, *158*, 53–60. [[CrossRef](#)] [[PubMed](#)]
121. He, Q.; Li, P.; Zhang, W.; Bi, Y. Cytosolic glucose-6-phosphate dehydrogenase plays an important role in the silicon-enhanced alkaline tolerance in highland barley. *Funct. Plant Biol.* **2021**, *48*, 119–130. [[CrossRef](#)]
122. Feng, R.; Wang, X.; He, L.; Wang, S.; Li, J.; Jin, J.; Bi, Y. Identification, Characterization, and Stress Responsiveness of Glucose-6-phosphate Dehydrogenase Genes in Highland Barley. *Plants* **2020**, *9*, 1800. [[CrossRef](#)]
123. Cardi, M.; Chibani, K.; Cafasso, D.; Rouhier, N.; Jacquot, J.-P.; Esposito, S. Abscisic acid effects on activity and expression of barley (*Hordeum vulgare*) plastidial glucose-6-phosphate dehydrogenase. *J. Exp. Bot.* **2011**, *62*, 4013–4023. [[CrossRef](#)]
124. Hu, Y.; You, J.; Li, J.; Wang, C. Loss of cytosolic glucose-6-phosphate dehydrogenase increases the susceptibility of *Arabidopsis thaliana* to root-knot nematode infection. *Ann. Bot.* **2018**, *123*, 37–46. [[CrossRef](#)]
125. Zhao, Y.; Cui, Y.; Huang, S.; Yu, J.; Wang, X.; Xin, D.; Li, X.; Liu, Y.; Dai, Y.; Qi, Z.; et al. Genome-Wide Analysis of the Glucose-6-Phosphate Dehydrogenase Family in Soybean and Functional Identification of GmG6PDH2 Involvement in Salt Stress. *Front. Plant Sci.* **2020**, *11*, 214. [[CrossRef](#)] [[PubMed](#)]

126. Liu, Y.; Wu, R.; Wan, Q.; Xie, G.; Bi, Y. Glucose-6-Phosphate Dehydrogenase Plays a Pivotal Role in Nitric Oxide-Involved Defense Against Oxidative Stress Under Salt Stress in Red Kidney Bean Roots. *Plant Cell Physiol.* **2007**, *48*, 511–522. [[CrossRef](#)] [[PubMed](#)]
127. Wang, H.; Hou, J.; Li, Y.; Zhang, Y.; Huang, J.; Liang, W. Nitric oxide-mediated cytosolic glucose-6-phosphate dehydrogenase is involved in aluminum toxicity of soybean under high aluminum concentration. *Plant Soil* **2017**, *416*, 39–52. [[CrossRef](#)]
128. Huang, J.; Han, R.; Ji, F.; Yu, Y.; Wang, R.; Hai, Z.; Liang, W.; Wang, H. Glucose-6-phosphate dehydrogenase and abscisic acid mediate programmed cell death induced by aluminum toxicity in soybean root tips. *J. Hazard. Mater.* **2022**, *425*, 127964. [[CrossRef](#)]
129. Tian, Y.; Peng, K.; Bao, Y.; Zhang, D.; Meng, J.; Wang, D.; Wang, X.; Cang, J. Glucose-6-phosphate dehydrogenase and 6-phosphogluconate dehydrogenase genes of winter wheat enhance the cold tolerance of transgenic *Arabidopsis*. *Plant Physiol. Biochem.* **2021**, *161*, 86–97. [[CrossRef](#)]
130. Zhang, Y.; Luo, M.; Cheng, L.; Lin, Y.; Chen, Q.; Sun, B.; Gu, X.; Wang, Y.; Li, M.; Luo, Y.; et al. Identification of the Cytosolic Glucose-6-Phosphate Dehydrogenase Gene from Strawberry Involved in Cold Stress Response. *Int. J. Mol. Sci.* **2020**, *21*, 7322. [[CrossRef](#)]
131. Lei, D.; Lin, Y.; Luo, M.; Zhao, B.; Tang, H.; Zhou, X.; Yao, W.; Zhang, Y.; Wang, Y.; Li, M.; et al. Genome-Wide Investigation of G6PDH Gene in Strawberry: Evolution and Expression Analysis during Development and Stress. *Int. J. Mol. Sci.* **2022**, *23*, 4728. [[CrossRef](#)]
132. Gong, H.; Chen, G.; Li, F.; Wang, X.; Hu, Y.; Bi, Y. Involvement of G6PDH in heat stress tolerance in the calli from *Przewalskia tangutica* and *Nicotiana tabacum*. *Biol. Plant.* **2012**, *56*, 422–430. [[CrossRef](#)]
133. Santiago, J.P.; Soltani, A.; Bresson, M.M.; Preiser, A.L.; Lowry, D.B.; Sharkey, T.D. Contrasting anther glucose-6-phosphate dehydrogenase activities between two bean varieties suggest an important role in reproductive heat tolerance. *Plant Cell Environ.* **2021**, *44*, 2185–2199. [[CrossRef](#)]