



HAL
open science

Effects of Electromagnetic Radiation on Neuropeptide Transcript Levels in the Synganglion of *Ixodes ricinus*

Lívia Šofrankova, Miroslav Baňas, Natália Pipova, Igor Majlá, Juraj Kurimský, Rom Cimbala, Marek Pavlík, Lourdes Mateos-Hernández, Ladislav Šimo, Viktória Majláthová

► **To cite this version:**

Lívia Šofrankova, Miroslav Baňas, Natália Pipova, Igor Majlá, Juraj Kurimský, et al.. Effects of Electromagnetic Radiation on Neuropeptide Transcript Levels in the Synganglion of *Ixodes ricinus*. Pathogens, 2023, 12 (12), pp.1398. 10.3390/pathogens12121398 . hal-04311327

HAL Id: hal-04311327

<https://hal.inrae.fr/hal-04311327>

Submitted on 28 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Article

Effects of Electromagnetic Radiation on Neuropeptide Transcript Levels in the Synganglion of *Ixodes ricinus*

Lívia Šofranková ¹, Miroslav Bañas ¹, Natália Pipová ¹, Igor Majláth ¹, Juraj Kurimský ², Roman Cimbala ², Marek Pavlík ², Lourdes Mateos-Hernández ³, Ladislav Šimo ³ and Viktória Majláthová ^{1,*}

¹ Department of Animal Physiology, Pavol Jozef Šafárik University in Košice, Šrobárova 2, 04180 Košice, Slovakia; livia.sofrankova@student.upjs.sk (L.Š.); miroslav.banas@student.upjs.sk (M.B.); natalia.kokosova@upjs.sk (N.P.); igor.majlath@upjs.sk (I.M.)

² Department of Electrical Power Engineering, Faculty of Electrical Engineering and Informatics, Technical University of Košice, Mäsiarska 74, 04120 Košice, Slovakia; juraj.kurimsky@tuke.sk (J.K.); roman.cimbala@tuke.sk (R.C.); marek.pavlik@tuke.sk (M.P.)

³ Laboratoire de Santé Animale, Unité Mixte de Recherche de Biologie Moléculaire et d'Immunologie Parasitaires (UMR BIPAR), Ecole Nationale Vétérinaire d'Alfort, INRAE, ANSES, F-94700 Maisons-Alfort, France; lourdes.mateos@vet-alfort.fr (L.M.-H.); ladislav.simo@vet-alfort.fr (L.Š.)

* Correspondence: viktoria.majlathova@upjs.sk

Abstract: Anthropogenic electromagnetic radiation is an important environmental factor affecting the functionality of biological systems. Sensitivity to various frequencies of electromagnetic radiation has been detected in ixodid ticks in the past. However, the physiological aspects of radiation effects have not yet been studied in ticks. In the presented experiment, 360 *Ixodes ricinus* ticks, 180 males and 180 females, were divided into 16 irradiated and 8 control groups. The irradiated groups were exposed to two different intensities of electromagnetic radiation with a frequency of 900 MHz at different lengths of exposure time. RT-PCR was utilized to determine the changes in mRNA levels in tick synganglia after irradiation. Four randomly selected neuropeptide genes were tested—allatotropin (*at*), FGLA-related allatostatins (*fgla/ast*), kinin, and arginine-vasopressin-like peptide (*avpl*). A significant decrease in transcript levels in all female groups exposed to higher intensity radiofrequency radiation for 1 to 3 h was found. After one hour of radiofrequency exposure, a significant downregulation in allatotropin expression in males was detected. A consistent downregulation of the *at* gene was detected in males irradiated with a higher intensity. Unfortunately, the specific functions of the studied neuropeptides in ticks are not known yet, so a more comprehensive study is necessary to describe the effects of EMF on observed neuropeptides. This study represents the first report on the effects of the abiotic environment on tick neurophysiology.

Keywords: ticks; neuropeptides; transcript levels; *Ixodes ricinus*



Citation: Šofranková, L.; Bañas, M.; Pipová, N.; Majláth, I.; Kurimský, J.; Cimbala, R.; Pavlík, M.; Mateos-Hernández, L.; Šimo, L.; Majláthová, V. Effects of Electromagnetic Radiation on Neuropeptide Transcript Levels in the Synganglion of *Ixodes ricinus*. *Pathogens* **2023**, *12*, 1398. <https://doi.org/10.3390/pathogens12121398>

Academic Editor: Ricardo Pérez-Sánchez

Received: 21 September 2023
Revised: 6 November 2023
Accepted: 21 November 2023
Published: 28 November 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Natural and anthropogenic electromagnetic fields (EMFs) are an important environmental factor affecting all living organisms. Among those, anthropogenic electromagnetic field, commonly labeled as electromagnetic smog, is a significant environmental pollutant affecting a wide range of cell and live system functions. Studies of the effects of EMFs and electromagnetic perception mechanisms have been conducted on in vitro cultures and in several invertebrate and vertebrate laboratory models [1–8]. It has been well documented that electromagnetic radiation affects the structure of the lipid bilayer, the polarity of membranes, and the sensitivity of ion channels [9–14]. The presence of EMFs can also cause the disturbance of the cell's electrochemical equilibrium, which subsequently induces the activation of the stress response [15]. Furthermore, electromagnetic fields influence the cell at the molecular level, which involves, for example, the gene expression and subsequent proteosynthesis. Among those, the upregulation of heat shock proteins (HSP) and

proteins involved in the immune response, cell proliferation, and cell apoptosis has been demonstrated [16–22]. In *Drosophila melanogaster*, radiation influenced the levels of biogenic amines in the central nervous system [23].

Pioneer studies describing the influence of electromagnetic radiation on ticks were published by Korotkov et al. (1996, 2000). Here the authors proved that microwave radiation exposure affected the hatching time, activity, and survival of *Hyalomma asiaticum* tick larvae. The experimental temperature during radiation exposure appeared to be an important factor of EMF influence in this study [24,25]. In addition, one-hour radiofrequency irradiation of *Dermacentor reticulatus* eggs resulted in the hatching of larvae with larger body dimensions in later studies [26]. The most recent studies investigating EMF-mediated changes in tick behavior have clearly demonstrated the ability of ticks to perceive electromagnetic fields and distinguish their different properties [27,28]. Interestingly, the presence of bacteria (*Rickettsia* spp.) in ticks was associated with enhancing the response of infected ticks to radiation [29]. The physiological basis of the behavioral responses observed in irradiated ticks has not been explained. However, as it is most likely mediated by the central nervous system, the levels of neurotransmitters and neuropeptides could have a role in the regulation of the behavior observed under irradiation.

Neuropeptides are small protein molecules acting as crucial regulators of arthropod physiology and behavior [30]. They are well known to be expressed in specific neuronal or neuroendocrine cells regulating a myriad of physiological processes, e.g., feeding, development, reproduction, homeostasis, growth, digestion, diuresis, sleeping, or stress. In addition, these signaling molecules are well known for acting as neuromodulators in the circuits of the central nervous system [31]. As of today, in the tick central nervous system, the synganglion, 38 different neuropeptide genes have been identified [32,33]. Among those, multiple have been immunolocalized in distinct neuronal cells within the synganglion as well as in their axons reaching several visceral organs [34–37]. Although the knowledge regarding the characterization and localization of tick neuropeptides along their receptors has accelerated in the last 15 years, most of their functions remain obscured.

In the presented study, we investigated the influence of different 900 MHz EMF intensities on selected neuropeptide transcripts—allatotropin (*at*), FGLa-related allatostatins (*fgla/ast*), kinin, and arginine-vasopressin-like peptide (*avpl*) in adult-stage male and female *I. ricinus* synganglia.

2. Materials and Methods

2.1. Experimental Ticks

Altogether, 360 adult-stage sheep ticks (*Ixodes ricinus*), 180 males and 180 females, were used. Ticks were collected via flagging near the village Dubovica, northeastern Slovakia (GPS coordinates 49°07'3.7" N 20°57'37.2" E). Ticks were collected in the spring (April–May) of 2022 on a meadow, its slope oriented towards the northeast. Ticks were collected in the shrubbery with many signs of animal tracks, located on the edge of a brook running along the perimeter of the meadow (Figure 1). The data on the intensity of electromagnetic fields were not available for Dubovica; there was a radio tower of a mobile phone provider (O₂ Slovakia) in Lipany, approximately 1.5 km from the collection site [38]. The intensity of the EMF in the nearest town, Sabinov (13 km distance), is, according to available data, 0.25–0.3 V/m [39]. Collected ticks were kept under a natural light regime in the laboratory desiccator to ensure proper constant humidity conditions.




Figure 1. A map of tick collection site. Ticks were collected in northeastern Slovakia (map scale = 10 km), near the village Dubovica (GPS: 49°07'3.7" N 20°57'37.2" E). The inset figure shows the village (insert figure scale = 1 km); the yellow circle marks the collection site. Source of maps: "<https://en.mapy.cz/>" (accessed on 26 October 2023).

2.2. Irradiation Procedure

Ticks were divided into 24 experimental groups, 5 ticks per group. To ensure the proper hydration of ticks, each group was placed in a plastic 2 mL tube with moistened filter paper before irradiation. The irradiation was conducted in an anechoic chamber (model 1710–100, Comtest Engineering, Leyde, The Netherlands), ensuring no external electromagnetic fields affected the experiment. A signal generator (N5183A, Agilent Technologies, Kuala Lumpur, Malaysia) was used as a source of radiofrequency electromagnetic field (RF EMF) connected to a Double-Ridged Waveguide Horn Antenna HF907 (Rohde and Schwarz, Munich, Germany). The antenna was placed approximately 2 m from the experimental apparatus containing ticks.

During the experiment, ticks were irradiated (Table 1) with a constant, polarized, unmodified electromagnetic field at a 900 MHz frequency and two different intensities—2 V/m and 40 V/m. The duration of the exposure was 10 min, 1 h, 3 h and 24 h. Together, 4 control groups of ticks were placed into the anechoic chamber for the designated times but without irradiation.

Table 1. Experimental groups created for the study.

Radiation Intensity	Biological Replications	Sex	Exposure Time	Number of Individuals
40 V/m	3	♀	10 min	5
			1 h	5
			3 h	5
			24 h	5
		♂	10 min	5
			1 h	5
			3 h	5
			24 h	5
2 V/m	3	♀	10 min	5
			1 h	5
			3 h	5
			24 h	5
		♂	10 min	5
			1 h	5
			3 h	5
			24 h	5
0 V/m	3	♀	10 min	5
			1 h	5
			3 h	5
			24 h	5
		♂	10 min	5
			1 h	5
			3 h	5
			24 h	5

2.3. Dissection Protocol

Immediately after irradiation, ticks were dissected live under a stereomicroscope. Ticks were individually placed on double-sided tape, ventral side down. Ticks were cut along the alloscutum using the tip of a scalpel, and the dorsal part of the cuticle was removed. A drop of ice-cold phosphate-buffer saline (PBS; 137 mM NaCl, 1.45 mM NaH₂PO₄, 20.5 mM Na₂HPO₄, pH 7.2) was added onto the exposed organs. Synganglia were surgically removed using precision tweezers with super-fine tips (type #5SF, DUMONT SWITZERLAND, Courtemaîche, Switzerland) and placed into a separate drop of PBS. Synganglia were cleaned from accessory tissues and washed in several drops of clean PBS. Tissues were flash-frozen in 1.5 mL tubes on dry ice.

2.4. RNA Isolation of Tick Synganglia and Quantitative Real-Time Reverse Transcriptase PCR (qRT-PCR)

A pool of 5 synganglia was used for each experimental condition. Total RNA was isolated using an RNeasy[®] Micro Kit (Qiagen, Venlo, The Netherlands). The concentration and purity of obtained RNA were measured using a NanoDrop ONE (Thermo Scientific, Waltham, MA, USA). The obtained RNA was reverse-transcribed to cDNA using a RevertAid H Minus First Strand cDNA Synthesis kit (Thermo Scientific), using oligo (dT) primers. To determine the change in the mRNA levels, qRT-PCR was performed using a LightCycler[®] 480 II (Roche, Meylan, France) thermocycler. LightCycler[®] 480 SYBR[®]

Green I Master mastermix (Roche) was used each 20 μ L reaction was prepared from 10 μ L of mastermix, 7 μ L of PCR-grade H₂O, 1 μ L of 10 μ M forward and 1 μ L of 10 μ M reverse primer, and 1 μ L of template cDNA. Primers for qRT-PCR are listed in Table 2. The nomenclature of studied neuropeptides and their genes is according to Coast and Schooley 2011 [40]. As a reference gene, the ribosomal protein S4 (*rps4*) was used [41].

Table 2. qRT-PCR primers used in this study.

Protein	Gene	Forward Primer (5'-3')	Reverse Primer (5'-3')	References
FGLA-related allatostatins	<i>fgla/ast</i>	AGCGGAGGTACAACCTTTGGC	CTCCTCTTCCAGCGCTCG	[42]
Allatotropin	<i>at</i>	GGCTTTGGCAAGAGAATGAG	GGCTATTTCTCCGTAACC	[42]
Arginine-vasopressin like peptide	<i>avpl</i>	TCTCACACATGCTGCTCCTG	GCATCGTTGAGGATGCACAT	This study
Kinin	<i>kinin</i>	ACCACGTTCTGATGAGCAT	ATGTTCCAGCGAATGAAGCT	This study
Ribosomal protein S4	<i>rps4</i>	GGTGAAGAAGATTGTCAAGCAGAG	TGAAGCCAGCAGGTTAGTTTG	[41]

Opaque 96-well plates were used for RT-PCR. The preincubation step of the cycling protocol was carried out at 95 °C for 5 min, followed by the amplification step at 95 °C (10 s), 60 °C (for *fgla/ast* and *at*) or 55 °C (for *kinin* and *avpl*) (10 s), and 72 °C (10 s). A total of 45 cycles were used for the amplification step. The melting curve was analyzed from 65 °C to 97 °C. The amplicons of targeted genes were verified via sequencing the qRT-PCR products (Eurofins, Luxembourg, Luxembourg).

Relative transcript levels were calculated utilizing the $\Delta\Delta$ Ct ratio method according to Livak and Schmittgen, 2001 using Microsoft Excel and are expressed as fold change value [43]. Graphs for each studied gene were created in GraphPad 5 Prism (GraphPad Software Inc., San Diego, CA, USA). Statistical significance was assessed using Student's *t*-test in GraphPad 5 Prism (GraphPad Software Inc.). This experiment was conducted in 3 biological replications and in two technical replications.

3. Results

Reduced mRNA levels of *at* were found in most of the irradiated groups, significantly after one hour of irradiation in both sexes. In females, the levels were reduced 0.53-fold and 0.21-fold after one-hour exposure. In male groups, the levels were reduced 0.26 and 0.14 folds after 1 h exposure under same experimental conditions. In females exposed to 40 V/m radiation, the transcript levels of *at* were significantly reduced not only after the already-mentioned one-hour irradiation but also after 3 (0.06 folds) and 24 h (0.04 folds). The only increase in *at* levels, although statistically non-significant, was noticed after 10 min of 2 V/m exposure in males and 3 h in females (1.44 and 1.38 folds, respectively) (Figure 2).

An elevation in mRNA levels of *fgla/ast* was found in all females irradiated with 2 V/m radiation, with fold changes ranging from 2.05 to 10.77. The 10.77-fold increase in the levels in females exposed to 2 V/m for 3 h was significant. In female groups exposed to 40 V/m for 1 h or more, *fgla/ast* levels were significantly reduced: 0.27 folds after 1 h, 0.21 after 3 h, and 0.14 folds after 24 h exposure. The levels of *fgla/ast* transcripts were reduced after 1 h (0.36-fold) and 3 h (0.45-fold) 2 V/m irradiation in males. In males exposed to a higher intensity, a decrease in mRNA levels after 1, 3, and 24 h of exposure was noted, but significantly only in the 1 h group (decrease of 0.05 folds) (Figure 3).




Figure 2. Effects on different EMF intensities on the allatotropin (*at*) transcripts in *I. ricinus* adult synganglia of (A) females; (B) males. Data were normalized using the ribosomal protein S4 transcript (*rps4*), and the expression levels in non-irradiated ticks were assigned a value of 1. Results shown are means and standard errors of means. Asterisks indicate the comparison of the mean to the value of the non-irradiated group. The statistical significance of the fold difference was calculated using an unpaired *t*-test and is expressed using *p* values: ** *p* < 0.01; *** *p* < 0.001.




Figure 3. Effects on different EMF intensities on the FGLA-related allatostatin (*fgla/ast*) transcripts in *I. ricinus* adult synganglia from (A) females; (B) males. Data were normalized using the ribosomal protein S4 transcript (*rps4*), and the expression levels in non-irradiated ticks were assigned a value of 1. Results shown are means and standard errors of means. Asterisks indicate the comparison of the mean to the value of the non-irradiated group. The statistical significance of the fold difference was calculated using an unpaired *t*-test and is expressed using *p* values: * *p* < 0.05; ** *p* < 0.01; *** *p* < 0.001.

The mRNA levels of *avpl* were significantly reduced 0.34-fold in the groups of females exposed to 2 V/m radiation for 24 h. The levels of *avpl* in the female groups exposed to

40 V/m reduced after 1 h 0.33 folds; after 3 h, 0.05 folds; and after 24 h of irradiation, 0.16 folds. In the rest of the female groups, the gene expression was upregulated, but the results were non-significant. A significant downregulation of transcripts was found in the group of males irradiated for one hour by 40 V/m (0.59-fold). The number of transcripts was slightly elevated in most male groups, however not significantly (Figure 4).

The mRNA levels of *kinin* were non-significantly increased in all groups of females irradiated with 2 V/m intensity (fold change from 1.56 to 11.44) and in the group exposed to 40 V/m radiation for 24 h (3.6-fold). Transcript levels were significantly reduced in the groups of females irradiated with the higher intensity for up to 3 h, specifically 0.06 folds after 10 min, 0.04 after 1 h, and 0.24 folds after 3 h of exposure. The levels of mRNA were slightly elevated in males during the first three hours of exposure to 2 V/m; however, these were non-significant results. When applying 40 V/m for 3 h, the levels of the *kinin* transcript were significantly reduced to 0.13-fold (Figure 5).




Figure 4. Effects on different EMF intensities on the arginine-vasopressin-like peptide (*avpl*) transcripts in *I. ricinus* adult synganglia from (A) females; (B) males. Data were normalized using the ribosomal protein S4 transcript (*rps4*), and the expression levels in non-irradiated ticks were assigned a value of 1. Results shown are means and standard errors of means. Asterisks indicate the comparison of the mean to the value of the non-irradiated group. The statistical significance of the fold difference was calculated using an unpaired *t*-test and is expressed using *p* values: * *p* < 0.05; *** *p* < 0.001.




Figure 5. Effects on different EMF intensities on the *kinin* transcripts in *I. ricinus* adult synganglia from (A) females; (B) males. Data were normalized using the ribosomal protein S4 transcript (*rps4*), and the expression levels in non-irradiated ticks were assigned a value of 1. Results shown are means and standard errors of means. Asterisks indicate the comparison of the mean to the value of the non-irradiated group. The statistical significance of the fold difference was calculated using an unpaired *t*-test and is expressed as *p* values: ** *p* < 0.01; *** *p* < 0.001.

4. Discussion

In our experiment, a 900 MHz frequency—the mainstream spectrum for mobile data and voice broadband—was used to study the effect of electromagnetic radiation on selected transcript levels in the synganglion of both sexes of *I. ricinus*. This frequency has proven bioactive effects [7,44] and has been utilized in other tick behavioral tests and studies in the past. Several species of ticks (*I. ricinus*, *D. reticulatus*) showed a significant response to it [27–29]. In the questing arena set-up, *D. reticulatus* ticks displayed jerking movements [27]. In the modified T-labyrinth arena (RST test), ticks preferred the arm exposed to irradiation when a 900 MHz frequency was used for 24 h [27,28]. Female *I. ricinus* ticks tested in the same manner did not display any significant affinity for either arm. However, tracking ticks' movement in the circular arena with irradiated and shielded parts revealed the difference in the movement dynamics [45].

We noted differences in the levels of neuropeptide mRNAs in the ticks' synganglion after exposure to electromagnetic radiation with different intensities. Previous studies on drosophila and honeybees have reported different effects of radiofrequency radiation on the organism depending on the radiation parameters used as well [44,46–48]. Modulation, pulsed fields, intensity, and specific absorption rate have been cited as factors that enhance the biological effects of exposure [49]. In the irradiation study on drosophila, all intensities (power densities) produced a change in gene expression and enzyme activity after

48 h exposure, and it seems that there is still bioactive effect if the irradiation is less than 0.0613 V/m [50]. The effect of radiation, according to studies, is most noticeable from 30 cm distance but can still have an effect from a 1 m distance, especially if the power density is higher than 1 $\mu\text{W}/\text{cm}$ [49].

In this study, an insignificant increase in the levels of transcripts was found in the lower-intensity radiation (2 V/m) female groups (Table 3). The 2 V/m intensity was utilized in this experiment, as it was found to be a common EMF intensity occurring in urban areas of eastern Slovakia [51]. It seems that the mRNA transcript levels are affected only slightly by lower-intensity radiation; however, utilizing higher intensity (40 V/m) produced an enhanced effect on the synganglion of *I. ricinus*. In females irradiated with 40 V/m, three out of four neuropeptide mRNA levels (*avpl*, *fgla/ast* and *at*) were significantly reduced after one hour of irradiation. Here, 40 V/m intensity was chosen for this experiment because it is close to the maximal 900 MHz radiation intensity allowed in Slovakia (which is 42 V/m [52]), although it can be rarely found in non-laboratory setups. As we hypothesize that EMF does affect neuropeptide levels, this intensity was chosen to simulate a super-normal stimulus. A much broader range of intensities and frequencies of EMF needs to be tested to acquire a more complex understanding of the effects of EMFs on the tick central nervous system in the future.

Table 3. Graphic depiction of the neuropeptide transcript levels in female experimental groups, comparison of the mean to the value of the non-irradiated group. The statistical significance of the fold difference was calculated using an unpaired *t*-test and is expressed with *p* values: grey color—no change, $p > 0.5$; green color—increase in mRNA level, $p < 0.5$; red color—decrease in mRNA level, $p < 0.5$; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

♀ Gene	2 V/m				40 V/m			
	10 min	1 h	3 h	24 h	10 min	1 h	3 h	24 h
<i>kinin</i>	Green	Green	Grey	Green	Red (***)	Red (***)	Red (**)	Grey
<i>avpl</i>	Grey	Green	Green	Red (*)	Grey	Red (*)	Red (***)	Red (***)
<i>at</i>	Grey	Red (***)	Green	Red	Grey	Red (**)	Red (***)	Red (***)
<i>fgla/ast</i>	Green	Green	Green (*)	Green	Grey	Red (*)	Red (***)	Red (***)

The results for both intensities varied in male ticks (Table 4). Although previous reports indicate sex-specific expression of certain transcripts among tick sexes [53,54], at this point, it is difficult to speculate which molecular or physiological mechanisms caused the differences observed in our study. Despite this fact, our results are consistent with the behavioral differences detected among male and female ticks exposed to irradiation in our previous experiments. Specifically, in *D. reticulatus*, females expressed a specific jerking movement with the whole body, while male ticks jerked mostly with their front legs, so the same radiation exposure parameters produced a stronger response in females. In the RST arena (radiation-shielded tube arena), however, there was no significant preference for exposed/shielded arm between the sexes in *I. ricinus* and *D. reticulatus* [27–29]. Taken together, more studies involving multiple tick species are necessary to figure out the sex-specific responsiveness of male and female ticks to EMFs.

The duration of radiation exposure was also considered in this study. Behavioral studies on *D. reticulatus* and *I. ricinus* in the past reported that ticks preferred the exposed arm of the labyrinth after 24 h of exposure [28,29]. In our experiment, irradiation seemed to have had a significant effect on tick mRNA levels after at least 1 h of exposure. Interestingly, changes in locomotor behavior and an influence on larvae development after approximately one hour of exposure to 900 MHz were reported in ticks before. In a behavioral study of *I. ricinus* in EMF, ticks were irradiated by GSM mobile signal for 24 h and, after one hour, started spending a longer time walking in the exposed part of the experimental arena [43]. One hour of exposure to the same frequency significantly enlarged the body dimensions of *D. reticulatus* larvae hatched from irradiated eggs. From the groups of eggs exposed to radiation for 30 and 90 min, the larvae were smaller than the control larvae

when hatched [27]. One hour of exposure therefore seems to be the minimal effective length of EMF irradiation for this frequency.

Table 4. Graphic depiction of the neuropeptide transcript levels in male experimental groups, comparison of the mean to the value of the non-irradiated group. The statistical significance of the fold difference was calculated using an unpaired *t*-test and is expressed with *p* values: grey color—no change, *p* > 0.5; green color—increase in mRNA level, *p* < 0.5; red color—decrease in mRNA level, *p* < 0.5; * *p* < 0.05; *** *p* < 0.001.

♂	2 V/m				40 V/m			
Gene	10 min	1 h	3 h	24 h	10 min	1 h	3 h	24 h
<i>kinin</i>	Green	Grey	Grey	Red	Grey	Red	Red (***)	Green
<i>avpl</i>	Grey	Grey	Grey	Green	Grey	Red (*)	Grey	Green
<i>at</i>	Grey	Red (***)	Red (***)	Red (***)	Red (***)	Red (***)	Red (***)	Red (***)
<i>fgla/ast</i>	Green	Red (***)	Red (*)	Green	Grey	Red (***)	Red (***)	Red (***)

Information about the effects of EMF on the arthropod nervous system is scarce. In previously published papers on drosophila, it is suggested that EMFs can affect the signaling pathways in ecdysteroid production [50] and the levels of biogenic amines in the brain [23]. Changes in biogenic amine levels after low-frequency irradiation were found in crickets as well [55]. The neuropeptide genes studied in this paper were selected randomly, as no previous studies on any neuropeptide genes in arthropods under influence of EMFs have been published so far. Here, a plausible approach for future studies would be mass sequencing, to obtain a more in-depth understanding of the changes in the synganglion transcripts under radiofrequency irradiation. Therefore, as our pioneer work investigated only four different neuropeptide transcripts, at this point, it is hard to ascertain the exact influence of irradiation on tick physiology and behavior that result from the orchestration of multiple gene products.

The increasing prevalence of ticks and tick-borne pathogens in urbanized areas like city parks and recreational woods near the cities is currently an urgent problem [56,57]. Green spaces in cities provide suitable environmental conditions for ticks, plenty of hosts, and possibly a constant dose of EMF exposure. However, it is hard to conclude if the omnipresent anthropogenic fields provide some kind of specific stimulation for ticks, as much more research needs to be conducted on this topic.

Since the first irradiation experiment published by Vargová et al. 2017, it was hypothesized that ticks could be able to perceive the inner electromagnetic field of their hosts [27]. Recently, it has been discovered that ticks likely utilize static electric fields to transport a very short distance through the air to latch onto their hosts while questing on vegetation. However, it was shown that this is a passive process of electrostatic attraction [58]. The question of the presence of some type of electromagnetic sense in ticks is still unanswered. Sensillae-like structures for infrared radiation in the capsule area of the Haller’s organ were described in *D. variabilis* [59]. Electromagnetic perception based on the presence of magnetite particles in the tissues found in bees and ants could also be a candidate mechanism of electromagnetic sensing in ticks [60,61].

5. Conclusions

To conclude this study, radiofrequency electromagnetic radiation does alter the levels of neuropeptide transcripts in the central nervous system of *Ixodes ricinus*. While both tested intensities affected the mRNA levels, the intensity of 40 V/m and 1 h of radiation exposure produced the most significant results. A more dramatic impact on female ticks was noted. Most EMF effects on transcript amount were gene-specific.

We present a completely novel research focus among studies on the effects of the abiotic environment on the physiology of ticks. As the neuropeptidergic system exerts various functions in the tick body, a more extensive study of gene expression under irradiation conditions is required in the future.

Author Contributions: Conceptualization, V.M., I.M. and L.Š. (Ladislav Šimo); Methodology, J.K., L.M.-H. and L.Š. (Lívia Šofranková); Software, L.Š. (Lívia Šofranková) and M.P.; Validation, L.Š. (Ladislav Šimo) and V.M.; Investigation, L.Š. (Lívia Šofranková), M.B., N.P. and M.P.; Resources, V.M., I.M., L.Š. (Ladislav Šimo) and R.C.; Formal Analysis, L.Š. (Lívia Šofranková); Data Curation, L.Š. (Lívia Šofranková) and L.Š. (Ladislav Šimo); Writing—Original Draft Preparation, L.Š. (Lívia Šofranková); Writing—Review and Editing, V.M., L.Š. (Ladislav Šimo) and I.M.; Visualization, L.Š. (Lívia Šofranková); Supervision, V.M., L.Š. (Ladislav Šimo) and I.M.; Funding Acquisition, V.M., L.Š. (Ladislav Šimo), J.K. and I.M. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Slovak Research and Development Agency (grant numbers APVV-17-0372, APVV-19-0440); Internal Scientific Grant System of the Pavol Jozef Šafárik University in Košice (grant numbers VVGS-2022-2192, VVGS-PF-2022-2135); Slovak Academic Information Agency (SAIA) via the the National Scholarship Programme of the Slovak Republic; French National Research Agency (ANR) (grant number: ANR-21-CE14-0012), project AxoTick. UMR BIPAR is supported by the French Government’s Investissement d’Avenir program, Laboratoire d’Excellence “Integrative Biology of Emerging Infectious Diseases” (grant number: ANR-10-LABX-62-IBEID) and the COST Action CA21170 “Prevention, anticipation and mitigation of tick-borne disease risk applying the DAMA protocol (PRAGMATICK)”.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data presented in this study are openly available in 10.6084/m9.figshare.24637908 licensed as CC BY 4.0.

Acknowledgments: Authors would like to thank to Institut français de Slovaquie and Campus France for providing support during the scientific internship in France.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Balmori, A. Electromagnetic Radiation as an Emerging Driver Factor for the Decline of Insects. *Sci. Total Environ.* **2021**, *767*, 144913. [[CrossRef](#)]
2. Singh, A.P.; Kaur, R. Electromagnetic Fields: Biological Implications on Various Life Forms. *Int. J. Bioassays* **2014**, *3*, 2030–2040.
3. De Iuliis, G.N.; Newey, R.J.; King, B.V.; Aitken, R.J. Mobile Phone Radiation Induces Reactive Oxygen Species Production and DNA Damage in Human Spermatozoa in Vitro. *PLoS ONE* **2009**, *4*, e6446. [[CrossRef](#)]
4. Ulubay, M.; Yahyazadeh, A.; Deniz, Ö.G.; Kivrak, E.G.; Altunkaynak, B.Z.; Erdem, G.; Kaplan, S. Effects of Prenatal 900 MHz Electromagnetic Field Exposures on the Histology of Rat Kidney. *Int. J. Radiat. Biol.* **2015**, *91*, 35–41. [[CrossRef](#)]
5. Molina-Montenegro, M.A.; Acuña-Rodríguez, I.S.; Ballesteros, G.I.; Baldelomar, M.; Torres-Díaz, C.; Broitman, B.R.; Vázquez, D.P. Electromagnetic Fields Disrupt the Pollination Service by Honeybees. *Sci. Adv.* **2023**, *9*, eadh1455. [[CrossRef](#)]
6. Kim, J.H.; Chung, K.H.; Hwang, Y.R.; Park, H.R.; Kim, H.J.; Kim, H.-G.; Kim, H.R. Exposure to RF-EMF Alters Postsynaptic Structure and Hinders Neurite Outgrowth in Developing Hippocampal Neurons of Early Postnatal Mice. *Int. J. Mol. Sci.* **2021**, *22*, 5340. [[CrossRef](#)]
7. Panagopoulos, D.J.; Chavdoula, E.D.; Nezis, I.P.; Margaritis, L.H. Cell Death Induced by GSM 900-MHz and DCS 1800-MHz Mobile Telephony Radiation. *Mutat. Res. Genet. Toxicol. Environ. Mutagen.* **2007**, *626*, 69–78. [[CrossRef](#)]
8. Wyszowska, J.; Shepherd, S.; Sharkh, S.; Jackson, C.W.; Newland, P.L. Exposure to Extremely Low Frequency Electromagnetic Fields Alters the Behaviour, Physiology and Stress Protein Levels of Desert Locusts. *Sci. Rep.* **2016**, *6*, 36413. [[CrossRef](#)]
9. Goodman, E.M.; Sharpe, P.T.; Greenebaum, B.; Marron, M.T. Pulsed Magnetic Fields Alter the Cell Surface. *FEBS Lett.* **1986**, *199*, 275–278. [[CrossRef](#)]
10. Carson, J.J.; Prato, F.S.; Drost, D.J.; Diesbourg, L.D.; Dixon, S.J. Time-Varying Magnetic Fields Increase Cytosolic Free Ca²⁺ in HL-60 Cells. *Am. J. Physiol. Cell Physiol.* **1990**, *259*, C687–C692. [[CrossRef](#)]
11. Lindström, E.; Lindström, P.; Berglund, A.; Mild, K.H.; Lundgren, E. Intracellular Calcium Oscillations Induced in a T-Cell Line by a Weak 50 Hz Magnetic Field. *J. Cell. Physiol.* **1993**, *156*, 395–398. [[CrossRef](#)]
12. Gaber, M.H.; Abd El Halim, N.; Khalil, W.A. Effect of Microwave Radiation on the Biophysical Properties of Liposomes. *Bioelectromagnetics* **2005**, *26*, 194–200. [[CrossRef](#)]
13. Ketabi, N.; Mobasheri, H.; Faraji-Dana, R. Electromagnetic Fields (UHF) Increase Voltage Sensitivity of Membrane Ion Channels; Possible Indication of Cell Phone Effect on Living Cells. *Electromagn. Biol. Med.* **2015**, *34*, 1–13. [[CrossRef](#)]

14. Altunkaynak, B.Z.; Altun, G.; Yahyazadeh, A.; Kaplan, A.A.; Deniz, O.G.; Türkmen, A.P.; Önger, M.E.; Kaplan, S. Different Methods for Evaluating the Effects of Microwave Radiation Exposure on the Nervous System. *J. Chem. Neuroanat.* **2016**, *75*, 62–69. [[CrossRef](#)]
15. Panagopoulos, D.J.; Karabarbounis, A.; Yakymenko, I.; Chrousos, G.P. Human-made Electromagnetic Fields: Ion Forced-oscillation and Voltage-gated Ion Channel Dysfunction, Oxidative Stress and DNA Damage (Review). *Int. J. Oncol.* **2021**, *59*, 92. [[CrossRef](#)]
16. Gerner, C.; Haudek, V.; Schandl, U.; Bayer, E.; Gundacker, N.; Hutter, H.P.; Mosgoeller, W. Increased Protein Synthesis by Cells Exposed to a 1800-MHz Radio-Frequency Mobile Phone Electromagnetic Field, Detected by Proteome Profiling. *Int. Arch. Occup. Environ.* **2010**, *83*, 691–702. [[CrossRef](#)]
17. Lin, H.; Blank, M.; Goodman, R. A Magnetic Field-Responsive Domain in the Human HSP70 Promoter. *J. Cell. Biochem.* **1999**, *75*, 170–176. [[CrossRef](#)]
18. Czyz, J.; Guan, K.; Zeng, Q.; Nikolova, T.; Meister, A.; Schönborn, F.; Schuderer, J.; Kuster, N.; Wobus, A.M. High Frequency Electromagnetic Fields (GSM Signals) Affect Gene Expression Levels in Tumor Suppressor P53-Deficient Embryonic Stem Cells. *Bioelectromagnetics* **2004**, *25*, 296–307. [[CrossRef](#)]
19. Zhao, T.-Y.; Zou, S.-P.; Knapp, P.E. Exposure to Cell Phone Radiation Up-Regulates Apoptosis Genes in Primary Cultures of Neurons and Astrocytes. *Neurosci. Lett.* **2007**, *412*, 34–38. [[CrossRef](#)]
20. Patruno, A.; Pesce, M.; Grilli, A.; Speranza, L.; Franceschelli, S.; De Lutiis, M.A.; Vianale, G.; Costantini, E.; Amerio, P.; Muraro, R.; et al. MTOR Activation by PI3K/Akt and ERK Signaling in Short ELF-EMF Exposed Human Keratinocytes. *PLoS ONE* **2015**, *10*, e0139644. [[CrossRef](#)]
21. Blank, M.; Goodman, R. Electromagnetic Fields Stress Living Cells. *Pathophysiology* **2009**, *16*, 71–78. [[CrossRef](#)]
22. Yalçın, S.; Erdem, G. Biological Effects of Electromagnetic Fields. *Afr. J. Biotechnol.* **2012**, *11*, 3933–3941.
23. Newland, P.L.; Al Ghamdi, M.S.; Sharkh, S.; Aonuma, H.; Jackson, C.W. Exposure to Static Electric Fields Leads to Changes in Biogenic Amine Levels in the Brains of *Drosophila*. *Proc. R. Soc.* **2015**, *282*, 20151198. [[CrossRef](#)]
24. Korotkov, I.; Burenkov, M.; Burenkova, L.; Pichugin, V.I.; Chunikhin, S.; Engovatov, V. The Reaction of the Tick *Hyalomma Asiaticum* (Acarina, Ixodidae) to 1-to 4-GHz Microwaves. *Med. Parazitol.* **1996**, *4*, 28–31.
25. Korotkov, I.S.; Burenkova, L.; Burenkov, M.; Pichugin, V.I. The Impact of Electromagnetic Radiation at Microwave Frequency (9.8 HhZ) on the Embryonic and Postembryonic Development of the Tick *Hyalomma Asiaticum* (Acarina, Ixodidae). *Med. Parazitol.* **2000**, 38–42.
26. Vargová, B.; Majláth, I.; Kurimský, J.; Cimbala, R.; Pipova, N.; Živčák, J.; Tryjanowski, P.; Peťko, B.; Džmura, J.; Ižariková, G.; et al. Morphometric Analysis—Effect of the Radiofrequency Interface of Electromagnetic Field on the Size of Hatched *Dermacentor Reticulatus* Larvae. *Ann. Agric. Environ. Med.* **2021**, *28*, 419–425. [[CrossRef](#)]
27. Vargová, B.; Kurimský, J.; Cimbala, R.; Kosterec, M.; Majláth, I.; Pipová, N.; Tryjanowski, P.; Jankowiak, Ł.; Majláthová, V. Ticks and Radio-Frequency Signals: Behavioural Response of Ticks (*Dermacentorreticulatus*) in a 900 MHz Electromagnetic Field. *Syst. Appl. Acarol.* **2017**, *22*, 683–693. [[CrossRef](#)]
28. Vargová, B.; Majláth, I.; Kurimský, J.; Cimbala, R.; Kosterec, M.; Tryjanowski, P.; Jankowiak, Ł.; Raši, T.; Majláthová, V. Electromagnetic Radiation and Behavioural Response of Ticks: An Experimental Test. *Exp. Appl. Acarol.* **2018**, *75*, 85–95. [[CrossRef](#)]
29. Frączak, M.; Vargová, B.; Tryjanowski, P.; Majláth, I.; Jerzak, L.; Kurimský, J.; Cimbala, R.; Jankowiak, Ł.; Conka, Z.; Majláthová, V. Infected *Ixodes Ricinus* Ticks Are Attracted by Electromagnetic Radiation of 900 MHz. *Ticks Tick Borne Dis.* **2020**, *11*, 101416. [[CrossRef](#)]
30. Nässel, D.R. Neuropeptides in the Nervous System of *Drosophila* and Other Insects: Multiple Roles as Neuromodulators and Neurohormones. *Prog. Neurobiol.* **2002**, *68*, 1–84. [[CrossRef](#)]
31. Yeoh, J.G.C.; Pandit, A.A.; Zandawala, M.; Nässel, D.R.; Davies, S.-A.; Dow, J.A.T. DIneR: Database for Insect Neuropeptide Research. *Insect Biochem. Mol. Biol.* **2017**, *86*, 9–19. [[CrossRef](#)]
32. Mateos-Hernández, L.; Pipová, N.; Allain, E.; Henry, C.; Rouxel, C.; Lagrée, A.-C.; Haddad, N.; Boulouis, H.-J.; Valdés, J.J.; Alberdi, P.; et al. Enlisting the *Ixodes Scapularis* Embryonic ISE6 Cell Line to Investigate the Neuronal Basis of Tick—Pathogen Interactions. *Pathogens* **2021**, *10*, 70. [[CrossRef](#)]
33. Gulia-Nuss, M.; Nuss, A.B.; Meyer, J.M.; Sonenshine, D.E.; Roe, R.M.; Waterhouse, R.M.; Sattelle, D.B.; de la Fuente, J.; Ribeiro, J.M.; Megy, K.; et al. Genomic Insights into the *Ixodes Scapularis* Tick Vector of Lyme Disease. *Nat. Commun.* **2016**, *7*, 10507. [[CrossRef](#)]
34. Šimo, L.; Slovák, M.; Park, Y.; Žitňan, D. Identification of a Complex Peptidergic Neuroendocrine Network in the Hard Tick, *Rhipicephalus Appendiculatus*. *Cell Tissue Res.* **2009**, *335*, 639–655. [[CrossRef](#)]
35. Kim, D.; Šimo, L.; Park, Y. Molecular Characterization of Neuropeptide Elevenin and Two Elevenin Receptors, IsElevR1 and IsElevR2, from the Blacklegged Tick, *Ixodes Scapularis*. *Insect Biochem. Mol. Biol.* **2018**, *101*, 66–75. [[CrossRef](#)]
36. Šimo, L.; Park, Y. Neuropeptidergic Control of the Hindgut in the Black-Legged Tick *Ixodes Scapularis*. *Int. J. Parasitol.* **2014**, *44*, 819–826. [[CrossRef](#)]
37. Šimo, L.; Žitňan, D.; Park, Y. Two Novel Neuropeptides in Innervation of the Salivary Glands of the Black-Legged Tick, *Ixodes Scapularis*: Myoinhibitory Peptide and SIFamide. *J. Comp. Neurol.* **2009**, *517*, spc1. [[CrossRef](#)]

38. © Voxo 2011–2023 Mapa Vysielačov v SR. Available online: <https://elektrosmog-info.voxo.eu/mapa-vysielačov> (accessed on 26 October 2023).
39. Electrical Power Engineering Department, Technical University in Košice; VUJE, Inc. Maps of Electromagnetic Fields. Available online: <http://www.emp.vuje.sk/EN4/#> (accessed on 26 October 2023).
40. Coast, G.M.; Schooley, D.A. Toward a Consensus Nomenclature for Insect Neuropeptides and Peptide Hormones. *Peptides* **2011**, *32*, 620–631. [[CrossRef](#)]
41. Koči, J.; Šimo, L.; Park, Y. Validation of Internal Reference Genes for Real-Time Quantitative Polymerase Chain Reaction Studies in the Tick, *Ixodes Scapularis* (Acari: Ixodidae). *J. Med. Entomol.* **2013**, *50*, 79–84. [[CrossRef](#)]
42. Hromníková, D. Use of Molecular and Transgenic Methods in Identification and Functional Analysis of Receptors. Ph.D. Thesis, Comenius University Bratislava, Bratislava, Slovakia, 2021.
43. Livak, K.J.; Schmittgen, T.D. Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the $2^{-\Delta\Delta CT}$ Method. *Methods* **2001**, *25*, 402–408. [[CrossRef](#)]
44. Panagopoulos, D.J.; Chavdoula, E.D.; Karabarbounis, A.; Margaritis, L.H. Comparison of Bioactivity Between GSM 900 MHz and DCS 1800 MHz Mobile Telephony Radiation. *Electromagn. Biol. Med.* **2007**, *26*, 33–44. [[CrossRef](#)]
45. Vargová, B.; Majláth, I.; Kurimský, J.; Cimbala, R.; Zbojovský, J.; Tryjanowski, P.; Majláthová, V. Locomotor Activity of *Ixodes Ricinus* Females in 900 MHz Electromagnetic Field. *Life* **2022**, *12*, 884. [[CrossRef](#)]
46. Panagopoulos, D.J. Comparing DNA Damage Induced by Mobile Telephony and Other Types of Man-Made Electromagnetic Fields. *Mutat. Res. Rev. Mutat. Res.* **2019**, *781*, 53–62. [[CrossRef](#)]
47. Vilić, M.; Tlak Gajger, I.; Tucak, P.; Štambuk, A.; Šrut, M.; Klobučar, G.; Malarić, K.; Žura Žaja, I.; Pavelić, A.; Manger, M.; et al. Effects of Short-Term Exposure to Mobile Phone Radiofrequency (900 MHz) on the Oxidative Response and Genotoxicity in Honey Bee Larvae. *J. Apic. Res.* **2017**, *56*, 430–438. [[CrossRef](#)]
48. Migdal, P.; Bieńkowski, P.; Cebrat, M.; Berbec, E.; Plotnik, M.; Murawska, A.; Sobkiewicz, P.; Łaszkiwicz, A.; Latarowski, K. Exposure to a 900 MHz Electromagnetic Field Induces a Response of the Honey Bee Organism on the Level of Enzyme Activity and the Expression of Stress-Related Genes. *PLoS ONE* **2023**, *18*, e0285522. [[CrossRef](#)]
49. Panagopoulos, D.J.; Chavdoula, E.D.; Margaritis, L.H. Bioeffects of Mobile Telephony Radiation in Relation to Its Intensity or Distance from the Antenna. *Int. J. Radiat. Biol.* **2010**, *86*, 345–357. [[CrossRef](#)]
50. Wang, Y.; Jiang, Z.; Zhang, L.; Zhang, Z.; Liao, Y.; Cai, P. 3.5-GHz Radiofrequency Electromagnetic Radiation Promotes the Development of *Drosophila Melanogaster*. *Environ. Pollut.* **2022**, *294*, 118646. [[CrossRef](#)]
51. Zbojovský, J.; Pavlík, M.; Cimbala, R.; Kurimský, J. Merania elektromagnetických polí v okolí vybraných telekomunikačných vysielačov v lokalitách Košice a okolie. In *Proceedings of the Zborník Príspevkov Prednesených na Vedeckom Seminári EMITICK22*; Technická Univerzita v Košiciach: Stará Lesná, Slovakia, 2022; pp. 33–38.
52. Act No. 355/2007 Z. z. on the Protection, Promotion and Development of Public Health and Amending Certain Acts, as Amended. Available online: <https://www.epi.sk/zz/2007-534> (accessed on 19 September 2023).
53. Tan, A.W.L.; Francischetti, I.M.B.; Slovak, M.; Kini, R.M.; Ribeiro, J.M.C. Sexual Differences in the Sialomes of the Zebra Tick, *Rhipicephalus Pulchellus*. *J. Proteom.* **2015**, *117*, 120–144. [[CrossRef](#)]
54. Bensaoud, C.; Nishiyama, M.Y.; Ben Hamda, C.; Lichtenstein, F.; Castro de Oliveira, U.; Faria, F.; Loiola Meirelles Junqueira-de-Azevedo, I.; Ghedira, K.; Bouattour, A.; M’Ghirbi, Y.; et al. De Novo Assembly and Annotation of *Hyalomma Dromedarii* Tick (Acari: Ixodidae) Sialotranscriptome with Regard to Gender Differences in Gene Expression. *Parasites Vectors* **2018**, *11*, 314. [[CrossRef](#)]
55. Wyszowska, J.; Kobak, J.; Aonuma, H. Electromagnetic Field Exposure Affects the Calling Song, Phonotaxis, and Level of Biogenic Amines in Crickets. *Environ. Sci. Pollut. Res.* **2023**, *30*, 93255–93268. [[CrossRef](#)]
56. Sormunen, J.J.; Kulha, N.; Klemola, T.; Mäkelä, S.; Vesilahti, E.-M.; Vesterinen, E.J. Enhanced Threat of Tick-Borne Infections within Cities? Assessing Public Health Risks Due to Ticks in Urban Green Spaces in Helsinki, Finland. *Zoonoses Public Health* **2020**, *67*, 823–839. [[CrossRef](#)] [[PubMed](#)]
57. Heylen, D.; Lasters, R.; Adriaensen, F.; Fonville, M.; Sprong, H.; Matthysen, E. Ticks and Tick-Borne Diseases in the City: Role of Landscape Connectivity and Green Space Characteristics in a Metropolitan Area. *Sci. Total Environ.* **2019**, *670*, 941–949. [[CrossRef](#)] [[PubMed](#)]
58. England, S.J.; Lihou, K.; Robert, D. Static Electricity Passively Attracts Ticks onto Hosts. *Curr. Biol.* **2023**, *33*, 3041–3047.e4. [[CrossRef](#)]
59. Mitchell, R.D.; Zhu, J.; Carr, A.L.; Dhammi, A.; Cave, G.; Sonenshine, D.E.; Roe, R.M. Infrared Light Detection by the Haller’s Organ of Adult American Dog Ticks, *Dermacentor Variabilis* (Ixodida: Ixodidae). *Ticks Tick Borne Dis.* **2017**, *8*, 764–771. [[CrossRef](#)]
60. Acosta-Avalos, D.; Wajnberg, E.; Oliveira, P.S.; Leal, I.; Farina, M.; Esquivel, D.M.S. Isolation of Magnetic Nanoparticles from *Pachycondyla Marginata* Ants. *J. Exp. Biol.* **1999**, *202*, 2687–2692. [[CrossRef](#)]
61. Lambinet, V.; Hayden, M.E.; Reigl, K.; Gomis, S.; Gries, G. Linking Magnetite in the Abdomen of Honey Bees to a Magnetoreceptive Function. *Proc. Royal Soc. B* **2017**, *284*, 20162873. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.