



HAL
open science

Pipecolate, specific biomarker of lysine deficiency

G. Roisé-Hamelin, C. Gaudichon, S. Devi, J.-C. Martin, C. Tardivel, A. Kurpad, D. Tomé, Delphine Jouan-Rimbaud Bouveresse, D. Azzout-Marniche

► To cite this version:

G. Roisé-Hamelin, C. Gaudichon, S. Devi, J.-C. Martin, C. Tardivel, et al.. Pipecolate, specific biomarker of lysine deficiency. International Symposium on “ Dietary Protein for Human Health ”, Sep 2023, Utrecht, Netherlands. . hal-04311376

HAL Id: hal-04311376

<https://hal.inrae.fr/hal-04311376>

Submitted on 28 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Pipecolate, specific biomarker of lysine deficiency

Gaëtan Roisné-Hamelin^a, Claire Gaudichon^a, Sarita Devi^b, Jean-Charles Martin^c, Catherine Tardivel^c, Anura Kurpad^b, Daniel Tomé^a, Delphine Jouan-Rimbaud Bouveresse^a and Dalila Azzout-Marniche^a

^a Université Paris-Saclay, AgroParisTech, INRAE, UMR PNCA, 91120, Palaiseau, France.


^b Division of Nutrition, St John's Reserach Institute, St John's National academy of health sciences, bangalore, India.

^c Aix Marseille université, INSERM, C2VN, Marseille, France.

Background & Objective

- The consumption of poor-quality protein increases the risk of essential amino acid (EAA) deficiency, particularly for lysine, threonine and methionine. Thus, it is necessary to be able to detect easily EAA deficiency.
- We have previously identified pipecolate and taurine as potential biomarkers for lysine and threonine deficiency, respectively (Moro et al. 2023, J nutr, 153:2571-2584).
- The purpose of this study was to develop metabolomic approaches to identify specific biomarkers for an EAA deficiency.


Methods



- Body weight and food intake (FI) were measured daily.
- 24h-urine was analyzed LC-MS metabolomic.
- Body composition was determined by EchoMRI.



- Body weight was analyzed by repeated measures, mixt model and other variables by one-way ANOVA.
- Metabolic features were analyzed by PLS-DA and individually test for diet effects by ANOVA.

Results



Our results confirmed that protein and EAA deficiency induced growth retardation and supplementation permits to restore growth, but a delay of length, lean body mass and specific organs weights remains after supplementation.

- The urinary metabolome allowed to discriminate between the deficient and no deficient diet, and we were able to identify specific signatures for methionine and lysine deficiency.
- The best model retained 4 latent variables with LV3 allowing the discrimination of methionine deficient diets and LV4 allowing the discrimination of lysine deficient diets.
- Further analyses are required to investigate the specific signature for threonine intake.



Urinary metabolites from lysine degradation pathway, particularly pipecolate and N-N-N-Trimethyllysine, signed lysine deficiency.

Conclusion

- Our results showed that EAA deficiencies influence the urinary metabolome.
- We identified specific urinary metabolomic for lysine and methionine deficiency.
- We confirmed that pipecolate is a urinary biomarker that specifically signs lysine deficiency.
- Our results showed that the deficiency/supplementation method could be applied to identify specific EAA biomarkers.
- The urinary biomarkers identified could be easily applied to detect EAA or protein status.

Contact

- dalila.azzout-marniche@agroparistech.fr
- gaetan.hamelin@agroparistech.fr
- delphine.bouveresse@agroparistech.fr