



Defining a dichotomous indicator for population-level assessment of dietary diversity among pregnant adolescent girls and women : a secondary analysis of quantitative 24-h recalls from rural settings in Bangladesh, Burkina Faso, India and Nepal

Eric O Verger, Sabrina Eymard-Duvernay, Dang Bahya-Batinda, Giles T Hanley-Cook, Alemayehu Argaw, Elodie Becquey, Loty Diop, Aulo Gelli, Helen Harris-Fry, Shivani Kachwaha, et al.

► To cite this version:

Eric O Verger, Sabrina Eymard-Duvernay, Dang Bahya-Batinda, Giles T Hanley-Cook, Alemayehu Argaw, et al.. Defining a dichotomous indicator for population-level assessment of dietary diversity among pregnant adolescent girls and women : a secondary analysis of quantitative 24-h recalls from rural settings in Bangladesh, Burkina Faso, India and Nepal. *Current Developments in Nutrition*, 2024, 8 (1), pp.102053. 10.1016/j.cdnut.2023.102053 . hal-04322954

HAL Id: hal-04322954

<https://hal.inrae.fr/hal-04322954>

Submitted on 5 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Defining a dichotomous indicator for population-level assessment of dietary diversity among pregnant adolescent girls and women: a secondary analysis of quantitative 24-h recalls from rural settings in Bangladesh, Burkina Faso, India and Nepal

Eric O. Verger, Sabrina Eymard-Duvernay, Dang Bahya-Batinda, Giles T. Hanley-Cook, Alemayehu Argaw, Elodie Becquey, Loty Diop, Aulo Gelli, Helen Harris-Fry, Shivani Kachwaha, Sunny S. Kim, Phuong Hong Nguyen, Naomi M. Saville, Lan Mai Tran, Rock R. Zagr , Edwige Landais, Mathilde Savy, Yves Martin-Prevel, Carl Lachat

PII: S2475-2991(23)26637-8

DOI: <https://doi.org/10.1016/j.cdnut.2023.102053>

Reference: CDNUT 102053

To appear in: *Current Developments in Nutrition*

Received Date: 19 September 2023

Revised Date: 21 November 2023

Accepted Date: 26 November 2023

Please cite this article as: E.O. Verger, S. Eymard-Duvernay, D. Bahya-Batinda, G.T. Hanley-Cook, A. Argaw, E. Becquey, L. Diop, A. Gelli, H. Harris-Fry, S. Kachwaha, S.S. Kim, P.H. Nguyen, N.M. Saville, L.M. Tran, R.R. Zagr , E. Landais, M. Savy, Y. Martin-Prevel, C. Lachat, Defining a dichotomous indicator for population-level assessment of dietary diversity among pregnant adolescent girls and women: a secondary analysis of quantitative 24-h recalls from rural settings in Bangladesh, Burkina Faso, India and Nepal, *Current Developments in Nutrition*, <https://doi.org/10.1016/j.cdnut.2023.102053>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Defining a dichotomous indicator for population-level assessment of dietary diversity among pregnant adolescent girls and women: a secondary analysis of quantitative 24-h recalls from rural settings in Bangladesh, Burkina Faso, India and Nepal

Author names and affiliations

Eric O. Verger^a, Sabrina Eymard-Duvernay^a, Dang Bahya-Batinda^a, Giles T. Hanley-Cook^b, Alemayehu Argaw^{b,c}, Elodie Becquey^d, Loty Diop^d, Aulo Gelli^e, , Helen Harris-Fry^f, Shivani Kachwaha^g, Sunny S. Kim^e, Phuong Hong Nguyen^e, Naomi M. Saville^h, Lan Mai Tranⁱ, Rock R. Zagréd^d, Edwige Landais^a, Mathilde Savy^a, Yves Martin-Prevel^a and Carl Lachat^b.

^a MoISA, Univ Montpellier, CIRAD, CIHEAM-IAMM, INRAE, Institut Agro, IRD, Montpellier, France; ^b Department of Food Technology, Safety and Health, Faculty of Bioscience Engineering, Ghent University, Ghent, Belgium; ^c Department of Population and Family Health, Institute of Health, Jimma University, Jimma, Ethiopia; ^d International Food Policy Research Institute (IFPRI), Dakar, Senegal; ^e International Food Policy Research Institute (IFPRI), Washington, DC, USA; ^f Department Population Health, London School of Hygiene & Tropical Medicine, London, UK; ^g Johns Hopkins University, Baltimore, Maryland, USA; ^h UCL Institute for Global Health, London, United Kingdom; ⁱ Emory University, Atlanta, Georgia, USA

Corresponding author

Eric O. Verger, MoISA, IRD, 911 avenue d'Agropolis, 34000 Montpellier, France. Email: eric.verger@ird.fr

Abstract (300 words maximum)

Background: The Minimum Dietary Diversity for Women of Reproductive Age (MDD-W) indicator was validated as a proxy of micronutrient adequacy among non-pregnant women in low- and middle-income countries (LMICs). At that time, indeed, there was insufficient data to validate the indicator among pregnant women, who face higher micronutrient requirements.

Objective: This study aimed to validate a minimum food group consumption threshold, out of the 10 food groups used to construct MDD-W, to be used as a population-level indicator of higher micronutrient adequacy among pregnant women aged 15-49 years in LMICs.

Methods: We used secondary quantitative 24-hour recall data from 6 surveys in 4 LMICs (Bangladesh, Burkina Faso, India and Nepal, total n=4909). We computed the 10-food group Women's Dietary Diversity Score (WDDS-10) and calculated the mean probability of adequacy (MPA) of 11 micronutrients. Linear regression models were fitted to assess the associations between WDDS-10 and MPA. Sensitivity, specificity and proportion of individuals correctly classified were used to assess the performance of MDD-W in predicting an MPA >0.60.

Results: In the pooled sample, median values (interquartile range) of WDDS-10 and MPA were 3 (1) and 0.20 (0.34), respectively, while the proportion of pregnant women with an MPA >0.60 was 9.6%. The WDDS-10 was significantly positively associated with MPA in each survey. Although the acceptable food group consumption threshold varied between 4 and 6 food groups across surveys, the threshold of 5 showed the highest performance in the pooled sample with good sensitivity (62%), and very good specificity (81%) and percentage of correctly classified individuals (79%).

Conclusions: The WDDS-10 is a good predictor of dietary micronutrient adequacy among pregnant women aged 15-49 years in LMICs. Moreover, the threshold of 5 or more food group for the MDD-W indicator may be extended to all women of reproductive age, regardless of their physiological status.

Teaser Text

This study aimed to validate whether the threshold of 5 or more food group for the MDD-W indicator can to be used among pregnant women aged 15-49 years in low- and middle-income countries.

Abbreviations:

BLUP: Best linear unbiased predictor

EAR: Estimated Average Requirement

LMIC: Low- and Middle-Income Country

MDD-W: Minimum Dietary Diversity for Women

MPA: Mean probability of adequacy

PA: Probability of adequacy

WDDS-10: 10-food group Women's Dietary Diversity Score

Keywords: dietary diversity; indicator; micronutrient adequacy; minimum dietary diversity for women; pregnant; resource-poor settings.

Introduction

Micronutrients are essential vitamins and minerals whose subclinical deficiencies contribute to an increased risk of morbidity and mortality (1). A recent analysis suggested that two-thirds of non-pregnant women of reproductive age have one or more micronutrient deficiencies worldwide, with higher prevalence in low- and middle-income countries (LMICs) (2). There are important changes in dietary requirements driven by physiological processes during pregnancy, including increased requirements for folate, iron, vitamin B12 and B6, and zinc (3). These deficiencies are exacerbated during pregnancy due to an additional demand for nutrients to support both fetal growth and development and maternal metabolism (4), and can result in adverse outcomes of pregnancy and birth (5), as well as maternal depression and cognitive impairment (6).

Dietary diversification is a food-based strategy that has been widely promoted to address micronutrient deficiencies (7). To help achieve healthy diets, eating a diversity of foods is needed to help achieve healthy diets (8,9) as recommended by most dietary guidelines (10). As a result, a large range of interventions and programmes to improve nutrition through dietary diversification has been developed, and has subsequently triggered a demand for a set of harmonized indicators to monitor progress. Subsequently, several simple indicators assessing dietary diversity were developed, primarily for use in global and national monitoring, and in survey contexts where more detailed dietary methods that include estimation of food quantities are infeasible.

In this context, the Women's Dietary Diversity Project developed and validated simple food group indicators with consistent and relevant meaning across different contexts and over time. The most recent example is the Minimum Dietary Diversity for Women of Reproductive Age (MDD-W), a simple population-level dichotomous indicator expressed as the proportion of non-pregnant women of reproductive age who consumed at least 5 out of 10 defined food

groups over the previous 24 hours (11). MDD-W was validated using nine datasets from 6 distinct LMICs as a proxy for a minimally acceptable level of intake adequacy of 11 micronutrients among non-pregnant women of reproductive age (12,13). While the initial MDD-W validation study was able to assess the performance of the indicator for non-pregnant non-lactating women and non-pregnant lactating women, this was not possible for pregnant women due to the lack of data. Recent studies have used the threshold of 5 or more food groups to determine whether pregnant women had more adequate micronutrient intakes but without further validation of this dichotomous indicator in this population group (14–16). However, pregnant women generally have higher micronutrient requirements than non-pregnant women (3), which may change the performance of food group indicators in predicting adequate micronutrient adequacy in this specific population. The only validation study among pregnant women we are aware of showed that an adapted 6 or more food group threshold markedly improved performance of the indicator in predicting micronutrient adequacy among pregnant girls and pregnant women in Bangladesh (17). Using secondary quantitative 24-hour recall data from 6 surveys in 4 LMICs, this study aimed to validate a minimum food group consumption threshold, out of the 10 food groups used to construct MDD-W, to be used as a population-level indicator of higher micronutrient adequacy among pregnant women aged 15-49 years in LMICs. We followed the methods used by previous studies on the development and validation of MDD-W to ensure comparability of the analysis and facilitate the interpretation of findings (12,13).

Methods

Selection of surveys

This study was based on a pre-identified set of datasets that was completed by a systematic review of studies which collected dietary intakes from pregnant adolescent girls and women

in LMICs, using one or multiple 24-hour dietary recalls. Inclusion criteria were: (i) food consumption data collected among pregnant women (15–49 y) in LMICs; (ii) quantitative dietary intake data collected through one or multiple 24-hour dietary recalls; (iii) use of relevant local food composition data with information on the 11 micronutrients included in the initial development and validation of MDD-W (vitamin A expressed in retinol activity equivalents (RAE), thiamin, riboflavin, niacin, vitamin B6, folate, vitamin B12, vitamin C, calcium, iron and zinc); (iv) minimum sample size of 100 pregnant adolescent girls and women and (v) repeated 24-hour dietary recalls from at least 10% of the study sample or being able to be matched with a relevant dietary intake survey with two non-consecutive days of recall to estimate external within-person variance.

Study design and participants

Six datasets with quantitative 24-hour recall data collected from rural areas in Bangladesh in 2015 (17), Burkina Faso in 2017/2019/2020 (BF1, (18)), 2020 (BF2, (19)) and 2019/2021 (BF3, (20)), India in 2019 (21) and Nepal in 2015 (22) were selected for analysis. Each dataset is described in more detail in **Supplemental table 1**, which includes their selection process. Briefly, there were 5 pre-identified datasets (Bangladesh, BF1, BF2, BF3 and India) and we undertook a literature research to identify others, leading to add the dataset from Nepal. The included studies' primary objectives were to assess the feasibility and impact of maternal nutrition packages or integrated agriculture-nutrition interventions (Bangladesh, BF1, BF3 and India), to assess the efficacy of fortified balanced energy-protein supplementation (BF2), or to characterize the status and determinants of intra-household food and nutrient allocation, and test the effect of pregnancy interventions upon dietary intake (Nepal). None of the study samples was nationally representative. Data quality control was carried out by the data providers, including the exclusion of outliers. The representativeness

of each sample has been discussed in the original articles and primary study protocols for all sites were approved by ethical review committees or institutional review boards (17–19,21–25).

Dietary data collection

In all studies, dietary data were collected using one to three quantitative multiple-pass 24-hour dietary recalls conducted by enumerators specially trained for this purpose (26). Participants were asked to describe all foods and beverages consumed during the preceding 24 hours. Recipes were usually collected from the household member who was responsible for cooking. Portion sizes were estimated using methods best suited to local foods and contexts (e.g. previously distributed plates and bowls, common household measures, water volume, rice, images, clay or wooden models, etc.). Only two datasets had repeated 24-hour dietary recalls on non-consecutive days, with two recalls for 19% of the sample (BF1) and three recalls for 87% of the sample (Nepal). Dietary data were converted into nutrient intakes using country specific food composition tables; the application of yields and nutrient retention factors was done by data providers according to their own practice and information is available from original studies (17–19,21–23).

MDD-W and WDDS-10

Among the various indicators with different food groupings developed and tested as part of the Women's Dietary Diversity Project I and II, the dichotomous MDD-W indicator has been shown to have a strong relationship to micronutrient adequacy and high consistency in terms of threshold which best discriminated higher versus lower micronutrient adequacy across various countries (12,27). The MDD-W was constructed considering 10 mutually exclusive food groups consisting of: 1) starchy staple foods, 2) pulses; 3) nuts and seeds; 4) dairy

products; 5) flesh foods; 6) eggs; 7) dark green leafy vegetables; 8) vitamin A-rich fruits and vegetables; 9) other vegetables; and 10) other fruits. The 10 food groups are summed into a score (WDDS-10) ranging from 0 to 10, starting with a score of 0 and adding 1 point per food group consumed (if the total consumption of the foods in the food group was at least 15 g/day)¹. The WDDS-10 was computed using a single day recall (the first day in case of repeated recalls). MDD-W was coded as 1 if WDDS-10 reached 5 food groups or more, and 0 if 4 or lower.

Micronutrient requirements, usual intakes and probability of adequacy

We used the Estimated Average Requirements (EAR) and coefficients of variations proposed by Nguyen *et al.* (17), that are based on the information from the WHO/FAO (29), the National Academy of Medicine (formerly the Institute of Medicine) (30,31) and the International Zinc Nutrition Consultative Group (IZiNCG) (32). These requirements were used regardless of the pregnancy trimester, age or country context of the participants (**Supplemental table 2**). These requirements were chosen rather than those proposed by Allen *et al.* (33) to enhance comparability and facilitate interpretation of findings with previous studies on the development and validation of MDD-W (12,13,17).

Analogous to previous studies on the development and validation of MDD-W (12,13), we used the probability approach to estimate the micronutrient adequacies of each of the 11 micronutrients (28). This approach is based on information or assumption about both the distribution of nutrient requirements in the population and the day-to-day variations (within-person) of nutrient intakes. We applied a Box-Cox transformation to the nutrient intake

¹ This is of course not easy to do in practice, when collecting data; therefore, what is recommended in the FAO MDD-W guidelines is to apply the 15g limit to each food. However, we decided here to stick to the methodology used for the validation of the MDD-W for the sake of comparability.

distribution of every micronutrient to obtain normal distributions. For each participant and micronutrient in each separate dataset, we calculated the best linear unbiased predictor (BLUP) of the individual's usual intake (34) which was then used to calculate the probability of adequacy for every micronutrient (see **Supplemental Methods**). All usual nutrient intakes have been calculated solely on the basis of food intakes, excluding intakes from food supplements (e.g. fortified balanced energy-protein supplementation in BF2). When datasets contained repeated 24-hour dietary recalls, the within person variance was defined as the mean of squared intra-individual SDs. When datasets contained only one 24-hour dietary recall, we used an external within-person variance estimate from a relevant dietary intake survey with two non-consecutive days of recall (35,36). We used the external within-person variance to between-person variance ratio multiplied by the between-person variance of our dataset as the within-person variance in the BLUP calculations. A relevant dietary intake survey was defined as a survey conducted in the same geographical and seasonal context among pregnant adolescent girls or women. For Bangladesh, we used the within-person variance estimate from a subsample of the baseline study (~20%) that also participated in the endline study conducted a year later (37). For BF2 and BF3, we used the within-person variance estimate from BF1 due to the fact that these three surveys were conducted in the same context (Boucle du Mouhoun, Centre-Ouest and Haut-Bassins for BF1, Haut-Bassins for BF2 and Boucle du Mouhoun for BF3) among pregnant adolescent girls and women. For India, we used the within-person variance estimate from repeated 24-hour dietary recall used to validate a Food Frequency Questionnaire among pregnant women living with or without HIV in Pune, India (38).

Probability of adequacy (PA) was calculated as the probability that a woman's usual intake was at or above the EAR during pregnancy (28). For each individual, we averaged the mean

of the individual PAs for the 11 micronutrients to form the mean probability of adequacy (MPA). Like individual PAs, the MPA has a possible range of 0–1.

Data analysis

Data were analyzed with Stata 17 (Statacorp, College Station, TX) and the Stata syntax that was used for MDD-W validation in non-pregnant women (12,13), with a few minor revisions to match the aims of our analyses. Descriptive statistics are reported as medians (interquartile ranges) due to skewness of the distributions, except for age, height, weight and energy intake, which are reported as means (SDs). Associations between the WDDS-10 and MPA (with or without adjustment for total energy intake) were assessed by fitting simple linear regressions. For the pooled sample, a mixed-effects regression model was used to examine the association between WDDS-10 and MPA, with random effect at dataset level to take into account the within-survey correlation. The MPA variable was previously transformed by BoX-Cox transformation for all the regression models.

We used receiver operating characteristic (ROC) analysis and area under the curve (AUC) to assess the diagnostic performance of WDDS-10 in predicting a MPA >0.60 , with an AUC >0.70 deemed acceptable for predictive capacity. We estimated sensitivity, specificity and percentage of correct classifications for MDD-W across datasets and in a pooled analysis. The MPA level of 0.60, as well as the interpretation thresholds, were selected to ensure comparability with the previous analysis used to validate the MDD-W (12,13). Sensitivity (i.e. ability to correctly detect a person with an MPA >0.60) is defined by the ratio between the true positives and the sum of true positives and false negatives. Specificity (i.e. ability to correctly detect a person with an MPA ≤ 0.60) is defined by the ratio between the true negatives and the sum of true negatives and false positives. A threshold was considered good when both sensitivity and specificity were >0.60 and it was considered fair enough if only one

test characteristic was >0.60 and the other >0.50 . Moreover, while we looked for the best balance between sensitivity and specificity, we favored specificity over sensitivity when trade-offs must be made, in order to be certain to identify the highest proportion of participants with a MPA ≤ 0.60 . The percentage of correct classifications is defined by the ratio between the sum of true positives and true negatives and the sum of true positives, false positives, true negatives and false negatives. A threshold was considered as good when the percentage of individuals correctly classified was >0.70 and it was considered fair enough if >0.60 .

In order to understand the implications of some methodological choices, we conducted additional robustness analyses to estimate sensitivity, specificity and the percentage of correct classifications for MDD-W across datasets and in a pooled analysis according to three distinct scenarios. In the first robustness analysis, we tested 3 scenarios (Sc1, Sc2 and Sc3) where only 1 of the 3 Burkinabe datasets was included in the pooled analysis (BF1, BF2 and BF3, respectively), in order to keep into account the potentially redundant nature of using three surveys from Burkina Faso. In the second robustness analysis, we used the same recommendations from WHO/FAO (29), the National Academy of Medicine (30,31) and the IZiNCG (32) but took into account pregnancy trimester, age and level of bioavailability of iron and zinc (see **Supplemental table 3**). In the third robustness analysis, we used the requirements proposed by Allen *et al.* (33) which take into account age and level of bioavailability of iron and zinc but not pregnancy trimester (see **Supplemental table 4**).

Results

Characteristics of participants

Data were available for 4909 pregnant adolescent girls and women (**Table 1**), with sample sizes of the datasets ranging from 452 (BF1) to 1912 (BF3). The mean (SD) age of

participants was 25.7 (6.2) years, with participants from Nepal being on average younger than pregnant women from other countries. The inclusion of adolescent girls (15-18 years) across studies varied from none (India) to up to 26% (Bangladesh), and was 7.1% in the pooled sample. The pregnancy trimester distribution was highly variable across datasets, with a near-even distribution in BF1, whereas almost all participants were in their third trimester in Nepal. Participants in their third trimester represented almost 60% of the pooled sample. Pregnant women in the Burkinabé datasets were on average taller and heavier than participants from other countries.

Dietary diversity

The median (interquartile range) WDDS-10 in the pooled sample was 3 (1), with higher median scores in the Bangladeshi, Nepalese, and Indian datasets compared to the three Burkinabe datasets (**Table 1**). **Figure 1** shows the percentage of pregnant adolescent girls and women consuming each of the 10 food groups used to construct MDD-W across the six datasets. Consistently across datasets, the diet of all participants was based on starchy staple foods. Most participants consumed other vegetables, but with large variations ranging from 55% in BF1 and BF3 to 91% in Nepal. The prevalence of participants consuming pulses and dairy products greatly differed across datasets: for pulses it was high in Nepal (over 80%), moderate in Bangladesh and India (59 and 46%, respectively), and low in the three Burkinabe datasets (27% in BF1, 14% in BF2 and 15% in BF3). As for the prevalence of consumption of dairy products, it was very high in India (over 80%), moderate in Nepal and Bangladesh (53 and 33%, respectively), and low in the three Burkinabe datasets (4% in BF1, 3% in BF2 and 11% in BF3). In contrast, the prevalence of participants consuming nuts and seeds, and dark green leafy vegetables was higher in the three Burkinabe datasets. The prevalence of

participants consuming flesh foods, eggs, and other fruits was higher in the Bangladeshi datasets.

Energy and nutrient intakes and the probability of adequacy

The mean (SD) energy intake of the pregnant adolescent girls and women was 2068 (969) kcal per day in the pooled sample (**Table 2**), ranging from 1816 (838) kcal in BF3 to 2473 (1482) kcal in BF2. For all micronutrients apart from zinc, median intakes in the pooled sample were below the EAR (**Supplemental table 5**). However, there were differences between datasets, with median intakes in the Nepalese and Bangladeshi datasets above the EAR for 5 and 4 micronutrients, respectively. Accordingly, PAs varied widely across datasets (**Table 2**). Across surveys, the PAs of vitamin A, riboflavin, folate, vitamin B12, calcium, and iron were <0.50. The median (IQR) MPA of the participants was 0.20 (0.34) in the pooled sample, ranging from 0.09 (0.21) in BF1 to 0.43 (0.32) in Nepal. The proportion of participants with MPA above the threshold of 0.60 was low, at 9.6% in the pooled sample and ranged from 2.4% (BF1) to 23.4% (Nepal).

Association between WDDS-10 and MPA

Figure 2 illustrates non-adjusted associations between WDDS-10 and MPA (see **Supplemental table 6** for details of the number of pregnant women consuming various numbers of food groups by dataset). The WDDS-10 was significantly and positively associated with the MPA in every dataset (all $P < 0.001$) (**Table 3**). Unadjusted regression coefficients ranged from 0.079 (95% CI: 0.070, 0.088) to 0.309 (95% CI: 0.250, 0.367) and was 0.168 (95% CI: 0.157, 0.178) for the pooled sample. The unadjusted models explained between 14% and 33% of the MPA variance, and 28% in the pooled sample. In models including total energy intake (kcal/d) as covariate, associations were attenuated in all datasets

but remained highly significant. Energy adjusted regression coefficients ranged from 0.038 (95% CI: 0.028, 0.050) to 0.166 (95% CI: 0.114, 0.218) and was 0.079 (95% CI: 0.069, 0.088) in the pooled sample. The energy adjusted models explained between 29% and 66% of the MPA variance, and 41% in the pooled sample.

Food group indicator performance and identification of thresholds

The AUC value in the pooled sample was 0.78 (95% CI: 0.75, 0.80), which indicates an acceptable predicting power, and ranged from 0.61 to 0.81 across datasets which indicates a low to good performance in predicting a MPA >0.60, except for BF1 where the 95% CI (0.43, 0.78) included 0.50 which indicates no statistically significant predictive power (**Table 4**). In the sensitivity and specificity analyses in the pooled sample, the threshold of WDDS-10 ≥ 5 food groups had the best performances in predicting an MPA >0.60 (i.e. both sensitivity and specificity >0.60 and percentage of individuals correctly classified >0.70) with good sensitivity (62%) and very good specificity (81%) and percentage of individuals correctly classified (79%). The threshold of ≥ 4 food groups showed slightly lower performances with very good sensitivity (84%), but fair enough specificity (55%) and a moderate percentage of correctly classified participants (58%). The threshold of ≥ 6 food groups had lower performances with low sensitivity (32%), but very good specificity (93%) and percentage of correctly classified participants (87%). The other thresholds had worse classification properties. However, findings were heterogeneous across datasets. In summary, when balancing sensitivity, specificity and percentage of correct classification, the most acceptable food group consumption threshold for predicting a MPA >0.60 was WDDS-10 ≥ 4 in BF1, BF2, and BF3, ≥ 5 in India and Nepal, and ≥ 6 in Bangladesh. The three distinct scenarios from our robustness analyses returned similar findings, confirming both the observed heterogeneity across countries and also that the threshold of

WDDS-10 ≥ 5 food groups had the best performance in predicting an MPA >0.60 in the pooled sample (data not shown).

Discussion

Following the approach used for developing and validating MDD-W among non-pregnant women (12,13), we analyzed six dietary datasets to determine the minimum number of food groups consumed, out of the 10 food groups of the MDD-W, which best discriminates between higher versus lower micronutrient adequacy among pregnant adolescent girls and women in four LMICs. At least half of the women in each dataset had PAs of six micronutrients at zero, highlighting the urgency of an emphasis on diet quality and nutrient adequacy population group. Consequently, pregnant adolescent girls and women had low nutrient adequacy, with median MPA values ranging from 0.09 to 0.43 across the datasets. These findings are consistent with those reported among lactating women, who also face higher nutrient requirements, where the MPA ranged from 0.23 to 0.50 in nine datasets from resource-poor settings (12,13). As with other population subgroups (12,13), the WDDS-10 was significantly and positively associated with MPA in each dataset. Similarly to the results found during the initial validation of MDD-W for non-pregnant women (12,13), our analyses showed that across the pooled sample a threshold of 5 or more food groups had the best performance in classifying pregnant adolescent girls and women as having a minimally acceptable level of dietary micronutrient adequacy (i.e., MPA >0.60). Nevertheless, we found evidence of heterogeneity across datasets, both in terms of dietary patterns and in the optimal threshold of WDDS-10 to predict a minimally acceptable level of micronutrient adequacy (which varied from 4 to 6). Pulses and dairy were more commonly consumed in South Asian countries, whereas nuts, seeds, and green leafy vegetables were more commonly consumed in Burkina Faso. This could be explained by geographical and

temporal differences, such as food availability, prices, budgets, and preferences. For example, each dataset only captured certain months of the year while seasonality could affect food availability and thus dietary diversity in these contexts (39,40). In terms of differences in thresholds, it should be noted that even in the validation study that led to adopt the MDD-W there were differences across datasets regarding the best threshold that predicted a $MPA > 0.60$ – which varied from 4 to 6 as in the present study (12). Various food (sub)groups contribute more or less to the MPA than others and/or can be consumed in larger or smaller quantities according to the context. This heterogeneity is not specific to pregnant adolescent girls and women. When recommending the threshold of 5 food groups, that work best in the pooled sample in this study as well as across the 9 datasets of the MDD-W validation study (12), we are pretty confident that this threshold would most likely minimize the gap to the true, context-specific and also probably season-specific optimal threshold that remains unknown in many contexts but was found in the range of 4 to 6 in most if not all published studies (12,17,41).

Measuring characteristics of diets and monitoring of their changes at global and national levels are needed to support governments in establishing policies and programmes to promote healthy diets, to assess the effectiveness of their actions and hold them accountable. This is the spirit behind the development of the MDD-W (12,13). Although MDD-W is already widely collected in large multi-topic surveys, such as Demographic and Health Surveys and Gallup World Poll, it only reflects dietary diversity which is one, albeit indispensable, subconstruct of healthy diets (42,43). Other promising metrics were recently designed to assess in a synthetic manner several subconstructs of healthy diets. The Global Diet Quality Score (GDQS), for example, is based on the consumption of 25 food groups that are globally important contributors to nutrient intake, on the one hand, and/or to non-communicable disease risk, on the other hand (44). Although it has been validated using several datasets

from various contexts, the validation was performed against several outcomes and by comparisons with the performance of other metrics and not directly to nutrient adequacy. In addition, the GDQS has not yet been widely used in large surveys, probably because some appraisal of quantities or portions consumed is needed for its construction. The Global Dietary Recommendations (GDR) score is another recently developed synthetic metric that was designed to assess the adherence to a dietary pattern respecting 11 global dietary recommendations from WHO, which include dietary factors protective against non-communicable diseases (45). Although the construction of the GDR score is based on a standardized Diet Quality Questionnaire that was validated against 24h-recalls in three different contexts, and has been used since in many other countries, as far as we know the GDR score itself was validated only with data from Brazil and the USA. Additional evidence are needed to establish its validity in various contexts and its equivalence across contexts (43). Thus, MDD-W arguably remains a statically robust and valid indicator, widely collected in large multi-topic surveys, to assess dietary diversity as a cornerstone of diet quality on a global and national scale. This work contributes to ongoing efforts to validate MDD-W in other populations such as adolescents and children (43).

The present analyses have some limitations. First, despite our efforts to obtain datasets from a diversity of contexts, our study only includes data from rural contexts in four LMICs among two regions (sub-Saharan Africa and South Asia). Although our findings are not globally representative, they are consistent with other analyses among non-pregnant women from more settings (12,13). Furthermore, the rural locations included in our study are settings where valid scores are arguably much needed, as they typically have a high burden of undernutrition and low dietary diversity (15,39,46,47). In the meantime, more datasets should be made available in settings where a reasonable proportion of pregnant adolescent girls and women reach an acceptable MPA, so that the best predictors of acceptable MPA can be further

studied. For example, in the BF1 sample of our study, only 11 (2.4%) pregnant women reached an MPA ≥ 0.60 , which strongly limits the search for the best dichotomous indicator predicting higher MPA. Another limitation concerns the use of an external within-person variance estimate to calculate the MPA in four of the six datasets. This results in more reliable prevalence estimates than when using a single day recall (36), but the use of within-person variance estimates from repeated measures within the samples is preferable (35). Although we tried to find and use an external estimate of within-person variance from a relevant food intake survey, we were limited in our ability to find studies with the same geographical (e.g. for India, the region of the external estimate study is 1500 km away from that of the dataset) or temporal (different seasonality between BF1 and BF3) characteristics. Future analyses from a wider variety of settings and with data containing repeated measures is recommended to confirm that a threshold of 5 or more groups is the best suited to indicate MPA > 0.60 . A last limitation is the use of a set of nutrient requirements which did not take into account the pregnancy trimester, the age of the participants or the level of bioavailability of iron and zinc. This simpler approach was preferred to take into account the fact that this information might not be accurately collected in large surveys. Nevertheless, taking these characteristics into account in three distinct robustness analyses did not affect our findings in terms of determining the threshold of WDDS-10 with the best classification characteristics.

In conclusion, our study suggests that the WDDS-10 is a good predictor of dietary micronutrient adequacy among pregnant adolescent girls and women in LMICs, as it was previously shown among non-pregnant and non-lactating women and lactating women (12,13). When a dichotomous indicator is preferred over a continuous measure, our results suggest that the MDD-W may be used as a proxy indicator for higher micronutrient adequacy in LMIC contexts in all women of reproductive age, regardless of physiological status. This might be particularly useful for international comparisons and when the physiological status

of women is unknown, which is the case in many large surveys. However, our findings suggest that context-specific thresholds might be more accurate and might therefore be preferred for research purposes. Given the low micronutrient adequacy in the populations studied, additional efforts are needed to enhance the diet of women of reproductive age. Even though the threshold of 5 or more groups might not accurately predict micronutrient adequacy in all contexts, the indicator allows tracking processes of such efforts over time and enables benchmarking between populations. However, there is a need to provide complementary assessment of other dimensions of diet quality, such as consumption of undesired foods, food safety aspects, and within food group contribution of foods. In addition, in food environments and diets with a considerable contribution of fortified foods, the validity of the 5 food group thresholds might require careful reconsideration.

Acknowledgements

We are grateful to Kripa Rajagopalan (Cornell University) and Rupak Shivakoti (Columbia University) for sharing the within-person variance estimate from their FFQ validation study among pregnant women living with or without HIV in Pune, India. The authors' responsibilities were as follows: EO, SED, DBB, GTH-C, EL, MS, YMP, and CL designed the study; AA, EB, LD, AG, GTH-C, HH-F, SK, SSK, PHN, NMS, LMT and RRZ provided the datasets; DBB harmonized the datasets; SBB analyzed data; EO and SBB drafted the figures, tables, and manuscript, and the other authors provided critical review; all authors read and approved the final manuscript.

Data Availability

Data described in the manuscript, code book, and analytic code will be made available upon request pending application and approval by authors of the current study.

439

440 **Funding**

441 This publication has been produced with the financial support of the European Commission,
 442 under the project “Knowledge and research for nutrition” implemented by Agrinatura EEIG. Its
 443 contents are the sole responsibility of the authors and do not necessarily reflect the views of the
 444 European Union and of Agrinatura EEIG.

445

446 **Author Disclosures**

447 The authors report no conflicts of interest.

448

References

1. Bailey RL, West Jr. KP, Black RE. The Epidemiology of Global Micronutrient Deficiencies. *Ann Nutr Metab* 2015;66:22–33.
2. Stevens GA, Beal T, Mbuya MNN, Luo H, Neufeld LM, Addo OY, Adu-Afarwuah S, Alayón S, Bhutta Z, Brown KH, et al. Micronutrient deficiencies among preschool-aged children and women of reproductive age worldwide: a pooled analysis of individual-level data from population-representative surveys. *Lancet Glob Health Elsevier*; 2022;10:e1590–9.
3. Marshall NE, Abrams B, Barbour LA, Catalano P, Christian P, Friedman JE, Hay WW, Hernandez TL, Krebs NF, Oken E, et al. The importance of nutrition in pregnancy and lactation: lifelong consequences. *Am J Obstet Gynecol* 2022;226:607–32.
4. Abu-Saad K, Fraser D. Maternal Nutrition and Birth Outcomes. *Epidemiol Rev* 2010;32:5–25.
5. Keats EC, Haider BA, Tam E, Bhutta ZA. Multiple-micronutrient supplementation for women during pregnancy. *Cochrane Database Syst Rev* 2019;3:CD004905.
6. Bourassa MW, Osendarp SJM, Adu-Afarwuah S, Ahmed S, Ajello C, Bergeron G, Black R, Christian P, Cousens S, de Pee S, et al. Review of the evidence regarding the use of antenatal multiple micronutrient supplementation in low- and middle-income countries. *Ann N Y Acad Sci* 2019;1444:6–21.
7. Adrianopoli M, D’Acapito P, Ferrari M, Mistura L, Toti E, Maiani G, Thompson B, Amoroso L. Improving diets and nutrition: food-based approaches. Rome Italy CABI 2014;230–45.
8. Verger EO, Port AL, Borderon A, Bourbon G, Moursi M, Savy M, Mariotti F, Martin-Prevel Y. Dietary Diversity Indicators and Their Associations with Dietary Adequacy and Health

- Outcomes: A Systematic Scoping Review. *Adv Nutr* [Internet] 2021 [cited 2021 Apr 22]; Available from: <https://doi.org/10.1093/advances/nmab009>
9. World Health Organization. WHO recommendations on antenatal care for a positive pregnancy experience [Internet]. Geneva: World Health Organization; 2016 [cited 2023 Jun 23]. 152 p. Available from: <https://apps.who.int/iris/handle/10665/250796>
 10. Herforth A, Arimond M, Álvarez-Sánchez C, Coates J, Christianson K, Muehlhoff E. A Global Review of Food-Based Dietary Guidelines. *Adv Nutr Bethesda Md* 2019;10:590–605.
 11. Food and Agriculture Organization. Minimum dietary diversity for women: An updated guide to measurement-from collection to action [Internet]. Rome: FAO; 2021. Available from: <https://doi.org/10.4060/cb3434en>
 12. Martin-Prével Y, Allemand P, Wiesmann D, Arimond M, Ballard T, Deitchler M, Dop M-C, Kennedy G, Lee WT, Moursi M. Moving forward on choosing a standard operational indicator of women's dietary diversity. FAO; 2015.
 13. Women's Dietary Diversity Project (WDDP) Study Group. Development of a Dichotomous Indicator for Population-Level Assessment of Dietary Diversity in Women of Reproductive Age. *Curr Dev Nutr* 2017;1:cdn.117.001701-cdn.117.001701.
 14. Gyimah LA, Annan RA, Apprey C, Edusei A, Aduku LNE, Asamoah-Boakye O, Azanu W, Lutterodt H. Dietary diversity and its correlates among pregnant adolescent girls in Ghana. *PLOS ONE Public Library of Science*; 2021;16:e0247979.
 15. Shrestha V, Paudel R, Sunuwar DR, Lyman ALT, Manohar S, Amatya A. Factors associated with dietary diversity among pregnant women in the western hill region of Nepal: A community based cross-sectional study. *PLOS ONE Public Library of Science*; 2021;16:e0247085.
 16. Yang J, Wang M, Tobias DK, Rich-Edwards JW, Darling A-M, Abioye AI, Noor RA, Madzorera I, Fawzi WW. Dietary diversity and diet quality with gestational weight gain and adverse birth outcomes, results from a prospective pregnancy cohort study in urban Tanzania. *Matern Child Nutr* 2022;18:e13300.
 17. Nguyen PH, Huybregts L, Sanghvi TG, Tran LM, Frongillo EA, Menon P, Ruel MT. Dietary Diversity Predicts the Adequacy of Micronutrient Intake in Pregnant Adolescent Girls and Women in Bangladesh, but Use of the 5-Group Cutoff Poorly Identifies Individuals with Inadequate Intake. *J Nutr* 2018;148:790–7.
 18. Becquey E, Diop L, Awonon J, Diatta AD, Ganaba R, Pedehombga A, Gelli A. A Poultry Value Chain Intervention Promoting Diversified Diets Has Limited Impact on Maternal and Child Diet Adequacy during the Lean Season in a Cluster Randomized Controlled Trial. *J Nutr* 2022;152:1336–46.
 19. de Kok B, Argaw A, Hanley-Cook G, Toe LC, Ouédraogo M, Dailey-Chwalibóg T, Diop L, Becquey E, Kolsteren P, Lachat C, et al. Fortified Balanced Energy-Protein Supplements Increase Nutrient Adequacy without Displacing Food Intake in Pregnant Women in Rural Burkina Faso. *J Nutr* 2021;151:3831–40.
 20. Kim SS, Zagré RR, Ouédraogo CT, Sununtnasuk C, Ganaba R, Zafimanjaka MG, Tharaney M, Sanghvi T, Menon P. Intensified Nutrition Interventions in Antenatal Care Services Increased Consumption of Iron and Folic Acid Supplements and Early Breastfeeding Practices in Burkina Faso: Results of a Cluster-Randomized Program Evaluation. *J Nutr* 2023;S0022-3166(23)72422-X.

21. Nguyen PH, Kachwaha S, Tran LM, Avula R, Young MF, Ghosh S, Sharma PK, Escobar-Alegria J, Forissier T, Patil S, et al. Strengthening Nutrition Interventions in Antenatal Care Services Affects Dietary Intake, Micronutrient Intake, Gestational Weight Gain, and Breastfeeding in Uttar Pradesh, India: Results of a Cluster-Randomized Program Evaluation. *J Nutr* 2021;151:2282–95.
22. Harris-Fry HA, Paudel P, Harrisson T, Shrestha N, Jha S, Beard BJ, Copas A, Shrestha BP, Manandhar DS, Costello AM de L, et al. Participatory Women's Groups with Cash Transfers Can Increase Dietary Diversity and Micronutrient Adequacy during Pregnancy, whereas Women's Groups with Food Transfers Can Increase Equity in Intrahousehold Energy Allocation. *J Nutr* 2018;148:1472–83.
23. Kim SS, Ouédraogo CT, Zagré RR, Ganaba R, Zafimanjaka MG, Tharaney M, Menon P. Multiple modifiable maternal, household and health service factors are associated with maternal nutrition and early breastfeeding practices in Burkina Faso. *Matern Child Nutr* 2023;19:e13457.
24. Harris-Fry H, Beard BJ, Harrisson T, Paudel P, Shrestha N, Jha S, Shrestha BP, Manandhar DS, Costello A, Saville NM. Smartphone tool to collect repeated 24 h dietary recall data in Nepal. *Public Health Nutr* England; 2018;21:260–72.
25. Harris-Fry H, Paudel P, Karn M, Mishra N, Thakur J, Paudel V, Harrisson T, Shrestha B, Manandhar DS, Costello A, et al. Development and validation of a photographic food atlas for portion size assessment in the southern plains of Nepal. *Public Health Nutr* Cambridge University Press; 2016;19:2495–507.
26. Gibson RS, Ferguson EL. An interactive 24-hour recall for assessing the adequacy of iron and zinc intakes in developing countries. [Internet]. Washington (DC) and Cali : International Food Policy Research Institute (IFPRI) and International Center for Tropical Agriculture (CIAT). 2008 [cited 2023 Jan 17]. 157 p. Available from: <https://www.ifpri.org/publication/interactive-24-hour-recall-assessing-adequacy-iron-and-zinc-intakes-developing-countries>
27. Arimond M, Wiesmann D, Becquey E, Carriquiry A, Daniels MC, Deitchler M, Fanou-Fogny N, Joseph ML, Kennedy G, Martin-Prevel Y, et al. Simple food group diversity indicators predict micronutrient adequacy of women's diets in 5 diverse, resource-poor settings. *J Nutr* 2010;140:2059S–69S.
28. National Research Council. Dietary Reference Intakes: Applications in Dietary Assessment [Internet]. Washington, DC: The National Academies Press; [cited 2014 Mar 7]. Available from: http://www.nap.edu/catalog.php?record_id=9956
29. World Health Organization. Vitamin and mineral requirements in human nutrition. World Health Organization; 2004.
30. Institute of Medicine. Dietary Reference Intakes: The Essential Guide to Nutrient Requirements [Internet]. The national academies press. Washington, D.C.; 2006 [cited 2023 Feb 3]. Available from: <https://www.nap.edu/read/11537/chapter/1>
31. Institute of Medicine (US) Committee to Review Dietary Reference Intakes for Vitamin D and Calcium. Dietary Reference Intakes for Calcium and Vitamin D [Internet]. Ross AC, Taylor CL, Yaktine AL, Del Valle HB, editors. Washington (DC): National Academies Press (US); 2011 [cited 2023 Feb 3]. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK56070/>
32. International Zinc Nutrition Consultative Group (IZiNCG), Brown KH, Rivera JA, Bhutta Z, Gibson RS, King JC, Lönnerdal B, Ruel MT, Sandtröm B, Wasantwisut E, et al. International

- Zinc Nutrition Consultative Group (IZiNCG) technical document #1. Assessment of the risk of zinc deficiency in populations and options for its control. *Food Nutr Bull* 2004;25:S99-203.
33. Allen LH, Carriquiry AL, Murphy SP. Perspective: Proposed Harmonized Nutrient Reference Values for Populations. *Adv Nutr Bethesda Md* 2020;11:469–83.
 34. Joseph ML, Carriquiry A. A measurement error approach to assess the association between dietary diversity, nutrient intake, and mean probability of adequacy. *J Nutr* 2010;140:2094S-101S.
 35. French CD, Arsenault JE, Arnold CD, Haile D, Luo H, Dodd KW, Vosti SA, Slupsky CM, Engle-Stone R, The Variance Components of Nutrient Intakes Data Working Group. Within-Person Variation in Nutrient Intakes across Populations and Settings: Implications for the Use of External Estimates in Modeling Usual Nutrient Intake Distributions. *Adv Nutr* 2021;12:429–51.
 36. Jahns L, Arab L, Carriquiry A, Popkin BM. The use of external within-person variance estimates to adjust nutrient intake distributions over time and across populations. *Public Health Nutr Cambridge University Press*; 2005;8:69–76.
 37. International Food Policy Research Institute (IFPRI). A&T Bangladesh Maternal Nutrition Endline Survey 2016: Households - Pregnant Women [Internet]. Harvard Dataverse; 2021 [cited 2023 May 17]. Available from: <https://dataverse.harvard.edu/dataset.xhtml?persistentId=doi:10.7910/DVN/PYLIY4>
 38. Rajagopalan K, Alexander M, Naik S, Patil N, Mehta S, Leu C-S, Bhosale R, Mathad JS, Caulfield LE, Gupta A, et al. Validation of New Interactive Nutrition Assistant - Diet in India Study of Health (NINA-DISH) FFQ with multiple 24-h dietary recalls among pregnant women in Pune, India. *Br J Nutr* 2021;126:1247–56.
 39. Hanley-Cook GT, Argaw A, de Kok B, Toe LC, Dailey-Chwalibóg T, Ouédraogo M, Kolsteren P, Huybregts L, Lachat C. Seasonality and Day-to-Day Variability of Dietary Diversity: Longitudinal Study of Pregnant Women Enrolled in a Randomized Controlled Efficacy Trial in Rural Burkina Faso. *J Nutr* 2022;152:2145–54.
 40. Saville NM, Cortina-Borja M, De Stavola BL, Pomeroy E, Marphatia A, Reid A, Manandhar DS, Wells JC. Comprehensive analysis of the association of seasonal variability with maternal and neonatal nutrition in lowland Nepal. *Public Health Nutr* 2021;25:1–16.
 41. Diop L, Becquey E, Turowska Z, Huybregts L, Ruel MT, Gelli A. Standard Minimum Dietary Diversity Indicators for Women or Infants and Young Children Are Good Predictors of Adequate Micronutrient Intakes in 24-59-Month-Old Children and Their Nonpregnant Nonbreastfeeding Mothers in Rural Burkina Faso. *J Nutr* 2021;151:412–22.
 42. Seligman HK, Levi R, Adebiyi VO, Coleman-Jensen A, Guthrie JF, Frongillo EA. Assessing and Monitoring Nutrition Security to Promote Healthy Dietary Intake and Outcomes in the United States. *Annu Rev Nutr* 2023;
 43. Verger EO, Savy M, Martin-Préve Y, Coates J, Frongillo E, Neufeld L, Saha K, Hayashi C, Holmes B, Vogliano C. Healthy diet metrics: a suitability assessment of indicators for global and national monitoring purposes [Internet]. Geneva: World Health Organization; 2023. Available from: <https://apps.who.int/iris/bitstream/handle/10665/371497/9789240072138-eng.pdf>
 44. Bromage S, Batis C, Bhupathiraju SN, Fawzi WW, Fung TT, Li Y, Deitchler M, Angulo E, Birk N, Castellanos-Gutiérrez A, et al. Development and Validation of a Novel Food-Based Global Diet Quality Score (GDQS). *J Nutr* 2021;151:75S-92S.

45. Herforth AW, Wiesmann D, Martínez-Steele E, Andrade G, Monteiro CA. Introducing a Suite of Low-Burden Diet Quality Indicators That Reflect Healthy Diet Patterns at Population Level. *Curr Dev Nutr* 2020;4:nzaa168.
46. Harris-Fry H, Azad K, Kuddus A, Shaha S, Nahar B, Hossen M, Younes L, Costello A, Fottrell E. Socio-economic determinants of household food security and women's dietary diversity in rural Bangladesh: a cross-sectional study. *J Health Popul Nutr* 2015;33:2–2.
47. Lander R, Hambidge K, Westcott J, Tejeda G, Diba T, Mastiholi S, Khan U, Garces A, Figueroa L, Tshefu A, et al. Pregnant Women in Four Low-Middle Income Countries Have a High Prevalence of Inadequate Dietary Intakes That Are Improved by Dietary Diversity. *NUTRIENTS* 2019;11.

Figures

Fig. 1. Percentage of participants having consumed the 10 food groups used to construct MDD-W in the previous 24-hours. BF1, rural Burkina Faso dataset (2017/2019/2020); BF2, rural Burkina Faso dataset (2020); BF3, rural Burkina Faso dataset (2019/2021).

Fig. 2. Average mean probability of adequacy by WDDS-10 score. BF1, rural Burkina Faso dataset (2017/2019/2020); BF2, rural Burkina Faso dataset (2020); BF3, rural Burkina Faso dataset (2019/2021). Error bars represent mean \pm standard error. Data points representing <10 participants are not shown. Details of the number of pregnant women by dataset are given in Supplemental Table 4.

Tables

Table 1. Characteristics of pregnant women¹

Dataset	<i>n</i>	Repeated recall, <i>n</i> (%) ²	Mean (SD) age, <i>y</i>	Adolescent, <i>n</i> (%)	First trimester, <i>n</i> (%)	Second trimester, <i>n</i> (%)	Third trimester, <i>n</i> (%)	Mean (SD) height, m	Mean (SD) weight, kg	Median (IQR) WDDS-10
Bangladesh	598	0 (0.0)	24.0 (5.6)	160 (26.0)	0 (0.0)	328 (54.8)	270 (45.2)	1.50 (0.06)	50.3 (8.1)	5 (2)
BF1	452	84 (18.6)	29.6 (5.3)	1 (0.2)	124 (27.4)	173 (38.3)	155 (34.3)	1.61 (0.07)	59.1 (8.0)	3 (2)
BF2	470	0 (0.0)	25.4 (6.4)	37 (7.9)	16 (3.4)	188 (40.0)	266 (56.6)	1.63 (0.06)	58.9 (8.7)	3 (2)
BF3 ³	1912	0 (0.0)	27.5 (6.6)	64 (3.4)	279 (14.7)	828 (43.8)	785 (41.5)	1.63 (0.01)	61.8 (2.5)	3 (2)
India	674	0 (0.0)	25.0 (4.0)	0 (0.0)	0 (0.0)	198 (29.4)	476 (70.6)	1.50 (0.06)	51.0 (8.5)	4 (2)
Nepal	803	745 (92.8)	21.5 (3.8)	88 (11.0)	0 (0.0)	1 (0.1)	802 (99.9)	1.51 (0.05)	52.1 (6.5)	4 (1)
Pooled ³	4909	N/A	25.7 (6.2)	350 (7.1)	419 (8.5)	1716 (34.9)	2754 (56.1)	1.58 (0.07)	56.7 (8.0)	3 (1)

¹ BF1, rural Burkina Faso dataset (2017/2019/2020); BF2, rural Burkina Faso dataset (2020); BF3, rural Burkina Faso dataset (2019/2021); SD, standard deviation; WDDS-10, 10-food group women dietary diversity score; IQR, Interquartile range. ² Women in the sample with more than one 24-hour dietary recall; ³ Information about the pregnancy trimester was missing for 20 participants.

Table 2. Energy intakes, probability of adequacy of individual micronutrients and mean probability of adequacy (MPA)¹

Dataset	Energy intakes, kcal/d ²	Vitamin A ³	Thiamin ³	Riboflavin ³	Niacin ³	Vitamin B6 ³	Folate ³	Vitamin B12 ³	Vitamin C ³	Calcium ³	Iron ³	Zinc ³	MPA ³	MPA >0.60, <i>n</i> (%)
Bangladesh	2330 (822)	0.00 (0.60)	1.00 (0.22)	0.00 (0.25)	1.00 (0.00)	1.00 (0.00)	0.00 (0.00)	0.00 (0.07)	1.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.18 (0.70)	0.40 (0.19)	94 (15.7)
BF1	1950 (939)	0.00 (0.10)	0.00 (0.04)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.67)	0.00 (0.00)	0.00 (0.49)	0.40 (0.95)	0.09 (0.21)	11 (2.4)
BF2	2473 (1482)	0.00 (0.48)	0.00 (0.76)	0.00 (0.01)	0.00 (0.22)	0.00 (0.67)	0.00 (0.15)	0.00 (0.00)	0.01 (1.00)	0.00 (0.05)	0.00 (0.95)	0.97 (0.79)	0.16 (0.34)	69 (14.7)
BF3	1816 (838)	0.00 (0.00)	0.00 (0.01)	0.00 (0.00)	0.00 (0.03)	0.00 (0.04)	0.00 (0.00)	0.00 (0.00)	0.00 (0.05)	0.00 (0.01)	0.07 (1.00)	0.63 (0.95)	0.13 (0.21)	73 (3.8)
India	2122 (924)	0.00 (0.00)	0.80 (0.99)	0.00 (0.23)	0.20 (0.92)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.04 (1.00)	0.00 (0.08)	0.00 (0.00)	0.62 (0.97)	0.20 (0.32)	35 (5.2)
Nepal	2254 (850)	0.05 (0.46)	0.96 (0.76)	0.06 (0.99)	0.64 (0.83)	1.00 (0.36)	0.00 (0.01)	0.00 (0.00)	1.00 (0.17)	0.00 (0.45)	0.00 (0.00)	0.99 (0.34)	0.43 (0.32)	188 (23.4)
Pooled	2068 (969)	0.00 (0.08)	0.03 (0.98)	0.00 (0.05)	0.03 (0.91)	0.00 (0.99)	0.00 (0.00)	0.00 (0.00)	0.02 (1.00)	0.00 (0.02)	0.00 (0.26)	0.69 (0.93)	0.20 (0.34)	470 (9.6)

¹BF1, rural Burkina Faso dataset (2017/2019/2020); BF2, rural Burkina Faso dataset (2020); BF3, rural Burkina Faso dataset (2019/2021). ² Values are means (SD) calculated from a single 24-hour dietary recall (the first one in case of repetitions). ³ Values are medians (interquartile range).

Table 3. Linear regression of WDDS-10 with mean probability of adequacy^{1,2}


Dataset	Unadjusted			Total energy (kcal/d) adjusted			
	WDDS-10	Constant	Adjusted R ²	WDDS-10	Energy intake, kcal/d	Constant	Adjusted R ²
Bangladesh	0.079 (0.070, 0.088)	-1.06 (-1.11, -1.01)	0.333	0.055 (0.046, 0.063)	0.0001 (0.0001, 0.0001)	-1.23 (-1.27, -1.18)	0.529
BF1	0.252 (0.195, 0.310)	-2.42 (-2.60, -2.24)	0.142	0.125 (0.067, 0.183)	0.0003 (0.0003, 0.0004)	-2.65 (-2.82, -2.48)	0.291
BF2	0.309 (0.250, 0.367)	-2.20 (-2.38, -2.01)	0.185	0.166 (0.114, 0.218)	0.0002 (0.0002, 0.0003)	-2.37 (-2.52, -2.21)	0.431
BF3	0.214 (0.194, 0.233)	-2.06 (-2.13, -2.00)	0.198	0.091 (0.074, 0.108)	0.0004 (0.0003, 0.0004)	-2.40 (-2.45, -2.34)	0.488
India	0.162 (0.139, 0.186)	-1.73 (-1.83, -1.63)	0.214	0.049 (0.032, 0.067)	0.0003 (0.0003, 0.0004)	-2.00 (-2.07, -1.94)	0.662
Nepal	0.082 (0.068, 0.095)	-0.93 (-0.99, -0.87)	0.149	0.038 (0.028, 0.050)	0.0002 (0.0001, 0.0002)	-1.11 (-1.16, -1.06)	0.465
Pooled ³	0.168 (0.157, 0.178)	-1.74 (-1.96, -1.51)	0.286	0.079 (0.069, 0.088)	0.0003 (0.0002, 0.0003)	-2.03 (-2.27, -1.78)	0.411


¹ Values are regression coefficients and (95% Confidence Intervals); WDDS-10, 10-food group women dietary diversity score; CI, confident interval; BF1, rural Burkina Faso dataset (2017/2019/2020); BF2, rural Burkina Faso dataset (2020); BF3, rural Burkina Faso dataset (2019/2021). ² The mean probability of adequacy after Box-Cox transformation was used as dependent variable in all the regression models. All *P*-values are <0.001. ³ A mixed-effects regression model, including a random intercept for survey, was fitted for the pooled sample.

Table 4. Test characteristics of food group indicators for classifying mean probability of adequacy >0.60 for pregnant adolescents and women¹

Dataset	AUC	WDDS-10 ≥ 4			WDDS-10 ≥ 5			WDDS-10 ≥ 6		
		Sensitivity	Specificity	PCC	Sensitivity	Specificity	PCC	Sensitivity	Specificity	PCC
Bangladesh	0.81 (95% CI: 0.77, 0.85)	98.9	19.6	32.1	97.9	41.5	50.3	78.7	67.3	69.1
BF1	0.61 (95% CI: 0.43, 0.78)	54.6	69.4	69.0	9.10	93.4	91.4	0.00	98.9	96.5
BF2	0.71 (95% CI: 0.65, 0.78)	55.1	77.3	74.0	21.7	96.8	85.7	2.9	99.8	85.5
BF3	0.74 (95% CI: 0.69, 0.79)	63.0	70.3	70.0	31.5	91.3	89.0	17.8	98.4	95.3
India	0.79 (95% CI: 0.73, 0.86)	97.1	37.7	40.8	77.1	71.5	71.8	34.3	91.6	88.6
Nepal	0.74 (95% CI: 0.71, 0.78)	94.7	30.6	45.6	70.2	70.6	70.5	26.6	92.7	77.2
Pooled	0.78 (95% CI: 0.75, 0.80)	84.0	54.9	57.7	61.7	80.6	78.8	32.1	93.2	87.4

¹ Values are percentages (except for the AUC values); AUC, area under the curve; CI, confident interval; BF1, rural Burkina Faso dataset (2017/2019/2020); BF2, rural Burkina Faso dataset (2020); BF3, rural Burkina Faso dataset (2019/2021). PCC, percentage correctly classified; WDDS-10, 10-food group Women Dietary Diversity Score.





Declaration of interests

☐ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☒ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Eric Verger reports financial support was provided by European Commission.