



HAL
open science

Intraspecific diversity of Microbial Anti-inflammatory Molecule (MAM) from *Faecalibacterium prausnitzii*

Sandrine Auger

► **To cite this version:**

Sandrine Auger. Intraspecific diversity of Microbial Anti-inflammatory Molecule (MAM) from *Faecalibacterium prausnitzii*. BactInflam Webinar, Jun 2022, Jouy-en-Josas, France. hal-04328397

HAL Id: hal-04328397

<https://hal.inrae.fr/hal-04328397>

Submitted on 7 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Intraspecific diversity of Microbial Anti-inflammatory Molecule (MAM) from *Faecalibacterium prausnitzii*

Sandrine Auger¹, Camille Kropp¹, Esther Borrás-Nogues¹, Wasaporn Chanput¹, Gwenaëlle Andre-Leroux², Oscar Gitton-Quent¹, Leandro Benevides¹, Natalia Breyner¹, Vasco Azevedo³, Philippe Langella¹ and Jean-Marc Chatel¹.

¹Université Paris-Saclay, INRAE, AgroParisTech, Micalis, 78352 Jouy-en-Josas, France

²Maiage, INRAE, 78350 Jouy-en-Josas, France

³Federal University of Minas Gerais (UFMG-ICB), Belo Horizonte, MG, Brazil

The commensal bacterium *Faecalibacterium prausnitzii* has unique anti-inflammatory properties, at least some of which have been attributed to its production of MAM, the Microbial Anti-inflammatory Molecule. Previous phylogenetic studies of *F. prausnitzii* strains have revealed the existence of various phylogroups. We address the question of whether MAMs from different phylogroups display distinct anti-inflammatory properties. We first performed wide-scale identification, classification, and phylogenetic analysis of MAM-like proteins encoded in different genomes of *F. prausnitzii*. When combined with a gene context analysis, this approach distinguished at least 10 distinct clusters of MAMs, providing evidence for functional diversity within this protein. We then selected 11 MAMs from various clusters and evaluated their anti-inflammatory capacities *in vitro*. A wide range of anti-inflammatory activity was detected. MAM from the M21/2 strain had the highest inhibitory effect (96% inhibition), while MAM from reference strain A2-165 demonstrated only 56% inhibition, and MAM from strain CNCM4541 was almost inactive. These results were confirmed *in vivo* in murine models of acute and chronic colitis. Our study provides insights into the family of MAM proteins and generates clues regarding the choice of *F. prausnitzii* strains as probiotics for use in targeting chronic inflammatory diseases.