



HAL
open science

Ability of black soldier fly larvae to bioaccumulate tocopherols from different substrates and measurement of larval tocopherol bioavailability in vitro

L. Morand-Laffargue, D. Vairo, C. Halimi, E. Chiarello, B. Creton, D. Sabatier, Patrick Borel

► To cite this version:

L. Morand-Laffargue, D. Vairo, C. Halimi, E. Chiarello, B. Creton, et al.. Ability of black soldier fly larvae to bioaccumulate tocopherols from different substrates and measurement of larval tocopherol bioavailability in vitro. *Journal of Insects as Food and Feed*, 2023, pp.1-13. 10.1163/23524588-20230183B . hal-04332134

HAL Id: hal-04332134

<https://hal.inrae.fr/hal-04332134>

Submitted on 2 Apr 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Research article

2

3 **Ability of black soldier fly larvae to bioaccumulate**
4 **tocopherols from different substrates and measurement of**
5 **larval tocopherol bioavailability *in vitro***

6 L. Morand-Laffargue¹, D. Vairo¹, C. Halimi¹, E. Chiarello², B. Creton³, D. Sabatier³, P.
7 Borel^{1*}

8

9 ¹C2VN, INRAE, Aix-Marseille Univ, INSERM, Marseille, France.

10 ²Department of Agri-Food Sciences and Technologies (DISTAL), University of Bologna,
11 Cesena, Italy.

12 ³BioMiMetiC, Boulbon, France.

13

14 *Corresponding author:

15 patrick.borel@univ-amu.fr

16 UMR C2VN "Center for CardioVascular and Nutrition Research of Marseille"

17 Faculté de Médecine

18 27, boulevard Jean Moulin

19 13005 Marseille, France

20 Phone: +33 (0)4 91 32 42 77

21

22 **Abstract**

23

24 Edible insects are an emerging approach to provide sustainable proteins in feed. Black
25 soldier fly larvae (BSFL) can also bioaccumulate micronutrients from various substrates. The
26 purpose of this study was to assess whether BSFL can bioaccumulate significant
27 concentrations of bioavailable α and γ -tocopherol (TOC) from vitamin E (VE) rich substrates.
28 BSFL were reared on VE rich substrates, e.g. wheat germ oil, bran, etc. α and γ -tocopherol
29 were quantified in larvae and substrates by HPLC. VE bioaccessibility was estimated using an
30 *in vitro* model of digestion. Uptake efficiency of micellarized VE by intestinal cell was
31 estimated using Caco-2 cells. BSFL were at least as rich in α -TOC, but not γ -TOC, as the
32 substrates they were reared on BSFL. VE bioaccessibility was almost always significantly
33 lower in BSFL than in corresponding substrates. Conversely, VE uptake efficiency was either
34 not significantly different or significantly higher in BSFLs than in substrates. Thus, VE
35 enrichment of BSFL from VE rich substrates, in particular co-products such as brans and oil
36 cakes, could be an innovative way to recycle VE and to provide significant amounts of
37 sustainable VE in farm animal feed.

38

39 **Keywords:** tocopherol; insect; bioaccessibility; co-products.

40

41 **Conflicts of interest:** BC and DS work in the BioMiMetiC company. This company conducts
42 research and development activities aimed at enhancing the value of entomo-conversion on a
43 wide variety of organic materials generated in the area at all levels of the food value chain. All
44 other authors have no conflict of interest to declare.

45

46 **Abbreviation:**

- 47 BSFL (black soldier fly larvae).
- 48 TOC (tocopherol)
- 49 VE (vitamin E)

50 1) Introduction

51 Vitamin E (VE) is a vital micronutrient that plays an important role in the health and
52 well-being of both humans and animals. It is a powerful antioxidant that helps to protect cells
53 from damages caused by free radicals and oxidative stress. In addition, it is important for
54 immune function, vision, reproduction, and skin health. In human it is supposed to play a
55 preventive role with respect to cardiovascular risk, immunomodulation, neuroprotection and
56 hepatoprotection (Fata *et al.*, 2014; Galli *et al.*, 2017; Gupta and Suh, 2016; Sozen *et al.*,
57 2019). When it comes to animal nutrition, it is particularly important for livestock, poultry,
58 and aquaculture. In livestock, VE supplementation can improve fertility, reduce the incidence
59 of mastitis in dairy cows (Politis, 2012), and enhance the immune response (Pinelli-Saavedra,
60 2003). It can also improve meat quality by reducing lipid oxidation (Monahan *et al.*, 1992;
61 Rey *et al.*, 2001) which can result in a longer shelf life and improved flavor. In poultry, it is
62 important for immune function, reproductive performance (Surai, 1999), and egg quality
63 (Cherian *et al.*, 1996). It is particularly important in aquaculture systems where fish and
64 shrimp are fed diets high in polyunsaturated fatty acids, which are prone to oxidation. In
65 addition to its benefits for animal health, VE is also important for food production. It is a key
66 ingredient in many food products, including baked goods, cereals, and oils. VE is often added
67 to these products as a natural preservative, as it helps prevent oxidation and spoilage. Overall,
68 the interest of VE for food and feed is clear. It plays a vital role in both human and animals
69 and it has additional interests in food and feed by promoting animal health and productivity
70 and by improving food quality.

71 VE is a fat-soluble vitamin gathering two families which are the tocopherols (TOCs)
72 (α , β , γ and δ) and the tocotrienols (α , β , γ and δ as well). Nevertheless, the main forms of VE
73 present in our diet are α - and γ -TOCs. The first is the most consumed in Europe while the
74 second is the most consumed in the US (Wagner *et al.*, 2004). One of the primary differences
75 between these two TOCs is their ability to scavenge free radicals. While α -TOC is considered
76 to be the most potent lipid antioxidant in our diet and in our body (Niki, 2014), γ -TOC has
77 been shown to have anti-inflammatory effects that are not observed with α -TOC (Jiang *et al.*,
78 2000; Jiang, 2014). Therefore, each TOC could be of interest depending on the biological
79 effect targeted.

80 α and γ -TOCs are mainly found in plant foods containing high concentrations of
81 unsaturated fatty acids, such as nuts, oilseeds and other common edible oils (de Camargo *et al.*
82 *et al.*, 2019; Shahidi *et al.*, 2021). They are also abundant in the by-products of these plant
83 foods, e.g. in brans and oil cakes. A completely innovative way of adding VE to farm animal
84 feed could be to provide it via edible insects containing significant amounts of this vitamin.
85 An advantage of this new way of providing this vitamin would be to provide a more
86 sustainable form of this vitamin and to be able to recycle in the food chain VE that would be
87 lost in waste or co-products rich in this vitamin and little or not used. Among the edible
88 insects used in animal feed, the black soldier fly (BSF, *Hermetia Illucens*) is of particular
89 interest. It has the ability to grow on a wide variety of organic substrates and it has been
90 shown that it can bioaccumulate high concentrations of another fat-soluble vitamin, i.e.
91 provitamin A (Borel *et al.*, 2021). It is therefore a perfect candidate for testing the possibility
92 of having a larva enriched in this vitamin by cultivating it on substrates rich in this vitamin
93 but which can be very different, e.g. vegetable oils, bran and oil cakes.

94 Although it has already been shown that BSF larvae (BSFL) contain α -TOC (Finke,
95 2013; Liu *et al.*, 2017; Shumo *et al.*, 2019), and can be enriched with α -TOC if it is added to
96 their diet (Liland *et al.*, 2017; Oonincx and Finke, 2021), it is not known whether there is an
97 effect of the substrate matrix in which VE is provided to the larvae on the efficiency of the
98 larvae to bioaccumulate this vitamin. It is also not known if the larvae have the same capacity

99 to bioaccumulate α and γ -TOC and if these two forms of VE are bioavailable when they are
100 incorporated into the insect matrix. To answer these many questions, we first raised different
101 groups of BSFL on substrates rich in α and/or γ -TOC. Then, the concentrations of these two
102 forms of VE were measured and compared in the larvae and in substrates they were reared on.
103 Finally, by using an *in-vitro* digestion model coupled with Caco-2 cell monolayers, we
104 compared the bioaccessibility and intestinal cell uptake efficiency of these two forms of VE
105 whether they came from the larval matrix or from the rearing substrate matrix.

106 **2) Material and methods**

107 *Chemicals*

108 Ethanol, *n*-hexane and methanol were purchased from Carlo Erba reagents (Peypin,
109 France). The TOC standards used to identify and quantify TOCs by HPLC, i.e. DL-all-rac- α -
110 TOC and (RRR)- γ -TOC (HPLC purity $\geq 96\%$), the internal standard used to calculate
111 extraction yield, i.e. retinyl acetate, and the enzymes and the chemicals used for the *in vitro*
112 digestions were from Sigma-Aldrich (Saint-Quentin-Fallavier, France). For cell culture,
113 Dulbecco's modified Eagle's medium (DMEM) containing 4.5 g/L glucose, trypsin-
114 ethylenediaminetetraacetic acid (EDTA), non-essential amino acids, penicillin/streptomycin,
115 and phosphate-buffered saline (PBS) were purchased from Life Technologies (Villebon-sur-
116 Yvette, France). Finally, fetal bovine serum (FBS) was from PAA (Vélizy-Villacoublay,
117 France).

118 *Experimental food*

119 We first chose a series of substrates among the richest in α and/or γ -TOC according to
120 food tables and data from the literature (**Table 1**). Then we chose co-products rich in these
121 two forms of VE because we had put forward the hypothesis that these co-products could be
122 valorized via their VE supply for the larvae. This led us to work on a total of 10 vitamin E-
123 rich substrates, and more specifically on two main types of different substrates, oils, e.g. corn
124 oil, and solid substrates, e.g. rice bran. Some of these substrates, wheat germ oil (Emile Noel
125 brand, France), corn oil (La Tourangelle brand, France), rice bran oil (Saldac brand, Italy),
126 rice bran (Moulin Des Moines brand, France) and wheat bran (Priméal brand, France) were
127 ordered via the internet. The brewer's spent grains were a kind gift from a local brewery
128 (Brasserie de la Plaine, Marseille, France). The turnip rapeseed and corn oil cakes came from
129 a local producer (Moulin Giraud, Ozan, France). Poultry feed (broiler grower, Evialis,
130 Vedène, France) was provided by BioMiMetiC that used it to rear their BSFL and thus which
131 was considered as a positive control. The other rearing substrates, i.e. rapeseed oil and white
132 mushrooms (which were used as a negative control because they do not contain detectable
133 amount of α and γ -TOC), as well as foods used in the *in vitro* digestion experiments, i.e.
134 potatoes, minced beef meat (with 5% of fat) and olive oil, were bought from local
135 supermarkets.
136

137 *BSFL farming*

138 As mentioned above, the BSFL were reared on either oils or solid substrates. As the
139 larvae could not be reared in oil we previously mixed the studied vegetable oils with hydrated
140 standard poultry feed to reach 10% of the total mass. For the by-products, only brewer's spent
141 grains did not need to be hydrated with water. The other substrates were hydrated so that the
142 relative humidity ranged from 60 to 70%, the ideal relative humidity for BSFL growth (Dzepe
143 *et al.*, 2020). The rearing procedure was carried out at the BioMiMetiC company in Boulbon
144 (France) as was described in detail in one of our previous studies (Borel *et al.*, 2021).
145 Following a 5-day fasting period, the larvae were collected and stored at $-80\text{ }^{\circ}\text{C}$.
146

147

148 *Assessment of the bioaccessibility of VE in rearing substrates and BSFL*

149 Before the experiment, 2 g of substrate or larva sample was ground with liquid
150 nitrogen in a mortar grinder (Pulverisette 2, FRITSCH GmbH, Idar-Oberstein, Germany). The
151 bioaccessibility of α and γ -TOC was measured using an *in vitro* digestion model, as was
152 described in detail in one of our previous studies (Borel et al., 2021). Briefly, samples were
153 mixed with 6.7 g mashed potato, 1.2 g minced meat and 200 mg olive oil. The mixture was
154 homogenized for 10 min at 37 °C in a rotary incubator (190 rpm) before adding 2.5 mL of
155 artificial saliva and the mixture was once again incubated 10 min at 37 °C. After adjusting the
156 pH to 4 ± 0.02 , 2 mL pepsin were added to the mixture before incubation during 30 min.
157 Then, pH was adjusted to 6 ± 0.02 prior to the addition of 9 mL pancreatin-bile extract
158 solution and 4 mL bile solution. The mixture was then incubated for 30 minutes and the
159 digestate aliquots were collected. The remaining mixture was centrifuged at $1,860 \times g$ for 1 h
160 12 min at 10 °C. Aliquots of filtered supernatant, containing the mixed micelles were stored at
161 80 °C until lipid extraction and quantification of vitamin E by HPLC.

162
163 *Cell culture of Caco-2 and method for measuring the efficiency of uptake of VE by these cells*

164 The cell culture protocol, as well as the cytotoxicity assay were the same as previously
165 described (Morand-Laffargue *et al.*, 2023). Regarding the measurement of the uptake
166 efficiency of VE from the micellar fraction by Caco-2 clone TC7 cells, 1.5 mL of the micellar
167 fractions obtained at the end of the *in vitro* digestion, previously diluted 5 times with DMEM
168 to avoid the toxicity of bile salts towards the cells, were added on the cells. Solutions
169 containing micellarized VE were incubated for 4 hours (Liu *et al.*, 2004) at 37 °C.

170
171 *VE extraction*

172 For the rearing substrates and the BSFL, 250 mg of samples finely ground with liquid
173 nitrogen in a mortar grinder (Pulverisette 2, FRITSCH GmbH, Idar-Oberstein, Germany)
174 were first homogenized in 250 μ L of distilled water and 500 μ L of an ethanol solution
175 containing retinyl acetate (as internal standard) at a concentration of 0.4 mg/L. In the case of
176 *in vitro* digestion samples, i.e. digestates and mixed micelles, and for the recovered medium
177 and scrapped cells, 500 μ L were added to 500 μ L of the same internal standard solution. Then
178 a double extraction with hexane was performed as described previously (Borel *et al.*, 2021).
179 Finally, the dry film was resolubilized in 200 to 2000 μ L of methanol, based on the theoretical
180 VE concentration of the sample.

181
182 *VE quantification*

183 The resolubilized extracts rich in VE were injected, the same day of their extraction,
184 into the HPLC system. After an initial injection, injection volumes were adjusted to obtain
185 signals within the calibration range. Injection volumes of 50 to 100 μ L for BSFL and rearing
186 substrate samples, 25 to 50 μ L for digestate samples, 50 to 100 μ L for micellar fractions and
187 180 μ L for recovered apical cell medium as well as scrapped cells were injected into the
188 HPLC system. All the analyses were performed using a 250×4.6 nm ZORBAX Eclipse
189 XDB-C18 5- μ m column and the corresponding analytical guard column (Agilent
190 Technologies, Santa Clara, CA, USA). The column was maintained at a constant temperature
191 (40 °C). VE analysis was conducted with a 100% methanol mobile phase and flow rate was
192 1.5 mL/min.

193 The HPLC system included an UltiMate 3000 system (Thermo Fisher Scientific, San
194 Jose, CA, USA) and a fluorimetric detector. For fluorimetric analysis of TOCs, detection was
195 set at 292 nm (excitation) and 325 nm (emission). Quantification was performed using
196 Chromeleon software comparing peak area with the curves of the TOC standards.

197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221

Calculations

VE bioaccessibility and uptake efficiency were calculated as:

$$\text{Bioaccessibility (\%)} = \frac{\text{amount of } \alpha \text{ or } \gamma \text{ TOC recovered in mixed micelles}}{\text{amount of } \alpha \text{ or } \gamma \text{ TOC recovered from the digesta}}$$

$$\text{Uptake efficiency (\%)} = \frac{\text{amount of } \alpha \text{ or } \gamma \text{ TOC recovered in cells}}{(\text{amount of } \alpha \text{ or } \gamma \text{ TOC recovered in cells}) + (\text{amount of } \alpha \text{ or } \gamma \text{ TOC in apical medium})}$$

Statistics

Results are expressed as means ± SD or SEM. The homogeneity of variances ($p > 0.05$) was assessed by the Levene’s test. In the case of variance heterogeneity, the data were log-transformed. Departures from normality were assessed using Q–Q plots of standardized residuals. Differences between groups were tested using several two-way ANOVA. When a significant effect of either the substrate matrix or the insect matrix was found by ANOVA, the comparison of means between the two matrices was carried out by pairwise comparisons using bilateral t-tests, with a Bonferroni adjustment. To compare the bioaccessibility of VE among the substrate groups or among the larval groups and to compare the VE amount in the micelles of larvae reared on the different substrates, one-way ANOVA were performed. Tukey-Kramer’s test was used as a post hoc test for pairwise comparisons. Pearson's correlation coefficients were calculated with 95% confidence intervals (CI) to assess possible correlations between certain variables. Values of $p < 0.05$ were considered significant. All statistical analyses were performed using R version 4.1.1 for Windows (R Core Team, 2021). The different ANOVA were performed using the “anova_test” function in the rstatix package v.0.7.2.

Table 1: Amount of α and γ -TOC (mg/kg) in the rearing substrates selected in the study.

	α -TOC	γ -TOC	Source
Rapeseed oil	181.0	383.5	(Wen <i>et al.</i> , 2020)
	189 – 240	370 – 510	(de Camargo <i>et al.</i> , 2019)
	202.9 ± 38.5	131.6 ± 24.2	Measured in this study
Corn oil	180 – 257	440 – 752	(de Camargo <i>et al.</i> , 2019)
	183.9	623.9	(Wen <i>et al.</i> , 2020)
	208.9 ± 27.0	339.9 ± 25.6	Measured in this study
Rice bran oil	151.8	64.4	(Wen <i>et al.</i> , 2020)
	7.3 – 159	2.6 – 80	(de Camargo <i>et al.</i> , 2019)
	91.2 ± 4.1	24.9 ± 2.7	Measured in this study
Wheat germ oil	1510 – 1920	traces – 523	(de Camargo <i>et al.</i> , 2019)
	1622.4	912.6	(Kumar and Krishna, 2015)
	1166.6 ± 343.0	152.9 ± 49.2	Measured in this study
Brewer’s spent grain (BSG)	8.8 – 11.0*	1.4 – 4.6*	(Badea <i>et al.</i> , 2018)
	1.7	3.6	(Bouillon, 2016)
	0.6 ± 0.05	0.2 ± 0.05	Measured in this study
Corn oil cake	12.9 ± 0.7	25.4 ± 1.5	Measured in this study
Poultry feed	7.0 ± 2.8	6.9 ± 1.3	Measured in this study
Rice bran	7 – 55*	4 – 23*	(Górnaś <i>et al.</i> , 2016)
	26.4 – 35.0*	20.9 – 41.0*	(Aguilar-Garcia <i>et al.</i> , 2007)
	27.4 ± 7.9	1.1 ± 0.2	Measured in this study
Turnip rapeseed cake	6.6 ± 0.3	12.6 ± 2.0	Measured in this study

	6 – 31*	0 – 1*	(Górnaś <i>et al.</i> , 2016)
Wheat bran	1.3 – 21.3*	0.9 – 6.9*	(Zhou <i>et al.</i> , 2004)
	n.d.	0.1 ± 0.02	Measured in this study
	0.75 ± 0.04	n.d.	(Barros <i>et al.</i> , 2008)
Mushrooms	n.d.*	n.d.*	(Glamočlija <i>et al.</i> , 2015)
	n.d.	n.d.	Measured in this study

222 Data are means ± SD. n.d.: not detected.

223 Data are expressed in fresh weight except when there is an asterisk which means that they are
224 expressed in dry weight.

225

226 3) Results

227 *Concentrations of α- and γ-TOC in VE rich substrates and in larvae reared on these*

228 *substrates*

229 **Figure 1** shows the different concentrations of α- (**Figure 1.A**) and γ-TOC (**Figure**
230 **1.B**), on a fresh weight basis, in substrates rich in these compounds and in BSFL reared on
231 these substrates. Firstly, note that both the mushrooms and the larvae reared on this substrate
232 did not contain detectable amounts of α- and γ-TOC. Looking at the graphs from left to right,
233 we first see that for the corn oil, rapeseed oil and rice bran oil groups, α-TOC concentrations
234 were not significantly different ($p > 0.3$) between the substrates and the larvae (72.4 ± 3.5 ,
235 71.7 ± 6.3 and 34.9 ± 0.5 mg/kg versus 76.9 ± 1.9 , 82.1 ± 9.2 and 38.7 ± 1.6 mg/kg,
236 respectively). Conversely, for the same groups, γ-TOC concentrations were significantly
237 ($p < 0.001$) lower in the larvae than in the substrates (17.2 ± 0.6 , 11.1 ± 1.0 and 4.9 ± 0.2
238 mg/kg versus 114.9 ± 3.3 , 48.1 ± 3.4 and 13.0 ± 0.3 mg/kg, respectively). The α- and γ-TOC
239 concentrations of the wheat germ oil substrate (719.1 ± 86.1 and 101.8 ± 12.0 mg/kg,
240 respectively) were significantly ($p < 0.001$) higher than those of the larvae reared on this
241 substrate (286.1 ± 4.6 and 29.6 ± 1.4 mg/kg, respectively). Larvae reared on brewer's spent
242 grains (BSG) contained significantly ($p < 0.001$) more α- (23.2 ± 0.3 mg/kg) and γ-TOC ($2.3 \pm$
243 0.2 mg/kg) than the brewer's spent grains (0.6 ± 0.02 and 0.2 ± 0.02 mg/kg, respectively). The
244 α-TOC concentrations of corn oil cake and turnip rapeseed cake (12.9 ± 0.4 and 6.6 ± 0.1
245 mg/kg respectively) were significantly lower than those of larvae reared on these cakes (41.5
246 ± 1.7 and 13.4 ± 1.7 mg/kg, respectively). However, the opposite was observed for the γ-TOC
247 concentration, since corn oil cake (25.4 ± 1.7 mg/kg) and turnip rapeseed cake (12.6 ± 1.0
248 mg/kg) contained significantly ($p < 0.001$) more γ-TOC than the larvae reared on these cakes
249 (15.3 ± 0.8 and 4.6 ± 0.6 mg/kg, respectively). Larvae reared on rice bran (59.6 ± 1.7 mg/kg)
250 contained significantly ($p < 0.001$) more α-TOC than rice bran (27.4 ± 4.0 mg/kg). The
251 concentration of α-TOC in wheat bran was not detectable but the concentration in wheat bran
252 fed larvae was 4.4 ± 0.7 mg/kg. The γ-TOC concentrations of rice (1.1 ± 0.1 mg/kg) and
253 wheat (0.1 ± 0.01 mg/kg) brans were significantly lower than those of larvae fed the same
254 brans (3.4 ± 0.4 and 4.4 ± 0.7 mg/kg respectively). The poultry feed contained significantly
255 more γ-TOC but less α-TOC than the larvae that consumed it (for γ-TOC: 6.9 ± 0.6 versus 2.6
256 ± 0.7 mg/kg, respectively, and for α-TOC: 7.0 ± 1.4 versus 16.7 ± 4.8 mg/kg, respectively).
257 Finally, positive correlations have been highlighted between the α-TOC and γ-TOC
258 concentrations of the substrates and those of the BSFL (Pearson's correlation coefficients = +
259 0.98 (CIr (95% Confidence Interval of r): [0.91-0.99]) and + 0.85 (CIr: [0.47-0.96]),
260 respectively, $p < 0.01$). Supplementary **figures S1A** and **S1B** show α-TOC and γ-TOC
261 concentrations on a dry weight basis.

262

263 **Figure 1. VE concentrations in rearing substrates and in larvae (mg/kg fresh weight).**

264

265 *Bioaccessibility of α - and γ -TOC from VE rich substrates and larvae reared on these*
 266 *substrates*

267 **Tables 2.A** and **2.B** respectively represent the bioaccessibility of α - and γ -TOC from
 268 the selected substrates and from the larvae fed on these substrates. Three types of comparisons
 269 are available: comparison of the bioaccessibility of each form of VE in different substrates,
 270 comparison of the bioaccessibility of each form of VE in larvae reared on different substrates,
 271 and comparison of the bioaccessibility of each form of VE when present in larvae versus
 272 when present in their corresponding substrate.

273 The first comparison shows that the bioaccessibility of these two forms of VE was
 274 significantly different between certain substrates. The maximum amplitude being 7 times for
 275 α -TOC (wheat bran vs corn oil), and 14 times for γ -TOC (rice bran vs corn oil). It is also
 276 interesting to note that the bioaccessibility of these two forms of VE was the highest in rice
 277 bran and wheat bran, and the lowest in oils, with the exception of wheat germ oil.

278 The comparison of the bioaccessibility of the two forms of VE in the larvae also shows
 279 significant differences. The maximum amplitude being about 35 times for α -TOC (larvae
 280 reared on rice bran vs larvae reared on corn oil), and 36 times for γ -TOC (larvae reared on
 281 rice bran vs larvae reared on corn oil as well).

282 The last comparison is that of the bioaccessibility of VE when it was present in the larvae and
 283 in their corresponding substrates. A striking overall observation is that the bioaccessibility of
 284 α and γ -TOC in the larvae was almost always significantly lower in the larvae than in their
 285 corresponding substrates, with an amplitude of up to about 34 times for α -TOC provided in
 286 wheat germ, and up to about 11 times for γ -TOC provide in wheat germ as well. There are
 287 however two exceptions: the turnip rapeseed cake for which the bioaccessibility of the two

Substrate	Bioaccess in substrates	Bioaccess in larvae	p-value**
Corn oil	8.8 ± 2.1^d	1.3 ± 0.4^c	0.002
Rapeseed oil	16.6 ± 5.3^{cd}	1.4 ± 0.3^c	<0.001
Rice bran oil	9.2 ± 2.3^d	1.7 ± 0.4^c	0.003

288 forms of VE was not significantly different between the larvae and the substrate, and the corn
 289 oil cake for which the bioaccessibility of the two forms of VE was significantly higher in the
 290 larvae than in the substrate.

291
 292 **Table 2.** Bioaccessibility of α - and γ -TOC¹ from larvae and their corresponding rearing
 293 substrates.

Wheat germ oil	58.1 ± 6.9 ^a	1.7 ± 0.1 ^c	<0.001
Brewers' spent grain	21.2 ± 3.6 ^{bc}	2.2 ± 0.6 ^c	<0.001
Corn oil cake	28.9 ± 1.9 ^{ac}	41.0 ± 0.9 ^a	0.002
Rice bran	59.7 ± 0.8 ^a	45.1 ± 3.5 ^a	0.008
Turnip rapeseed cake	42.4 ± 4.1 ^{ab}	37.1 ± 5.1 ^a	0.4
Wheat bran	61.9 ± 1.0 ^a	7.6 ± 0.2 ^b	<0.001
Poultry feed	40.6 ± 8.5 ^{ab}	18.6 ± 2.0 ^{ab}	0.02

294 **A. α-TOC**

295 **B. γ-TOC**

Substrate	Bioaccess in substrates	Bioaccess in larvae	p-value**
Corn oil	3.6 ± 0.5 ^c	1.3 ± 0.4 ^f	0.009
Rapeseed oil	5.3 ± 1.4 ^c	1.8 ± 0.4 ^{ef}	0.01
Rice bran oil	5.3 ± 0.7 ^c	2.9 ± 0.4 ^{de}	0.02
Wheat germ oil	25.1 ± 4.2 ^b	2.2 ± 0.4 ^{ef}	<0.001
Brewers' spent grain	16.8 ± 2.6 ^b	2.6 ± 0.7 ^{df}	0.002
Corn oil cake	26.4 ± 2.1 ^{ab}	38.2 ± 1.9 ^{ab}	0.009
Rice bran	52.0 ± 1.3 ^a	46.9 ± 3.6 ^{ab}	0.2
Turnip rapeseed cake	36.0 ± 2.7 ^{ab}	37.0 ± 5.1 ^a	1.0
Wheat bran	49.1 ± 1.6 ^{ab}	8.2 ± 0.3 ^c	<0.001
Poultry feed	36.9 ± 7.3 ^{ab}	19.0 ± 1.8 ^{bc}	0.04

296 ¹Percent of α- or γ-TOC recovered in the micelle fraction relative to the quantity of these
 297 compounds found in the digestate at the end of the *in vitro* digestion. Values are means ±
 298 SEM (n=4). **p-value of a two-way ANOVA for each TOC, followed by pairwise
 299 comparisons using bilateral t-tests. In each column, means that bear different superscript
 300 letters are significantly different (*p* < 0.05; ANOVA followed by Tukey's HSD test).
 301

302 *Uptake efficiency of micellized VE by Caco-2 cells*

303 **Table 3** shows the absorption efficiency of α- and γ-TOC contained in the mixed
 304 micelle fractions which were obtained after the *in vitro* digestions. The absorption efficiency
 305 of α-TOC from micelles of larvae reared on wheat germ oil was about twice as high as that
 306 from micelles from this substrate. In the case of rice bran, the absorption efficiency of α-TOC
 307 from micelles of larvae was not significantly different from that of the substrate. The
 308 absorption efficiencies of γ-TOC from *in vitro* digestions of larvae reared on wheat germ oil
 309 or rice bran were not significantly different from their corresponding substrates, but note that
 310 the means were higher in larvae than in the substrates (about +50%).

311 **Table 3.** VE uptake efficiency (%) by Caco-2 cells. ¹

	Micellar fraction from rearing substrate digestion	Micellar fraction from larva digestion	Substrate vs larvae comparison **p-value
α-TOC			
Wheat germ oil	16.2 ± 2.1	34.2 ± 3.6	0.005
Rice bran	24.1 ± 1.3	28.8 ± 3.8	0.29
γ-TOC			
Wheat germ oil	12.4 ± 2.3	19.7 ± 5.0	n.s****

Rice bran	20.3 ± 2.0	30.1 ± 5.6	n.s
-----------	------------	------------	-----

312 [†]Percentage of α - or γ -TOC that was recovered in the Caco-2 cells relative to the quantity of
313 these compounds in the mixed micelle-rich fractions deposited on the cells before incubation
314 for 4 hours at 37°C. Values are means ± SEM (n=4). ** For α -TOC, *p*-value of a two-way
315 ANOVA followed by pairwise comparisons using bilateral *t*-tests. ***For γ -TOC, the result
316 of the two-way ANOVA was non-significant (*p* = 0.7) thus post-hoc tests were not carried
317 out.

318 **Figure 2** shows the VE amount absorbed by the Caco-2 cells on a fresh weight basis. This
319 value is of great interest to compare the theoretical amount of VE that each matrix can bring
320 to the intestinal cell because, in addition to considering the cell uptake efficiency, it also
321 considers the bioaccessibility of VE in the substrate or the larvae. The cellular amount of VE
322 after digestion and cell uptake of micelles from wheat germ oil was significantly higher than
323 that of larvae reared on wheat germ oil (for α -TOC (**Figure 2A**), 30.3 ± 7.8 µg and 3.3 ± 0.5
324 µg, respectively, and for γ -TOC (**Figure 2B**) 3.7 ± 1.2 µg and 0.2 ± 0.04 µg, respectively).
325 Conversely to what was observed for wheat germ oil, cells incubated with micelles from
326 digestion of larvae reared on rice bran had higher concentrations of α and γ -TOC than cells
327 incubated with micelles from digestion of rice bran, although this was not significantly
328 different (*p* = 0.08 for α -TOC and *p* = 0.3 for γ -TOC). Supplementary **figures S2A** and **S2B**
329 show α -TOC and γ -TOC cellular concentrations on a dry weight basis.

330
331 **Figure 2. Amounts of VE taken up by Caco-2 cells incubated with micelle-rich fractions**
332 **obtained after *in vitro* digestions of rearing substrates or larvae.**

333 4) Discussion

334 The first objective of this study was to assess the ability of BSFL to bioaccumulate α -
335 and γ -TOC from substrates rich in these compounds but made up of very different matrices
336 (de Camargo *et al.*, 2019; Franke *et al.*, 2010; Shahidi *et al.*, 2021). Indeed, these substrates
337 were either solid, e.g. bran or cake, or liquid, i.e. vegetable oils. The first interesting result of
338 this study is the fact that when BSFL were grown on white mushrooms, they did not contain
339 VE. This result not only suggests that BSFL reared on a substrate without VE can apparently
340 grow like larvae that have access to VE, but also shows that BSFL do not have the ability to
341 synthesize VE. With regard to the substrates rich in VE, the first observation is that the
342 contents of α and γ -TOC that we measured agreed with the data in the literature, except in
343 wheat bran where we found no α -TOC and a very low concentration of γ -TOC compared to
344 what has been reported in the literature (**Table 1**). This suggests that our extraction protocol
345 was not adapted for this substrate. Regarding α -TOC concentrations measured in the BSFL,
346 they differed strongly depending on the rearing substrate, since they ranged from 18.1 ± 2.3
347 mg/kg for BSFL reared on turnip rapeseed cake to 286.1 ± 4.6 mg/kg for BSFL reared on
348 wheat germ oil. This very high concentration, which is moreover largely higher than the α -
349 TOC concentration of all the VE-rich substrates, with the exception of wheat germ, has never
350 been observed. Indeed, what has been reported in whole larvae is 9.6 mg/kg if reared on
351 chicken manure and 19.3 mg/kg if reared on spent grain (Shumo *et al.*, 2019).

352 If we now compare the concentrations of α -TOC in the larvae and in the substrates on which
353 they were raised, we see that they are always higher in the larvae than in the substrates,
354 sometimes significantly, with the exception of the larvae raised on the wheat germ oil. To
355 explain this exception, we hypothesize that the very high amount of α -TOC provided by
356 wheat germ oil led the larvae to reach the maximum concentration of α -TOC they can reach.
357 Concerning γ -TOC, what was immediately striking was that the concentrations in the larvae
358 are, 7 times out of 10, lower than in their corresponding substrate. In fact, it is likely that the

359 result should have been the same with the 3 other substrates, made of bran, because we
360 deduced that our method of extracting the two VE species from these substrates was not very
361 effective which must have led to underestimating their VE content. To support this
362 hypothesis, we rely on the results of bioaccessibility which show that we can detect α and γ -
363 TOC in the micelles resulting from the digestion of wheat bran (**Table 2**) whereas we do not
364 detect α -TOC, and we detect very low concentration of γ -TOC, in wheat bran when we have
365 extracted this substrate with organic solvents (**Figure 1**), even though wheat bran contains
366 both forms of VE (**Table 1**). This difference in bioaccumulation of these two forms of VE in
367 BSFL is quite remarkable. The most likely hypothesis is that, as has been described in
368 *Drosophila*, which is a dipteran like BSF, this is due to a preferential degradation of γ -TOC,
369 as compared to α -TOC, by cytochrome P450 TOC- ω -hydroxylase (Parker and McCormick,
370 2005). If this hypothesis is true, this suggests that α -TOC has an essential function in the BSF
371 and that it is important to provide it in its diet during its breeding, as is the case for other
372 production animals.

373 After demonstrating that larvae can bioaccumulate very high concentrations of α -TOC
374 and that they can also be enriched in γ -TOC, the question arose of the bioavailability of these
375 two forms of VE in the insect matrix. The first key step in the bioavailability of lipid
376 molecules is their incorporation into the micelles during digestion. The rate of incorporation is
377 called bioaccessibility. When we look at **Table 2** we can clearly see that, with one exception
378 which is corn oil cake, the bioaccessibility of the two forms of VE was significantly lower
379 when they were in the insect matrix than when they were in the plant matrix. We do not know
380 the mechanism but we can propose two hypotheses. The first is that, as the insect's
381 cuticle/exoskeleton contains a variety of compounds, e.g. chitin, proteins and lipids (Kramer
382 *et al.*, 1995) , it is possible that a fraction of the VE was associated with one or several of
383 these compounds and is therefore not bioaccessible. The second hypothesis, knowing that all
384 these substrates contain not only VE but also phytosterols, which are well known to inhibit
385 the micellization of cholesterol and fat-soluble vitamins (Fardet *et al.*, 2017), is that the
386 phytosterols have strongly bioaccumulated in the larvae and/or are much more bioaccessible
387 in the larvae than in their corresponding substrates. This effect of phytosterols could also
388 explain why the bioaccessibility of VE in the oils was significantly lower than in the solid
389 matrices. It can indeed be assumed that the bioaccessibility of phytosterols present in oils is
390 greater than that of these compounds when they are present in plant matrices in which they
391 can interact with dietary fibers. These hypotheses obviously need to be verified by dedicated
392 studies.

393 Bioaccessibility gives an idea of the extractability of a bioactive from a matrix in
394 which it is present. But, from a nutritional point of view, it is important to consider both this
395 parameter and the content of bioactive of interest in the matrix. It is therefore relevant to
396 compare the VE concentrations in the micelles which come from the digestions of the larvae
397 having been reared on the different substrates (Supplemental **figures S3 and S4**). This
398 comparison clearly shows that the larvae which give the highest amount of α and γ -TOC in
399 micelles are the two cakes, the rice bran and the chicken feed, substrates which were not the
400 richest in VE.

401 The amount of VE in micelles is obviously a very important parameter of
402 bioavailability, but we know that everything that is present in micelles is not necessarily
403 absorbed by the intestinal cells, or it can be absorbed and then resecreted in the digestive
404 lumen and is therefore not bioavailable to the body, as is the case for phytosterols (Ling and
405 Jones, 1995). To get a better idea of what the in vivo bioavailability might be it is therefore
406 essential to measure the absorption efficiency of VE present in micelles. And that's what was
407 done with the Caco-2 model by choosing the oil and the solid substrate giving the greatest
408 amount of α -TOC in the micelles (**Table 3**). When we look at this table we see that the only

409 significant difference was that the VE uptake efficiency was twice as high from the micelles
 410 resulting from the digestion of larvae reared on the substrate containing wheat germ oil than
 411 from the micelles resulting from the digestion of this same substrate. Unfortunately, we do not
 412 have a solid hypothesis to explain this difference.

413 But what is important from a nutritional point of view is the
 414 quantity of VE which is found in the cells, i.e. which is actually bioavailable. This quantity
 415 depends on the absorption efficiency by the cells but also on the amount of VE present in the
 416 micelles. When we look at these data (**Figures 2A** and **2B**) we can clearly see that the
 417 amounts of VE found in the cells which have been incubated with micelles from the digestion
 418 of larvae which have been reared on the substrate containing wheat germ oil was 3 to 4 times
 419 lower than the amount of VE found in cells that were incubated with micelles from larvae
 420 reared on rice bran. This suggests that rice bran is a better source of bioavailable VE than
 421 wheat germ oil, although the latter was the richest source of VE among all the studied
 422 substrates.

423 Finally, in order to get an overview of the capacity of VE-rich BSFL to be used in the
 424 animal feed industry as a significant source of VE, the amount of VE that could potentially be
 425 provided by VE-rich BSFL to different livestock was calculated according to the VE needs of
 426 the different animal species and to the recommended quantities of BSFL that are usually
 427 incorporated in their diet (**Table 4**). This table suggests that these vitamin-rich larvae would
 428 be of particular interest in poultry as they could cover all the vitamin requirements of laying
 429 hens, broilers and turkeys.

430 In conclusion, this study first shows that BSFL can bioaccumulate both α - and γ -TOC
 431 from VE-rich substrates. Larvae α -toc concentrations are mostly equal to or higher than those
 432 of their rearing substrates. Conversely larvae γ -toc concentrations are generally lower than
 433 those of their substrates suggesting a different metabolism of these two forms of VE in the
 434 larvae. Regarding VE bioaccessibility it was generally lower in the insect matrix than in the
 435 corresponding plant matrices. Nevertheless, most BSFL reared on by-products (brans and
 436 press oil cakes) contained significant amounts of micellarized VE, which could lead to a
 437 sustainable VE valorization of these by-products. Finally, adding VE rich BSFL in the diet of
 438 some farm animals could provide a significant proportion of the needs of these animals in this
 439 vitamin. The enrichment of larvae with VE could make it possible to recycle part of the VE
 440 present in waste or co-products, it would make it possible to provide the natural forms of this
 441 vitamin and would increase the economic value of these larvae.

442
 443 **Table 4.** Proportion of the recommended VE dose that could come from the incorporation of
 444 VE-rich larvae in animal feed.

Species	VE concentration that was added in the diet (mg/kg)	BSFL fraction provided in the diet	Recommended proportion of the BSFL fraction added in the diet (weight%)	mg of VE* provided by BSFL incorporated at the recommended proportion	VE % provided by BSFL
Broiler chicken (<i>Gallus gallus domesticus</i>)	7 (National Research Council, 1994)	Prepupae	15% (Pieterse <i>et al.</i> , 2019)	9	128

Laying hens (<i>Gallus gallus domesticus</i>)	3 (National Research Council, 1994)	Prepupae	20% (Tahamtani <i>et al.</i> , 2021)	12	400
Salmon (<i>Salmo salar</i>)	40 (National Research Council, 1993)	Prepupae	50% (Sealey <i>et al.</i> , 2011)	30	75
Tilapia (<i>Oreochromis niloticus</i>)	40 (National Research Council, 1993)	Prepupae	50% (Taufek <i>et al.</i> , 2021)	30	75
Turkey (<i>Meleagris gallopavo</i>)	7 (National Research Council, 1994)	Prepupae	50% (Chia <i>et al.</i> , 2021)	30	428
Dog (<i>Canis lupus familiaris</i>)	20 (National Research Council, 2006)	Partially defatted black soldier fly	20 % (Freel <i>et al.</i> , 2021)	12	60

445 *Using the theoretical α -TOC content of BSFL reared on rice bran, the BSFL group that
446 provided the highest amount of α -TOC in micelles.
447

448 **Supporting information:**

449 The costs of this project were covered equally by the own budget of P. Borel's research team,
450 which came mainly from INRAE endowments, and by the BioMiMetiC company.

451 For the purpose of Open Access, a CC-BY public copyright licence ([Creative Commons — Attribution 4.0 International — CC BY 4.0](https://creativecommons.org/licenses/by/4.0/))
452 has been applied by the authors to the present document and will be applied to all subsequent versions up to the Author Accepted
453 Manuscript arising from this submission.
454



455
456

457 **5) References**

458 Aguilar-Garcia, C., Gavino, G., Baragaño-Mosqueda, M., Hevia, P. and Gavino, V.C., 2007.
459 Correlation of tocopherol, tocotrienol, γ -oryzanol and total polyphenol content in rice bran
460 with different antioxidant capacity assays. *Food Chemistry* 102: 1228–1232.
461 <https://doi.org/10.1016/j.foodchem.2006.07.012>.

462 Badea, A., Carter, A., Legge, W.G., Swallow, K., Johnston, S.P. and Izydorczyk, M.S., 2018. Tocols
463 and oil content in whole grain, brewer's spent grain, and pearling fractions of malting, feed,
464 and food barley genotypes. *Cereal Chemistry* 95: 779–789.
465 <https://doi.org/10.1002/cche.10093>.

466 Barros, L., Cruz, T., Baptista, P., Estevinho, L.M. and Ferreira, I.C.F.R., 2008. Wild and commercial
467 mushrooms as source of nutrients and nutraceuticals. *Food and Chemical Toxicology* 46:
468 2742–2747. <https://doi.org/10.1016/j.fct.2008.04.030>.

- 469 Borel, P., Hammaz, F., Morand-Laffargue, L., Creton, B., Halimi, C., Sabatier, D. and Desmarchelier,
470 C., 2021a. Using black soldier fly larvae reared on fruits and vegetables waste as a sustainable
471 dietary source of provitamin a carotenoids. *Food Chemistry* 359: 129911.
472 <https://doi.org/10.1016/j.foodchem.2021.129911>.
- 473 Bouillon, M., 2016. Brewers Spent Grain: its chemical composition, potential applications and source
474 of value-added products. <https://doi.org/10.13140/RG.2.2.31341.82400>.
- 475 Cherian, G., Wolfe, F.W. and Sim, J.S., 1996. Dietary Oils with Added Tocopherols: Effects on Egg
476 or Tissue Tocopherols, Fatty Acids, and Oxidative Stability. *Poultry Science* 75: 423–431.
477 <https://doi.org/10.3382/ps.0750423>.
- 478 Chia, S.Y., Tanga, C., Osuga, I., Alaru, P., Mwangi, D.M., Githinji, M., Dubois, T., Ekesi, S., van
479 Loon, J. and Dicke, M., 2021. Black Soldier Fly Larval Meal in Feed Enhances Growth
480 Performance, Carcass Yield and Meat Quality of Finishing Pigs. *Journal of Insects as Food
481 and Feed* 7: 1–16. <https://doi.org/10.3920/JIFF2020.0072>.
- 482 de Camargo, A.C., Franchin, M. and Shahidi, F., 2019. Tocopherols and Tocotrienols: Sources,
483 Analytical Methods, and Effects in Food and Biological Systems. *Encyclopedia of Food
484 Chemistry*. Elsevier, , pp. 561–570. <https://doi.org/10.1016/B978-0-08-100596-5.22358-5>
- 485 Dzepe, D., Nana, P., Fotso, A., Tchuinkam, T. and Djouaka, R., 2020. Influence of larval density,
486 substrate moisture content and feedstock ratio on life history traits of black soldier fly larvae.
487 *Journal of Insects as Food and Feed* 6: 133–140. <https://doi.org/10.3920/JIFF2019.0034>.
- 488 Fardet, A., Morise, A., Kalonji, E., Margaritis, I. and Mariotti, F., 2017. Influence of phytosterol and
489 phytostanol food supplementation on plasma liposoluble vitamins and provitamin A
490 carotenoid levels in humans: An updated review of the evidence. *Critical Reviews in Food
491 Science and Nutrition* Taylor & Francis, 57: 1906–1921.
492 <https://doi.org/10.1080/10408398.2015.1033611>.
- 493 Fata, G.L., Weber, P. and Mohajeri, M.H., 2014. Effects of Vitamin E on Cognitive Performance
494 during Ageing and in Alzheimer’s Disease. *Nutrients Multidisciplinary Digital Publishing
495 Institute*, 6: 5453–5472. <https://doi.org/10.3390/nu6125453>.
- 496 Finke, M.D., 2013. Complete Nutrient Content of Four Species of Feeder Insects: Nutrient Content of
497 Feeder Insects. *Zoo Biology* 32: 27–36. <https://doi.org/10.1002/zoo.21012>.
- 498 Franke, S., Fröhlich, K., Werner, S., Böhm, V. and Schöne, F., 2010. Analysis of carotenoids and
499 vitamin E in selected oilseeds, press cakes and oils. *European Journal of Lipid Science and
500 Technology* 112: 1122–1129. <https://doi.org/10.1002/ejlt.20090025>.
- 501 Freel, T.A., McComb, A. and Koutsos, E.A., 2021. Digestibility and safety of dry black soldier fly
502 larvae meal and black soldier fly larvae oil in dogs. *Journal of Animal Science* 99: skab047.
503 <https://doi.org/10.1093/jas/skab047>.
- 504 Galli, F., Azzi, A., Birringer, M., Cook-Mills, J.M., Eggersdorfer, M., Frank, J., Cruciani, G.,
505 Lorkowski, S. and Özer, N.K., 2017. Vitamin E: Emerging aspects and new directions. *Free
506 Radical Biology & Medicine* 102: 16–36.
507 <https://doi.org/10.1016/j.freeradbiomed.2016.09.017>.
- 508 Glamočlija, J., Stojković, D., Nikolić, M., Ćirić, A., Reis, F.S., Barros, L., Ferreira, I.C.F.R. and
509 Soković, M., 2015. A comparative study on edible *Agaricus* mushrooms as functional foods.
510 *Food & Function* 6: 1900–1910. <https://doi.org/10.1039/c4fo01135j>.

- 511 Górnaś, P., Radenkova, V., Pugajeva, I., Soliven, A., Needs, P.W. and Kroon, P.A., 2016. Varied
512 Composition of Tocochromanols in Different Types of Bran: Rye, Wheat, Oat, Spelt,
513 Buckwheat, Corn, and Rice. *International Journal of Food Properties* Taylor & Francis, 19:
514 1757–1764. <https://doi.org/10.1080/10942912.2015.1107843>.
- 515 Gupta, S.D. and Suh, N., 2016. Tocopherols in cancer: an update. *Molecular nutrition & food research*
516 60: 1354–1363. <https://doi.org/10.1002/mnfr.201500847>.
- 517 Jiang, Q., 2014. Natural forms of vitamin E: metabolism, antioxidant, and anti-inflammatory activities
518 and their role in disease prevention and therapy. *Free Radical Biology and Medicine* 72: 76–
519 90. <https://doi.org/10.1016/j.freeradbiomed.2014.03.035>.
- 520 Jiang, Q., Elson-Schwab, I., Courtemanche, C. and Ames, B.N., 2000. gamma-tocopherol and its
521 major metabolite, in contrast to alpha-tocopherol, inhibit cyclooxygenase activity in
522 macrophages and epithelial cells. *Proceedings of the National Academy of Sciences of the*
523 *United States of America* 97: 11494–11499. <https://doi.org/10.1073/pnas.200357097>.
- 524 Kramer, K.J., Hopkins, T.L. and Schaefer, J., 1995. Applications of solids NMR to the analysis of
525 insect sclerotized structures. *Insect Biochemistry and Molecular Biology* 25: 1067–1080.
526 [https://doi.org/10.1016/0965-1748\(95\)00053-4](https://doi.org/10.1016/0965-1748(95)00053-4).
- 527 Kumar, G.S. and Krishna, A.G.G., 2015. Studies on the nutraceuticals composition of wheat derived
528 oils wheat bran oil and wheat germ oil. *Journal of Food Science and Technology* 52: 1145–
529 1151. <https://doi.org/10.1007/s13197-013-1119-3>.
- 530 Liland, N.S., Biancarosa, I., Araujo, P., Biemans, D., Bruckner, C.G., Waagbø, R., Torstensen, B.E.
531 and Lock, E.-J., 2017. Modulation of nutrient composition of black soldier fly (*Hermetia*
532 *illucens*) larvae by feeding seaweed-enriched media. *PLOS ONE Public Library of Science*,
533 12: e0183188. <https://doi.org/10.1371/journal.pone.0183188>.
- 534 Ling, W.H. and Jones, P.J., 1995. Dietary phytosterols: a review of metabolism, benefits and side
535 effects. *Life Sciences* 57: 195–206. [https://doi.org/10.1016/0024-3205\(95\)00263-6](https://doi.org/10.1016/0024-3205(95)00263-6).
- 536 Liu, C.-S., Glahn, R.P. and Liu, R.H., 2004. Assessment of carotenoid bioavailability of whole foods
537 using a Caco-2 cell culture model coupled with an in vitro digestion. *Journal of Agricultural*
538 *and Food Chemistry* 52: 4330–4337. <https://doi.org/10.1021/jf040028k>.
- 539 Liu, X., Chen, X., Wang, H., Yang, Q., Ur Rehman, K., Li, W., Cai, M., Li, Q., Mazza, L., Zhang, J.,
540 Yu, Z. and Zheng, L., 2017. Dynamic changes of nutrient composition throughout the entire
541 life cycle of black soldier fly. *PLOS ONE* 12: e0182601.
542 <https://doi.org/10.1371/journal.pone.0182601>.
- 543 Monahan, F.J., Buckley, D.J., Morrissey, P.A., Lynch, P.B. and Gray, J.I., 1992. Influence of dietary
544 fat and α -tocopherol supplementation on lipid oxidation in pork. *Meat Science* 31: 229–241.
545 [https://doi.org/10.1016/0309-1740\(92\)90042-3](https://doi.org/10.1016/0309-1740(92)90042-3).
- 546 Morand-Laffargue, L., Hirschberg, J., Halimi, C., Desmarchelier, C. and Borel, P., 2023. The
547 zeaxanthin present in a tomato line rich in this carotenoid is as bioavailable as that present in
548 the food sources richest in this xanthophyll. *Food Research International* 168: 112751.
549 <https://doi.org/10.1016/j.foodres.2023.112751>.
- 550 National Research Council, 1993. *Nutrient Requirements of Fish*. National Academies Press, 132 pp.
551 <https://doi.org/10.17226/2115>.
- 552 National Research Council, 1994. *Nutrient Requirements of Poultry*. National Academies Press, 174
553 pp. <https://doi.org/10.17226/2114>.

- 554 National Research Council, 2006. Nutrient Requirements of Dogs and Cats. National Academies
555 Press, Washington, D.C. <https://doi.org/10.17226/10668>.
- 556 Niki, E., 2014. Role of vitamin E as a lipid-soluble peroxy radical scavenger: in vitro and in vivo
557 evidence. *Free Radical Biology & Medicine* 66: 3–12.
558 <https://doi.org/10.1016/j.freeradbiomed.2013.03.022>.
- 559 Oonincx, D. g. a. b. and Finke, M. d., 2021. Nutritional value of insects and ways to manipulate their
560 composition. *Journal of Insects as Food and Feed Wageningen Academic Publishers*, 7: 639–
561 659. <https://doi.org/10.3920/JIFF2020.0050>.
- 562 Parker, R.S. and McCormick, C.C., 2005. Selective accumulation of α -tocopherol in *Drosophila* is
563 associated with cytochrome P450 tocopherol- ω -hydroxylase activity but not α -tocopherol
564 transfer protein. *Biochemical and Biophysical Research Communications* 338: 1537–1541.
565 <https://doi.org/10.1016/j.bbrc.2005.10.124>.
- 566 Pieterse, E., Erasmus, S.W., Uushona, T. and Hoffman, L.C., 2019. Black soldier fly (*Hermetia*
567 *illucens*) pre-pupae meal as a dietary protein source for broiler production ensures a tasty
568 chicken with standard meat quality for every pot. *Journal of the Science of Food and*
569 *Agriculture* 99: 893–903. <https://doi.org/10.1002/jsfa.9261>.
- 570 Pinelli-Saavedra, A., 2003. Vitamin E in immunity and reproductive performance in pigs.
571 *Reproduction, Nutrition, Development* 43: 397–408. <https://doi.org/10.1051/rnd:2003034>.
- 572 Politis, I., 2012. Reevaluation of vitamin E supplementation of dairy cows: bioavailability, animal
573 health and milk quality. *Animal* 6: 1427–1434. <https://doi.org/10.1017/S1751731112000225>.
- 574 Rey, A.I., Kerry, J.P., Lynch, P.B., López-Bote, C.J., Buckley, D.J. and Morrissey, P.A., 2001. Effect
575 of dietary oils and alpha-tocopheryl acetate supplementation on lipid (TBARS) and cholesterol
576 oxidation in cooked pork. *Journal of Animal Science* 79: 1201–1208.
577 <https://doi.org/10.2527/2001.7951201x>.
- 578 Sealey, W., Gaylord, G., Barrows, F., Tomberlin, J., Mcguire, M., Ross, C. and St-Hilaire, S., 2011.
579 Sensory Analysis of Rainbow Trout, *Oncorhynchus mykiss*, Fed Enriched Black Soldier Fly
580 Prepupae, *Hermetia illucens*. *Journal of the World Aquaculture Society* 42
581 <https://doi.org/10.1111/j.1749-7345.2010.00441.x>.
- 582 Shahidi, F., Pinaffi-Langley, A.C.C., Fuentes, J., Speisky, H. and de Camargo, A.C., 2021. Vitamin E
583 as an essential micronutrient for human health: Common, novel, and unexplored dietary
584 sources. *Free Radical Biology and Medicine* 176: 312–321.
585 <https://doi.org/10.1016/j.freeradbiomed.2021.09.025>.
- 586 Shumo, M., Osuga, I.M., Khamis, F.M., Tanga, C.M., Fiaboe, K.K.M., Subramanian, S., Ekesi, S., van
587 Huis, A. and Borgemeister, C., 2019b. The nutritive value of black soldier fly larvae reared on
588 common organic waste streams in Kenya. *Scientific Reports* 9: 10110.
589 <https://doi.org/10.1038/s41598-019-46603-z>.
- 590 Sozen, E., Demirel, T. and Ozer, N.K., 2019. Vitamin E: Regulatory role in the cardiovascular system.
591 *IUBMB Life* 71: 507–515. <https://doi.org/10.1002/iub.2020>.
- 592 Surai, P. F., 1999. Vitamin E and avian reproduction/Poultry and avian biology reviews. V,
593 10, 3-60, ISSN: 1357-048X.
- 594 Tahamtani, F., Ivarsson, E., Wiklicky, V., Lalander, C., Wall, H., Rodenburg, B., Tuytens, F. and
595 Hernandez, C., 2021. Feeding live Black Soldier Fly larvae (*Hermetia illucens*) to laying hens:

- 596 effects on feed consumption, hen health, hen behavior, and egg quality. *Poultry Science* 100:
597 101400. <https://doi.org/10.1016/j.psj.2021.101400>.
- 598 Taufek, N., Lim, J. and Bakar, N., 2021. Comparative evaluation of *Hermetia illucens* larvae reared
599 on different substrates for red tilapia diet: effect on growth and body composition. *Journal of*
600 *Insects as Food and Feed Wageningen Academic Publishers*, 7: 79–88.
601 <https://doi.org/10.3920/JIFF2019.0058>.
- 602 Wagner, K.-H., Kamal-Eldin, A. and Elmadfa, I., 2004. Gamma-Tocopherol – An Underestimated
603 Vitamin? *Annals of Nutrition and Metabolism Karger Publishers*, 48: 169–188.
604 <https://doi.org/10.1159/000079555>.
- 605 Wen, Y., Xu, L., Xue, C., Jiang, X. and Wei, Z., 2020. Assessing the Impact of Oil Types and Grades
606 on Tocopherol and Tocotrienol Contents in Vegetable Oils with Chemometric Methods.
607 *Molecules* 25: 5076. <https://doi.org/10.3390/molecules25215076>.
- 608 Zhou, K., Su, L. and Yu, L. (Lucy), 2004. Phytochemicals and Antioxidant Properties in Wheat Bran.
609 *Journal of Agricultural and Food Chemistry American Chemical Society*, 52: 6108–6114.
610 <https://doi.org/10.1021/jf049214g>.
- 611