



HAL
open science

The influence of preschoolers' emotional and behavioural problems on obesity treatment outcomes: Secondary findings from a randomized controlled trial

Elodie Eiffener, Karin Eli, Anna Ek, Pernilla Sandvik, Maria Somaraki, Stef Kremers, Ester Sleddens, Paulina Nowicka

► To cite this version:

Elodie Eiffener, Karin Eli, Anna Ek, Pernilla Sandvik, Maria Somaraki, et al.. The influence of preschoolers' emotional and behavioural problems on obesity treatment outcomes: Secondary findings from a randomized controlled trial. *Pediatric Obesity*, 2019, 14 (11), 10.1111/ijpo.12556 . hal-04334624

HAL Id: hal-04334624

<https://hal.inrae.fr/hal-04334624>

Submitted on 11 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Abstract**

2

3 **Background:** Few studies have explored the influence of preschoolers' behavioral problems on
4 obesity treatment.

5 **Objectives:** To assess emotional and behavioral problems before and after an obesity
6 intervention and examine relationships between changes in child behavior and changes in weight
7 status.

8 **Method:** The study included 77 children (4-6 years, 53% girls, mean Body Mass Index (BMI)
9 z-score 3.0 (SD 0.6)) who participated in the More and Less Study, a randomized controlled trial.
10 Families were randomized to a parenting program or to standard treatment. The children's
11 heights and weights (BMI z-score, primary outcome) were measured at baseline and 12 months
12 post-baseline. Parents rated their children's behaviors (secondary outcome) on the Child
13 Behavior Checklist (CBCL) for ages 1.5-5 years, a questionnaire that measures psychosocial
14 health and functioning, encompassing emotional and behavioral problems. Changes in child
15 behavior during treatment were examined through paired samples t-tests; the influence of child
16 behavior on treatment effects was examined through linear regressions.

17 **Results:** Child emotional and behavioral problems significantly improved after obesity
18 treatment. Lower scores were found for Emotional Reactivity, Sleep Problems, Affective
19 Problems, Aggressive Behavior, Externalizing Behaviors, Oppositional Defiant Problems, and
20 Total Problems. Child behavior significantly affected obesity treatment results: Attention
21 Problems and ADHD at baseline contributed to increasing BMI z-scores while Oppositional
22 Defiant Problems, Externalizing Behaviors and a higher number of behavioral problems
23 predicted decreasing BMI z-scores.

24

25 **Conclusions:** Child behaviors at baseline influenced treatment results. Child emotional and
26 behavioral problems improved post-treatment. The results suggest that obesity treatment may
27 help in reducing emotional distress among preschoolers.

28

29 **Introduction**

30 Emotional and behavioral problems are common among children worldwide, with global
31 prevalence estimates as high as 13% (1). Most common are anxiety disorders (7%), followed by
32 disruptive disorders (6%), depressive disorders (4%), oppositional defiant disorder (4%),
33 attention-deficit-hyperactivity-disorder (ADHD) (3%) and conduct disorders (2%), alongside
34 combinations of the above and other, rarer disorders (1). Research studies have shown that
35 children's emotional and behavioral problems may be associated with psychosocial difficulties
36 and other health problems later in life (2).

37

38 Among child behavior problems, externalizing and internalizing behaviors (including those
39 associated with neurodevelopmental disorders such as ADHD and autism spectrum disorder) are
40 particularly prevalent among children with obesity, affecting about one-third of children referred
41 to obesity treatment (3). In children, ADHD may lead to abnormal eating patterns and
42 consequently obesity, although studies have not found consistent associations between child
43 ADHD and Body Mass Index (BMI) (4). Child temperament has also been associated with
44 obesity. A systematic review found that child temperament (i.e. poor self-regulation, distress to
45 limitations, low/high soothability, low negative affectivity) was associated with higher BMI in
46 infants and preschool-age children (5). More recently, a study showed that 5-year-old children
47 with externalizing temperament (e.g. impulsiveness, low self-regulation) are more likely to be
48 overweight (6). Another study showed that certain subgroups of preschoolers with difficult
49 temperament (easily distressed, inhibited, cry often and have low soothability) might be at risk

50 for eating and weight regulation problems later in life (7). According to a meta-analysis, children
51 with overweight and obesity tend to be more impulsive than children with normal weight (8),
52 and children with overweight are more likely to experience multiple associated psychological
53 problems, such as co-existing depression, anxiety or behavioral difficulties, compared to children
54 with normal weight (9). The associations described above suggest that children's emotional and
55 behavioral problems need to be considered in obesity treatment, particularly because obesity
56 treatment often entails structured and regulated eating routines that may run counter to some
57 children's temperament (9, 10).

58
59 Most studies on associations between childhood obesity and behavioral problems have focused
60 on school-aged children and adolescents. However, a recent population-based study has found
61 that overweight at the age of 6 is associated with more total behavioral problems (11).
62 Additionally, a longitudinal analysis of a prospective cohort study has shown that externalizing
63 behavior at 2 years of age is associated with a higher BMI from age 2 to age 12 (12). However,
64 evidence on how behavioral problems influence obesity treatment in children is limited (13-16)
65 and no study has examined associations between obesity treatment and behavioral problems
66 among children under the age of 6 years.

67
68 The present study reports on secondary outcomes of a Randomized Controlled Trial (RCT) based
69 in Stockholm County, Sweden, which assessed a new childhood obesity intervention for
70 preschool-aged children (defined as children between the ages of 4 and 6 years). The RCT
71 evaluated a parent support program (whose main focus was evidence-based parenting practices)
72 in comparison to standard treatment (whose main focus was food and physical activity) (17). The
73 RCT's primary outcome was changes in child weight status 12-months post baseline. Parents
74 filled out questionnaires at baseline and after 12 months. Among these, the Child Behavior

75 Checklist (CBCL), a validated instrument with two versions (one for preschool-age children (1.5-
76 5 years old) and one for school-age children (6-18 years old)), enabled us to assess the influence
77 of child emotional and behavioral problems on obesity treatment outcomes over time.

78
79 The CBCL is a robust and widely used instrument (18, 19), several studies have examined
80 associations between child behavior and child obesity using the CBCL. However, all studies but
81 two have focused on children older than 5 years of age, using the CBCL for school-age children
82 (CBCL 6-18). A study based in Greece used the CBCL among school-age children (mean age
83 11, n=141) to assess the prevalence of Internalizing and Externalizing Behaviors in children with
84 and without obesity and found that both Internalizing (e.g. anxiety problems, mood changes,
85 somatic complaints) and Externalizing Behaviors (e.g. aggressive, antisocial and/or delinquent)
86 were more prevalent in children with obesity (20). A large study in Chile (n= 11 207) found that
87 children aged 6 years with overweight or obesity had higher CBCL-measured behavioral
88 problems compared to their peers with normal weight (11). Similarly, a study conducted in South
89 Korea among children aged 10-12 years (n=504) found that children with overweight were more
90 likely to present patterns of emotional and behavioral problems (i.e. more social, delinquent and
91 total problems) compared to children with normal weight (21). Additionally, a study conducted
92 in Turkey that compared emotional and behavioral problems among adolescents aged 12-16
93 years with obesity (n=60) or normal weight (n=30) found that adolescents with obesity had
94 higher rates of emotional and behavioral problems, as measured by scores on the Anxiety-
95 Depression and Withdrawal scales, and by rates of social and total problems (22). Of note, the
96 subsample of adolescents with obesity who were recruited through an endocrinology clinic
97 (n=30) had higher emotional and behavioral problem scores than the subsample of adolescents
98 with obesity who were recruited through a local high school (n=30) (22). Moreover, a Dutch
99 research group found that among children aged 8-14 years (n=248), lower BMI z-score at

100 baseline and higher social competence were associated with obesity treatment success at a
101 younger age and with less somatic complaints (23). In contrast, a cross-sectional study based in
102 the United States with mothers of children aged 3 years of Latino and low-income background
103 (n=174) found no association between emotional problems and obesity (24). To further
104 understandings of how emotional and behavioral problems may be associated with childhood
105 obesity treatment, we decided to include the CBCL among the measurements in our RCT.

106
107 The overall aim of the present study is to examine preschool-aged children's emotional and
108 behavioral problems in relation to changes in weight status during obesity treatment. First, we
109 examined changes in problem behaviors from baseline to 12 months post-baseline. Second, we
110 examined baseline behaviors as predictors of obesity treatment outcomes (child BMI z-score)
111 and how changes in these behaviors may contribute to BMI z-score change 12 months post-
112 baseline.

113
114 We hypothesized that emotional and behavioral problems would be lessened at 12 months post
115 obesity treatment. We hypothesized that the effect would be more pronounced in the parent
116 support treatment, since it emphasized evidence-based parenting practices suitable for addressing
117 child behavioral problems. In addition, we expected that children with higher levels of baseline
118 emotional and behavioral problems would exhibit a smaller change in their BMI z-score, given
119 that managing eating situations might be more challenging for them. As for changes in emotional
120 and behavioral problems from baseline to 12 months post obesity treatment, we hypothesized
121 that greater reductions in emotional and behavioral problems would predict greater reductions in
122 child BMI z-score.

123

124 **Methods**

125 Study design and intervention

126 The present paper reports on secondary data from the More and Less (ML) Study, a parallel
127 open-label RCT. The overall aim of the ML Study was to evaluate a parent support program
128 compared to standard treatment for childhood obesity, with BMI z-score defined as the primary
129 outcome. Families with children aged 4 to 6 years (hereafter called 'preschool-age') diagnosed
130 with obesity were randomized to one of the two treatment approaches. The parent support
131 program (10 weekly sessions à 90 minutes) was delivered by trained dietitians. The sessions
132 centered on evidence-based parenting practices (e.g. encouragement, monitoring, positive
133 involvement, limit setting and problem solving along with regulation of emotional expression),
134 which were discussed and practiced through role playing. The sessions emphasized specific
135 lifestyle components (e.g. healthy eating, physical activity) to support a healthy home
136 environment (17). Standard treatment was delivered to individual families by pediatricians (the
137 first visit) and pediatric nurses (follow-up visits) in outpatient pediatric clinics.

138 The present paper presents data on child emotional and behavioral problems at baseline and 12
139 months post-baseline in relation to weight status. The primary outcomes of the RCT have been
140 published elsewhere; these findings show that children whose families were randomized to the
141 parent support group had a larger decrease in BMI z-score compared to standard treatment 12
142 months post-baseline (25). The protocol for the study was registered with the Clinical Trials
143 Registry, clinicaltrials.gov on February 14, 2013, (ID: NCT01792531).

144

145

146 Study participants

147 Participant recruitment was conducted through primary child health care, school health care and
148 outpatient pediatric clinics between 2012 and 2016 in Stockholm County, Sweden. The study
149 was approved by the Regional Ethical Board in Stockholm (dnr: 2011/1329-31/4; 2012/1104-32;

150 2012/2005-32; 2013/486-32; 2016/80-32). Informed consent forms were signed by all
151 participating parents. The ML study included children who were 4-6 years old, had obesity as
152 defined by international age- and sex-specific cut-offs (26), were not diagnosed with any chronic
153 disease or developmental problem, and whose parents were sufficiently proficient in Swedish to
154 fill out questionnaires and participate in discussion-centered treatment (17). The sample size was
155 determined based on BMI z-score (primary outcome) adjusting for a 21% dropout rate (17).

156

157 Measurements

158 *The CBCL*

159 The CBCL for the ages of 1.5 to 5 years was filled in by one of the parents at baseline and after
160 12 months. On the CBCL 1.5-5, child behavior is rated by parents in 99 questions plus one open
161 question, encompassing emotional, behavioral and social problems that, together with the total
162 score, allow for the classification of child behavior according to different scales (27). When
163 rating, parents use a 3-Point Likert scale, ranging from 0 (*not true*) to 1 (*somewhat or sometimes*
164 *true*) and 2 (*very often or often true*), and for the open-ended question parents can mention any
165 problem not covered by the preceding 99 questions. Examples include ‘*My child doesn’t want to*
166 *sleep alone*’ (part of the scales Sleep Problems and DSM-Anxiety Problems) and ‘*My child is*
167 *overeating*’ (part of the scale DSM-Affective Problems). The questions can be arranged to
168 classify child behavior according to different scales.

169

170 The CBCL consists of two scale systems, the *syndrome scale* and the *DSM scale*. The *syndrome*
171 *scale* categorizes child behavior into seven syndromes: Emotionally Reactive,
172 Anxious/Depressed, Somatic Complaints, Withdrawn, Attention Problems, Aggressive Behavior
173 and Sleep Problems (27). The syndromes can be further divided into two subscales. The
174 Internalizing Behaviors subscale includes the first four syndromes and describes conflicts within

175 the inner self. The Externalizing Behaviors subscale consists of the syndromes Attention
176 Problems and Aggressive Behavior, and describes problems with other people and their
177 expectations. Sleep Problems constitute a stand-alone subscale. Additionally, a Total Problem
178 score can be calculated using all 99 question items (27).

179
180 The *DSM scale* refers to the 4th edition of the American Psychiatric Association's *Diagnostic*
181 *and Statistical Manual of Mental Disorders (DSM-IV)* (28). In this scale, child behavior is
182 classified into five subscales: Affective Problems, Anxiety Problems, Pervasive Developmental
183 Problems, Attention Deficit/Hyperactivity Problems (ADHD) and Oppositional Defiant
184 Problems (27). High scores on a CBCL DSM-oriented scale do not correspond to a DSM
185 diagnosis, but help identify children who might meet the criteria for clinically relevant
186 symptoms. In Sweden, the CBCL was validated in clinical and non-clinical population (n=2088).
187 The face validity was assessed with several methods, including pediatric screening interview,
188 Life Stress Score (50 items), Edinburg Postal Depression Scale and Sense of Coherence (19).

189 In this study, we compare our results to reference values from a Danish sample (29) and from a
190 large international sample (30) to place the scores from the Swedish sample of children with
191 obesity in context. The international pool includes both the syndrome scale and the DSM-
192 oriented scale. The Danish sample leaves out the scores for the seven syndromes and includes
193 only Internalizing and Externalizing Behaviors, Total Problems and the DSM-oriented scale.

194

195 *BMI z-score*

196 The children's weight and height were measured to the nearest 0.1 kg/cm by trained healthcare
197 professionals with calibrated measuring tools and in outpatient pediatric clinics at baseline and
198 after 12 months (17). Children were measured in light clothing. Each measurement was taken
199 three times, and a mean value was calculated. The BMI was calculated as follows: (weight in kg)

200 / (height in m)². The BMI z-score was derived using Swedish age- and sex-specific reference
201 values (31).

202 Statistical Analyses

203 To assess and evaluate children's behavior before and after the intervention, the mean scores of
204 the subscales of the syndrome scale and the DSM scale and the Total Problem score were
205 computed. The mean scores at baseline and at 12 months post-baseline were then compared by
206 using dependent paired samples t-tests. Linear regression analyses were conducted to examine
207 (1) the relationship between baseline child behaviors and baseline BMI z-score; (2) the
208 relationship between baseline child behaviors and changes in child BMI z-score after 12 months
209 (post BMI z-scores minus pre BMI z-scores); (3) the relationship between changes in child
210 behavior after 12 months (post scores minus baseline scores) and changes in child BMI z-score
211 after 12 months. Each of these analyses was carried out using four different models, which
212 focused on the influence of child behavior according to: (1) the seven syndromes, (2)
213 Internalizing and Externalizing Behaviors, (3) the Total Problem score, and (4) the DSM-
214 oriented sub-scales. All models were adjusted for child age and gender, as well as parents'
215 income, foreign background and BMI self-reported at baseline. The software package 'IBM
216 SPSS Statistics 24' was used for all statistical analyses. P-values <0.05 were regarded as
217 significant in all analyses.

218

219 **Results**

220 Complete questionnaires (with less than 9 items missing) were present for 121 children at
221 baseline and for 83 children at 12 months post-baseline. Among these, 77 children had complete
222 CBCL data both at baseline and at 12 months. The data for the 77 children who had complete
223 CBCL data at both time points form the basis for the present analysis. The background
224 characteristics of the sample are presented in Table 1. Children in the total sample were on

225 average 5.2 years old (*SD* 0.8), with the children in the parent support program being slightly
226 younger (5.1 years, *SD* 0.8) than in the standard treatment (5.3 years, *SD* 0.7). Half of the sample
227 were girls (53%), with more girls in the standard treatment (67%) compared to the parent support
228 program (40%). The average BMI z-score in the total sample was 3.0 (*SD* 0.6). The mothers
229 were on average 36.8 years old (*SD* 5.6) and the fathers 39.4 years old (*SD* 6.7). Most parents
230 had overweight (mothers: 28%; fathers: 42%) or obesity (mothers: 36%; fathers: 49%).
231 Approximately half of the parents in the total sample were of foreign origin; more mothers than
232 fathers had a university degree (mothers: 46%; fathers: 40%). The study population, of families
233 who provided CBCL data, differed from the total population of families enrolled in in the ML
234 in two aspects: parents who did not fill out any of the CBCL questionnaires were more likely to
235 be of foreign origin (mothers: 81%; fathers: 62%) compared to the parents who filled out both
236 CBCL questionnaires at baseline and at 12 months (mothers: 51%; fathers: 52%); these parents
237 were also less likely to have a university degree (mothers: 28%; fathers: 29%), compared to the
238 parents who provided complete CBCL data (mothers: 46%; fathers: 40%) (data not shown).

239
240 Mean scores on the different CBCL scales for the total sample, parent support program and the
241 standard treatment are presented in Table 2a, in comparison to reference values from an early
242 validation study among Danish preschoolers and the international reference values (19, 20). The
243 mean scores of our population are similar to those of a non-clinical population (18) and to a
244 Danish population (30). Table 2b presents differences related to changes in CBCL scores
245 between baseline and follow-up. For the total sample, significant decreases were observed from
246 baseline to 12 months in Aggressive Behavior, Externalizing Behaviors, Total Problems and
247 Oppositional Defiant Problems. Children in the parent support program had no significant
248 changes in CBCL scores from baseline to 12 months. In the standard treatment group there was

249 a significant decrease from baseline to 12 months in Emotional Reactivity, Sleep Problems, Total
250 Problem score, Affective Problems and Oppositional Defiant Problems.

251
252 Linear regression analyses of associations between child behavior and BMI z-score before
253 childhood obesity treatment found that child behavior at baseline had no association with child
254 BMI z-score at baseline (data not shown). However, significant associations were found between
255 child behavior at baseline and changes in child BMI z-score (Table 3). In the parent support
256 program, children who scored high on ADHD at baseline were more likely to experience
257 increases in BMI z-score after 12 months, whereas children who scored high on Oppositional
258 Defiant Problems at baseline were more likely to experience decreases in BMI z-score. In the
259 standard treatment group, children with more baseline Externalizing Behaviors, Total Problems
260 and/or Oppositional Defiant Problems were more likely to experience decreases in BMI z-score.
261 For the whole sample, children who scored high on Attention Problems and ADHD at baseline
262 were more likely to experience increases in BMI z-score, while children who scored high on
263 Oppositional Defiant Problems were more likely to experience decreases in BMI z-scores.

264
265 Table 4 demonstrates how changes in child behavior were associated with changes in BMI z-
266 scores. Only one significant association was identified, in the standard group, between reductions
267 in symptoms of Depression and Anxiety and increasing BMI z-scores.

268
269 **Discussion**
270 This is the first study to assess how preschool-age children’s emotional and behavioral problems
271 evolve over the course of obesity treatment and to identify the influence of specific behaviors on
272 obesity treatment effects. Fewer problematic behaviors were reported by parents at 12 months
273 post-baseline in the total sample of children and in the standard treatment group; no significant

274 changes were found in the parent support group. Examining baseline emotional problems as
275 predictors of treatment results, we found that for the total sample and for the parent support
276 group, children with Attention Problems and ADHD were more likely to have increased BMI z-
277 scores. For all groups, children with Oppositional Defiant Problems were more likely to
278 experience a decrease in BMI z-scores. Finally, when examining how changes in child behaviors
279 influenced changes in weight status during treatment, we found that children in standard
280 treatment with reduced Anxiety and Depression scores were more likely to experience an
281 increase in their BMI z-scores.

282
283 It is promising that obesity treatment was associated with reduced levels of emotional and
284 behavioral problems in preschool-age children. Previous childhood obesity intervention studies,
285 which involved parents of older children, have shown favorable results with respect to child
286 behavioral problems (15, 32). Obesity interventions that target preschool-age children are still
287 rare (33, 34), with few RCTs; of the extant RCTs, the secondary outcomes are rarely reported,
288 especially with regard to child psychosocial functioning. In the present study, the positive gains
289 in emotional and behavioral wellbeing post-treatment may indicate that parents were empowered
290 through the program, and were better able to respond to their children's emotional distress and
291 behavioral problems. Although the study had low statistical power due to the small sample sizes
292 in each treatment arm, and as such warrants cautious interpretations, it is interesting that, contrary
293 to our hypothesis, children's emotional and behavioral problems improved more significantly
294 after standard treatment. This is particularly surprising because the evidence-based practices
295 taught in the parent support group have been successful in improving difficult child behavior
296 (35). However, as child behaviors were reported by parents, it is possible that the parent group
297 approach made parents more attuned to children's emotional and behavioral expressions,
298 resulting in higher behavioral problem scores compared to the standard treatment group.

299

300 The reported degree of behavioral problems was very low in comparison with rates in the
301 international validation study, with the exception of Sleep Problems and Affective Problems
302 (30). While the prevalence of Sleep Problems among preschoolers is known to differ among
303 populations (36), it should be noted that the parents in this study constituted a treatment-seeking
304 clinical population. Hence, the high prevalence of Sleep Problems could be an indicator of a
305 challenging or stressful home environment. Poor sleep quality has also been associated with
306 obesity in children and adolescents (37). For Affective Problems, the scores in this study were
307 higher than the international and Danish scores (29, 30). While few studies have explored
308 affective problems in young children, stress is an important risk factor for the development of
309 affective problems, especially in high-risk groups (38). Although stress was not measured in this
310 study, elevated stress in the home environment might underlie the high reported scores of
311 Affective Problems. In a previous study, a significant association between maternal depression
312 and child emotional and behavioral problems was found in a subsample of the ML population
313 (39). Maternal depression is also a major risk factor for the development of distress and
314 depression among children (40), and the high scores of reported Affective Problems might be
315 due, in part, to maternal affective problems.

316

317 The study's findings of a relationship between baseline ADHD and Attention Problems and an
318 increase in BMI z-score after treatment are in line with previous research (41). Especially in
319 young children, Attention Problems or ADHD symptoms predict increased body fat mass later
320 in childhood (42). While the association between ADHD and obesity is not yet understood,
321 ADHD may affect obesity treatment through an interplay of environmental, biological and
322 lifestyle factors (e.g. physical activity, eating patterns, genetics, impulsivity, parental behavior)
323 (43). In particular, impulsivity may influence obesity treatment, as children who display

324 impulsivity have been found to be food-reactive and experience difficulties in self-control (44,
325 45). It is important to note that although Attention Problems were associated with increased BMI
326 z-scores after obesity treatment, BMI z-scores might have increased even further without any
327 treatment.

328
329 Compared to potential associations between ADHD and obesity, previous research has not paid
330 as much attention to potential associations between Oppositional Defiant Problems and
331 childhood obesity. We found that a higher baseline degree of Oppositional Defiant Problems was
332 associated with decreases in children's BMI z-score over the course of obesity treatment. These
333 findings stand in contrast to a study conducted in Germany, which found that Oppositional
334 Defiant Disorder, together with conduct disorder, were strong predictors of overweight and
335 obesity, even after controlling for comorbid ADHD (46). A possible explanation is that the data
336 used in the German study included older children (6 to 12 years old) referred to psychiatric
337 treatment (46). We also found that higher scores at baseline in Externalizing Behaviors were
338 associated with a decrease in BMI z-score in the standard treatment group. Although we did not
339 find an association between changes in Externalizing behaviors and changes in BMI z-score, we
340 found that these behaviors decreased for both groups during the 12-month intervention. A study
341 conducted in the U.S. also found a co-occurrence for Externalizing Behaviors and obesity among
342 5-year-old children (47). Neither the German study nor the U.S. study reported on obesity
343 interventions, such that their findings are not directly comparable to the present study. Our
344 findings are in line with a recent study that evaluated a clinical population of 662 children
345 (ranging in age from 4 to 20 years) who participated in a family-based obesity program focusing
346 on nutrition over one year (16). The authors highlighted the role of impulsivity in externalizing
347 behaviors, suggesting that the structure provided by the program was particularly beneficial for
348 children exhibiting those behaviors (16). This explanation might also apply to the standard

349 treatment group in our study. Further analysis is needed to understand whether the parent support
350 group's reduced focus on lifestyle components such as eating patterns and physical activity might
351 have contributed to the lack of significant reduction in externalizing behaviors in this group.

352
353 Finally, while we found that changes in child emotional and behavioral problems were associated
354 with changes in BMI z-score in the standard treatment group, it remains unclear why children
355 whose parents reported reductions in depression/anxiety also experienced increases in BMI z-
356 score levels. Though depression/anxiety results for the parent support program did not reach
357 statistical significance, they pointed to reductions over time, which is in accordance with
358 previous research. The evidence from previous research on associations between depression and
359 weight status presents mixed results. Some studies have indicated that weight increases with
360 depressive symptoms in adolescence (48), while other studies found that adolescents with
361 depressive symptoms have a lower BMI (49), but no studies have assessed how affective
362 symptoms might related to body weight among young children. Hence, more research is needed
363 to evaluate associations between affective problems among preschool-age children and obesity.

364

365 *Strengths and limitations*

366 A notable strength of this study was the use of high-quality measurements. The CBCL has been
367 validated in 23 countries in Asia, Australasia, Europe (including Sweden), the Middle East and
368 South America, including 19,000 children of preschool age, and was found to be valid and
369 reliable in measuring children's emotional, behavioral and social problems across cultures (18).
370 Moreover, the child anthropometrics were measured by trained health care professionals, thus
371 removing possible self-reporting bias concerning height and weight (17). Another strength of the
372 study was its pre- and post-treatment design, which allowed us to examine changes in CBCL
373 scores following an obesity intervention. The study was limited by the absence of blinding for

374 participants and researchers, who were all aware of which treatment was being administered.
375 Further, for ethical reasons, this RCT did not include a control group with a no-treatment
376 approach; given that all recruited children were diagnosed with clinical obesity, they were
377 offered treatment. Additionally, the study's power, established on primary outcomes (the BMI
378 z-score and not on the CBCL), may have influenced results close to significance although they
379 had a high standardized beta. While the sociodemographic variation among parents with regards
380 to educational level and foreign background was wide, children with missing CBCL data were
381 more likely to have parents with a foreign background and lower educational levels. This may
382 have affected the results and could be due to the nature of the comprehensive CBCL, which
383 includes 100 items. Finally, the clinical relevance of the reported changes in child behavior
384 problems is not clear.

385

386 *Conclusions*

387 This study was the first to assess how preschool-age children's behavior changes following
388 obesity treatment, and to identify the influence of specific behavioral patterns on obesity
389 treatment effects. After 12 months of treatment, children in the total sample experienced
390 reductions in scores for Emotional Reactivity, Sleep and Affective Problems, Aggressive and
391 Externalizing Behaviors, Oppositional Defiant Problems, and Total Problems. Attention
392 Problems were more likely to contribute to increasing BMI z-scores, while Oppositional Defiant
393 Problems, Externalizing Behaviors and Total Problems contributed to decreasing BMI z-scores.
394 The results suggest that obesity treatment may provide the necessary structure to support children
395 with emotional and behavioral problems and their parents. Future childhood obesity
396 interventions should incorporate the measurement of child emotional and behavioral problems
397 to assess the impact of these interventions on children's emotional and behavioral wellbeing.

398

399 **Conflicts of interest statement**

400 The authors have no financial relationships relevant to this article to disclose and no
401 potential conflicts of interest to disclose.

402

403 **Funding**

404 This study was supported by the Swedish Research Council (2014-02404), Karolinska
405 Institutet Doctoral Funds, the Swedish Society of Medicine, VINNOVA (2011-3443),
406 Jerring Foundation, Samariten Foundation, Magnus Bergvall Foundation, Ingrid and
407 Fredrik Thuring Foundation, Helge Ax:son Foundation, Crown Princess Lovisa
408 Foundation, Foundation Frimurare Barnhuset in Stockholm, Foundation Pediatric Care,
409 Foundation Martin Rind, Jane and Dan Olsson Foundation, Groschinsky Foundation,
410 Sigurd and Elsa Golje Memory Foundation and iShizu Matsumurais Donation. The
411 funding sources had no role in the study design, collection, analysis or interpretation of
412 the data, writing the manuscript, or the decision to submit the paper for publication.

413

414 **Acknowledgements**

415 We want to thank all participating families, child health care and school nurses and all personnel
416 involved in the standard treatment offered in the pediatric outpatient clinics. We also thank Sofia
417 Ljung, Jonna Nyman, Mahnoush Etminan Malek, Karin Nordin, Kathryn Lewis Chamberlain,
418 Jan Ejderhamn, Philip A. Fisher, Patricia Chamberlain and Claude Marcus who were involved
419 in design or in data collection in the More and Less Study. PN conceived the idea of this study
420 in collaboration with EE, AE, ES and SK. EE drafted the paper together with KE and PS. EE
421 conducted the main analyses. PN was responsible for the data collection and critically reviewed
422 all analyses together with PS, MS, AE and ES. All authors made substantial contributions to
423 conception and design, data collection and to interpretation of the data. All authors contributed
424 to reviewing and approving the final manuscript.

425 **REFERENCES**

- 426 1. Polanczyk GV, Salum GA, Sugaya LS, Caye A, Rohde LA. Annual research
427 review: A meta-analysis of the worldwide prevalence of mental disorders in children
428 and adolescents. *Journal of child psychology and psychiatry, and allied disciplines*.
429 2015;56(3):345-65.
- 430 2. Layard R DJ. *A good childhood: Searching for values in a competitive age*.
431 London: Penguin; 2009. 256 p.
- 432 3. Wentz E, Bjork A, Dahlgren J. Neurodevelopmental disorders are highly over-
433 represented in children with obesity: A cross-sectional study. *Obesity (Silver Spring,*
434 *Md)*. 2017;25(1):178-84.
- 435 4. Tong L, Shi H, Li X. Associations among ADHD, Abnormal Eating and
436 Overweight in a non-clinical sample of Asian children. *Scientific reports*.
437 2017;7(1):2844.
- 438 5. Bergmeier H, Skouteris H, Horwood S, Hooley M, Richardson B. Associations
439 between child temperament, maternal feeding practices and child body mass index
440 during the preschool years: a systematic review of the literature. *Obesity reviews : an*
441 *official journal of the International Association for the Study of Obesity*. 2014;15(1):9-
442 18.
- 443 6. Skogheim TS, Vollrath ME. Associations of Child Temperament with Child
444 Overweight and Breakfast Habits: A Population Study in Five-Year-Olds. *Nutrients*.
445 2015;7(12):10116-28.
- 446 7. Messerli-Burgy N, Stulb K, Kakebeeke TH, Arhab A, Zysset AE, Leeger-
447 Aschmann CS, et al. Emotional eating is related with temperament but not with stress
448 biomarkers in preschool children. *Appetite*. 2018;120:256-64.

- 449 8. Thamotharan S, Lange K, Zale EL, Huffhines L, Fields S. The role of
450 impulsivity in pediatric obesity and weight status: a meta-analytic review. *Clinical*
451 *psychology review*. 2013;33(2):253-62.
- 452 9. Rankin J, Matthews L, Cobley S, Han A, Sanders R, Wiltshire HD, et al.
453 Psychological consequences of childhood obesity: psychiatric comorbidity and
454 prevention. *Adolescent health, medicine and therapeutics*. 2016;7:125-46.
- 455 10. Goldschmidt AB, Sinton MM, Aspen VP, Tibbs TL, Stein RI, Saelens BE, et al.
456 Psychosocial and familial impairment among overweight youth with social problems.
457 *International journal of pediatric obesity : IJPO : an official journal of the International*
458 *Association for the Study of Obesity*. 2010;5(5):428-35.
- 459 11. Kagawa RMC, Fernald LCH, Behrman JR. Weight Status and Behavioral
460 Problems among Very Young Children in Chile. *PLOS ONE*. 2016;11(9):e0161380.
- 461 12. Anderson SE, He X, Schoppe-Sullivan S, Must A. Externalizing behavior in
462 early childhood and body mass index from age 2 to 12 years: longitudinal analyses of a
463 prospective cohort study. *BMC pediatrics*. 2010;10:49-.
- 464 13. Stavrou S, Nicolaides NC, Papageorgiou I, Papadopoulou P, Terzioglou E,
465 Chrousos GP, et al. The effectiveness of a stress-management intervention program in
466 the management of overweight and obesity in childhood and adolescence. *Journal of*
467 *molecular biochemistry*. 2016;5(2):63-70.
- 468 14. de Niet J, Timman R, Jongejan M, Passchier J, van den Akker E. Predictors of
469 participant dropout at various stages of a pediatric lifestyle program. *Pediatrics*.
470 2011;127(1):e164-70.
- 471 15. Myers MD, Raynor HA, Epstein LH. Predictors of Child Psychological Changes
472 During Family-Based Treatment for Obesity. *Archives of Pediatrics & Adolescent*
473 *Medicine*. 1998;152(9):855-61.

- 474 16. Blomquist K, Griffin S, Hays E, Schmalz D, Sease K, Reeves C. Behavioral and
475 Psychological Predictors of BMI Reduction in Children Seeking Treatment at a Hospital
476 Clinics Family Based Pediatric Weight Management Program. *Journal of Childhood*
477 *Obesity*. 2018;03(2).
- 478 17. Ek A, Chamberlain KL, Ejderhamn J, Fisher PA, Marcus C, Chamberlain P, et
479 al. The More and Less Study: a randomized controlled trial testing different approaches
480 to treat obesity in preschoolers. *BMC public health*. 2015;15:735.
- 481 18. Ivanova MY, Achenbach TM, Rescorla LA, Harder VS, Ang RP, Bilenberg N,
482 et al. Preschool psychopathology reported by parents in 23 societies: testing the seven-
483 syndrome model of the child behavior checklist for ages 1.5-5. *Journal of the American*
484 *Academy of Child and Adolescent Psychiatry*. 2010;49(12):1215-24.
- 485 19. Höök B, Cederblad M. *Prövning av CBCL för förskolebarn (ASEBA)*. Umeå,
486 Sweden: Umeå University, Child and Adolescent Psychiatry DoCS; 2008.
- 487 20. Pervanidou P, Bastaki D, Chouliaras G, Papanikolaou K, Kanaka-Gantenbein C,
488 Chrousos G. Internalizing and externalizing problems in obese children and adolescents:
489 associations with daily salivary cortisol concentrations. *Hormones (Athens, Greece)*.
490 2015;14(4):623-31.
- 491 21. Hwang JW, Lyoo IK, Kim BN, Shin MS, Kim SJ, Cho SC. The relationship
492 between temperament and character and psychopathology in community children with
493 overweight. *Journal of developmental and behavioral pediatrics : JDBP*. 2006;27(1):18-
494 24.
- 495 22. Erermis S, Cetin N, Tamar M, Bukusoglu N, Akdeniz F, Goksen D. Is obesity a
496 risk factor for psychopathology among adolescents? *Pediatrics international : official*
497 *journal of the Japan Pediatric Society*. 2004;46(3):296-301.

- 498 23. de Niet J, Timman R, Rokx C, Jongejan M, Passchier J, van Den Akker E.
499 Somatic complaints and social competence predict success in childhood overweight
500 treatment. *International journal of pediatric obesity : IJPO : an official journal of the*
501 *International Association for the Study of Obesity*. 2011;6(2-2):e472-9.
- 502 24. Nagata JM, Hagan MJ, Heyman MB, Wojcicki JM. No Association between
503 Obesity and Behavior in Low-income, Preschool Latino Children. *Journal of health care*
504 *for the poor and underserved*. 2015;26(2):410-20.
- 505 25. Ek A, Lewis Chamberlain K, Sorjonen K, Hammar U, Etminan Malek M,
506 Sandvik P, et al. A parent treatment program for preschoolers with obesity: A
507 randomized controlled trial. *Pediatrics*. Provisionally accepted March 19 2019.
- 508 26. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition
509 for child overweight and obesity worldwide: international survey. *BMJ (Clinical*
510 *research ed)*. 2000;320(7244):1240-3.
- 511 27. Achenbach TM, & Rescorla, L. A. *Manual for the ASEBA School-Age Forms*
512 *and Profiles*. Burlington, VT: University of Vermont Research Center for Children,
513 Youth, & Families; 2000.
- 514 28. Association AP. *Diagnostic and statistical manual of mental disorders: DSM-IV-*
515 *TR*. Washington, DC: American Psychiatric Association; 2000.
- 516 29. Kristensen S, Henriksen TB, Bilenberg N. The Child Behavior Checklist for
517 Ages 1.5-5 (CBCL/1(1/2)-5): assessment and analysis of parent- and caregiver-reported
518 problems in a population-based sample of Danish preschool children. *Nordic journal of*
519 *psychiatry*. 2010;64(3):203-9.
- 520 30. Rescorla LA, Achenbach TM, Ivanova MY, Harder VS, Otten L, Bilenberg N, et
521 al. International comparisons of behavioral and emotional problems in preschool
522 children: parents' reports from 24 societies. *Journal of clinical child and adolescent*

523 psychology : the official journal for the Society of Clinical Child and Adolescent
524 Psychology, American Psychological Association, Division 53. 2011;40(3):456-67.

525 31. Karlberg J, Luo ZC, Albertsson-Wikland K. Body mass index reference values
526 (mean and SD) for Swedish children. *Acta paediatrica* (Oslo, Norway : 1992).
527 2001;90(12):1427-34.

528 32. Vignolo M, Rossi F, Bardazza G, Pistorio A, Parodi A, Spigno S, et al. Five-
529 year follow-up of a cognitive-behavioural lifestyle multidisciplinary programme for
530 childhood obesity outpatient treatment. *European journal of clinical nutrition*.
531 2008;62(9):1047-57.

532 33. Loveman E, Al-Khudairy L, Johnson RE, Robertson W, Colquitt JL, Mead EL,
533 et al. Parent-only interventions for childhood overweight or obesity in children aged 5 to
534 11 years. *The Cochrane database of systematic reviews*. 2015(12):Cd012008.

535 34. Colquitt JL, Loveman E, O'Malley C, Azevedo LB, Mead E, Al-Khudairy L, et
536 al. Diet, physical activity, and behavioural interventions for the treatment of overweight
537 or obesity in preschool children up to the age of 6 years. *The Cochrane database of*
538 *systematic reviews*. 2016;3:Cd012105.

539 35. Chamberlain P, Price J, Leve LD, Laurent H, Landsverk JA, Reid JB. Prevention
540 of behavior problems for children in foster care: outcomes and mediation effects.
541 *Prevention science : the official journal of the Society for Prevention Research*.
542 2008;9(1):17-27.

543 36. Mindell JA, Sadeh A, Kwon R, Goh DY. Cross-cultural differences in the sleep
544 of preschool children. *Sleep medicine*. 2013;14(12):1283-9.

545 37. Fatima Y, Doi SA, Mamun AA. Sleep quality and obesity in young subjects: a
546 meta-analysis. *Obesity reviews : an official journal of the International Association for*
547 *the Study of Obesity*. 2016;17(11):1154-66.

- 548 38. Ostiguy CS, Ellenbogen MA, Linnen AM, Walker EF, Hammen C, Hodgins S.
549 Chronic stress and stressful life events in the offspring of parents with bipolar disorder.
550 Journal of affective disorders. 2009;114(1-3):74-84.
- 551 39. Hedman J. Psychological problems in preschoolers with obesity and the link to
552 depression in their parents [Bachelor Thesis]. Stockholm, Sweden: Karolinska Institutet;
553 2013.
- 554 40. Beardslee WR, Bemporad J, Keller MB, Klerman GL. Children of parents with
555 major affective disorder: a review. The American journal of psychiatry.
556 1983;140(7):825-32.
- 557 41. Quesada D, Ahmed NU, Fennie KP, Gollub EL, Ibrahimou B. A Review:
558 Associations Between Attention-deficit/hyperactivity Disorder, Physical Activity,
559 Medication Use, Eating Behaviors and Obesity in Children and Adolescents. Archives
560 of psychiatric nursing. 2018;32(3):495-504.
- 561 42. Bowling AB, Tiemeier HW, Jaddoe VWV, Barker ED, Jansen PW. ADHD
562 symptoms and body composition changes in childhood: a longitudinal study evaluating
563 directionality of associations. Pediatric obesity. 2018;13(9):567-75.
- 564 43. Hanc T, Cortese S. Attention deficit/hyperactivity-disorder and obesity: A
565 review and model of current hypotheses explaining their comorbidity. Neuroscience and
566 biobehavioral reviews. 2018;92:16-28.
- 567 44. Graziano PA, Calkins SD, Keane SP. Toddler self-regulation skills predict risk
568 for pediatric obesity. International journal of obesity (2005). 2010;34(4):633-41.
- 569 45. Nederkoorn C, Braet C, Van Eijs Y, Tanghe A, Jansen A. Why obese children
570 cannot resist food: the role of impulsivity. Eating behaviors. 2006;7(4):315-22.
- 571 46. Pauli-Pott U, Neidhard J, Heinzl-Gutenbrunner M, Becker K. On the link
572 between attention deficit/hyperactivity disorder and obesity: Do comorbid oppositional

573 defiant and conduct disorder matter? *European child & adolescent psychiatry*.
574 2013;23(7):531-7.

575 47. Suglia SF, Duarte CS, Chambers EC, Boynton-Jarrett R. Social and behavioral
576 risk factors for obesity in early childhood. *Journal of developmental and behavioral*
577 *pediatrics : JDBP*. 2013;34(8):549-56.

578 48. Hillman J, Dorn L, Huang B. Association of Anxiety and Depressive Symptoms
579 and Adiposity Among Adolescent Females, Using Dual Energy X-ray Absorptiometry.
580 *Clinical pediatrics*. 2010;49(7):671-7.

581 49. Matziou V, Hatira K, Perdikaris P, Zachos I, Mellou K, Bakoula C. Weight
582 status and depressive symptoms in 18 year-old Greek adolescents. *Pediatric reports*.
583 2010;2(2):e16.

584

585 **Table 1.** Background characteristics of the study sample

	Whole sample (n=77)	Parent-group treatment (n=38)	Standard treatment (n=39)
	Mean (sd) or %	Mean (sd) or %	Mean (sd) or %
Child			
Age (years)	5.2 (0.7)	5.1 (0.8)	5.3 (0.7)
Sex (girl)	53	40	67
First born	49	40	58
Live with both parents	78	76	77
Number of siblings	1.1 (0.8)	1.1 (0.8)	1.0 (0.9)
BMI z-score	2.9 (0.6)	3.0 (0.6)	2.9 (0.6)
Mother			
Age (years)	36.8 (5.6)	38.1 (5.5)	36.6 (5.9)
BMI	28.4 (6.3)	28.7 (6.9)	28.1 (5.7)
Weight category			
<i>Normal weight</i>	36	38	34
<i>Overweight</i>	28	27	29
<i>Obesity</i>	36	35	37
Foreign origin	51	50	53
University degree	46	50	42
Income (SEK/month)			
< 10 000	18	11	26
10 000 < 20 000	40	29	50
20 000 < 30 000	33	45	21
30 000 < 40 000	8	13	3
40 000 < 50 000	0	0	0
> 50 000	1	3	0
Fathers			
Age (years)	39.4 (6.7)	39.8 (6.8)	38.9 (6.6)
BMI	30.3 (4.7)	30.4 (4.8)	30.2 (4.6)
Weight category			
<i>Normal weight</i>	9	9	8
<i>Overweight</i>	42	42	42
<i>Obesity</i>	49	49	50
Foreign origin	52	55	50
University degree	40	34	44
Income level (SEK/month)			
< 10 000	9	7	11
10 000 < 20 000	21	19	22
20 000 < 30 000	54	52	56
30 000 < 40 000	13	16	11
40 00 < 50 000	2	3	0
> 50 000	2	3	0

586 **Note:** Parents weight categories according to the World Health Organization's BMI cut-off criteria (Normal
587 weight BMI=18.5-24.9; Overweight BMI ≥ 25.0; Obesity BMI ≥ 30.0). **Abbreviations:** BMI, Body Mass Index;
588 SEK, Swedish kronor (mean monthly income level in 2015: 33 600 SEK (men), 26 400 SEK (women)); Foreign
589 origin, parent and both grandparents born in other country than Sweden or parent born in Sweden and
590 grandparents born abroad. **Missing values:** 1 child for 'First born' and 'Number of siblings', 1 mother for all

591 maternal variables + 1 mother for 'BMI' and 'weight category', 8 fathers missing for all variables + 1 for 'Age'
592 and 'University degree' + 2 for 'income level'

Table 2.a. Child Behavior Checklist for Ages 1.5-5. Total scores for all subscales at baseline and 12-months with mean values (*SD*) for the whole sample, Parent-group treatment and Standard treatment including reference values

	Whole Sample (n=77)		Parent-group treatment (n=38)		Standard treatment (n=39)		Danish Reference values**	International Reference values*
	Baseline	12 months	Baseline	12 months	Baseline	12 months		
Syndrome Scale (No of items)								
Emotionally Reactive (9 items)	2.30 (2.76)	1.91 (2.92)	2.08 (2.47)	1.97 (3.16)	2.51 (3.03)	1.85 (2.70)	n.a.	2.7 (0.7)
Anxious/Depressed (8 items)	1.47 (1.95)	1.43 (2.27)	1.39 (1.82)	1.76 (2.67)	1.54 (2.10)	1.10 (1.78)	n.a.	3.1 (1.0)
Somatic Complaints (11 items)	1.32 (1.82)	1.19 (1.38)	1.24 (1.83)	1.03 (1.32)	1.41 (1.83)	1.36 (1.44)	n.a.	2.3 (0.6)
Withdrawn (8 items)	1.08 (1.29)	1.12 (1.34)	1.39 (1.32)	1.26 (1.48)	0.77 (1.20)	0.97 (1.20)	n.a.	1.7 (0.7)
Sleep Problems (7 items)	2.88 (2.22)	2.57 (2.62)	2.87 (2.18)	3.37 (2.91)	2.90 (2.29)	1.79 (2.05)	n.a.	2.7 (0.3)
Attention Problems (5 items)	1.78 (1.68)	1.51 (1.84)	2.00 (1.75)	1.66 (1.79)	1.56 (1.60)	1.36 (1.89)	n.a.	2.3 (0.6)
Aggressive Behavior (19 items)	6.90 (5.44)	5.66 (5.96)	7.03 (5.11)	6.00 (5.53)	6.77 (5.80)	5.33 (6.41)	n.a.	9.7 (1.8)
Total Internalizing Behaviors (36 items)	6.17 (6.55)	5.65 (6.68)	6.11 (6.21)	6.03 (7.45)	6.23 (6.95)	5.28 (5.91)	3.9 (5.1)	9.6 (2.6)
Total Externalizing Behaviors (24 items)	8.68 (6.55)	7.17 (7.33)	9.03 (6.31)	7.66 (6.75)	8.33 (6.83)	6.69 (7.91)	6.7 (6.0)	12.0 (2.3)
Total Problems score (100 items)	25.7 (18.34)	22.29 (19.27)	25.82 (16.72)	24.13 (19.26)	25.59 (20.01)	20.49 (19.37)	17.3 (15.3)	33.3 (7.1)
DSM Scale (No of items)								
Affective Problems (10 items)	2.86 (2.14)	2.60 (2.31)	2.74 (2.03)	2.82 (2.60)	2.97 (2.26)	2.38 (1.99)	1.1 (1.6)	2.3 (0.7)
Anxiety Problems (10 items)	2.68 (2.41)	2.58 (2.52)	2.61 (2.40)	2.87 (2.48)	2.74 (2.46)	2.31 (2.56)	1.6 (1.9)	3.8 (1.2)
Pervasive Developmental Problems (13 items)	2.27 (2.61)	2.09 (2.35)	2.79 (2.50)	2.42 (2.46)	1.77 (2.66)	1.77 (2.21)	1.6 (2.3)	3.2 (0.9)
ADHD (6 items)	3.04 (2.34)	2.66 (2.46)	3.26 (2.42)	2.97 (2.42)	2.82 (2.27)	2.36 (2.50)	2.3 (2.2)	4.7 (1.0)
Oppositional Defiant Problems (6 items)	2.83 (2.42)	2.13 (2.37)	2.71 (2.34)	2.08 (2.25)	2.95 (2.52)	2.18 (2.51)	2.3 (2.1)	3.3 (0.6)

* International Reference Values from Rescorla et al. (2011)

** Danish Reference Values from Kristensen, Henriksen & Bilenberg (2010)

Abbreviations: ADHD, Attention Deficit/Hyperactivity Problems; BMI, Body Mass Index (weight in kilograms/ height in meters squared); DSM, Diagnostic and Statistical Manual of Mental Disorders (4th edition, 1994), Parent-group treatment = child obesity treatment focused on general parenting; Standard treatment = child obesity treatment focused on lifestyle

Table 2.b. Child Behavior Checklist for Ages 1.5-5. Changes in total scores for all subscales at 12 months compared to baseline with mean values (*SD*) for the whole sample, Parent-group treatment and Standard treatment

	Whole sample (n=77)			Parent-group treatment (n=38)			Standard treatment (n=39)		
	Mean (SD)	p-value	95% CI	Mean (SD)	p-value	95% CI	Mean (SD)	p-value	95% CI
Syndrome Scale (No of items)									
Emotionally Reactive (9 items)	-0.39 (2.06)	0.10	(-0.85;0.07)	-0.10 (2.05)	0.75	(-0.77;0.56)	-0.66 (2.06)	0.05	(-1.33;0.00)
Anxious/Depressed (8 items)	-0.03 (2.02)	0.86	(-0.49;0.42)	0.36 (1.79)	0.21	(-0.22;0.95)	-0.43 (2.17)	0.21	(-1.14;0.26)
Somatic Complaints (11 items)	-0.13 (1.72)	0.51	(-0.52;0.26)	-0.21 (1.63)	0.43	(-0.74;0.32)	-0.05 (1.83)	0.86	(-0.64;0.54)
Withdrawn (8 items)	-0.03 (1.47)	0.81	(-0.29;0.37)	-0.13 (1.63)	0.62	(-0.66;0.40)	0.20 (1.30)	0.33	(-0.21;0.62)
Sleep Problems (7 items)	-0.31 (2.04)	0.18	(-0.77;0.15)	0.50 (2.02)	0.13	(-0.16;1.16)	-1.10 (1.75)	0.00	(-1.67;-0.53)
Attention Problems (5 items)	-0.27 (1.48)	0.11	(-0.60;0.06)	-0.34 (1.52)	0.17	(-0.84;0.16)	-0.20 (1.45)	0.38	(-0.67;0.26)
Aggressive Behavior (19 items)	-1.23 (4.82)	0.02	(-2.32;-0.13)	-1.02 (5.03)	0.21	(-2.68;0.63)	-1.43 (4.66)	0.06	(-2.94;-0.07)
Total internalizing Behaviors (36 items)	-0.51 (5.29)	0.39	(-1.72;0.68)	-0.07 (4.93)	0.92	(-1.70;1.54)	-0.94 (5.65)	0.30	(-2.78;0.88)
Total externalizing Behaviors (24 items)	-1.50 (5.62)	0.02	(-2.78;-0.23)	-1.36 (5.64)	0.14	(-3.22;0.48)	-1.64 (5.67)	0.07	(-3.48;0.19)
Total Problem score (100 items)	-3.41 (14.50)	0.04	(-6.70;-0.12)	-1.68 (13.82)	0.45	(-6.22;2.85)	-5.10 (15.12)	0.04	(-10.00;-0.19)
DSM Scale (No of items)									
Affective Problems (10 items)	-0.26 (1.94)	0.24	(-0.70;0.18)	0.07 (2.01)	0.81	(-0.58;0.74)	-0.59 (1.83)	0.05	(-1.18;0.00)
Anxiety Problems (10 items)	-0.09 (2.30)	0.73	(-0.61;0.43)	0.26 (2.02)	0.42	(-0.40;0.92)	-0.43 (2.52)	0.28	(-1.25;0.38)
Pervasive Developmental Problems (13 items)	-0.18 (1.94)	0.41	(-0.62;0.26)	-0.36 (2.09)	0.28	(-1.05;0.32)	0.00 (1.79)	1.00	(-0.58;0.58)
ADHD (6 items)	-0.37 (1.92)	0.09	(-0.81;0.06)	-0.28 (1.90)	0.35	(-0.91;0.33)	-0.46 (1.97)	0.15	(-1.10;0.17)
Oppositional Defiant Problems (6 items)	-0.70 (2.10)	0.00	(-1.17;-0.22)	-0.63 (2.21)	0.08	(-1.35;0.09)	-0.76 (2.01)	0.02	(-1.42;-0.11)

Significant associations are shown in **bold**.

Abbreviations: ADHD, Attention Deficit/Hyperactivity Problems, BMI, Body Mass Index; DSM, Diagnostic and Statistical Manual of Mental Disorders (4th edition, 1994)

Notes: Parent-group treatment = child obesity treatment focused on general parenting; Standard treatment = child obesity treatment focused on lifestyle.

Table 3. Estimates from linear regression models for associations between Child Behavior Checklist scores at baseline and change in child BMI Z-score from baseline to 12 months follow-up after obesity treatment

	Whole sample (n=77)				Parent-group treatment (n=38)				Standard treatment (n=39)			
	R ²	p	β	95% CI	R ²	p	β	95% CI	R ²	p	β	95% CI
Syndrome Scale												
<i>Model 1</i>	0.30				0.52				0.51			
Emotionally Reactive		0.15	-0.35	(-0.78;0.13)		0.46	-0.39	(-1.74;0.84)		0.47	-0.34	(-0.88;0.43)
Anxious/Depressed		0.28	0.29	(-0.23;0.77)		0.72	0.21	(-1.24;1.73)		0.10	0.67	(-0.09;0.99)
Somatic Complaints		0.23	0.22	(-0.14;0.55)		0.30	0.40	(-0.42;1.25)		0.46	0.20	(-0.27;0.56)
Withdrawn		0.78	-0.05	(-0.43;0.33)		0.58	-0.23	(-1.20;0.71)		0.94	0.02	(-0.48;0.52)
Sleep Problems		0.63	-0.08	(-0.42;0.26)		0.66	0.16	(-0.68;1.05)		0.64	-0.13	(-0.53;0.33)
Attention Problems		0.03	0.40	(0.03;0.72)		0.16	0.48	(-0.24;1.26)		0.29	0.31	(-0.21;0.67)
Aggressive Behavior		0.12	-0.36	(-0.77;0.09)		0.54	-0.24	(-1.22;0.67)		0.13	-0.79	(-1.23;1.72)
<i>Model 2</i>	0.14				0.40				0.32			
Total internalizing Behaviors		0.88	0.03	(-0.34;0.39)		0.90	0.04	(-0.67;0.75)		0.39	0.30	(-0.28;0.68)
Total Externalizing Behaviors		0.53	-0.12	(-0.48;0.25)		0.51	0.18	(-0.44;0.85)		0.04	-0.72	(-0.96;0.10)
<i>Model 3</i>	0.14				0.43				0.27			
Total Problem score		0.48	-0.09	(-0.32;0.15)		0.15	0.30	(-0.14;0.84)		0.03	-0.44	(-0.55;0.02)
DSM Scale												
<i>Model 4</i>	0.29				0.74				0.42			
Affective Problems		0.63	-0.09	(-0.45;0.28)		0.39	0.18	(-0.29;0.70)		0.30	-0.41	(-0.83;0.27)
Anxiety Problems		0.68	0.09	(-0.36;0.54)		0.70	0.10	(-0.50;0.73)		0.46	0.30	(-0.40;0.85)
Pervasive Developmental Problems		0.66	-0.09	(-0.49;0.31)		0.42	0.23	(-0.41;0.91)		0.86	-0.06	(-0.59;0.50)
Attention Deficit/Hyperactivity Problems		0.01	0.42	(0.09;0.72)		0.00	0.68	(0.29;1.21)		0.24	0.29	(-0.16;0.58)
Oppositional Defiant Problems		0.00	-0.45	(-0.72;-0.11)		0.00	-0.83	(-1.46;-0.40)		0.05	-0.53	(-0.72;0.01)
Covariates¹												
	0.13				0.37				0.12			
Child age		0.88	-0.01	(-0.28;0.24)		0.54	0.12	(-0.30;0.55)		0.59	-0.10	(-0.45;0.27)
Child sex		0.76	0.04	(-0.22;0.30)		0.45	0.15	(-0.30;0.65)		0.30	-0.20	(-0.49;0.16)
Mother BMI		0.24	0.15	(-0.10;0.40)		0.04	0.41	(0.01;0.85)		0.60	-0.10	(-0.42;0.25)
Mother foreign background		0.36	0.14	(-0.16;0.43)		0.17	0.27	(-0.15;0.77)		0.77	0.09	(-0.44;0.58)
Mother income		0.69	-0.05	(-0.34;0.23)		0.35	0.17	(-0.26;0.69)		0.66	-0.08	(-0.48;0.31)
Father BMI		0.03	0.27	(0.02;0.51)		0.05	0.35	(-0.01;0.77)		0.32	0.20	(-0.16;0.45)
Father foreign background		0.70	-0.05	(-0.35;0.24)		0.57	0.10	(-0.31;0.54)		0.66	-0.15	(-0.62;0.40)
Father income		0.83	-0.02	(-0.27;0.22)		0.86	-0.03	(-0.41;0.34)		0.98	0.00	(-0.33;0.34)

Significant associations are shown in **bold**. Abbreviations: p, p-value; β, standardized beta; CI, confidence intervals; BMI, Body Mass Index; DSM, Diagnostic and Statistical Manual of Mental Disorders (4th edition, 1994) ¹Covariates included as step one in all models. Notes: Parent-group = child obesity treatment focused on evidence-based parenting practices; Standard treatment group = child obesity treatment focused on lifestyle.

Table 4. Estimates from linear regression models for the associations between changes in Child Behavior Checklist scores and change in weight status (BMI z-scores) after an obesity treatment.

	Whole sample (n=77)				Parent-group treatment (n=38)				Standard treatment (n=39)			
	R ²	p	β	95% CI	R ²	p	β	95% CI	R ²	p	β	95% CI
Syndrome Scale												
<i>Model 1</i>	0.17				0.49				0.46			
Emotionally Reactive		0.86	0.03	(-0.38;0.45)		0.53	-0.23	(-1.71;0.93)		0.51	0.18	(-0.28;0.54)
Anxious/Depressed		0.45	-0.15	(-0.54;0.25)		0.57	0.26	(-0.87;1.51)		0.01	-0.90	(-1.08;0.11)
Somatic Complaints		0.53	-0.10	(-0.41;0.22)		0.87	-0.06	(-1.06;0.91)		0.93	0.02	(-0.41;0.44)
Withdrawn		0.43	0.12	(-0.19;0.43)		0.70	0.13	(-0.63;0.90)		0.09	-0.45	(-0.78;0.07)
Sleep Problems		0.71	-0.06	(-0.39;0.27)		0.73	0.10	(-0.59;0.81)		0.34	0.29	(-0.28;0.77)
Attention Problems		0.79	-0.04	(-0.35;0.27)		0.79	-0.10	(-1.00;0.78)		0.15	-0.38	(-0.67;0.11)
Aggressive Behavior		0.31	0.20	(-0.19;0.58)		0.62	0.18	(-0.70;1.12)		0.20	0.48	(-0.20;0.90)
<i>Model 2</i>	0.14				0.41				0.20			
Total internalizing Behaviors		0.64	-0.09	(-0.46;0.29)		0.39	0.26	(-0.51;1.23)		0.18	-0.45	(-0.73;0.15)
Total Externalizing Behaviors		0.49	0.13	(-0.24;0.50)		0.89	-0.03	(-0.75;0.66)		0.71	0.11	(-0.36;0.52)
<i>Model 3</i>												
Total Problem score		0.63	0.07	(-0.22;0.35)		0.36	0.20	(-0.31;0.80)		0.46	-0.18	(-0.46;0.22)
DSM Scale												
<i>Model 4</i>												
Affective Problems		0.75	0.06	(-0.30;0.41)		0.73	-0.11	(-0.82;0.59)		0.42	0.24	(-0.30;0.67)
Anxiety Problems		0.66	0.08	(-0.28;0.44)		0.07	0.47	(-0.08;1.47)		0.23	-0.36	(-0.63;0.16)
Pervasive Developmental Problems		0.94	-0.01	(-0.36;0.33)		0.86	-0.05	(-0.78;0.67)		0.86	-0.05	(-0.51;0.43)
Attention Deficit/Hyperactivity Problems		0.33	-0.16	(-0.47;0.16)		0.22	-0.30	(-0.95;0.24)		0.62	-0.13	(-0.48;0.29)
Oppositional Defiant Problems		0.34	0.16	(-0.18;0.50)		0.32	0.31	(-0.38;1.07)		0.85	-0.04	(-0.44;0.37)
Covariates¹	0.13				0.37				0.12			
Child age		0.88	-0.01	(-0.28;0.24)		0.54	0.12	(-0.30;0.55)		0.59	-0.10	(-0.45;0.27)
Child sex		0.76	0.04	(-0.22;0.30)		0.45	0.15	(-0.30;0.65)		0.30	-0.20	(-0.49;0.16)
Mother BMI		0.24	0.15	(-0.10;0.40)		0.04	0.41	(0.01;0.85)		0.60	-0.10	(-0.42;0.25)
Mother foreign background		0.36	0.14	(-0.16;0.43)		0.17	0.27	(-0.15;0.77)		0.77	0.09	(-0.44;0.58)
Mother income		0.69	-0.05	(-0.34;0.23)		0.35	0.17	(-0.26;0.69)		0.66	-0.08	(-0.48;0.31)
Father BMI		0.03	0.27	(0.02;0.51)		0.05	0.35	(-0.01;0.77)		0.32	0.20	(-0.16;0.45)
Father foreign background		0.70	-0.05	(-0.35;0.24)		0.57	0.10	(-0.31;0.54)		0.66	-0.15	(-0.62;0.40)
Father income		0.83	-0.02	(-0.27;0.22)		0.86	-0.03	(-0.41;0.34)		0.98	0.00	(-0.33;0.34)

Significant associations are shown in **bold**. Abbreviations: p, p-value; β, standardized beta; CI, confidence intervals; BMI, Body Mass Index; DSM, Diagnostic and Statistical Manual of Mental Disorders (4th edition, 1994) ¹Covariates included as step one in all models. Notes: Parent-group treatment = child obesity treatment focused on general parenting; Standard treatment = child obesity treatment focused on lifestyle.