



HAL
open science

Future challenges for using in silico molecular typology for risk assessment of pesticides metabolites - the example of Typol

Pierre Benoit, Laure Mamy, Yoce Aprianto, Dominique Patureau, Eric Latrille, Virginie Rossard, Rémi Servien, Fabienne Bessac, Sophie Hoyau, Christelle Margoum, et al.

► **To cite this version:**

Pierre Benoit, Laure Mamy, Yoce Aprianto, Dominique Patureau, Eric Latrille, et al.. Future challenges for using in silico molecular typology for risk assessment of pesticides metabolites - the example of Typol. 11th European Modelling Workshop EMW2023 "Exposure and effect modelling - Linking the domains", INRAE; Organizing Committee and Scientific committee: Carola Schriever (BASF), Bernhard Gottesbueren (make-sense consulting), Bas Buddendorf (Wageningen University Research), Thomas Preuss (Bayer AG), Marc Voltz (INRAE), Cécile Dages (INRAE), Sep 2023, Montpellier, France. hal-04334634

HAL Id: hal-04334634

<https://hal.inrae.fr/hal-04334634v1>

Submitted on 11 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Future challenges for using in silico molecular typology for risk assessment of pesticide metabolites - the example of Typol

Benoit P.^(1*), Mamy L.⁽¹⁾, Aprianto Y.⁽¹⁾, Patureau D.⁽²⁾, Latrille E.⁽²⁾, Rossard V.⁽²⁾, Servien R.⁽²⁾, Bessac F.⁽³⁾, Hoyau S.⁽³⁾, Margoum C.⁽⁴⁾, Rocco R.⁽⁴⁾, Martin-Laurent F.⁽⁵⁾


(1) UMR ECOSYS, INRAE, AgroParisTech, Université Paris-Saclay, 91120 Palaiseau, France

(2) UR LBE, INRAE, 11100 Narbonne, France


(3) LCPQ, UMR 5626, Université de Toulouse, CNRS, 31062 Toulouse, France

(4) UR RiverLy, INRAE, 69625 Villeurbanne, France

(5) UMR Agroécologie, INRAE, 21065 Dijon, France



➤ Environmental Risk Assessment of Pesticide Transformation Products (TPs)




Escher & Fenner, 2011

➤ Pesticide TPs : What do we know ?

The immersed face of the iceberg

Large number of potential TPs are still unknown but

- Increasing knowledge related to analytical methods improvement (non target methods)



Changes in molecular structures induce changes in :

- Environmental fate
- Toxicity and ecotoxicity

Analytical methods

GC-MS, LC-MS, GC-MS & LC-MS




Saaidi et al., 2022

➤ Pesticide TPs : What do we know ?


Knowledge Gaps

About identification, quantification



Countries

Anagnostopoulou et al., 2022



Increasing number of scientific publications

- Occurrence in environmental compartments
 - Water resources
 - Less information on soil, air, biota

About Fate and Effects (toxicity, ecotoxicity)

- Mostly via in silico approaches


➤ In Silico Approach applied on pesticide TPs

- **Computational tools and softwares to predict degradation pathways**

The pathway prediction system provides different relative reasoning models to predict likely biotransformation pathways and products

Examples :

- **EnviPath*** (<https://envipath.org>)
- **Eawag_BBD*** (<http://eawag-bbd.ethz.ch/predict/>)
- **PathPred*** (<https://www.genome.jp/tools/pathpred/>)
- **Chemical Transformation Simulator** (<https://qed.epa.gov/cts/>)



- **Computational tools and softwares to calculate fate and ecotoxicity endpoints**


From the molecular structure knowledge – QSARs

Examples

- **ECOSAR™** (Ecological Structure Activity Relationships) : based on a linear relationships between the predicted log Kow values (**EPI Suite** KOWWIN prediction) and the associated log of the evaluated toxicity values (mmol/L) for fish, daphnids, and green algae


➤ Predicting ecotoxicological endpoints

ECOSAR : chronic toxicity for aquatic organisms



Anagnostopoulou et al., 2022

➤ Clustering approaches





Hypothesis

The structure of molecules is **quantitatively correlated** with molecular descriptors and reflect their physicochemical properties, which enable to predict the behavior of pesticide TPs in the environment respecting their parent compounds (PCs)


➤ TyPol : Typology of Organic Contaminants

An operational tool for mapping and choosing "model" molecules to study their fate and impact in environment


- ❑ To select “model” molecules according to phenomenological (different behaviors) and operational (intensity of current or future risks) criteria
- ❑ Achieve a mutualizable tool (database) which makes possible to argue the choice of model contaminants to carry out studies concerning their fate and their ecotoxicological effects
- ❑ Test novel molecules (including transformation products) *in silico* , extrapolate (behaviours and impacts)




Biodegradation: DT50




TyPol - Basic principles



TyPol - A new methodology for organic compounds clustering based on their molecular characteristics and environmental behavior 


Rémi Servien^{a,b,*}, Laure Mamy^c, Ziang Li^d, Virginie Rossard^b, Eric Latrille^b, Fabienne Bessac^{e,f,g}, Dominique Patureau^b, Pierre Benoit^d

Servien et al., 2014



Issues	Process
Atmospheric release (Post application)	Volatilization
Atmospheric dissemination (polluted site)	Adsorption
Soil purification capacity	Absorption (plants)
Transfer to organisms & plants (Biavailability)	Biodegradation
Biological effects	Abiotic degradation
Persistence in wastewater treatment plant	Dissolution
Groundwater transfer	Ecotoxicity
Surface water transfer	

Parameters
Pvap, K _H
Koc
DT50 _a
DT50 _b
Kow
Sw, Kow
BCF, EC50, LC50, DL50 (various organisms)
ADI, AOEL



K_H : Henry's constant
DT50: Half-life time


BCF: Bioconcentration factor
EC50: Concentration producing 50 % effect

LC50: Lethal concentration
LD50: Lethal dose 50

ADI : Acceptable daily intake
AOEL : Acceptable operator exposure level

Environmental and ecotoxicological parameters retrieved from existing databases and literature


➤ TyPol - Basic principles



in silico and statistical approaches


TyPol







TyPol – A new methodology for organic compounds clustering based on their molecular characteristics and environmental behavior 

Rémi Servien^{a,b,*}, Laure Mamy^c, Ziang Li^d, Virginie Rossard^b, Eric Latrille^b, Fabienne Bessac^{e,f,g}, Dominique Patureau^b, Pierre Benoit^d

Servien et al., 2014



Descriptor type	Molecular descriptors
Constitution	Number and types atoms and bonds Molecular weight 
Geometry	Surface 
Topology	Connectivity Index 
Quantum	Polarizability Dipolar moment Orbital energies (homo, lumo) Ionization potential 

	August 2023
Molecules	568 parents/214 TPs
Descriptors	32939
Parameters	10569

➡ 40 most generic descriptors *Mamy et al., 2015*

➤ Questions addressed

- To what extent can the TyPol tool effectively classify pesticide parent compounds and their corresponding TPs and be used to highlight the relationships between chemical structures and the environmental fate or effects of putative pesticide TPs?
- To what extent can the clusters obtained with TyPol tool be used for a first approach in the risk assessment of pesticide TPs?

☰ Hypothesis





TyPol tool allows clustering pesticide parent compounds and their TPs **based on similarities** in molecular attributes, physicochemical parameters, environmental fate, and ecotoxicological effects

➤ Filiation parent compound → transformation products

➤ Screening chlordecone putative metabolites

Chlordecone dehalogenation products and potential congeners * (Dolfing et al., ES&T 2012)

- 6-hydrochlordecone
- 8-hydrochlordecone*
- 9-hydrochlordecone
- 10-hydrochlordecone
- 3,7-dihydrochlordecone*
- cis 8,10-dihydrochlordecone*
- 3,7,10-trihydrochlordecone*
- 8,10,10-trihydrochlordecone*
- 3,7,10,10-tetrahydrochlordecone*
- 2,3,7,8,10,10-hexahydrochlordecone
- 2,3,7,8,9,10,10-heptahydrochlordecone
- 1,2,3,7,8,9,10,10-octahydrochlordecone
- 1,2,3,4,6,7,8,9,10-nonahydrochlordecone
- 1,2,3,6,7,8,9,10,10-nonahydrochlordecone
- 1,3,4,6,7,8,9,10,10-nonahydrochlordecone*
- 2,3,4,6,7,8,9,10,10-nonahydrochlordecone
- décahydrochlordecone




Benoit et al. STOTEN 2017

- Visualization of class change for certain metabolites
- Potential consequences in terms of behavior and therefore risk

Projet Biodechlor « Search for the biological signature of chlordecone degradation in the soils of the French West Indies »
Coord : F. Martin Laurent (INRAE, Agroécologie, Dijon)

Combining TyPol to suspect screening approach




Short communication
 Identification and characterization of tebuconazole transformation products in soil by combining suspect screening and molecular typology



Veronika Storck ^{a,b,c}, Luigi Lucini ^{c,**}, Laure Mamy ^d, Federico Ferrari ^b, Evangelia S. Papadopoulou ^e, Sofia Nikolaki ^f, Panagiotis A. Karas ^e, Remi Servien ^g, Dimitrios G. Karpouzas ^e, Marco Trevisan ^c, Pierre Benoit ^d, Fabrice Martin-Laurent ^{a,*}

^a INRA, Mixed Research Unit 1347 Agroecology, Dijon, France
^b Aefjoria srl, Spinoff Catholic University of the Sacred Heart, Fidenza, Italy
^c Catholic University of the Sacred Heart, Department of Agronomy and Environmental and Chemistry, Piacenza, Italy
^d INRA, Mixed Research Unit 1402 ECOSYS, Thiverval-Grignon, France
^e University of Thessaly, Department of Biochemistry and Biotechnology, Larissa, Greece
^f University of Patras, Department of Environmental and Natural Resources Management, Agrinio, Greece
^g INRA, Mixed Research Unit 1331 Taxalim, Toulouse, France

- 76 tebuconazole metabolites
- Complementary diagnostic from *in silico* clustering by TyPol



Storck V., Lucini L., Mamy L., Ferrari F., Papadopoulou E.S., Nikolaki S., Karas P.A., Servien R., Karpouzas D.G., Trevisan M., Benoit P., Martin-Laurent F. 2016. *Env. Pollution*, 208, 537-545.

➤ Characterizing the Environmental Fate and Ecotoxicological Effects of Pesticide Transformation Products


Target organisms

Their physicochemical properties

Feasibility for using analytical instruments

Extensive use at EU level


Occurrence



Aprianto 2023

➤ Data preparation for Typol Database

Journal of Hazardous Materials 440 (2022) 129706




Enhanced database creation with *in silico* workflows for suspect screening of unknown tebuconazole transformation products in environmental samples by UHPLC-HRMS

Kevin Rocco*, Christelle Margoum, Loïc Richard, Marina Coquery

INRAE, UR RiverLy, 69625 Villeurbanne, France

Transformation Predictors



Input data:
chemical identifier of
TBZ, TBT, FNX (**SMILES**)

Output data:
Various format (depends
on predictors)

Output converter: OpenBabel (V2.4.1)
<https://sourceforge.net/projects/openbabel/>

Output harmonization:
InChiKey format, one-
way readable identifier

O U T P U T	Tebuconazole	Terbuthylazine	Fenoxycarb
	257 TPs	21 TPs	11 TPs
	In total 289 TPs		

List of TPs was retrieved from Rocco *et al.* (2022; 2023), INRAE-UR RiverLy, Villeurbanne

➤ Data preparation for Typol Database

Descriptors Calculation

40 molecular descriptors

Dragon 7.0

Constitutional

- ❖ No of atoms
- ❖ Molecular weight
- ❖ No of rings
- ❖ No of bonds
- ❖ Etc..

Topologica

- ❖ Connectivity index (CI)
- ❖ Valence CI

ChemOffice Ultra 12.0+Excel

Geometric

- ❖ Connolly molecular surface area

Computational Chemistry (DFT)

3D Quantum-chemical

- ❖ Dipole moment
- ❖ HOMO energy
- ❖ LUMO energy
- ❖ Polarizability
- ❖ Ionization energy

- Using Gaussian 16
- Executed in terminal macOS Ventura 13
- Access to CALMIP supercomputing

Environmental & Ecotoxicological Parameters

EPI Suite (v4.11)

Environmental parameter

Environmental Process

Dissolution

Volatilization

Adsorption

Degradation

Bioaccumulation

Ecotoxicity

ECOSAR (v.2.0.2)

Ecotoxicological parameter (acute & chronic endpoints)

Parameter

Sw (mg/L); log Kow

Pvap (mPa); K_H (Pa m³/mol)

Koc (L/kg)

DT₅₀ (days)

BCF

LC₅₀ / EC₅₀, NOEC (mg/L)

fish, daphnia, & green algae


Most values were retrieved from Rocco *et al.* (2022; 2023), except DT₅₀ & K_H

➤ Example of results


217 compounds (3 Parent Compounds Tebuconazole, Terbutylazine et Fenoxycarb and 214 TPs)

Clustering 1 : Based on molecular descriptors


Clustering 2 : Based on molecular descriptors & environmental parameters



≡ TPs that **are clustered away** from parent compounds **may have different behavior** due to marked changes in molecular structure




➤ Example of results



217 compounds (3 Parent Compounds Tebuconazole, Terbutylazine et Fenoxycarb and 214 TPs)

Clusters main characteristics





Molecular descriptors

Environmental parameters

Aprianto 2023

➤ Identifying molecular descriptors driving the fate of TPs



217 compounds (3 Parent Compounds Tebuconazole, Terbutylazine et Fenoxycarb and 214 TPs)



➤ Relating cluster changes to pathways, fate and impacts

217 compounds (3 Parent Compounds Tebuconazole, Terbutylazine et Fenoxycarb and 214 TPs)

Understanding why some TPs shifted from one cluster to another



Cluster 3 was characterized by higher log Kow (lower Sw), higher P_{vap}, higher K_H, high persistence

➔ Higher Risk to bioaccumulate in food webs compared to cluster 2

Pose risk to aquatic organisms

Some TPs have been detected in water samples in France (Rocco et al., 2022)


Some TPs have been identified in laboratory experiment (Richard, 2021)

Aprianto 2023

➤ Conclusions

- TyPol allows to classify the pesticides and their TPs in different clusters characterized by distinct properties: molecular descriptors, environmental parameters (log Kow, Sw, Koc, BCF, Pvpap, and KH) & some ecotoxicity endpoints (LC50 and NOEC)
- Some TPs can be assumed to have the same behavior as their parent compounds while others would be more persistent and/or toxic or, on the contrary, less persistent and/or toxic
- TyPol relies on the robustness of Input Data
 - > Increased knowledge from **suspect screening approach**
 - > Increased knowledge from predictive tools for exposure
 - > Increased knowledge from predictive tools for effects
 - > **Also integrating** more experimental values
- Potential use of TyPol for Pesticide TPs Environmental Risk Assessment
 - > **Screening and prioritization framework**
 - > **Selecting compounds for experimental testing**
 - > **Selecting compounds for exposure and effects modeling scenarios**

➤ Thanks for your attention



➤ References

- Servien R., Mamy L., Li Z., Rossard V., Latrille E., Bessac F., Patureau D., Benoit P. 2014. TyPol – A new methodology for organic compounds clustering based on their molecular characteristics and environmental behavior . *Chemosphere* , 111, 613-622.
- Mamy , L., Patureau , D., Barriuso , E., Bedos , C., Bessac , F., Louchart , X., Martinlaurent , F., Miege , C., Benoit, P. 2015. Prediction of the Fate of Organic Compounds in the Environment From Their Molecular Properties: A Review. *Critical Reviews in Environmental Science and Technology* . [doi: 10.1080/10643389.2014.955627](https://doi.org/10.1080/10643389.2014.955627)
- Storck V., Lucini L., Mamy L., Ferrari F., Papadopoulou ES, Nikolaki S., Karas PA, Servien R., Karpouzias DG, Trevisan M., Benoit P., Martin-Laurent F., 2016. Identification and characterization of tebuconazole transformation products in soil by combining suspect screening and molecular typology. *Environmental Pollution*, 208, 537-545. [doi:10.1016/j.envpol.2015.10.027](https://doi.org/10.1016/j.envpol.2015.10.027)
- Benoit P., Mamy L., Servien R., Li Z., Latrille E., Rossard V., Bessac F., Patureau D., Martin-Laurent F. 2017. Categorizing chlordecone potential degradation products to explore their environmental fate. *Science of the Total Environment* , 574, 781-795. <http://dx.doi.org/10.1016/j.scitotenv.2016.09.094>
- Traoré H, Crouzet O., Mamy L., Sireyjol C., Rossard V., Servien R., Latrille E., Martin-Laurent F., Patureau D., Benoit P. 2018. Clustering pesticides according to their molecular properties and their impacts by considering additional ecotoxicological parameters in the TyPol method. 2017. *Environmental Science and Pollution Research* **25**: 4728–4738 . <https://doi.org/10.1007/s11356-017-0758-8>
- Aemig , Q., Hélias , A., Patureau , D., 2021. Impact assessment of a large panel of organic and inorganic micropollutants released by wastewater treatment plants at the scale of France, *Water Research*, 188, 116524, <https://doi.org/10.1016/j.watres.2020.116524> .
- Bonnot K. 2021. In silico approach to analyze and predict the transformation of emerging organic contaminants in the environment. Thesis Montpellier SupAgro - Ademe -INRAE. Defended December 2021. 232pp.
- Mamy L., Bonnot K., Benoit P., Bockstaller C., Latrille E., Rossard V., Servien R., Patureau D., Prevost L., Pierlot F., Bedos C. 2021 Assessment of pesticide emission potential to atmosphere from their molecular properties using the TyPol tool . *Journal of Hazardous Materials* , 415, 125613 <https://doi.org/10.1016/j.jhazmat.2021.125613>
- Bonnot K., Benoit P., Mamy L., Patureau P. 2022. Transformation products of PPCPs in the environment: review of existing knowledge and classification of formation pathways according to parent molecule chemical structures. *Critical Reviews in Environmental Science and Technology*, in press Jan 2022.
- Bonnot K., Benoit P., Hoyau S., Mamy L., Patureau D., Servien R., Rapacioli M., Bessac F. 2022. Accuracy of computational chemistry methods to calculate organic contaminant molecular properties. *Submitted to Journal of Computational Chemistry Jan 2022*
- Servien R, Latrille E, Patureau D, Hélias A (2021) Machine learning models based on molecular descriptors to predict human and environmental toxicological factors in continental freshwater. *bioRxiv* , 2021.07.20.453034, ver. 6 peer-reviewed and recommended by Peer Community in Ecotoxicology and Environmental Chemistry. <https://doi.org/10.1101/2021.07.20.453034>
- Servien , R., Leenknecht , C., Bonnot , K., Rossard , V., Latrille , E., Mamy , L., Benoit, P., Hélias , A., Patureau , D., 2022. Improved impact assessment of micropollutants release from WWTPs, *Case Studies in Chemical and Environmental Engineering*, 100172. <https://doi.org/10.1016/j.cscee.2021.100172> .