



HAL
open science

Benchmarking the north atlantic salmon stock assessment: a new life cycle model to evaluate salmon mixed stock status and fisheries management scenarios across the north atlantic basin

Maxime Olmos, Remi Lemaire-Patin, Pierre-yves Hervann, James Ounsley, Maud Queroue, Geir Bolstad, Peder Fiske, Gérald Chaput, Etienne Prévost, Marie Nevoux, et al.

► **To cite this version:**

Maxime Olmos, Remi Lemaire-Patin, Pierre-yves Hervann, James Ounsley, Maud Queroue, et al.. Benchmarking the north atlantic salmon stock assessment: a new life cycle model to evaluate salmon mixed stock status and fisheries management scenarios across the north atlantic basin. 2023. hal-04337713v1

HAL Id: hal-04337713

<https://hal.inrae.fr/hal-04337713v1>

Preprint submitted on 12 Dec 2023 (v1), last revised 15 Jan 2024 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

BENCHMARKING THE NORTH ATLANTIC SALMON STOCK ASSESSMENT: A NEW LIFE CYCLE MODEL TO EVALUATE SALMON MIXED STOCK STATUS AND FISHERIES MANAGEMENT SCENARIOS ACROSS THE NORTH ATLANTIC BASIN

ICES WKBSalmon 2023

15 October 2023

Authors

Maxime Olmos^{1,2,*}, Remi Lemaire-Patin^{1,2}, Pierre Yves Hervann^{1,2}, James Ounsley,
Maud Queroue^{1,2}, Geir Bolstad, Peder Fiske, Gérald Chaput³, Etienne Prevost^{1,2}, Marie
Nevoux^{1,2}, Dauphin Guillaume³, Etienne Rivot^{1,2,#}

¹ DECOD Ecosystem Dynamics and Sustainability, Institut Agro, INRAE, Ifremer, France

² MIAME, Management of Diadromous Fish in their Environment, OFB, INRAE, L'Institut Agro, E2S UPPA, Rennes, France

³ Fisheries and Oceans, Moncton, Canada

*Email: maxime.olmos@ifremer.fr

#

Email: etienne.rivot@institut-agro.fr

Version: 10.13.2023

Recommended format for citation:

Olmos, M., Lemaire-Patin, R., Hervann, P-Y., Ounsley, J., Queroue, M., Bolstad, G., Fiske, P., Chaput, G., Prevost, E., Nevoux, M., Dauphin, G., Rivot, E. 2023. Benchmarking the north atlantic salmon stock assessment: a new life cycle model to evaluate salmon mixed stock status and fisheries management scenarios across the north atlantic basin. ICES WKBSalmon 2023 Working Paper, October 2023, 92 pp.

<https://doi.org/10.17895/ices.pub.24752079>

1 Acknowledgements

2

3 This work was made possible by the investment of the numerous people from various institutions
4 who collect and compile the data used by the ICES Working Group on North Atlantic Salmon.

5 The project has received funding from the Office Français de la Biodiversité under grant
6 agreement INRAE-OFB SalmoGlob-ToolBox, and from the European Regional Development
7 Fund through the Interreg Channel VA Programme, project SAMARCH Salmonid Management
8 Round the Channel.

9

10

11

12 Important note

13

14 This WP is based the version of the Life Cycle model developed in 2021. The structure of the
15 LCM is based on the one described in Olmos et al. (2019). The catch allocation rule is different
16 between the LCM and the PFA models. In particular, in the PFA models, the catch allocation rule
17 at West Greenland allocates catches among SU within the same complex in proportion to the pre
18 fishery abundance. In e LCM, catches at WG are allocated using proportions based on genetic
19 data.

20

21

22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54

TABLE OF CONTENT

1 INTRODUCTION 1

2 MATERIAL AND METHODS..... 5

 2.1 Outlines of the hierarchical Bayesian life cycle model..... 5

 2.1.1 Spatial structure integrating all SUs within the North Atlantic basin..... 5

 2.1.2 A harmonized life cycle model accounting for variability of life history traits..... 6

 2.1.3 Covariation among the 25 Sus 7

 2.2 Toward a new stock assessment model for the WGNAS 8

 2.2.1 Limits of PFA models and improvement through the life cycle model..... 8

 2.2.2 Likelihood function and data..... 8

 2.2.3 Forecasting and risk analysis framework 9

 2.2.3.1 Management objectives- Conservations Limits (CLs)..... 10

 2.2.3.2 Risk analysis framework for the Western Greenland and the Faroes fishery..... 11

 2.2.3.3 Expected differences in forecast and risk analysis between life cycle and PFA models 13

 2.2.4 MCMC simulations and model checking..... 14

3 RESULTS 17

 3.1 A harmonized life cycle to estimate temporal variations and spatial coherence of demographic parameters across the North Atlantic basin 17

 3.1.1 A widespread decline of abundances in all CSG 17

 3.1.2 Post-smolt survival rate 17

 3.1.3 Proportion of fish maturing as 1SW 18

 3.2 Comparisons between the new life cycle and currently used PFA models to assess and forecast salmon population dynamic and to evaluate catch options for mixed stock marine fisheries 19

 3.2.1 Hindcast and Forecast analysis..... 19

 3.2.1.1 Europe : NE and SE 20

 Hindcast 20

 3.2.1.2 NA 21

 3.2.2 Risk analysis: Evaluating catch options for mixed stock marine fisheries 22

 3.2.2.1 Eggs deposition compared to CLs..... 22

55	3.2.2.2	The West Greenland mixed stock fishery	23
56	3.2.2.3	The Faroes mixed stock fisheries	24
57	3.2.2.4	Evaluating catch options for both Faroes and West Greenland Fisheries	25
58	4	CONCLUSION.....	26
59	5	REFERENCES	27
60	6	SUPP.MAT.1. - Detailed process and observation equations of the Bayesian life cycle	
61		model.....	43
62	1.1	Population dynamics	43
63	1.1.1	Simplified life history.....	43
64	1.1.2	Eggs deposition.....	43
65	1.1.3	Egg-to-smolt transition.....	43
66	1.1.3.1	Egg-to-smolt survival.....	43
67	1.1.3.2	Distribution according to smolt ages.....	44
68	1.1.4	Marine phase.....	44
69	1.1.4.1	Post-smolt survival and proportion of fish maturing as 1SW.....	44
70	1.1.4.2	Sequential marine fisheries and natural mortality.....	45
71	1.1.4.3	From returns to spawners (homewater catches).....	46
72	1.2	Observation equations	47
73	1.2.1	Returns.....	47
74	1.2.1.1	The case of Northern NEAC SU.....	47
75	1.2.2	Homewater catches.....	48
76	1.2.3	Catches at sea for sequential distant marine fisheries operating on mixed stocks	48
77	7	SUPP.MAT.II : Comparison between the PFA and life cycle modelling framework	67
78	8	Supp.Mat.III: RESULTS	73
79			

80 1 INTRODUCTION

81 Atlantic salmon (*Salmo salar*) (hereafter A. salmon) figures prominently in emblematic
82 migratory species of the Atlantic Ocean and is characterized by a highly evolved homing ability
83 that is the root of its structure into individual river populations. After spending their first years in
84 rivers of the eastern North America and the western Europe, juvenile A. salmon migrate to
85 common feeding grounds in the North Atlantic, especially aggregating off West Greenland, in the
86 Labrador Sea, and the Faroes Islands and Norwegian Sea (Aas et al., 2010; D. Mills, 1989), where
87 they grow over multiple winters before heading back to their home waters once mature. A. salmon
88 are susceptible to be harvested at several stages in their life cycle. While some fisheries affect
89 individual salmon populations in coastal areas before entering their natal river or in freshwater
90 (final stages of spawning migration), mixed stock fisheries harvest populations originating from
91 various continental habitats in high seas foraging areas, in West Greenland or in the vicinity of the
92 Faroe Islands (Chaput, 2012; ICES, 2017). These areas are also where salmon populations are
93 exposed to common marine environmental conditions (Beaugrand & Reid, 2003, 2012; Friedland
94 et al., 2014; K. E. Mills et al., 2013).

95 The regulation of mixed stock high seas fishery was of sufficient concern that an international
96 body (the North Atlantic Salmon Conservation Organization (NASCO; <http://www.nasco.int/>))
97 was formed in 1982 and a treaty subsequently signed by participating countries to manage the
98 marine fisheries impacting different stock of A. salmon (Windsor and Hutchinson, 1994). The
99 annual stock status reports developed by the Working Group North Atlantic Salmon of the
100 International Council for the Exploration of the sea (ICES/CIEM WGNAS) and the subsequent
101 scientific advices provided to the NASCO have formed the basis for the negotiations and

102 subsequent management of these fisheries. West Greenland and Faroe high sea fisheries are
103 managed following a fixed escapement strategy aiming to achieve the spawner (or egg)
104 requirements for the contributing stocks on both sides of the Atlantic Ocean (Chaput, 2012; W.
105 Crozier et al., 2003; W. W. Crozier et al., 2004; Potter et al., 2004). Until 2020, ICES has advised
106 this management by recommending maximum catch levels based on a forecast of A. salmon
107 abundance prior to the high seas fisheries exploitation (Pre Fishery Abundance, PFA). The so-
108 called PFA models used for stock assessment of Atlantic salmon have been developed based on
109 data aggregated at the scale of regional or national stock units (SU) over the North Atlantic area
110 within three continental stock groups (CSG), i.e. eastern North America (NA), Southern European
111 (SE) and Northern European (NE) (Chaput, 2012; W. W. Crozier et al., 2004; Potter et al., 2004).
112 Each of these three independent models reconstructs long-term series of abundance at sea before
113 marine fisheries for a given CSG and forecasts the returns of adults in their natal rivers of the
114 matching SUs (homewaters). PFA models have been incorporated in a risk analysis framework to
115 assess the consequences of mixed stock marine fisheries at West Greenland and Faroes on the
116 returns (Friedland et al., 2005; ICES, 2015) and assess compliance of realized spawning
117 escapement to conservation limits (biological references point below which the stock should not
118 pass) at both the SU and CSG scales.

119 However, PFA models suffer from three weaknesses (here after [W]) that hinder their
120 relevance for analysing the demographic processes driving the population's dynamics of European
121 and American A. salmon populations. [W1] PFA models rely on a constructed stock-recruitment
122 dynamic. Since they do not explicitly represent the population dynamics as a life cycle, there is no
123 dynamic link between PFA and subsequent egg deposition. Thus, the productivity required by the
124 models to forecast the returns, which is roughly inferred from reconstructions, may be susceptible

125 to time series bias. In addition, the lack of flexibility in the modeling framework restricts the
126 hypotheses that can be explored on drivers and mechanisms of changes in *A. salmon* demographics
127 and population dynamics. [W2] The PFA modeling framework actually works as a combination
128 of two models. A first model, the run reconstruction model relies on estimates of the abundance
129 of fish returning to spawn and biological parameters (sex ratio, fecundity and mean proportion of
130 smolts ages) to estimate the potential number of spawners or eggs (measure of the Stock) for each
131 year of the time series. The same model is used to estimate the abundance of fish at the PFA stages
132 (measure of the Recruitment), through a back-calculation procedure (similar to a Virtual
133 Population Analysis) using data on catches at sea and hypothesis on natural mortality rates at sea.
134 Hence, the measure of the stock and the recruitment are derived from the same data, whilst they
135 are considered independent in the rest of the process. A second part of the modelling framework
136 consists of estimating the productivity parameters between the Stock and the Recruitment for all
137 years of the historical time series, and uses time series hypothesis (random walk) to forecast the
138 evolution of the productivity parameter during three years after the last year of the assessment.
139 This forecast of the productivity parameters serves as a basis to forecast the PFA and the number
140 of fish that returns to homewater based on catches scenarios at sea. [W3] The three PFA models
141 are independent and some of their core demographic hypotheses are not harmonized. The fact that
142 SE and NE models consider the dynamics of both one-sea-winter and two-sea-winter fish while
143 the NA model only considers that of the latter precludes the simultaneous analysis of the
144 population dynamics among all SU in the North Atlantic (Chaput et al., 2005). Therefore, these
145 structural differences rule out any covariance structure in the dynamics of the SUs even though
146 the SUs may share common environments at sea and be jointly exploited in sea fisheries.

147 This paper aims to bring a major contribution to improve the scientific basis for A. salmon
148 stock assessment. We then address weaknesses [W1]-[W3] from PFA models by developing a
149 Bayesian life cycle modelling framework for the combined analysis of A. salmon population
150 dynamics across a large number of SU in the North Atlantic Ocean. To this goal we extend the
151 framework developed by Olmos *et al.*, (2019) for North American and Southern European CSG
152 to include the dynamics of all SU of the three CSG representing the all Atlantic salmon distribution
153 (Northern Europe, Southern Europe and North America) within a single harmonized and unified
154 hierarchical Bayesian life cycle approach with populations following a similar life history process.
155 The final goal of this paper is to provide a benchmark for the assessment and forecast models
156 currently used by ICES for Atlantic salmon stock assessment in the North Atlantic.

157 To this end, using the same dataset, we compare previous PFA assessment models from ICES
158 (ICES, 2018) with our Bayesian life cycle model to illustrate how our new approach improve the
159 scientific basis for Atlantic salmon stock assessment. We illustrated how the life cycle model
160 outperformed PFA models by investigating how harmonizing life histories and considering
161 covariation in demographic parameters may (i) provide new insight about the life history
162 parameters driving the dynamic of all North Atlantic salmon population; (ii) impact hindcast
163 analysis, (iii) forecast analysis and (iv) risk analysis.

164

165 2 MATERIAL AND METHODS

166 In this section we focus on two main aspects of our modelling exercise. First, we present the
167 outlines of the Bayesian life cycle modelling framework for the combined analysis of A. salmon
168 population dynamics across a large number of SU in the North Atlantic. Then, we present the
169 general framework used to compare PFA and LCM models.

170 2.1 Outlines of the hierarchical Bayesian life cycle model

171 The life cycle model is formulated in a Bayesian hierarchical state-space framework (Buckland et
172 al., 2004; Cressie et al. 2009; Parent & Rivot, 2012; Rivot et al., 2004) that incorporates
173 stochasticity in population dynamics as well as observation errors. We extended the framework
174 developed by Olmos et al. (2019) (all SU from North America and Southern Europe) to include
175 the dynamics of all SU of the three CSG (North America, Southern Europe and Northern Europe).
176 To keep the presentation concise, below we resume the important characteristics of the model and
177 all model equations and data sources are detailed in Supp.Mat.I.

178 2.1.1 Spatial structure integrating all SUs within the North Atlantic basin

179 The model considers the dynamics of 25 SU (subscript $r = 1, \dots, N$ with $N=25$) (Fig. 1):

- 180 • 6 SU from NA CSG, indexed by $r = 1, \dots, 6$: 1 = Newfoundland, 2 = Gulf, 3 = Scotia-Fundy,
181 4 = USA, 5 = Quebec and 6 = Labrador;
- 182 • 8 SU from the SE CSG, indexed by $r = 7, \dots, 14$: 7 = France, 8 = UK England and Wales, 9
183 = Ireland, 10 = UK Northern Ireland - FO, 11 = UK Northern Ireland - FB, 12 = UK Scotland
184 East, 13 = UK Scotland West, 14 = Iceland South-West;

- 185 • 11 SU from NE CSG, indexed by $r=15, \dots, 25$: 15 = Iceland North-East, 16 = Sweden, 17 =
186 Norway South-East, 18 = Norway South-West, 19 = Norway Middle, 20 = Norway North, 21
187 = Finland, 22 = Russia Kola Barents, 23 = Russia Kola White Sea, 24 = Russia Arkhangelsk
188 Karelia and 25 = Russia River Pechora.

189 SU are defined on the basis of freshwater areas. All salmon within a SU are assumed to have the
190 same demographic parameters and to undertake a similar migration route at sea.

191 2.1.2 A harmonized life cycle model accounting for variability of life history traits

192 The model is built in discrete time on a yearly basis (subscript $t = 1, \dots, n$ with $n=47$ in this present
193 application based on data from 1971 to 2017). The population dynamic of each of the 25 SU is
194 represented by an age- and stage-structured life cycle model (Fig. 2). The model incorporates
195 variations in the age of out-migrating juveniles from freshwater (i.e., smolt ages) and the sea-age
196 of returning adults among SUs. Smolts migrate to sea after 1 to 6 years in freshwater (depending
197 on SU). Following the approach used by ICES for catch advice purposes (ICES 2022), only two
198 sea-age classes are considered in the model: maiden salmon that return to homewaters to spawn
199 after one year at sea, referred to as one-sea-winter (1SW) salmon, or grilse, and maiden salmon
200 that return after two winters at sea (2SW). This is a simplification of the larger diversity of life
201 history traits as some maiden fish may spend more than two winters at sea before returning to
202 spawn, and some salmon return as repeat spawners. However, maiden spawners older than 2SW
203 are relatively rare in North America and Southern Europe and the six smolt-age by two sea-age
204 combinations represent the essence of life history variation.

205 The model tracks the abundance of fish ($N_{s,t,r}$) for each SU (r) by year (t) and life stage (s),
 206 sequentially from eggs (N_1) to 1SW (N_7) or 2SW (N_{10}) spawners for the period considered
 207 (starting in 1971, year of return to rivers) (Fig. 2; Supp.Mat.I). Spawners are fish that contribute
 208 to reproduction and that survived all sources of natural and fishing mortality. All transition rates
 209 between stages s for each SU (r) in year t are denoted $\theta_{s,t,r}$.

210 2.1.3 Covariation among the 25 Sus

211 The model explicitly incorporates two components of temporal covariation among all SUs (Fig.
 212 2). First, the post-smolt survival (called productivity in PFA models, - *so in this manuscript post-*
 213 *smolt survival and productivity defined the same variable-* denoted $\theta_{3,t,r}$) and the proportion of fish
 214 maturing as 1SW (denoted $\theta_{4,t,r}$) are modelled as multivariate random walks in the logit scale
 215 which captures spatial covariation associated with environmental stochasticity. Random variations
 216 are drawn from multivariate Normal distributions in the logit scale with variance-covariance
 217 matrices Σ_{8_3} and Σ_{8_4} (Ripa and Lundberg, 2000; Minto *et al.*, 2014) :

218 (1)
$$(\text{logit}(\theta_{3,t+1,r}))_{r=1:N} \sim \text{MVNormal} ((\text{logit}(\theta_{3,t,r}))_{r=1:N}, \Sigma_{8_3})$$

219 (2)
$$(\text{logit}(\theta_{4,t+1,r}))_{r=1:N} \sim \text{MVNormal} ((\text{logit}(\theta_{4,t,r}))_{r=1:N}, \Sigma_{8_4})$$

220 with N = the number of SU in the model (here $N=25$).

221 The pairwise correlation matrix ρ can be calculated from the variance-covariance matrix:

222 (3)
$$\rho = \text{jdiag}(\Sigma)^{-1} \times \Sigma \times \text{jdiag}(\Sigma)^{-1}$$

223 The second source of covariation among SU is the harvest dynamics of the sequential marine
224 fisheries that operate on mixtures of SUs, with the portfolio of SU available for each fishery
225 dependent on marine migration route hypotheses (Fig. 2 and Fig. Supp.Ma.I).

226 **2.2 Toward a new stock assessment model for the WGNAS**

227 In this section we present how the hierarchical life cycle model may improve the assessment of
228 north Atlantic salmon by (i) highlighting the main differences between the currently used stock
229 assessment (PFA models) and the hierarchical life cycle model (Fig.3, Table.1, Supp.Mat.II), (ii)
230 presenting the data we used and (iii) defining the risk analysis framework we used to compare both
231 approaches.

232 2.2.1 Limits of PFA models and improvement through the life cycle model

233 We highlight the limits of the PFA models and how the life cycle model may address and solve
234 those limits in Table 1. Details about the currently used PFA models can be found in Supp. Mat.
235 II and in previous studies (Rago et al., 1993; Crozier et al., 2004; Chaput et al., 2005; ICES, 2018)

236 2.2.2 Likelihood function and data

237 Building an integrated model (Maunder & Punt, 2013; Rivot et al., 2004; Schaub & Abadi, 2011)
238 that explicitly integrates complicated observation models would dramatically increase the
239 complexity of the full model. Therefore for, a sequential approach (Michielsens et al., 2008;
240 Staton, Catalano, & Fleischman, 2017) is used that consists of (i) processing observation models
241 separately to reconstruct probability distributions that synthesize observation uncertainty around
242 the time series of catches and returns for the 13 SUs and (ii) using those distributions as likelihood
243 approximations in the population dynamics state-space model. Probability distributions for returns

244 and catches are derived from a variety of raw data and observation models, specific to each SU
245 (except for the mixed-stock fisheries at sea) as originally developed by ICES to provide input for
246 PFA models (see ICES, 2015b and Supporting Information S1 for further details).

247 For both PFA and life cycle models these consist of the following in time series (1971–2017) of
248 estimates (approximated as log-normal distributions) of the number of maturing anadromous
249 Atlantic salmon that return to homewaters for each of the 13 SUs by 1SW and 2SW maiden sea-
250 age classes (Supp.Mat.I, Figure.1).

251 Because the life cycle is expandable and provides an opportunity to assimilate new sources of
252 information to improve the ecological and biological realism of the model, other times series are
253 implemented in the likelihood component of the model such as (ii) time series of estimates
254 (approximated as log-normal distributions) of homewater catches for each SU by sea-age class;
255 and (iii) time series of estimates (approximated as log-normal distributions) of catches for the
256 mixed-stock fisheries at sea operating sequentially on combinations of SUs, and using additional
257 data on the SU origin of the catches. More details about how the data are implemented in the PFA
258 and life cycle model can be found in (Supp.Mat.II. Table SII.1).

259 2.2.3 Forecasting and risk analysis framework

260 For both PFA and life cycle models we used probabilistic forecasts from the models to evaluate
261 the probability that egg deposition fall below management objectives for different catch options in
262 the Western Greenland and the Faroes fisheries.

263 The forecasted abundance of eggs deposition after all marine distant fisheries (but before
264 homewater fisheries) is then compared to the management objectives defined below. The same
265 models are used for fitting the historical time series and forecasting. In the case of the life cycle

266 model, an exception are the transitions that involve a fishing mortality, modelled by directly
267 retrieving catches to the abundance (this is because scenarios are defined by fixing catches and not
268 harvest rates).

269 For the life cycle model, the post-smolt marine survival and the proportion maturing are forecasted
270 following the multivariate random walks defined at equations (1)-(2). For the NA PFA, only the
271 post-smolt marine survival is forecasted following the multivariate random walks (the model
272 tracking only 2SW). For the NE PFA model the post-smolt marine survival and the proportion
273 maturing are forecasted following random walks with no covariance structure among SUs.

274 Because of the random walk hypothesis, the forecasted marine survival and proportion maturing
275 during the forecasting period will remain at the same average level than the last year of the fitted
276 time series, but with an uncertainty that increases with time due to error propagation through the
277 random walk. Parameters uncertainty is integrated using Monte Carlo simulation, by simulating
278 multiple population trajectories with parameters randomly drawn in the posterior MCMC sample.

279 *2.2.3.1 Management objectives- Conservations Limits (CLs)*

280 In managing Atlantic salmon fisheries, NASCO (1998) has adopted a fixed escapement strategy
281 (Potter, 2001), in recognition of the importance of the spawning stock to subsequent recruitment.

282 Management objectives are based on Conservation Limits (CLs) as defined by ICES and NASCO.
283 CLs are defined as the quantity of eggs that should be deposited by spawners to produce a desired
284 production of smolts (Table 2). Following the principles adopted by NASCO (1998) CLs for North
285 Atlantic salmon have been defined by ICES WGNAS as limit reference points, in the sense that
286 having abundance of eggs spawned falls below these limits should be avoided with high
287 probability.

288 Management objectives in SE and NE are to reach or exceed CLs for both 1SW and 2SW fish.
289 However, in NA management objectives currently defined by ICES consider the 2SW fish
290 component of the returns only. However, in this paper, CLs for North America have been defined
291 as the total required egg deposition for both sea age classes (1SW+2SW) for all SU including
292 North America. CLs used by ICES are only available at a more aggregated spatial scale than SU
293 defined in our life cycle model (Table 2). Specifically, one CL is available for Scotland (sum of
294 Eastern Scotland and Western Scotland in life cycle model), one CL for Norway (sum of 4 SU in
295 life cycle model, South-East Norway, South-West Norway, Middle Norway and North Norway)
296 and one CL for Russia (sum of 4 SU in life cycle model, Russia Kola Barents Russia Kola White
297 Sea, Russia Arkhangelsk Karelia and Russia River Pechora). To be compared to the CLs defined
298 by ICES, eggs deposition in the life cycle model were then summed to match with the spatial scale
299 considered for CLs. Finally, abundances defined at different life stages, demographic parameters
300 and fisheries exploitation rates posterior estimated from the life cycle model were also aggregated
301 over the PFA spatial scale (representing 17 SU or aggregation of SU) to be compared to PFA
302 posterior estimates.

303 *2.2.3.2 Risk analysis framework for the Western Greenland and the Faroes fishery*

304 SUs from NA, SE and NE are all potentially harvested by the West Greenland fishery (although
305 the proportion of fish originating from Northern Europe is very low in West Greenland catches).
306 A risk framework for the provision of catch advice for the West Greenland fishery has been applied
307 since 2003 by NASCO and ICES (ICES, 2013).
308 Only fish from SE and NE are potentially harvested at the Faroes fisheries. There is currently no
309 agreed framework for the provision of catch advice for the Faroes fishery adopted by NASCO.
310 However, NASCO has asked ICES, for a number of years, to provide catch options or alternative

311 management advice with an assessment of risks relative to the objective of exceeding stock
312 conservation limits for salmon in the European area (NE and SE complexes).

313 As an important contribution, our new life cycle model provides a unique framework for evaluating
314 catch options for the Faroes and West Greenland separately or simultaneously and for all SU
315 separately or simultaneously.

316 For the purpose of demonstration, in this paper 36 scenarios that consider both Faroes and West-
317 Greenland catches were then built by crossing 6 combinations of catches from 0 to 2500 tons (0,
318 500, 1000, 1500, 2000, 2500) for both the Western Greenland and Faroes fisheries. For each
319 scenario, catches options were converted to number of fish really caught using mean weight of fish
320 following ICES (2015a). Population dynamics was simulated with homewater catches and
321 proportions to allocate catches at West Greenland and Faroes fisheries to the different SU fixed to
322 the average of the last five years of the time series of data (2013-2017), and 0 catches for other
323 distant fisheries.

324 For the West Greenland fishery, the catch of 1SW salmon of North American and European origins
325 is further discounted by the fixed sharing fraction (F_{na}) historically used in the negotiations of the
326 West Greenland fishery, that is a 40%:60% West Greenland:(North America & Europe) split. For
327 instance, in a scenario with a 100t quotas, a total of 250t are actually caught, 150t are reserved for
328 the Western Greenland fishery and 100t are reserved for the North American and European
329 commercial fishery (note that the scientific advice given by ICES since several years is a quotas
330 of 0t).

331 For each scenario, we provide forecasts during three years (in this application, 2018- 2020) starting
332 after the last year of our assessment model (2017). Monte Carlo simulations are run to integrate
333 over both process' errors and parameters' uncertainty. Parameters uncertainty is integrated by

334 randomly sampling the parameters in the joint Bayesian posterior distribution probability around
335 parameters, which captures the covariance structure among the parameters. For a given set of
336 parameters, the population dynamics is simulated including process error (i.e., inter-annual
337 variability).

338 The probability of each SU (or aggregation of SU as defined in Table 2) achieving its CL
339 individually and the probability of this being achieved by all management units simultaneously
340 within a same CSG (i.e. in the same given year) are calculated from Monte Carlo trials. This allows
341 managers to evaluate both individual and simultaneous achievement of management objectives in
342 making their management decisions.

343 *2.2.3.3 Expected differences in forecast and risk analysis between life cycle and PFA models*

344 We expect that most of the differences we might observe between PFA and life cycle model will
345 rely on two main reasons:

- 346 1. The demographic structure between PFA and life cycle models can be different. PFA
347 models rely on different life history hypothesis depending on CSG: NE and SE PFA models
348 consider both 1SW and 2SW life histories while NA PFA model considers only 2SW life
349 history
- 350 2. The way the PFA and life cycle models handle differently the uncertainty can lead to major
351 differences, and have an impact on hindcast, forecast and risk analysis.
 - 352 ○ In the life cycle model all latent variables are correlated through the life cycle
353 structure whereas in the PFA model, all latent variables are not linked (similar to a
354 stock recruitment dynamic). This will change how the uncertainty is propagated
355 between the different parameters and latent variables.

- 356 ○ In the PFA models, the stock abundances (lagged eggs for NEAC and lagged
357 spawners for NAC) is defined in the model through a prior distribution which is not
358 updated within the model and so the uncertainty is not propagated through the other
359 latent variables and parameters. Here again, this will create a big difference in term
360 of how the uncertainty is quantified and propagated between the life cycle and PFA
361 models
- 362 ○ Finally, in PFA models, the only likelihood function is defined for the distribution
363 of returns whereas in the life cycle models the likelihood function include the
364 distribution of returns, the distributions of both freshwater and fisheries catches and
365 the proportion to allocate the catches to each stock units.
- 366 ○ To sum up, we argue that those differences in how PFA and life cycle models
367 account for uncertainty will generate strong differences in the approximation of
368 Bayesian posterior distributions.

369 2.2.4 MCMC simulations and model checking

370 The life cycle model and the PFA models have different statistical structures. So the number of
371 iterations to make the model converged and to generate Bayesian posterior distributions are
372 different.

373 *Life cycle model*

374 Bayesian posterior distributions were approximated using Monte Carlo Markov Chain (MCMC)
375 methods using Nimble (<https://r-nimble.org>) (de Valpine et al., 2017). The Nimble code for our
376 model is available on GitHub: [https://github.com/MaxOlmos/SALMOGLOB-Life-Cycle-](https://github.com/MaxOlmos/SALMOGLOB-Life-Cycle-Model/blob/master/model_nimble_compPFA.r)
377 [Model/blob/master/model_nimble_compPFA.r](https://github.com/MaxOlmos/SALMOGLOB-Life-Cycle-Model/blob/master/model_nimble_compPFA.r). Two independent MCMC chains with dispersed

378 initialization values were used. The level of autocorrelation of MCMC chains is very high (still
379 significant at lag 30). The first 200 000 iterations were used as a burn-in period. For each chain, to
380 reduce the autocorrelation in the MCMC sample used for final inferences, one out of 400 iterations
381 post burn-in was kept and the resulting sample of 4000 iterations per chain was used to characterize
382 the posterior distribution (total iteration is 3 200 000). Convergence was assessed using the
383 Gelman-Rubin statistic (Brooks & Gelman, 1998) as implemented in the R Coda package
384 (`gelman.diag()`). Following the methodology developed in Olmos et al. (2019), the model fit to
385 each data source was assessed by checking that the 90% credibility envelope of the posterior
386 predictive distribution of each variable contained the observation. In addition, Bayesian p-values
387 calculated from chisquare discrepancy tests (Gelman et al., 2014a) were calculated to check the
388 ability of the model to replicate a posteriori data similar to those observed. As in Olmos et al.
389 (2019, 2020), posterior predictive distributions show that the model fits well to all observations,
390 and posterior predictive checks do not indicate strong inconsistencies between the model a
391 posteriori and the data. Those results are not developed further in this paper (see Olmos et al.
392 (2019) for more details)

393 *NAC PFA model*

394 Two independent MCMC chains with dispersed initialization values were used. The first 10 000
395 iterations were used as a burn-in period. For each chain to reduce the autocorrelation in the MCMC
396 sample used for final inferences, one out of 100 iterations post burn-in was kept and the resulting
397 sample of 50000 iterations per chain was used to characterize the posterior distribution (total
398 iteration is 80 000)

399 *NEAC PFA model*

400 Two independent MCMC chains with dispersed initialization values were used. The first 3000
401 iterations were used as a burn-in period. For each chain to reduce the autocorrelation in the MCMC
402 sample used for final inferences, one out of 100 iterations post burn-in was kept and the resulting
403 sample of 7 000 iterations per chain was used to characterize the posterior distribution (total
404 iteration is 20 000).

405

406 **3 RESULTS**

407 In this section we first present the new results provided by the life cycle model for Atlantic salmon
408 across the North Atlantic basin. Then we present the results of the comparison between the PFA
409 and LCM models.

410 **3.1 A harmonized life cycle to estimate temporal variations and spatial coherence of** 411 **demographic parameters across the North Atlantic basin**

412 3.1.1 A widespread decline of abundances in all CSG

413 The model estimates time series of all key life stages in the model for all SU or for any aggregation
414 of SU at the scale of SU or countries (Supp.Mat.III. Fig. SIII.1). Time series of total PFA in each
415 CSG show very similar continuous declines by a factor 3, between the 1970s and the 2010s (Fig.4)
416 with a stronger decline for the NA CSG. The decline in PFA is marked by a strong decrease in
417 abundances in the 1990s.

418 3.1.2 Post-smolt survival rate

419 The time-series of post-smolt survival for the 25 SU show a common decreasing trend over years
420 (Fig.5a -d). The trends averaged over all SU of the same CSG exhibit slightly different tendencies
421 over the years. The post-smolt survival in NA exhibit a strong decline by a factor 3 in the period
422 1985-1995. This decline is also observable in SE with a sharp decline by a factor 1.8 in 1987. The
423 sharp decline in the late 80's-early 90's is less visible in NE. Trend in NE shows a continuous and
424 smoothed decline over the period.

425 The majority of pairwise correlations are positive, with a median correlation among all SU of 0.084
426 ± 0.139 (correlations are calculated in the logit scale; Fig.5e and Fig.5f). In general, correlations
427 are stronger between geographically close SU. The results show strong correlations for SU within
428 NA (0.333), followed by SE (0.138) and NE (0.083). Correlations between the NE SU are stronger
429 for the block of SU going from Sweden (East) to Russia-KB (West).

430 3.1.3 Proportion of fish maturing as 1SW

431 Time trends in the proportion of fish that mature as 1SW also show a strong coherence among SU.
432 These are in accordance with the expectation of higher correlations between SU of the same CSG.
433 Overall, there is an increasing trend from the 1970s to the 1990s that corresponds to declines in
434 the proportions of 2SW fish in the returns followed by a levelling off or even a decline from the
435 2000s (Fig.6a -d). All time trends are consistent with the average trend, except for France which
436 shows a consistent decline during the entire period. Consistently with the low proportion of 1SW
437 observed in the returns, the two most eastern SU, Russia-AK and Russia-RP, and US differ from
438 the others SU with a very low probability of maturing. As observed for the post-smolt survival,
439 most of the pairwise correlations are positive across the 24 SU, with an average correlation of 0.1
440 (correlations are calculated in the logit scale; Fig.6e and Fig.6f). In general, the correlations are
441 stronger for geographically close SU. The results show strong correlations for SU within NA
442 (0.409), followed by SE (0.149) and NE (0.087).

443 **3.2 Comparisons between the new life cycle and currently used PFA models to assess and**
444 **forecast salmon population dynamic and to evaluate catch options for mixed stock**
445 **marine fisheries**

446 Most of the differences we might observe in the following sections between PFA and life cycle
447 models will rely on two main reasons defined previously in section 2.2.2.3: the difference in
448 demographic structures between PFA and life cycle model and how the uncertainty is considered
449 and propagated in the model by the PFA and the life cycle frameworks.

450 Overall, the life cycle model generates more accurate posterior estimates of abundances,
451 demographic parameters and fishery exploitation rates for both hindcast and forecast analysis (Fig.
452 7, Fig.8 and Supp.Mat.III. Fig.SIII.2-18.). In the following sections we analyzed results per CSG
453 for hindcast and forecast analysis and finally we explored how both PFA and life models
454 performed to evaluate catch options for mixed stock marine fisheries.

455 3.2.1 Hindcast and Forecast analysis

456 In this section we provide a comparison of the key variables and parameters constituting both PFA
457 models and life cycle models: Productivity (1SW and/or 2SW), Proportion of fish maturing as
458 1SW, Eggs deposition, Pre-Fishery Abundance (1SW and/or 2SW), Returns (1SW and/or 2SW).

459 3.2.1.1 *Europe : NE and SE*

460 *Hindcast*

461 Because most of the results are consistent between all stock units within SE and NE CSGs, we
462 propose in this section to present the results for only one SU, Scotland. We provided results for
463 other stock units in Supp.Mat.III.

464 Posterior estimates of abundances (Eggs deposition, PFA, Returns) and demographic parameters
465 (Productivity and proportion of fish maturing as 1SW) are consistent between the life cycle and
466 PFA models over the time-serie from 1978 to 2020 (Fig.7 and Supp. Mat.III). For both PFA and
467 LCM models trends in Returns (1SW and 2SW), PFA (1SW and 2 SW) and eggs deposition (total
468 1SW + 2SW) show a general temporal decreasing trend; trend in productivity shows a decline
469 while proportion in fish maturing as 1SW show an increasing trend over the time-series.

470 For SE and NE stock units, results highlight few differences in posterior distributions for PFA and
471 LCM models. Those differences mainly concern times series of PFA non maturing and
472 Productivity for FR and EW stock units. For FR and EW stock units, differences in posterior
473 distributions for PFA and LCM result in how the predictions might fit the data in the life cycle
474 model, as already highlighted in Olmos et al. (2019) (see Supp. Mat in Olmos et al. 2019).

475 *Forecast*

476 Overall, for both PFA and LCM models, forecasted years (2018-2019-2020) show a common
477 pattern. LCM showed more accurate predictions. Uncertainties in forecast from PFA models can
478 be really big. As a consequence, when considering the forecast for the egg deposition, even if PFA
479 and life cycle predict similar median, they show different Bayesian posterior distributions due to

480 how the models account and propagate differently the uncertainty. Such a difference might have
481 an impact on the risk analysis (see section 3.2.2).

482 3.2.1.2 NA

483 *Hindcast*

484 In this section we presented the results for only two SUs, GF and NF, which illustrate the main
485 differences between posterior estimates from PFA and life cycle models for NA SUs. We provided
486 results for other stock units in Supp.Mat.III.

487 Posterior estimates of abundances (Eggs deposition, PFA, Returns) and demographic parameters
488 (Productivity) are not fully consistent between the life cycle and PFA models over the time-serie
489 from 1971 to 2020 (Fig.8, Supp. Mat. III). The reason is because PFA models for NA CSG
490 considers only fish non-maturing after the first winter at sea. Then, for stock units having returns
491 largely dominated by 2SW (US, Scotia-Fundy, Quebec, Gulf), posterior estimates of abundances
492 (Fig.8.e, Fig.8.f and Fig.8.h, Supp. Mat. III) and productivity show a similar trends and values for
493 both PFA and life cycle models (see purple 2SW productivity from the LCM and blue PFA
494 productivity Fig.8.g). However, for SUs having returns largely dominated by 1SW (Labrador and
495 Newfoundland), consistency between PFA and life cycle models will depend on how parameters
496 and variables are accounting for only non-maturing fish or for both maturing and non maturing
497 fish. Variables and parameters specific to non-maturing fish as, 2SW Returns (Fig.8.g) and PFA
498 of fish non maturing the first year at sea (Fig.8.h) show consistent temporal pattern and value
499 between PFA and life cycle model. However, in the life cycle model eggs deposition is defined as
500 the sum of eggs from both 1SW and 2SW fish while PFA model only track 2SW fish. Posterior

501 estimates of eggs deposition from the life cycle model were estimated to be 5 times larger than
502 estimates from PFA models.

503 *Forecast*

504 Overall, for both PFA and LCM models, forecasted years (2018-2019-2020) show differences.
505 LCM showed more accurate predictions. Uncertainties in forecast from PFA models can be really
506 big.

507 *Note : In this version (13.10.2023) of the manuscript we are facing some issues with how the PFA*
508 *model forecast the egg deposition. Indeed, PFA models built by the WGNAS are defined to forecast*
509 *Pre-fishery abundances only. But to be able to compare PFA and life cycle models we had to model*
510 *transitions between PFA stages and eggs stages for the PFA models. For the hindcast it worked*
511 *well but for the forecast years 2019-2020 present higher values than hindcasted years, which does*
512 *not make sense for now. We are still investigating the reason and hope to provide a response for*
513 *the benchmark. However, we argue that it does not impact our demonstration showing that the life*
514 *cycle model is an operational to assess salmon population across the North Atlantic basin*

515 3.2.2 Risk analysis: Evaluating catch options for mixed stock marine fisheries

516 3.2.2.1 *Eggs deposition compared to CLs*

517 Results show how uncertainty in the forecasts increases with forecasting horizon. This is mostly
518 the consequence of uncertainty propagation through time in forecasts of the post-smolts survival
519 and proportion maturing modelled as multivariate random walks. Results show that the life cycle
520 model generates more accurate posterior estimates than PFA (Fig. 7 and 8).

521 3.2.2.2 *The West Greenland mixed stock fishery*

522 The probability that the eggs deposition achieves the CLs under any fishing scenarios is directly
523 quantified through Monte Carlo draws. The probabilities of achieving management objectives are
524 higher for the stocks in Northern and Southern Europe (Fig. 9). Stocks from Northern Europe have
525 the highest probabilities of achieving their management objectives. In Southern Europe, Northern
526 Ireland, Southwest Iceland, Scotland and England and Wales and Northern Ireland have the highest
527 probabilities of achieving their management objectives. In contrast, Ireland and France from SE,
528 and stocks from NA such as US, Scotia-Fundy and Gulf have very low probabilities of achieving
529 their management objectives. As expected, different catch options at West Greenland have
530 minimal influence on the probability of achieving management objectives for stocks that represent
531 only a very low proportion of the catches at West Greenland, such as all stocks of NE (that
532 represent less than 5% of the total fish harvested in West Greenland) and most of the stocks of SE.
533 Because they present the highest exploitation rate at WG, stocks from NA such as Labrador,
534 Quebec, and Gulf have their probability of achieving management objectives decreasing when
535 catch options increase.

536 When assessed at the scale of stock complexes (i.e. by comparing the total egg deposition at the
537 scale of stock complex to the total CL at the scale of stock complex), probabilities to achieve
538 conservation limits are the highest for Northern Europe, the lowest for southern Europe, and
539 intermediate for North America (Fig 9). The North American stock complex logically revealed the
540 most sensitive to catches scenarios at WG (Fig. 9)

541 When comparing PFA and LCM models, results are pretty consistent. Differences exists for SU
542 from NA CSG like NF that considers only fish non-maturing after the first winter at sea Fig. 9). In
543 NE and SE CSG, differences exist for NO and RU stocks between PFA and LCM (Fig. 9). Those

544 differences can be explained by the fact that in the life cycle models the risk analysis is carried out
545 for all stock units constituting NO (NO.MI, NO.NO, NO.SE,; NO.SW) and RU (RU.AK, RU.KB;
546 RU.KW, RU.RP) by summing a posteriori the abundances of eggs deposition whereas in PFA
547 models, the risk analysis is conducted from aggregated data, where the calculated productivity is
548 the aggregated productivity of a given stock unit. Differences in uncertainties generated by each
549 model can also explain some observed differences in Fig.9. For example, for IR SU (Fig. 9 and
550 Supp.Mat.III Fig.SIII.8), the probability to reach the conservation limit is higher is higher with the
551 PFA model than with the life cycle model. But both PFA and life cycle model predict the same
552 abundance in term of median (Supp.Mat.III Fig.SIII.8). The only difference is that for the life cycle
553 model the posterior distribution is narrower (uncertainty is lower) and so only a small part of the
554 posterior distribution overlaps the conservation limits (meaning that for the life cycle model fewer
555 Monte Carlo draws are above the conservation limits), which generates a lower probability of
556 reaching the conservation limit.

557 3.2.2.3 *The Faroes mixed stock fisheries*

558 As expected, SU from North America are insensitive to catch options at Faroes as fish from North
559 America are not supposed to be caught there (Fig. 10). Catch options at Faroes influence the
560 probability of achieving management objectives for European SU only (Fig. 10). Southwest
561 Iceland, Northeast Iceland, England and Wales, Northern Ireland, Norway and Russia have the
562 highest probabilities of achieving their management objectives (Fig. 10). By contrast France,
563 Scotland and Sweden have low probabilities of achieving their management objectives. Logically,
564 Sweden and Norway that represent the highest proportion of catches at Faroes are the most
565 sensitive to the catch options at Faroes.

566 When assessed at the scale of stock complexes (i.e. by comparing the total egg deposition at the
567 scale of stock complex to the total CL at the scale of stock complex), probabilities to achieve
568 conservation limits are the highest for Northern Europe and the lowest for southern Europe (Fig
569 10). The Northern European stock complex logically revealed the most sensitive to catches
570 scenarios at Faroes. (Fig. 10).

571 Results between PFA and LCM are consistent. The mains differences occurs again for Norway
572 and Russia stock units which have the specificity to have their productivity calculated from
573 aggregated stock units whereas in the life cycle model productivity and eggs depositions are
574 estimated for each stock unit constituting Norway and Russia and then aggregated to be compared
575 to PFA outputs. And the main differences can also be explained by how the uncertainty is
576 quantified by the PFA and life cycle model.

577 *3.2.2.4 Evaluating catch options for both Faroes and West Greenland Fisheries*

578 The new life cycle model allows for the first time to evaluate simultaneously catch options in
579 Faroes and West Greenland (Fig. 11). As already shown with the independent assessment of
580 Faroes and West Greenland fisheries, Southwest Iceland have the highest probabilities of
581 achieving their management objectives and Ireland the lowest.

582 Interestingly, eggs deposition in France England&Wales and in Norway are more sensitive to
583 catch options in the Faroes fisheries than in the West Greenland fishery. Indeed, those three
584 stock units (or aggregated stock units) represent 50% of the catches in the 1SW maturing Faroes
585 fishery, and a few amounts of fish harvested in the West Greenland fishery.

586

587 **4 CONCLUSION**

588 The new life cycle model provides a singular harmonized framework to simultaneously assess
589 two sea-classes of Atlantic salmon for all SU in North America and Europe. This represent a
590 paradigm shift from the stock assessment and forecasting approach currently used by ICES
591 considers the North American and European (Southern and Northern) continental stock groups
592 separately and these models have different demographic structures. It also allows for modelling
593 covariations among all SU and for partitioning the effects of fisheries from the effects of
594 environmental factors at a hierarchy of spatial scales, including at the level of the North Atlantic,
595 of each CSG, and for each SU within a CSG.

596 Moving from PFA models to life cycle model to assess, forecast and carry out risk analysis for
597 Atlantic salmon stock unit across the all North Atlantic basin will change the workflow of the
598 WGNAS (Fig.12).

599

7 REFERENCES

- 601 Buckland, S. T., Newman, K. B., Thomas, L., and Koesters, N. B. 2004. State-space models for
602 the dynamics of wild animal populations. *Ecological Modelling*, 171: 157–175.
- 603 Chaput, G., Legault, C., Reddin, D., Caron, F., and Amiro, P. 2005. Provision of catch advice
604 taking account of non-stationarity in productivity of Atlantic salmon (*L.*) in the
605 Northwest Atlantic. *ICES Journal of Marine Science*, 62: 131–143.
- 606 Clark, J. S. 2004. Why environmental scientists are becoming Bayesians: Modelling with Bayes.
607 *Ecology Letters*, 8: 2–14.
- 608 Cressie, N., Calder, C. A., Clark, J. S., Hoef, J. M. V., and Wikle, C. K. 2009. Accounting for
609 uncertainty in ecological analysis: the strengths and limitations of hierarchical statistical
610 modeling. *Ecological Applications*, 19: 553–570.
- 611 Crozier, W. W., Schön, P.-J., Chaput, G., Potter, E. C. E., Maoiléidigh, N. Ó., and MacLean, J.
612 C. 2004. Managing Atlantic salmon (*Salmo salar* L.) in the mixed stock environment:
613 challenges and considerations. *ICES Journal of Marine Science: Journal du Conseil*, 61:
614 1344–1358.
- 615 ICES. 2018. Report of the Working Group on North Atlantic Salmon (WGNAS), 4–13 April
616 2018, Woods Hole, MA, USA. ICES CM 2018/ACOM:21. 386 pp.
- 617 Minto, C., Mills Flemming, J., Britten, G. L., Worm, B., and Rose, K. 2014. Productivity
618 dynamics of Atlantic cod. *Canadian Journal of Fisheries and Aquatic Sciences*, 71: 203–
619 216.
- 620 Olmos, M., Massiot-Granier, F., Prévost, E., Chaput, G., Bradbury, I. R., Nevoux, M., and Rivot,
621 E. 2019. Evidence for spatial coherence in time trends of marine life history traits of
622 Atlantic salmon in the North Atlantic. *Fish and Fisheries*, 20: 322–342.
- 623 Parent, E., and Rivot, E. 2012. Introduction to hierarchical Bayesian modeling for ecological
624 data. Chapman & Hall/CRC applied environmental statistics. CRC Press, Boca Raton.
625 405 pp.
- 626 Potter, T. 2001. Past and present use of reference points for Atlantic salmon. In: Prévost, E. &
627 Chaput, G., eds. Stock, recruitment and reference points: assessment and management of
628 Atlantic salmon. Paris: Hydrobiologie et Aquaculture, INRA Editions, pp. 195–223.
- 629 Rago, P. J., Reddin, D. G., Porter, T. R., Meerburg, D. J., Friedland, K. D., and Potter, E. C. E.
630 1993. A continental run reconstruction model for the non-maturing component of North
631 American Atlantic salmon: analysis of fisheries in Greenland and Newfoundland-
632 Labrador, 1974-1991. ICES, COPENHAGEN(DENMARK). 1993.
- 633 Ripa, J., and Lundberg, P. 2000. The route to extinction in variable environments. *Oikos*, 90: 89–
634 96.
- 635 Rivot, E., Prévost, E., Parent, E., and Baglinière, J. L. 2004. A Bayesian state-space modelling
636 framework for fitting a salmon stage-structured population dynamic model to multiple
637 time series of field data. *Ecological Modelling*, 179: 463–485.
- 638 Windsor, M. L., and Hutchinson, P. 1994. International management of Atlantic salmon, *Salmo*
639 *salar* L., by the North Atlantic Salmon Conservation Organization, 1984-1994. *Fisheries*
640 *Management and Ecology*, 1: 31–44.
- 641

Table 1 :Weaknesses of the currently used stock assessment (PFA models) and proposed improvements through the hierarchical life cycle model.

	Improvement through the life cycle model
<p>[W1] A coarsely constructed stock-recruitment dynamic</p> <ul style="list-style-type: none"> • Forecasts of the returns are based on forecasts of productivity between a spawning potential and abundance at the PFA stage (measure of the recruitment) • But the dynamic link between PFA and subsequent egg depositions is not represented; so statistical inferences on productivity is susceptible to time series bias • Lack of flexibility in the statistical modelling framework restricts the integration of the large amount of available data and knowledge 	<ul style="list-style-type: none"> • A life cycle to represent all stages and life histories. • This integrated life cycle framework is expandable • Assimilates new sources of information to improve the ecological and biological realism of the model
<p>[W2] Measure of the stock and the recruitment are derived from the same data</p> <ul style="list-style-type: none"> • A combination of three models <ol style="list-style-type: none"> i. The same model is used to estimate the abundance of fish at the PFA stages (recruitment) and to estimate the potential number of spawners or eggs (stock) ii. A model to forecast the evolution of the productivity parameter iii. The forecast model serves as a basis to forecast the PFA and the number of fish that return to homewater based on catches scenarios at sea 	<ul style="list-style-type: none"> • The same model is used for both the inferences hindcasting and forecasting phases • All the model properties and sources of uncertainties are readily integrated into the forecast process.
<p>[W3] Three different and independent PFA models for each complex with different life histories modelled</p> <ul style="list-style-type: none"> • The two European models explicitly consider 1SW and 2SW fish in the population dynamics, • The model for NA only considers the dynamics of 2SW <ol style="list-style-type: none"> i. implicitly assumes that 2SW spawners only produce 2SW fish in future cohorts, and excludes contributions of 1SW ii. Temporal variations of productivities are therefore not comparable to the PFA models built for the European CSG considering both 1SW and 2SW productivity. • Cannot evaluate the commonality in temporal trends between all SU in the North Atlantic. • Ignores any covariance structure in the dynamics of the SU <ol style="list-style-type: none"> i. that may share common environments at sea ii. that are exploited in sea fisheries • Precludes evaluation of the consequences of scenarios on multiple stock complexes simultaneously • Hypotheses on drivers and mechanisms of changes cannot be easily tested 	<ul style="list-style-type: none"> • A single unified life cycle approach with all populations following a similar life history process • A framework to enhance the ecosystem approach. This framework analyzes the mechanisms that shape population responses to variations in marine ecosystems <ol style="list-style-type: none"> i. by modelling covariations among all SU ii. by partitioning the effects of fisheries from the effects of environmental factors iii. at a hierarchy of spatial scales • Evaluate catch options for the Faroes and West Greenland separately or simultaneously and for all SU separately or simultaneously.

Table 2: Management objectives relevant to the development of catch options for the stocks units in NA, SE and NE.

North America		
Stock Units	CLs	References
Labrador	243660000	O'Connel et al. 1997
Newfoundland	267780000	Reddin et al. 2009
Quebec	50380000	Atlantic salmon management plan 2016, Ministère des Forêts, de la Faune et des Parcs (2016).
Gulf	248680000	Cameron et al. 2009, Breau et al. 2009, Chaput et al. 2010, Cairns et al. 2015
Scotia Fundy	224140000	Gibson et al. 2014, Bowlby et al. 2013, Jones et al. 2014
US	435369000	Baum, E.T. 1995
Southern Europe		
Stock Units	CLs	References
Iceland (south+west)	64273104	
Scotland	1609542000	
Northern Ireland	56281942	
Ireland	710711690	ICES, 2015a
England&Wales	211419850	
France	55165500	
Northern Europe		
Stock Units	CLs	References
Iceland (north+east)	23889096	
Sweden	13997100	
Norway	444064980	ICES, 2015a
Finland	104278220	
Russia	357856550	




Figure 2: Spatial repartition of the 25 stock units considered in North Atlantic. Stock units of North America: NFDL=Newfoundland, GF=Gulf, SF=Scotia-Fundy, US=USA, QB=Quebec and LB=Labrador ; Stock units in Southern Europe: IR=Ireland, E&W=England&Wales, FR=France, E.SC=Eastern Scotland, W.SC=Western Scotland, N.IR=Northern Ireland, IC.SW=South-West Iceland ; Stocks units in Northern Europe: FI=Finland, IC.NE=North-East Iceland, NO.MI=Middle Norway, NO.NO=North Norway, NO.SE=South-East Norway, NO.SW=South-West Norway, RU.AK=Russia Arkhangelsk Karelia, RU.KB=Russia Kola Barents Sea, RU.KW=Russia Kola White Sea, RU.RP=River Pechora, SWD=Sweden. **UPDATE NI SUS**




Figure 3: Structure of the age- and stage-based life cycle model for the 25 SU. Grey boxes: different stages during the marine (grey) and freshwater (black) phases. Green circles belong to NA SU, Red circles belong to SE SU and blue circles belong to NE SU. Cylinder: sources of covariations among the 24 SU. Orange cylinders: key parameters (post-smolt survival and maturing probability). Purple cylinders: fisheries operating on mixture of SE and NE SU. Green cylinders: fisheries operating on mixture of NA SU. Grey cylinders: fisheries operating on mixture of NA, SE and NE SU.




Figure 4: Outline of the modelling flows for PFA (blue) and life cycle (red) models. Run reconstruction (in green) are processing observation models that reconstruct probability distributions synthesizing observation uncertainty around the time series of catches and returns for the 25 SUs. Outputs from the Run Reconstruction models are used as pseudo observation for both the PFA models and the life cycle model. PFA models are defined for each complex whereas life cycle model is a unique framework integrating all complexes with the same statistical model.




Figure 5 Time series of estimated abundances at the PFA stage (maturing + non maturing PFA) for all SU for the three continental stock groups: (a) summed by CSG and all CSG (Atlantic basin), (b) North America, (c) Southern Europe; (d) Northern Europe. Thick lines: median of the marginal posterior distributions. PFA are standardized to the first year values.




Figure 6: Post-smolt survival : (a) Time series of smolt-PFA survival (also called productivity) (median of the marginal posterior distributions plotted in the natural scale) for the 25 SU (thin grey lines), averaged over the three continental stock groups (thick color lines) and averaged across all CSG (Atlantic basin, thick black line). (b) –(c) –(d) Time series of post-smolt survival (median of the marginal posterior distributions plotted in the natural scale) for the 25 SU grouped by continental stock groups (North American, Southern Europe and Northern Europe respectively). (e) Pairwise correlations calculated between all SUs (median of the posterior distribution from the variance-covariance matrix in the logit scale). (f) Pairwise correlations (calculated in the logit scale) averaged over all SUs, over SU within the same CSG (NA, SE, NE) and over pairs of SU that belong to two different CSG (NA-SE, NA-NE and SE-NE).




Figure 7: Proportion of maturing as 1SW : (a) Time series of proportion of maturing as 1SW (median of the marginal posterior distributions plotted in the natural scale) for the 25 SU (thin grey lines), averaged over the three continental stock groups (thick color lines) and averaged across all CSG (Atlantic basin, thick black line). (b) –(c)–(d) Time series of proportion of maturing as 1SW (median of the marginal posterior distributions plotted in the natural scale) for the 25 SU grouped by continental stock groups (North American, Southern Europe and Northern Europe respectively). (e) Pairwise correlations calculated between all SUs (median of the posterior distribution from the variance-covariance matrix in the logit scale). (f) Pairwise correlations (calculated in the logit scale) averaged over all SUs, over SU within the same CSG (NA, SE, NE) and over pairs of SU that belong to two different CSG (NA-SE, NA-NE and SE-NE).




Figure 8: Probability distributions of the number of egg potentially spawned, PFA (total PFA.m + PFA.nm), PFA.maturing (PFA.m, PFA.non maturing (PFA.nm)), Productivity, Proportion of fish maturing as 1SW, Returns 1SW, and Returns (2SW) for Scotland. Thick point is the median and line represent the 95% posterior credibility interval. Blue: PFA model; Red : Life cycle model (LCM); Dark color: historical time series; Light colors: Forecasted years (under the scenario of no catches at West Greenland and Faroes)




Figure 9: Probability distributions of (a, e) 2SW Productivity, (b, f) the number of egg potentially spawned by 2SW fish, (c, g) 2SW Returns, and (d, h) PFA.nm for Quebec (a - f) and Newfoundland (e-h). Thick point is the median and line represent the 95% posterior credibility interval. Blue: PFA model; Red : Life cycle model (LCM); Dark color: historical time series; Light colors: Forecasted years (under the scenario of no catches at West Greenland)




Figure 10: Probability to reach Conservation Limits obtained under different catches options at West Greenland. Catches options: 0, 500, 1000, 1500, 2000 and 2500 tons (3 years projections). (LB, NF, QB, GF, SF, US) North America; (EW, IR, N.IR, SC, IC.SW) Southern Europe; (IC.NE, SW, NO, FI, RU) Northern Europe. Panels (NA, SE, NE) give probabilities to simultaneously achieving the management objectives for all SU of North America (NA), Southern Europe (SE) and Northern Europe (NE). Models : circle=Life cycle model (LCM), triangle= PFA




Figure 11: Probability to reach Conservation Limits obtained under different catches options at Faroes. Catches options: 0, 500, 1000, 1500, 2000 and 2500 tons (3 years projections). (EW, IR, N.IR, SC, IC.SW) Southern Europe; (IC.NE, SW, NO, FI, RU) Northern Europe. Panels (NA, SE, NE) give probabilities to simultaneously achieving the management objectives for all SU of North America (NA), Southern Europe (SE) and Northern Europe(NE). Models : circle=Life cycle model (LCM), triangle=PFA. Stock Units of North America are not impacted by Faroes fisheries and are not represented.




Figure 12a: Probability to reach Conservation Limits simultaneously under different catches options at West Greenland and Faroes: 0, 500, 1000, 1500, 2000 and 2500 tons (3 years projections) for SU of the Southern European complex potentially impacted by both mixed stock fisheries.




Figure 11b: Probability to reach Conservation Limits simultaneously under different catches options at West Greenland and Faroes: 0, 500, 1000, 1500, 2000 and 2500 tons (3 years projections) for SU of the Northern European complex potentially impacted by both mixed stock fisheries.




Figure 13: Conceptual representation about how the new life cycle model impacts the workflow of the WGNAS in term of human resource and interactions.

6 SUPP.MAT.1. - Detailed process and observation equations of the Bayesian life cycle model

1.1 Population dynamics

1.1.1 Simplified life history

The age- and stage-structured life cycle model has a similar structure for each SU. It includes variation in the age of juveniles out-migrating from freshwater (i.e. smolts) and the sea-age of returning adults. Smolts migrate seaward after 1 to 6 years spent in freshwater (depending on SU). Two sea-age classes are considered in the model: Maiden salmon that return and reproduce after one year at sea, referred to as one-sea-winter (1SW) salmon or grilse, and maiden salmon that return after two winters spent at sea (2SW). This is a simplification of the variety of life history as some maiden fish may spent more than two winters at sea before returning to spawn, or some may be repeat spawners. However, those fish are rare and the 6 smolt-ages \times 2 sea-ages combinations capture the essence of life history variations.

1.1.2 Eggs deposition

The total number of eggs potentially spawned in year t for SU r is calculated from the number of 1SW ($N_{7,t,r}$) and 2SW ($N_{10,t,r}$) spawners escaping the homewater fisheries and the average number of eggs potentially spawned per 1SW and 2SW salmon, denoted $eggs_{1,t,r}$ and $eggs_{2,t,r}$ (Table 2):

$$(A1.1) \quad N_{1,t,r} = N_{7,t,r} \times eggs_{1,t,r} + N_{10,t,r} \times eggs_{2,t,r}$$

1.1.3 Egg-to-smolt transition

The egg-to-smolt transition consists of two steps: the survival from egg-to-smolt per cohort, and the distribution of the surviving smolts according to their age at downstream migration.

1.1.3.1 Egg-to-smolt survival

Because no smolt production data is available at the scale of SU, it is difficult to separate the variability of the egg-to-smolt survival from that of the post-smolt survival, and parameters of the egg-to-smolt transitions have to be fixed. The egg-to-smolt survival is density independent, with average survival rate θ_1 arbitrarily fixed to 0.007 (Hutchings & Jones, 1998; Massiot-Granier et al. 2014) for all years and all SU (Table 2). Environmental stochasticity is modelled by logNormal random noise with variance $\sigma_{\theta_1}^2$ fixed to an arbitrarily value corresponding to $CV_{\theta_1}=0.4$ ($\sigma_{\theta_1}^2 = \log(CV_{\theta_1}^2 + 1)$) which is a median values for the inter-annual variability found in

the literature (Prevost et al., 2003; Pulkkinen et al., 2013). The total number of smolts produced in the cohort c (corresponding to egg deposition of year c), denoted $N_{2,c,r}$ is then modelled as:

$$(A1.2) \quad \log(N_{2,c,r}) \sim \text{Normal}(\log(\theta_{2,c,a,r} * N_{1,c,r}) - \frac{1}{2}\sigma_{1,c,r}^2, \sigma_{2,c,r}^2)$$

This model configuration only allows for random stochasticity in the egg-to-smolt survival and does not account for any compensation neither (but see Olmos et al. 2019 for a sensitivity analysis to inclusion of density dependence). This implicitly assumes that any trends in the stock productivity over time are a response to changes in the marine phase, what may inflate the importance of trends in the post-smolt survival.

1.1.3.2 Distribution according to smolt ages

The probabilities of a smolt in the cohort c migrating at age $a = 1, \dots, 6$ at year $t = c + a + 1$, denoted $\theta_{2,c,a,r}$ are fixed to their averaged proportions $psm_{1:6,r}$ specific to each SU (Table 2).

Given $\theta_{2,c,a,r}$ the number of smolts from the cohort c that migrate at age a year $t = c + a + 1$ is modelled as:

$$(A1.4) \quad N_{2,c,a,t=c+a+1,r}^1 = \theta_{2,c,a,r} \times N_{2,c,r}$$

Last, the number of smolts migrating in the spring of year t is the sum of all smolts of different ages (and therefore of different cohorts) migrating in year t :

$$(A1.5) \quad N_{3,t,r} = \sum_{a=1}^{a=6} N_{2,c=t-a-1,a,t,r}^1$$

Note:

In a previous model version, smolt ages distribution ($\theta_{2,c,a,r}$) was estimated. Proportion of smolt ages were set a tight Dirichlet prior distribution. However, this transition revealed serious bottleneck for computational time needed to reach convergence. A simpler solution using fixed proportions is then adopted to keep reasonable model execution time.

1.1.4 Marine phase

The marine phase is modelled as a sequence of three blocks of transitions: survival from smolts to the PFA stage, the maturation of fish at the PFA stage, and the fishing and natural mortality between PFA and returns.

1.1.4.1 Post-smolt survival and proportion of fish maturing as 1SW

Time series of post-smolt survivals ($\theta_{3,t,r}$) and the proportion of fish maturing as 1SW ($\theta_{4,t,r}$) are modelled as multivariate random walks in the logit scale. Random variations are drawn from multivariate Normal distributions with variance-covariance matrix Σ_{θ_3} and Σ_{θ_4} that define the covariations among the SU (Minto et al., 2014; Ripa and Lundberg, 2000):

$$(A1.6) \quad \begin{aligned} & \text{First year (for } r = 1:N): \text{logit}(\theta_{3_{t,r}}) \sim \text{Normal}(0,1) \\ & \text{Then } (\text{logit}(\theta_{3_{t+1,r}}))_{r=1:N} \sim \text{MVNormal}((\text{logit}(\theta_{3_{t,r}}))_{r=1:N}, \Sigma_3) \end{aligned}$$

$$(A1.7) \quad \begin{aligned} & \text{First year (for } r = 1:N): \text{logit}(\theta_{4_{t,r}}) \sim \text{Normal}(0,1) \\ & \text{Then } (\text{logit}(\theta_{4_{t+1,r}}))_{r=1:N} \sim \text{MVNormal}((\text{logit}(\theta_{4_{t,r}}))_{r=1:N}, \Sigma_4) \end{aligned}$$

Then, given the number of smolts migrating in year t ($N_{3_{t,r}}$) and the post-smolt survival ($\theta_{3_{t,r}}$), the number of posts-smolts that survive to the PFA stage ($N_{4_{t+1,r}}$) in January of year $t + 1$ is modelled as:

$$(A1.8) \quad N_{4_{t+1,r}} = \theta_{3_{t,r}} \times N_{3_{t,r}}$$

Given the number of fish at the PFA stage ($N_{4_{t+1,r}}$) and the maturation rate ($\theta_{4_{t+1,r}}$), mature ($N_{S_{t+1,r}}$) and non mature fish ($N_{8_{t+1,r}}$) at the PFA stage are modelled as:

$$(A1.9) \quad N_{S_{t+1,r}} = \theta_{4_{t+1,r}} \times N_{4_{t+1,r}}$$

$$(A1.10) \quad N_{8_{t+1,r}} = (1 - \theta_{4_{t+1,r}}) \times N_{4_{t+1,r}}$$

Note: For this 2018 version, those transitions were modelled as stochastic, using lognormal distribution with standard deviation fixed to a very low value. In the current version (2023). Those transitions are now modelled as deterministic to keep reasonable model execution time.

1.1.4.2 Sequential marine fisheries and natural mortality

After the PFA stage, both maturing and non-maturing fish are subject to natural mortality and sequential fisheries mortalities operating on mixed stocks (Tables 4 & 5). The following modelling structure applies for each of those transitions. For any marine fishery f , operating in year t on a number of fish $N_{f_{t,r}}$ originated from the stock unit r with an exploitation rate $h_{f_{t,r}}$, the catches $C_{f_{t,r}}$ (unknown states) and the number of fish that escape the fishery $N_{f,esc_{t,r}}$ are modelled as:

$$(A1.11) \quad C_{f_{t,r}} = h_{f_{t,r}} \times N_{f_{t,r}}$$

$$(A1.12) \quad N_{f,esc_{t,r}} = (1 - h_{f_{t,r}}) \times N_{f_{t,r}}$$

Exploitation rates $h_{f_{t,r}}$ are modelled as variable over time but their variability across SU is modelled differently depending on the data available to allocate catches to each SU and on expert knowledge about migration routes. Exploitation rates of the West Greenland fishery (WG; operating on a mixture of SU from North America and Europe) and of the Faroes fishery (FA; operating on SU from Europe only) were all supposed to vary across years and SU (Tables 5 & 6).

For the fisheries specific to the SU from NA (Table 5), because no proportion data are available to allocate catches among SU, catches were allocated to each SU by considering a single h homogeneous for all SU. There is however an exception to this general rule (Prévost et al., 2009): for the Labrador/Newfoundland (LAB/NFDL and Saint-Pierre et Miquelon (SPM) fisheries on 1SWm and 2SW fish, a separate h is estimated for Labrador SU, and a single h is considered for the five other North American SU, corresponding to catch allocation proportional to abundance.

All fisheries at sea are separated by periods of time where only natural mortality occurs (ICES, 2018; Potter, 2016; Prévost et al., 2009). Fish that escape the fishery f at year t hence suffer natural mortality rate $\theta_{S_{t,f}} = e^{-M \times \Delta_{t,f}}$ where the monthly mortality rate M is fixed, constant across years and SU's ($M = 0.03 \cdot \text{month}^{-1}$; Table 2) and the duration $\Delta_{t,f}$ (in months) are assumed known and constant across years but with some variations among SU to account for variability in migration routes (Tables 4 & 5):

$$(A1.13) \quad N_{f+1_{t,r}} = (1 - \theta_{S_{t,f}}) \times N_{f.\text{esc}_{t,r}}$$

1.1.4.3 From returns to spawners (homewater catches)

Fish that escape all marine mortality and return as 1SW fish ($N_{6_{t,r}}$) or 2SW fish ($N_{9_{t,r}}$), are subject to homewater fisheries that operate locally on each SU. Homewater fisheries are modelled with exploitation rates $h_{\text{HW}f_{t,r}}$ that are assumed to vary with years and SU and for the two sea-age classes separately (Tables 5 & 5). Homewater fishery harvest rates are estimated. After homewater fishery, a proportion of fish may potentially delay spawning to the next year. The proportion of delayed spawners are supposed known but varies with SU, years and sea-age classes and are denoted $p_{\text{delSp}_{t,r}}$ (specific for 1SW and 2SW). Fish that do not delay spawning are potentially subject to an additional fishery with harvest rate denoted $h_{\text{sup}_{t,r}}$ (specific for 1SW and 2SW). Fish that delay spawning to the next year may then be subject to a specific fishery with (estimated) harvest rates $h_{\text{delSp}_{t,r}}$ (specific to 1SW and 2SW). An additional survival rate ($\theta_{6_{t,r}}$ or $\theta_{9_{t,r}}$ for 1SW and 2SW, respectively) is then finally applied on all remaining fish before spawning. In practice, the proportion of delayed spawners is non-zero only for Russian stock units. The supplementary harvest rate $h_{\text{sup}_{t,r}}$ (and associated catches) are non null only for Scotland East and West. But these transitions are modelled uniformly for all stock units with zero proportion of delayed spawners in the data and zero additional catches for almost all SU. Last, the number of 2SW spawners in the US stock unit is also supplemented by stocking. The transition is also modelled uniformly for all SU but the number of fish stocked $n_{\text{Stock}.2\text{SW}_{t,r}}$ is null for all SU except USA. Finally, the number of fish that escape the homewater fishery and potentially spawn as 1SW ($N_{7_{t,r}}$) and 2SW ($N_{10_{t,r}}$) are modelled as:

$$(A1.14) \quad N_{7_{t,r}} = ((1 - h_{\text{HW}f.1\text{S}_{t,r}}) \times (1 - p_{\text{delSp}.1\text{SW}_{t,r}}) \times (1 - h_{\text{sup}.1\text{S}_{t,r}}) \times N_{6_{t,r}} + (1 - h_{\text{HW}f.1\text{SW}_{t-1,r}}) \times p_{\text{delSp}.1\text{SW}_{t-1,r}} \times (1 - h_{\text{delSp}.1\text{SW}_{t,r}}) \times N_{6_{t-1,r}}) \times \theta_{6_{t,r}}$$

$$(A1.15) \quad N_{10_{t,r}} = ((1 - h_{\text{HW}f.2\text{SW}_{t,r}}) \times (1 - p_{\text{delSp}.2\text{SW}_{t,r}}) \times N_{9_{t,r}} + (1 - h_{\text{HW}f.2\text{SW}_{t-1,r}}) \times p_{\text{delSp}.2\text{SW}_{t-1,r}} \times (1 - h_{\text{delSp}.2\text{SW}_{t,r}}) \times N_{9_{t-1,r}}) \times \theta_{9_{t,r}} + n_{\text{Stock}.2\text{SW}_{t,r}}$$

1.2 Observation equations

The model incorporates observation errors for all time series of returns and catches. A sequential approach (Michielsens et al., 2008; Staton et al., 2017) is used that consists of two steps:

- In a first step, observation models are processed separately to reconstruct probability distributions that synthesize observation uncertainty around catches and returns for each year and each of the 25 SU. Probability distributions for returns and catches are derived from a variety of raw data and observation models, specific to each SU and each year and originally developed by ICES to provide input for PFA models for NA (Rago et al., 1993) and SE (Potter et al., 2004b) stock units.
- In a second step, those distributions are used to approximate likelihoods in the population dynamics state-space model.

1.2.1 Returns

Returns are estimated for each year, each SU and for the two sea-age classes separately. Raw data used to estimate return essentially consist in homewater catches available at the scale of rivers or regional fishery jurisdictions, scaled by harvest and declaration rates and then aggregated at the scale of larger stock units. Uncertainties then essentially arise from a numerical (Monte Carlo) integration of uncertainties about harvest and declaration rates. Other fishery independent information like counting fences or mark and recapture data can also be used. Detailed description of the raw data and models used in each SU is provided in the WGNAS Stock Annex for Atlantic salmon (Crozier et al., 2003; ICES, 2002, 2015b; Potter et al., 2004b; Rago et al., 1993).

1.2.1.1 The case of Northern NEAC SU

ICES provides a shorter time series of data for Northern NEAC SU because some data are missing for Norway for the first time of the time series before 1982. The Norwegian data for the period 1971-1982 were completed using the following hypotheses (*Com pers.* Geir Bolstad and Peder Fiske, NINA):

- Homewater catches - Catch data for Norway (homewater catches, 1SW and 2SW separately) for the period 1971-1982 were extracted from the ICES WGNAS report of year 2002 (table 3.3.3.1f. Allocations of catches among the four regions of Norway was done using averages proportions calculated from the first five years for which data are available 1983-1987.
- Returns – The probability distribution of returns (1SW and 2SW, separately) was estimated by dividing the catches by guesstimates of exploitation rates and unreported catches for the period 1982-1971. Harvest rates and unreported catches were extrapolated backwards in time from year 1983. Uncertainty about those rates was bumped by 20% to account for the additional uncertainty due to extrapolation.

- Note that all MSW were considered as 2SW as for all other European SU.

The resulting probability distributions of returns are shown in Fig. 1. Numerical integration of uncertainty support the hypothesis that the returns are logNormaly distributed, allowing to approximate the likelihood for the returns as follows. For any year t and SU r , the expected mean of the distribution derived from the observations models for 1SW (respectively, 2SW) returns in log scale, denoted $\mathbb{E}_{\log(R_{1SW_{t,r}})}$ (resp. $\mathbb{E}_{\log(R_{2SW_{t,r}})}$), is considered as an observed realization of a Normal distribution of non-observed returns (in log-scale) $N_{6_{t,r}}$ (resp. $N_{9_{t,r}}$), with known variance $\sigma_{1SW_{t,r}}^2$ (resp. $\sigma_{2SW_{t,r}}^2$) set to the value derived from the observation errors models. These observation errors are considered independent across years, SU and sea-age classes.

$$(A1.16) \quad \mathbb{E}_{\log(R_{1SW_{t,r}})} \sim Normal(\log(N_{6_{t,r}}), \sigma_{1SW_{t,r}}^2)$$

$$(A1.17) \quad \mathbb{E}_{\log(R_{2SW_{t,r}})} \sim Normal(\log(N_{9_{t,r}}), \sigma_{2SW_{t,r}}^2)$$

1.2.2 Homewater catches

The homewater fisheries take adult fish that are mainly returning to the natal rivers to spawn. Point estimates of total catches reported by ICES (ICES 2018) pool all homewater fisheries capturing returning fish in coastal areas, estuaries and freshwater, for each SU, each year and each sea-age class separately. They are here denoted $\mathbb{E}_{\log(C_{HW,1SW_{t,r}})}$ and $\mathbb{E}_{\log(C_{HW,2SW_{t,r}})}$ for 1SW and 2SW fish, respectively. The likelihood term for homewater catches is built from logNormal observation errors with known observation error. Available knowledge support that homewater catches are known with only few errors. Relative error is then arbitrarily fixed to CV=0.05 for both sea-ages, for all years and all SU (but note this value can be changed to acknowledge for greater observation errors). Observation errors are considered independent across years, SU and sea-age classes. The likelihood terms associated with homewater catches are:

$$(A1.18) \quad \mathbb{E}_{\log(C_{HW,1SW_{t,r}})} \sim Normal(\log(h_{HWf,1SW_{t,r}} \times (1 - p_{delSp,1SW_{t,r}}) \times N_{6_{t,r}}), \sigma_{HW,1SW}^2)$$

$$(A1.19) \quad \mathbb{E}_{\log(C_{HW,2SW_{t,r}})} \sim Normal(\log(h_{HWf,2SW_{t,r}} \times (1 - p_{delSp,2SW_{t,r}}) \times N_{9_{t,r}}), \sigma_{HW,2SW}^2)$$

with $\sigma_{HW,1SW}^2 = \sigma_{HW,2SW}^2$ the variance corresponding to CV=0.05.

Observation model for the delayed catches are modelled using the same hypothesis and the same CV of observation errors.

1.2.3 Catches at sea for sequential distant marine fisheries operating on mixed stocks

For any marine fishery f operating on a mixture of SU, likelihood equations consist in logNormal observation errors on the total catches summed over all SU (still based on the same likelihood

approximation method), eventually supplemented by Dirichlet likelihood terms to adjust the proportion of catches allocated to each SU when proportion data are available (Table 7 and Fig. 2 - 6). Observation errors on the total catches and on the proportions are considered independent across fisheries, years and SU.

Observation models based on ICES (2018) data are built independently from the state-space model to estimate logNormal probability distributions of total catches at sea for each fishery f and each year t , with expected mean and variance (in log-scale) denoted $\mathbb{E}_{\log(C_f)}$ and $\sigma^2_{f_t}$, respectively. Variances $\sigma^2_{f_t}$ are derived by integrating uncertainty in the catch declaration rates, the proportions of fish of wild origin in the catches, and sampled biological characteristics of the catches including average weight of a fish used to convert catches in weights to number of fish, and scale samples used to separate the two sea-age classes in the catches. An exception is for the WG fishery for which observation errors are considered to be low (ICES 2005b) and fixed to $CV = 0.1$.

By denoting $C_{f_t} = \sum C_{f_t,r}$ the total catches from the state process summed over all SU, the likelihood term for the total catch is modelled as:

$$(A1.20) \quad \mathbb{E}_{\log(C_{f_t})} \sim Normal(\log(C_{f_t}), \sigma^2_{f_t})$$

Proportion of catches allocated to each SU are available for the West Greenland fishery (European and North American continental stock groupings) and for the Faroes fishery (1SWm and 1SWnm, and 2SW, for the European continental stock groupings only).

Proportions used to allocate West Greenland catches to each of the 25 SU in North America and Europe (Fig. 5) are derived from a compilation of individual assignment data from scale reading and genetic analyses. Proportions of the total catches at WG are first attributed to European and North American based on scales (1971-1999) and genetics samples (2000-2017) (ICES 2017a; ICES 2017b). Then, proportions attributed to each SU within the European stock group are fixed through time as compiled from ICES (2017b). Within the North American continental stock group, proportions are based on Bradbury et al. (2016a,2016b) that provide estimates of the proportion of fish originated from North American SU for 13 years based on genetic samples. The average value of the 13 years are used for the years without available data.

Proportions used to allocate Faroes catches to European SU are derived from a compilation of assignment data from scale reading (to separate fish from Southern and Northern Europe origin) and genetics data to allocate to each SU (ICES, 2018). Data are not informative enough to account for annual variability and those proportions are considered constant over the time series (Table 7, Fig. 2-4).

When available, observed proportion of each SU in the total catches, denoted $p_{f_t,r}^{obs}$ enters into a Dirichlet likelihood modelled as:

$$(A1.21) \quad (p_{f_t,r=1}^{obs}, \dots, p_{f_t,r=\Omega}^{obs}) \sim Dirichlet(\eta_{sample} \times (p_{f_t,r=1}, \dots, p_{f_t,r=\Omega}))$$

where $p_{f_t,r} = \frac{C_{f_t,r}}{C_{f_t}}$ is the proportion of fish from SU r in the total catches calculated from the state process.

When no proportions data are available, only the logNormal likelihood on total catches is used. The hypothesis of a homogeneous exploitation rate among SU replaces the Dirichlet likelihood. As a direct consequence, the proportions of any SU in the catches are set in pro-rata to the abundance among the SU just before the fishery.

Table 1. Summary of the main life stages and transitions of the life cycle model.

Stages	Transitions	Parameters	Parameters	Observation equations
N1: Eggs	$N7_t \rightarrow N1_t$ $N10_t \rightarrow N1_t$	Sex-ratio and fecundity	Fixed	No
N2: Total number of Smolt	$N1_t \rightarrow N2_t$	Freshwater survival (θ_1) - Average value - Lognormal noise	Fixed Fixed = 0.007 Fixed CV = 0,4	No
N3: Number of smolts in each age class (6 age classes)	$N3_{t+1+1}$ $N2_t \rightarrow N3_{t+1+a}$ $N3_{t+1+6}$	Proportion of smolt age (P_{smolt})	Fixed	No
N3tot: Total number of smolts migration year t	$N3_{tot_t}$			No
N4 : PFA (Pre Fishery Abundance)	$N3_{tot_t} \rightarrow N4_{t+1}$	Post-smolt survival (θ_3)	Estimated (Multivariate random walk with covariation among SU)	No
N5 : PFA maturing N8 : PFA non maturing	$N4_t \rightarrow N5_t$ $N4_t \rightarrow N8_t$	<i>Proportion maturing PFA</i> (θ_4)	Estimated (Multivariate random walk with covariation among SU)	No

N5.1 1SW maturing (1SW m) Faroes fishery	$N5_t \rightarrow N5.1_t$	Natural mortality (M) Harvest rates	Fixed Estimated (non informative prior)	Catches Faroes 1SW m observed with LogNormal errors and known variance
N8.1 : 1SW non maturing (1SW nm) Faroes fishery	$N8_t \rightarrow N8.1_t$	Natural mortality (M) Harvest rates	Fixed Estimated (non informative prior)	Catches Faroes 1SW nm observed with LogNormal errors and known variance
N8.2 : 2SW Faroes fisheries	$N8.1_t \rightarrow N8.2_{t+1}$	Natural mortality (M) Harvest rates	Fixed Estimated (non informative prior)	Catches Faroes 2SW observed with LogNormal errors and known variance
N6 : Returns 1SW N9 : Returns 2SW	$N5.1_t \rightarrow N6_t$ $N8.2_t \rightarrow N9_t$	Natural mortality (M)	Fixed Fixed	Returns 1SW and 2SW observed with LogNormal errors and known variance
N7 : Spawners 1SW N10 : Spawners 2SW	$N6_t \rightarrow N7_t$ $N9_t \rightarrow N10_t$	Harvest rates	Estimated (non informative prior)	1SW and 2SW homewater catches observed with LogNormal errors and fixed variance

Table 2. Parameters fixed or drawn in tight informative priors for the 25 stock units (Source: ICES 2018). (*) Number of eggs per fish. The number of eggs per fish includes the proportion of females in spawners. Fecundity can vary over time in the model, although these are fixed values in the data for most of the SU (except Scotland for which a time series is provided that accounts for the variation of the fecundity induced by the variation of the average length and weight of returning fish).

		NAC						S.NEAC							
		LB	NF	QB	GF	SF	US	FR	E&W	IR	NLFO	NLFB	SC.W	SC.E	IC.SW
Egg to smolts survival		$\theta_{1_{tr}} \sim \text{Lognormaly distributed with average value } E_1 = 0.007 \text{ and inter-annual variability } CV_1 = 0.4$													
	<i>psm</i> _{1,r}	0	0	0	0	0	0.377	0.917	0.23	0.05	0.38	0.38	0.2	0.05	0
	<i>psm</i> _{2,r}	0	0.041	0.058	0.398	0.6	0.52	0.083	0.75	0.75	0.59	0.59	0.5	0.45	0.05
Proportion of smolt ages	<i>psm</i> _{3,r}	0.077	0.598	0.464	0.573	0.394	0.103	0	0.02	0.2	0.03	0.03	0.3	0.45	0.73
	<i>psm</i> _{4,r}	0.542	0.324	0.378	0.029	0.006	0	0	0	0	0	0	0	0.05	0.21
	<i>psm</i> _{5,r}	0.341	0.038	0.089	0	0	0	0	0	0	0	0	0	0	0
	<i>psm</i> _{6,r}	0.04	0	0.01	0	0	0	0	0	0	0	0	0	0	0
Natural mortality rate (per month) after the PFA stage (for 1SW and 2SW fish)		$M = 0.03 \cdot \text{month}^{-1}$													
Migration duration between stages		See Table 5 and 6													
Number of eggs per fish (*)	<i>eggs</i> _{1,r}	1500	3000	468	547	917	200	1552	1350	2040	1972	1972	2000(*)	2000(*)	2501
	<i>eggs</i> _{2,r}	5500	4000	6402	5956	6107	5500	5520	4550	5950	4069	4069	6000(*)	6000(*)	6149

Table 2. (continuing)

		N.NEAC										
		IC.NE	SW	NO.SE	NO.SW	NO.MI	NO.NO	FI	RU.KB	RU.KW	RU.AK	RU.RP
Egg to smolts survival		$\theta_{1,r,t} \sim \text{Lognormally distributed with average value } \mathbb{E}_1 = 0.007 \text{ and interannual variability } CV_1 = 0.4$										
	<i>psm</i> _{1,r}	0	0.07	0	0	0	0	0	0	0	0	0
	<i>psm</i> _{2,r}	0.09	0.65	0.379	0.379	0.057	0.003	0	0.05	0.1	0.05	0
Proportion of smolt ages	<i>psm</i> _{3,r}	0.37	0.25	0.524	0.524	0.608	0.263	0.26	0.4	0.6	0.55	0.6
	<i>psm</i> _{4,r}	0.49	0.03	0.094	0.094	0.316	0.583	0.59	0.4	0.3	0.4	0.4
	<i>psm</i> _{5,r}	0.05	0	0.004	0.004	0.019	0.138	0.14	0.1	0	0	0
	<i>psm</i> _{6,r}	0	0	0	0	0	0.012	0.01	0.05	0	0	0
Natural mortality rate (per month) after the PFA stage (for 1SW and 2SW fish)		$M = 0.03 \cdot \text{month}^{-1}$										
Migration duration between stages		See Table 5 and 6										
Number of eggs per fish (*)	<i>eggs</i> _{1,r}	1974	1500	887	887	1050	450	600	350	2700	450	450
	<i>eggs</i> _{2,r}	7350	4200	4944	4944	5128	6673	10010	10000	4200	9600	10500

Table 3. Parameters of the marine phase drawn in non-informative prior and for which update from the data is expected. Note that harvest rates for the West Greenland and Faroes fishery are parameterized on the logit scale and written as the sum of a grand mean and yearly, region-specific and residual effects. Such a parameterization revealed needed to avoid miss fit of the proportions used to allocate the total catches to the different SU.

Non diagonal (plain) N×N variance-covariance matrix (N=25)	$\Sigma_8 = \begin{matrix} \sigma^2_{8_{1,1}} & \dots & \sigma^2_{8_{1,i}} \\ \dots & \dots & \dots \\ \sigma^2_{8_{i,1}} & \dots & \sigma^2_{8_{i,i}} \end{matrix}$
Note: Two different matrices for the post-smolt survival (Σ_{θ_3}) and for the proportion of fish maturing as 1SW (Σ_{θ_4})	$\Sigma_8^{-1} \sim \text{Wishart}(\Omega, \delta)$ with scale matrix Ω set as the N×N identity matrix and δ the degree of freedom set to N
Exploitation rate of West Greenland and Faroes mixed stock fisheries (for any year t and stock unit r)	$\begin{aligned} \text{logit}(h_{f,t,r}) &\sim \text{logit_h_mu}_f + \text{logit_h_time}_{f,t} + \text{logit_h_su}_{f,r} + \text{logit_h_res}_{f,t,r} \\ \text{logit_h_mu}_f &\sim N(-5,4) \\ \text{logit_h_time}_{f,t} &\sim N(0,4) \\ \text{logit_h_su}_{f,r} &\sim N(0,4) \\ \text{logit_h_res}_{f,t,r} &\sim N(0,4) \end{aligned}$
Exploitation rate of other fisheries f (NA marine fisheries and homewater fisheries) (for any year t and stock unit r)	$h_{f,t,r} \sim \text{Beta}(1,2)$

Table 5. Summary of the duration among stages and the sequential fisheries (operating on mixed stocks at sea and homewater fisheries) for stock units in the North American continental stock grouping (Source: ICES 2018; Prévost et al., 2009).


North American continental stock grouping		
Stages/Fisheries	Migration duration	Exploitation rate
PFA maturing		
↓	7 months	
1SWm NFDL/LB/SPM Fisheries		Variable among years (NFDL zone 3-7) Homogeneous among SU (NFDL zone 8-14, LAB, SPM) Specific estimate for SU = Labrador + homogeneous among all other SU
↓	1 month	
Returns 1SW		
↓	0	
1SW homewater Fishery		Variable among SU
↓	0	
Spawners 1SW		
PFA non maturing		
↓	7 months	
1SWnm NFDL/LB Fisheries		Variable among years Homogeneous among SU
↓	2 months	
1SWnm West Greenland Fishery		Variable among years and SU + data to allocate catches among SU. Allocation in two steps: 1) prop. to allocate fish from North America / Europe ; 2) prop. to allocate North American fish among SU in NA
↓	8 months	
2SWm NFDL/LB/SPM Fisheries		Variable among years (NFDL zone 3-7): Homogeneous among SU (NFDL zone 8-14, LAB, SPM): Specific estimate for SU = Labrador + homogeneous among all other SU
↓	1 month	
Returns 2SW		
↓	0	
2SW homewater Fishery		Variable among years and SU
↓	0	
Spawners 2SW		

Table 6. Summary of the duration among stages and the sequential fisheries (operating on mixed stocks at sea and homewater fisheries) for stock units in the Southern and Northern European continental stock groupings (Source: ICES 2018; Potter, 2016).


Southern and Northern Europe continental stock groupings		
Stages/Fisheries	Migration duration	Exploitation rate
PFA maturing		
↓	0.5 months	
1SWm Faroes Fishery		Variable among years and SU + data to allocate catches among SU
↓	7.5 months	
Returns 1SW		Variable among years and SU + data to allocate catches among SU
↓	0	
1SW homewater Fishery		Variable among years and SU
↓	0	
Spawners 1SW		
PFA non maturing		
↓	0.5 months	
1SWnm Faroes Fishery		Variable among years and SU + data to allocate catches among SU
↓	8.5 months	
1SWnm West Greenland Fishery		Variable among years and SU + data to allocate catches among SU. Allocation in two steps: 1) prop. to allocate fish from Europe / North America; 2) prop. to allocate European fish among SU in Europe
↓	5 months	
2SWm Faroes Fishery		Variable among years and SU + data to allocate catches among SU
↓	3.5 months	
Returns 2SW		
↓	0	
2SW homewater Fishery		Variable among years and SU
↓	0	
Spawners 2SW		

Table 7. Proportions to allocate the total catches among different SU from Southern and Northern Europe in the Faroes fishery. Proportions sum to 1 for each fishery and are considered constant over time (Source: ICES 2018). Fish originated from North America are not harvested in the Faroes fishery.

	S.NEAC								N.NEAC										
	FR	E&W	IR	NI.FO	NI.FB	SC.W	SC.E	IC.SW	IC.NE	SW	NO.SE	NO.SW	NO.MI	NO.NO	FI	RU.KB	RU.KW	RU.AK	RU.RP
1SW maturing	0.021	0.083	0.328	0.053	0.013	0.103	0.268	0.014	0.006	0.001	0.015	0.003	0.026	0.018	0.010	0.008	0.028	0.001	0.001
1SW non maturing	0.007	0.054	0.027	0.004	0.001	0.064	0.129	0.005	0.006	0.011	0.097	0.033	0.185	0.136	0.05	0.022	0.055	0.024	0.083
2SW	0.007	0.054	0.027	0.004	0.001	0.064	0.129	0.005	0.006	0.011	0.097	0.033	0.185	0.136	0.05	0.022	0.055	0.024	0.083




1
2



3
4 Figure 14: Probability distributions (median, quantiles 5% and 95%) of the number of fish returning as 1SW and 2SW for SU of North
5 America, Southern Europe and Northern Europe for PFA (blu) and LCM (red) (Source: ICES 2018).

6




7

8

9

10

Figure 2 : (a) Time series of total catches of the 1SW maturing stage in the Faroes fishery (Source: ICES 2015b); (b) proportions of the catches attributed to South European and North European stock units (Source: ICES 2015b). (Proportions attributed to SU from NA are 0).




11

12

13

14

Figure 3: (a) Time series of total catches of the 1SW maturing stage in the Faroes fishery (Source: ICES 2015b); (b) proportions of the catches attributed to South European and North European stock units (Source: ICES 2015b). (Proportion attributed to SU from NA are 0).



15


16

17

18

Figure 4: (a) Time series of total catches of the 2SW maturing stage in the Faroes fishery (Source: ICES 2015b); (b) proportions of the catches attributed to South European and North European stock units (Source: ICES 2015b). (Proportion attributed to SU from NA are 0).

19



20

21

22

23


24

25

26

Figure 5: (a) Time series of total catches of 1SW non-maturing in the West Greenland fishery (Source: ICES 2018); (b) Proportions of the catches attributed to North American / European stock complex (top), attributed to North American stock units within North American stock complex (bottom left) and attributed to European stock units within European stock complex (bottom right). Genetics data (Bradbury et al (2016)) provide estimates of the proportion of fish originated from the 7 SU of North America (Olmos et al 2019). Total catches and proportions were updated from Olmos et al. (2019) to account for fish of NE SU.

27



28

29

30


31

32

33

Figure 6. Time series of point estimates (median of logNormal distributions) of catches for the sequential fisheries at sea occurring on mixed stocks of North American 1SW maturing fish (Source: ICES 2018). (a) catches of all SU in north-eastern Newfoundland (Salmon Fishing Areas 3 to 7); (b) Labrador origin catches in Labrador Fisheries; (c) catches of SU 1-5 in Labrador fisheries, South-west Newfoundland Fisheries (Salmon Fishing Areas 8 to 14A) and in the Saint Pierre and Miquelon fisheries.

34



35

36

37 Figure 7. (Continuing). Time series of distribution (median and quantiles 5% and 95% of logNormal distributions) of catches for the sequential
 38 fisheries at sea occurring on mixed stock fisheries, on North American 1SW non-maturing fish (Source: ICES 2018). (a) 1SW catches for all SUs in
 39 north and eastern Newfoundland (Salmon Fishing Areas 3 to 7); (b) 2SW catches for all SUs in north and eastern Newfoundland (Salmon Fishing
 40 Areas 3 to 7); (c) 2SW Labrador (SU 6) origin catches in the Labrador fisheries; (d) 2SW catches of SU 1-5, in Labrador fisheries, south and west
 41 Newfoundland Fisheries (Salmon Fishing Areas 8 to 14A) and in the Saint Pierre and Miquelon fisheries.


42

43 **7 SUPP.MAT.II : Comparison between the PFA and life cycle modelling framework**

44

45

46



47

48

49

Figure SII.15: Structure of the stage-based life cycle model (red) and the PFA models (green). Run reconstructions models (green) are observation models generating input data for both PFA and life cycle models

50

51 Table SII.2: Comparison of the model resolution, the likelihood, the hypothesis of the parameters to estimate, and the information priors used in PFA and life cycle model

		PFA NA FORECAST	PFA Europe FORECAST		LIFE CYCLE	
			Southern Europe	Northern Europe		
Model Resolution	Temporal resolution [t]	1978-2017	1978-2017	1991-2017	1971-2017	
	Spatial resolution [r]	6 SUs	6 Regions: Fr, E&W, SC, IR, IC.SW, N.IR	5 Regions: IC.NE, SW, NO,RU,FI	25 SUs	
	Life History at sea [a]	1SW	1SW & 2SW	1SW & 2SW	1SW & 2SW	
Likelihood	Catches at sea	Abundances of Returns [t,r,a]	LogNormal Distribution	LogNormal Distribution	LogNormal Distribution	LogNormal Distribution
		HomeWater Catches (and delayed Spawners) [t,r,a]	-	-	-	LogNormal Distribution
		Catches in the Faroes fisheries (1SWm, 1SWnm, 2SW) [t,a]	-	No Likelihood Prior Distributions	No Likelihood Prior Distributions	LogNormal Distribution
		Prop. of catches attributed to SUs in the Faroes fisheries (1SWm, 1SWnm, 2SW) [t,r,a]	-	Fixed	Fixed	Dirichlet Distribution
		Catches in the LAB/NFLD/SPM fisheries (1SWm, 1SWnm, 2SW) [t,a]	No Likelihood Prior Distributions	-	-	LogNormal Distribution
		Prop. of catches attributed to SUs in the LAB/NFLD/SPM fisheries (1SWm, 1SWnm, 2SW)	Homogeneous exploitation rates (no data, except LB) based on proportion of regional abundance	-	-	Homogeneous exploitation rates (no data)
		Catches in the West Greenland fisheries [t,a]	No Likelihood Prior Distributions	No Likelihood Prior Distributions	No Likelihood Prior Distributions	Dirichlet Distribution
		Prop. of catches attributed to SUs in the 1SWm West Greenland fisheries [t,r,a]	Homogeneous exploitation rates (no data) based on proportion of regional abundance	Fixed and stationnary	Fixed and stationnary	SU-specific genetic information data to allocate catches accross year Dirichlet Distribution

52

53 Table SII.3 (continued): Comparison of the model resolution, the likelihood, the hypothesis of the parameters to estimate, and the information priors used in PFA and life cycle model

		PFA NA FORECAST	PFA Europe FORECAST		LIFE CYCLE
			Southern Europe	Northern Europe	
Estimated parameters	Productivity[t,r]	Multivariate random walks in the log scale, with covariations among all stock units	Random walks in the log scale. No covariance structure among SUs	Random walks in the log scale. No covariance structure among Sus	Multivariate random walks in the log scale. Random variations are drawn from multivariate normal distributions with variance-covariance
	Proportion of fish maturing as 1SW [t,r]	-	Random walks in the logit scale. No covariance structure among SUs	Random walks in the logit scale. No covariance structure among Sus	Multivariate random walks in the log scale. Random variations are drawn from multivariate normal distributions with variance-covariance
	Fisheries exploitation rates[t,r,a]	derived	derived	derived	Estimated with a prior distribution
	Pre-Fishery abundance[t,r]	Random effect	derived	derived	derived
Variables and Parameters fixed or drawn in tight informative priors	Number of eggs per fish	Fixed	Fixed	Fixed	Fixed
	Proportion of smolt ages	-	-	-	Informative prior
	Natural mortality rate (per month) after the PFA stage (for 1SW and 2SW fish)	Fixed	Fixed	Fixed	Informative prior
	Migration duration between stages	Fixed	Fixed	Fixed	Fixed
	Lagged Eggs/Spawners	Normal distribution	Normal distribution	Normal distribution	<i>Information from Lagged Eggs/Spawners is accounted by the likelihood of homewater catches</i>

54


55

56 Table SII.4 : Comparison between the hypothesis used to forecast and carry out the risk analysis using PFA and life cycle models

	PFA NA FORECAST	PFA Europe FORECAST		LIFE CYCLE
		Southern Europe	Northern Europe	
Risk Analysis	Probability of meeting region specific 2SW eggs deposition CLs	Probability of meeting region specific 1SW and 2SW eggs deposition CLs	Probability of meeting region specific 1SW and 2SW eggs deposition CLs	Probability of meeting region specific 1SW and 2SW eggs deposition CLs
Forecasted years	2018-2020	2018-2020	2018-2020	2018-2020
Fishing Scenarios in Faroes Fisheries	-	TAC = 0, 500, 1000, 1500, 2000, 2500	TAC = 0, 500, 1000, 1500, 2000, 2500	TAC = 0, 500, 1000, 1500, 2000, 2500
Fishing Scenarios in Faroes Fisheries	TAC = 0, 500, 1000, 1500, 2000, 2500	TAC = 0, 500, 1000, 1500, 2000, 2500	TAC = 0, 500, 1000, 1500, 2000, 2500	TAC = 0, 500, 1000, 1500, 2000, 2500
Fisheries Catches at sea and proportion to allocate catches to Sus for years 2018-2020	Fixed to the average of the last five years of the time series of data (2013-2017)	Fixed to the average of the last five years of the time series of data (2013-2017)	Fixed to the average of the last five years of the time series of data (2013-2017)	Fixed to the average of the last five years of the time series of data (2013-2017)
Homewater catches	-	-	-	Fixed to the average of the last five years of the time series of data (2013-2017)

57

58



59

60


Figure SII.16 : Comparison of the West Greenland catch allocation assumptions between the PFA and the life cycle model. For the life cycle model see data in Supp. Mat. I, Fig.2 – Fig7.

61 **8 Supp.Mat.III: RESULTS**

62

63

— North America — Northern Europe — Southern Europe




64

Figure SIII.17: Time series of estimated abundances aggregated per CSG at four stages in the life cycle. (a) total returns to homewater (1SW + 2SW); (b) proportion of 1SW in returns; (c) total spawners (1SW + 2SW); (d) proportion of 1SW in spawners; (e) total egg deposition by spawners; (f) proportion of eggs spawned by 1SW. Trend lines are medians of marginal posterior distributions. Abundances are standardized to the first year values.

66

67


68



70

71 Figure SIII.18 UK England and Whales - Probability distributions of (a) the number of egg potentially spawned, (b) PFA (total PFA.m + PFA.nm), (c)
 72 PFA.m, (d) PFA.nm, (e) Productivity, (f) Proportion of fish maturing as 1SW, (g) Returns 1SW, and Returns (2SW). Thick point is the median and line
 73 represent the 95% posterior credibility interval. Blue: PFA model; Red : Life cycle model (LCM); Dark color: historical time series; Light colors:
 74 Forecasted years (under the scenario of no catches at West Greenland and Faroes)

75



76

77

78


79

80

81

82

Figure SIII.19 Finland - Probability distributions of (a) the number of egg potentially spawned, (b) PFA (total PFA.m + PFA.nm), (c) PFA.m, (d) PFA.nm, (e) Productivity, (f) Proportion of fish maturing as 1SW, (g) Returns 1SW, and Returns (2SW). Thick point is the median and line represent the 95% posterior credibility interval. Blue: PFA model; Red : Life cycle model (LCM); Dark color: historical time series; Light colors: Forecasted years (under the scenario of no catches at West Greenland and Faroes)



83

84

85


86

87

88


89

Figure SIII.20 France - Probability distributions of (a) the number of egg potentially spawned, (b) PFA (total PFA.m + PFA.nm), (c) PFA.m, (d) PFA.nm, (e) Productivity, (f) Proportion of fish maturing as 1SW, (g) Returns 1SW, and Returns (2SW). Thick point is the median and line represent the 95% posterior credibility interval. Blue: PFA model; Red : Life cycle model (LCM); Dark color: historical time series; Light colors: Forecasted years (under the scenario of no catches at West Greenland and Faroes)



90
91
92
93
94
95

Figure SIII.21 Gulf Region - Probability distributions of 2SW Productivity, the number of egg potentially spawned by 2SW fish, 2SW Returns, and PFA.nm. Thick point is the median and line represent the 95% posterior credibility interval. Blue: PFA model; Red: Life cycle model (LCM); Dark color: historical time series; Light colors: Forecasted years (under the scenario of no catches at West Greenland)



96
 97 Figure SIII.22 Iceland North-east - Probability distributions of (a) the number of egg potentially spawned, (b) PFA (total PFA.m + PFA.nm), (c) PFA.m,
 98 (d) PFA.nm, (e) Productivity, (f) Proportion of fish maturing as 1SW, (g) Returns 1SW, and Returns (2SW). Thick point is the median and line
 99 represent the 95% posterior credibility interval. Blue: PFA model; Red : Life cycle model (LCM); Dark color: historical time series; Light colors:
 100 Forecasted years (under the scenario of no catches at West Greenland and Faroes)

96

97

98

99

100

101

102





Figure SIII.23 Iceland South West - Probability distributions of (a) the number of egg potentially spawned, (b) PFA (total PFA.m + PFA.nm), (c) PFA.m, (d) PFA.nm, (e) Productivity, (f) Proportion of fish maturing as 1SW, (g) Returns 1SW, and Returns (2SW). Thick point is the median and line represent the 95% posterior credibility interval. Blue: PFA model; Red : Life cycle model (LCM); Dark color: historical time series; Light colors: Forecasted years (under the scenario of no catches at West Greenland and Faroes)



110

111

112


113

114

115


116

Figure SIII.24 Ireland - Probability distributions of (a) the number of egg potentially spawned, (b) PFA (total PFA.m + PFA.nm), (c) PFA.m, (d) PFA.nm, (e) Productivity, (f) Proportion of fish maturing as 1SW, (g) Returns 1SW, (h) and Returns (2SW). Thick point is the median and line represent the 95% posterior credibility interval. Blue: PFA model; Red : Life cycle model (LCM); Dark color: historical time series; Light colors: Forecasted years (under the scenario of no catches at West Greenland and Faroes)



117
118
119
120
121
122

Figure SIII.25 Labrador - Probability distributions of (a) the number of egg potentially spawned by 2SW fish 2SW Returns, (b) and PFA.nm (c) productivity of 2SW fish, (d) Returns 2SW. Thick point is the median and line represent the 95% posterior credibility interval. Blue: PFA model; Red : Life cycle model (LCM); Dark color: historical time series; Light colors: Forecasted years (under the scenario of no catches at West Greenland)



123

124

125


126

127

128

129

Figure SIII.26 UK Northern Ireland - Probability distributions of (a) the number of egg potentially spawned, (b) PFA (total PFA.m + PFA.nm), (c) PFA.m, (d) PFA.nm, (e) Productivity, (f) Proportion of fish maturing as 1SW, (g) Returns 1SW, and Returns (2SW). Thick point is the median and line represent the 95% posterior credibility interval. Blue: PFA model; Red : Life cycle model (LCM); Dark color: historical time series; Light colors: Forecasted years (under the scenario of no catches at West Greenland and Faroes)



130

131

132

133

134

135

Figure SIII.27 Newfoundland - Probability distributions of (a) the number of egg potentially spawned by 2SW fish 2SW Returns, (b) and PFA.nm (c) productivity of 2SW fish, (d) Returns 2SW. Thick point is the median and line represent the 95% posterior credibility interval. Blue: PFA model; Red : Life cycle model (LCM); Dark color: historical time series; Light colors: Forecasted years (under the scenario of no catches at West Greenland)




Figure SIII.28 Norway - Probability distributions of (a) the number of egg potentially spawned, (b) PFA (total PFA.m + PFA.nm), (c) PFA.m, (d) PFA.nm, (e) Productivity, (f) Proportion of fish maturing as 1SW, (g) Returns 1SW, and Returns (2SW). Thick point is the median and line represent the 95% posterior credibility interval. Blue: PFA model; Red : Life cycle model (LCM); Dark color: historical time series; Light colors: Forecasted years (under the scenario of no catches at West Greenland and Faroes)

136

137


138

139

140

141

142



143

144


145

146

147

148

Figure SIII.29 Quebec - Probability distributions of (a) the number of egg potentially spawned by 2SW fish 2SW Returns, (b) and PFA.nm (c) productivity of 2SW fish, (d) Returns 2SW. Thick point is the median and line represent the 95% posterior credibility interval. Blue: PFA model; Red : Life cycle model (LCM); Dark color: historical time series; Light colors: Forecasted years (under the scenario of no catches at West Greenland)



149

150

151

152

153

154

155

Figure SIII.30 Russia - Probability distributions of (a) the number of egg potentially spawned, (b) PFA (total PFA.m + PFA.nm), (c) PFA.m, (d) PFA.nm, (e) Productivity, (f) Proportion of fish maturing as 1SW, (g) Returns 1SW, and Returns (2SW). Thick point is the median and line represent the 95% posterior credibility interval. Blue: PFA model; Red : Life cycle model (LCM); Dark color: historical time series; Light colors: Forecasted years (under the scenario of no catches at West Greenland and Faroes)






Figure SIII.31 Scotland - Probability distributions of (a) the number of egg potentially spawned, (b) PFA (total PFA.m + PFA.nm), (c) PFA.m, (d) PFA.nm, (e) Productivity, (f) Proportion of fish maturing as 1SW, (g) Returns 1SW, and Returns (2SW). Thick point is the median and line represent the 95% posterior credibility interval. Blue: PFA model; Red : Life cycle model (LCM); Dark color: historical time series; Light colors: Forecasted years (under the scenario of no catches at West Greenland and Faroes)



163
164
165
166
167
168

Figure SIII.32 Scotia-Fundy - Probability distributions of 2SW Productivity, the number of egg potentially spawned by 2SW fish, 2SW Returns, and PFA.nm. Thick point is the median and line represent the 95% posterior credibility interval. Blue: PFA model; Red: Life cycle model (LCM); Dark color: historical time series; Light colors: Forecasted years (under the scenario of no catches at West Greenland)



169

170

171


172

173

174

175

Figure SIII.33 Sweden - Probability distributions of (a) the number of egg potentially spawned, (b) PFA (total PFA.m + PFA.nm), (c) PFA.m, (d) PFA.nm, (e) Productivity, (f) Proportion of fish maturing as 1SW, (g) Returns 1SW, and Returns (2SW). Thick point is the median and line represent the 95% posterior credibility interval. Blue: PFA model; Red : Life cycle model (LCM); Dark color: historical time series; Light colors: Forecasted years (under the scenario of no catches at West Greenland and Faroes)



176

177

178

179

Figure SIII.34: USA - Probability distributions of 2SW Productivity, the number of egg potentially spawned by 2SW fish, 2SW Returns, and PFA.nm. Thick point is the median and line represent the 95% posterior credibility interval. Blue: PFA model; Red: Life cycle model (LCM); Dark color: historical time series; Light colors: Forecasted years (under the scenario of no catches at West Greenland)

180