

Best practices for consistent and reliable life cycle assessments of urban agriculture

Erica Dorr, Benjamin Goldstein, Christine Aubry, Benoit Gabrielle, Arpad Horvath

▶ To cite this version:

Erica Dorr, Benjamin Goldstein, Christine Aubry, Benoit Gabrielle, Arpad Horvath. Best practices for consistent and reliable life cycle assessments of urban agriculture. Journal of Cleaner Production, 2023, 419, pp.138010. 10.1016/j.jclepro.2023.138010. hal-04341695

HAL Id: hal-04341695 https://hal.inrae.fr/hal-04341695

Submitted on 20 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 How to do life cycle assessments of urban agriculture

- 2 Erica Dorr^{1,2}, Benjamin Goldstein³, Christine Aubry¹, Benoit Gabrielle^{2,*}, Arpad Horvath⁴
- 3
- ⁴ Université Paris-Saclay, INRAE, AgroParisTech, UMR SAD-APT, Palaiseau, France
- ⁵ Université Paris-Saclay, INRAE, AgroParisTech, UMR ECOSYS, Palaiseau, France
- 6 ³McGill University, Faculty of Agricultural and Environmental Sciences, Canada
- ⁴ Department of Civil and Environmental Engineering, University of California, Berkeley,
- 8 USA
- 9 *Corresponding author

10

- 11 Highlights
- Life cycle assessment is inconsistently applied to urban agriculture
- Identified key challenges of doing life cycle assessment of this unique activity
- Made practical recommendations for how to address these challenges
- Outlined research directions and scientific questions for this maturing topic
- Following recommendations will strengthen and clarify body of research

17

- 18 Submitted manuscript
- 19 Final version (still under embargo) appeared in the Journal of Cleaner Production
- 20 <u>https://dx.doi.org/10.1016/j.jclepro.2023.138010</u>

- 20 Abstract
- 21 There is increasing interest in evaluating the environmental performance of urban agriculture
- 22 (UA), especially using life cycle assessment (LCA). However, LCA has been applied to UA
- 23 inconsistently, making it difficult to confidently compare or draw conclusions from existing
- 24 studies. Here, we outline the key challenges of applying LCA to UA and recommend concrete
- 25 steps to help bring consistency and comprehensiveness to the topic. First, we clarify the
- 26 research questions that can be addressed with LCA. We then provide practical
- 27 recommendations for performing LCAs of UA, considering several of its unique aspects that
- 28 require special attention by practitioners. These include crop diversity, data availability,
- 29 modeling compost, soil carbon sequestration, producing growing media, distribution of crops,
- and variability and uncertainty. Next, we propose future research areas that will benefit LCA
- 31 generally and its application to UA, such as framing UA as urban green infrastructure,
- 32 evaluations at the city-scale, accounting for ecosystem services, and including social
- 33 dimensions of UA. By following these recommendations, future LCAs of UA can be more
- 34 consistent, comparable, and holistic, and will help build knowledge and inform policy making
- 35 and practices around UA.
- 36 Keywords: LCA guidelines, farms, gardens, social LCA, ecosystem services, compost

37

1 Introduction

38

Urban agriculture (UA) is a multifunctional activity with many assumed and demonstrated 39 benefits for cities and their inhabitants. These social, economic, and environmental benefits 40 position UA as a powerful tool to improve urban environments, contribute to more sustainable 41 42 of urban food systems, and enhance wellbeing of urban dwellers (Azunre et al., 2019; Gómez-Villarino et al., 2021). Food grown in cities can have lower environmental burdens than food 43 from conventional farms for a variety of indicators, including site-specific pollution to diffuse 44 greenhouse gas (GHG) emissions (Nicholls et al., 2020; O'Sullivan et al., 2019). While 45 pollution at farms can be measured on site, environmental footprinting methods, such as life 46 cycle assessment (LCA), are needed to capture impacts across the food value chain. Although 47 the LCA method is standardized, findings from available LCAs of UA are highly variable 48 because of inconsistencies in how the method has been employed (Dorr et al., 2021a). We 49 50 lack reliable answers to important questions surrounding the environmental performance of 51 UA. What types of UA have lower impacts than others? What are the main sources of impacts in UA? Can UA help reduce the environmental impacts of the food system? Researchers 52 53 require guidance to more consistently make decisions regarding system modeling, system 54 boundaries, and reporting so that LCAs of UA can help answer these and other questions. 55 General LCA frameworks and guides have been proposed to improve the rigor and 56 comparability of LCAs, and include the International Organization for Standardization (ISO) 57 framework (ISO, 2006a, 2006b), the ILCD handbook (European Commission, 2010a), and the 58 Product Environmental Footprint Category Rules Guidance (European Commission, 2017). For LCA of specific sectors, methodological guidelines have highlighted unique aspects that 59 require special attention. Failure to account for these aspects can skew results and hamper 60 decision making. For instance, the inclusion of direct and indirect land use change in biofuel 61 production fundamentally altered the carbon calculus of this technology, and caused a 62 reappraisal of government policies to support first-generation biofuels (Searchinger et al., 63 64 2008).

65 To avoid similar mistakes in other fields, researchers have produced LCA guidelines for diverse industries and technologies ranging from waste management (Laurent et al., 2014) to 66 67 bioplastics (Bishop et al., 2021). In the area of food, best practices exist for LCAs of crop production (Adewale et al., 2018), organic agriculture (van der Werf et al., 2020), fruit 68 orchards (Cerutti et al., 2014), vegetables (Perrin et al., 2014), climate-smart agriculture 69 (Acosta-Alba et al., 2019) and agricultural use of microbial inoculants (Kløverpris et al., 70 2020). Other work has evaluated the combination of agricultural LCAs with circular economy 71 72 (Stillitano et al., 2021) or ecosystem service assessments (Tang et al., 2018). Not to mention 73 the large body of work reviewing the methodological choices, challenges, and best practices of agricultural LCAs in general (Audsley et al., 1997; Brentrup et al., 2004; Caffrey and Veal, 74 2013; Cucurachi et al., 2019; Dijkman et al., 2018; Mclaren, 2010; Nemecek and Gaillard, 75 76 2010; Notarnicola et al., 2017, 2012). Such methodological reflections and recommendations have not yet been done for UA. 77 78 This study intends to fill this gap by providing a guideline for how to assess UA using LCA. It is applicable to all forms of UA in its most general definition of "food production in and 79 around cities" (Mougeot, 2000). It builds on observations from a previous literature review 80 81 and meta-analysis of the environmental impacts of UA (Dorr et al., 2021a) to provide 82 practical recommendations when applying LCA to UA, and takes a more comprehensive 83 approach to both UA and LCA. This guideline was also tested and iteratively refined during a 84 recent LCA of a diverse set of urban farms in France and the United States (Dorr et al., 2022b). 85 This paper begins by reflecting on the goals and expectations of LCAs of UA, followed by 86 practical recommendations to make LCAs of UA more consistent, and research directions to 87 improve LCAs of UA. In doing so, this paper identifies the challenges of including certain 88 aspects of UA in LCA, reviews how these aspects are currently treated in LCAs of UA, and 89 recommendations for how to best treat them going forward. This guideline is intended to 90

complement existing frameworks for agricultural LCAs, and some issues relevant to both

91

LCA of conventional agriculture and UA were included here. Our hope is that by outlining clear rules for dealing with the unique challenges of applying LCA to UA, future work can be done in an a consistent, transparent, and comprehensive manner. Such consistency is needed to determine under what conditions and in what forms UA can meaningfully contribute to urban sustainability.

2 Why do LCAs of urban agriculture?

Since there are diverse framings of UA, it is useful to clarify why we should study it with LCA, by defining both the goals and the larger questions they aim to answer. Reflecting on these questions is especially timely as UA LCAs evolve from an early stage with relatively simple goals of assessing impacts of a farm or garden, to a more mature stage assessing more complex topics. The goal of an LCA dictates how the assessment is set up. All decisions regarding system boundaries, functional unit (FU), and interpretations should be consistent with the defined goal(s) of the study, which should reflect the pursuit of an overarching question(s). Table 1 highlights some key, largely unanswered questions around UA that LCA can address. Goals of existing UA LCAs include evaluating the environmental impacts of urban food production at the farm-scale, identifying ways to reduce these impacts, comparing UA to rural agriculture or to other urban land uses, comparing types of UA, and evaluating the consequences of developing UA (such as reduced lawn management, or municipal treatment of organic waste) ((Dorr et al., 2021a). A more detailed review of UA LCAs that addressed each question, with goal, scope, and FU recommendations, is in Appendix A.

Table 1 The goal of a life cycle assessment should answer a larger question. Some relevant questions for life cycle assessment of urban agriculture are presented here, along with a description/justification for each question and possible functional units (FU). Some questions are already prevalent in the literature, and some are our original suggestions and have not been addressed before.

Question	Description	FU
Is UA an environmentally positive way to feed the city, relative to the status quo of conventional food systems?	In light of new urban food planning strategies, and initiatives to reduce impacts of public food procurement, we should investigate if UA is a useful strategy.	Single crop, mixed crops, cost/revenue, individual diet, citywide food flows
Is UA an environmentally positive type of green	Green infrastructure is promoted in cities, and many types are possible.	Land area, cost/revenue

infrastructure to implement in a city?
How does UA affect the

How does UA affect the GHG emission or other environmental impacts of a city?

What are potential trade-offs of socially motivated UA projects?

Which type of UA should be developed/promoted for a given motivation (indoor or outdoor, hydroponics or soil-based, commercial or non-profit, professional or volunteer-based...)?

How can UA be designed or managed to minimize environmental impacts?

116

117

118

119

120

121

122

123

124

125

City leaders must decide which types to implement.

Cities have pledged to reduce GHG emissions, which UA may address through land use, replacing other food sources, changing consumers' behaviors, or altering organic waste treatment.

Many UA projects do not claim to have environmental motivations or particularly low impacts, but they are promoted based on other merits (often social). Are there important tradeoffs between the social and environmental dimensions? Can we justify an environmentally harmful activity if it delivers social benefits?

Developers, city leaders, and stakeholders may have land that they want to dedicate to UA. With the vast diversity of types of UA, they may need support deciding which type to develop, according to environmental and other dimensions.

In many cases, UA will be practiced regardless of the above questions. Then, we should inform practitioners of the best practices to minimize their impact.

Urban metabolism, land area, operation of other sectors (i.e. waste treatment)

Single crop, mixed crops, land area, cost/revenue, total operations of urban farm, social functions (i.e. hours of education, number of participants)

Single crop, mixed crops, land area, cost/revenue, total operations of urban farm, social functions (i.e. hours of education, number of participants)
Any

3 Challenges and practical recommendations for UA LCAs

Below, we describe unique aspects of UA that present methodological challenges for LCAs, and our recommendations for addressing them. Each section includes an explanation of the challenge, examples of how it has been treated in previous urban or rural agriculture LCAs, and recommendations for future work. Section 3.3, on compost, includes additional subsections because there are numerous challenges, and to the best of our knowledge its inclusion in agricultural LCAs has not been reviewed before. A summary of key recommendations is provided in Table 2, which draws from both the practical recommendations here and the research directions presented in section 4.

126 3.1 Crop diversity

127 Challenge:

Mass-based FUs are most common in crop production LCAs (Notarnicola et al., 2017). For monoculture farms, there are no allocation issues: all inputs and impacts are assigned to one crop. For farms growing multiple crops either with temporal diversity (crop rotation) or spatial diversity (polyculture/intercropping), allocation between crops is needed (Adewale et al., 2018). For polycultures, rural/professional farmers can often specify which inputs were used on various farm parcels, and fixed inputs can be allocated by mass, revenue or other measure (Caffrey and Veal, 2013). For crop rotations, allocation principals have been proposed (Brankatschk and Finkbeiner, 2015). Such allocation is difficult for UA, where crop diversity is often exceedingly high: urban farms may cultivate on average 20-30 crops per year, with extremes of 80-130 (Gregory et al., 2016; Kirkpatrick and Davison, 2018; Pourias et al., 2016). It is unreasonable to expect urban farmers to distinguish inputs for so many crops, so LCA practitioners often contend with the challenge of including many crops in one FU. This issue is not unique to UA—it is also relevant for diversified rural farms and community-supported agriculture (CSA) (Caffrey and Veal, 2013; Christensen et al., 2018)—but is more pronounced with UA.

143 Examples:

LCA stage	No.	Recommendation				
Goal and scope	1	Be transparent, thorough, and critical when evaluating compost, substrate, and other organic inputs. They are especially important for UA, and are not usually the focus in agricultural LCAs.				
	2	Use multiple FUs—at least land and product-based. Include post-farm transport of products—especially the (near) zero impacts of transport by bike or on foot.				
	3					
	4	Account for seasonality, local context, and (where relevant) last mile transport for more precise comparisons to rural agriculture.				
Life cycle inventory	5	Collect primary data from functioning urban farms, because UA may not operate as expected or as measured under ideal, controlled conditions.				
Life cycle impact assessment	6	Use sensitivity analyses for important parameters with high uncertainty or variability to obtain a range or distribution of results. Such parameters may be related to:				
		Infrastructure lifetime				
		 Substrate lifetime 				
		 Compost emission factors 				
		 Delivery logistics 				
	7	Present results with and without major avoided burdens and carbon sequestration benefits.				
Interpretation	8	Provide more holistic descriptions of UA case studies and their urban contexts, because UA is diverse and vaguely defined. This includes the motivations, management/farming structures, or innovative status of a case study.				
	9	Compare impacts with an area-based FU to other urban green infrastructure or urban land use options.				
	10	Include social, economic, and ecosystem service-related measures, even if they not life-cycle based.				

Table 2 Ten key recommendations for performing UA LCAs are summarized according to their position along the 4-step LCA process.

Most UA LCAs with high crop diversity chose FUs covering total annual operations of a farm or impacts per unit area (Martinez et al., 2018; Pérez-Neira and Grollmus-Venegas, 2018; Sanyé-Mengual et al., 2018a). This avoids highly uncertain allocations, considers additional functions of agriculture, and facilitates cross-farm comparisons. However, results are difficult to extrapolate since they represent production of varied crops which are usually not functionally comparable, and sometimes the crops grown are not communicated. Another strategy uses published data or farmer estimates to estimate an life cycle inventory for each

crop (Caputo et al., 2020; Kulak et al., 2013; Liang et al., 2019). This allows for crop-level analysis for polycultures, but accuracy is inevitably lost when equating UA to other systems. For instance, when these data come from rural agriculture, representativeness of UA is likely sacrificed. Other researchers have allocated between many crops based on mass, area, calorie content, nutritional index, or time of cultivation of each crop, to generate results per crop (Pennisi et al., 2019; Rufí-Salís et al., 2020a; Sanyé-Mengual et al., 2015b). Finally, some researchers used a simplified FU covering a basket of crops (i.e. 1 kilogram of mixed lettuce, tomato, and pepper) (Boneta et al., 2019; Hu et al., 2019). These results are difficult to use elsewhere since unique mixes of crops are not precisely comparable, and authors may not include which crops are included in the mix or in what proportions. LCAs of rural farms with many crops have also used a FU of kilogram of mixed crop (Christensen et al., 2018; Pepin, 2022), which complicates interpretation.

Recommendations:

The main options for dealing with multi-crop UA systems are to evaluate a basket of products (by mass or by converting to calories or nutritional indexes), allocate between products, or choose a FU that is not based on food production. It is impossible to universally recommend a FU for LCAs of such diverse systems aiming to answer different questions, and ultimately the choice of FU depends on the goal of the LCA, but we can recommend some best practices. When a FU other than single crop is used, a breakdown of how much of each crop was grown should be provided, to give some indication of what the food outputs of the system were. Ideally UA LCAs should aim for crop-specific inventories within urban farms, to allow for a FU of production of a single crop, but due to high crop biodiversity this may not be feasible. Finally, providing results across multiple FUs can illuminate tradeoffs and compensate for the opaque nature of mixed-product FUs such as mass of mixed crops.

3.2 Data availability

Challenge:

Data collection in LCA is often highly labor-intensive. For an agricultural LCA, data on farm inputs and outputs are needed. In conventional agriculture, primary data come from farmer interviews, receipts, or informed estimates/calculations (Christensen et al., 2018). Secondary data, such as the UC Davis Crop Budgets (Caffrey and Veal, 2013), can address data gaps or create entire inventories. Similar quality data are rare for UA because urban farmers usually keep limited records (Cleveland, 1997; Egerer et al., 2018; Whittinghill and Sarr, 2021). Inputs and food production in UA (especially informal UA) can be extremely variable and difficult to predict, casting doubt on the applicability of secondary data for UA (Dorr et al., 2022c). Collective and community-based UA may have many participants who harvest and use agricultural inputs, which further complicates record keeping. Self-reporting and participatory methods face issues of reliable, consistent data collection and participant fatigue (CoDyre et al., 2015).

Examples:

The available UA LCAs are based on both primary and secondary data. Data for UA LCAs come from many different sources, including directly measured data, operations records, farmer interviews and surveys, and secondary data from urban or rural agriculture (Dorr et al., 2021a). Data sources and data collection difficulties are largely discussed in research on UA practices in general, but not so much in UA LCAs (McDougall et al., 2019; Pollard et al., 2018).

Recommendations:

Due to the variability and lack of data regarding UA practices, collecting primary data from case studies should be prioritized. Past records of operation may be used, although it is unlikely that urban farmers have records of all necessary information for an LCA. A data collection campaign, with commitment from farmers, may be necessary. Researchers should discuss data needs with farmers early and often to identify the most feasible methods to collect data, create a data collection plan, and regularly follow up to ensure reliability. This is a crucial step because unclear or overly burdensome data collection efforts may be abandoned

or unusable. Researchers should consider the types of data that may already be collected at urban farms (i.e. level of detail, time frame, units), and adapt the data collection plan accordingly. Surveys, growing logs, and harvest notebooks should be co-designed with farmers to track harvest and inputs (Nicholls et al., 2020). Water use should be measured using water meters or calculated using the number of buckets or watering cans used and their volume (Pollard et al., 2018). Researchers should periodically check for leaks in irrigation systems, which may be substantial (Dorr et al., 2022c). Soil amendments, such as compost and fertilizers, should be tracked through the amount applied, or the amount purchased/delivered (although this may require temporal allocation to growing season). The detailed description of our data collection methods with UA case studies in the appendix of (Dorr et al., 2022b) provides concrete examples of how to collect data across diverse systems.

3.3 Compost

Compost is the main input to many urban farms (see detailed review in the Appendix B) (Cofie et al., 2006; Dobson et al., 2021; Edmondson et al., 2014). A proposed environmental advantage of UA is its potential to grow food and reduce landfill burdens by applying compost from urban organic waste (Mohareb et al., 2017; Specht et al., 2014). Compost is thus central to UA despite infrequent and inconsistent quantification in UA LCAs (Dorr et al., 2021a). Even for rural agriculture LCAs, compost is often omitted, or its inclusion is inconsistent and unclear (Bartzas et al., 2015). Surprisingly, compost is not explicitly mentioned in reviews of LCAs of organic agriculture, where it is expected to be extensively used (Meier et al., 2015; van der Werf et al., 2020). LCAs focusing on compost use in agriculture found that the GHGs emitted from microbial decomposition (CH4 and N_2O) are a major contributor to climate change impacts, and avoided burdens (i.e. subtracting emissions from avoided processes, such as avoided incineration of organic waste) and allocation have large effects on the results for rural agriculture (Bartzas et al., 2015; Christensen et al., 2018; Martínez-Blanco et al., 2009) and for UA (Dorr et al., 2022b, 2017; Liang et al., 2019; Martin et al., 2019). Therefore, compost is given extra attention for this section.

3.3.1 Off-farm compost system modeling

Challenge:

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

Off-farm compost refers to compost purchased from municipal or industrial composting facilities, as opposed to on-farm compost, described below. In the authors' experiences, the majority of compost used in UA is purchased, because urban farms do not have the capacity to make sufficient quantities of compost on-farm. Off-farm compost used in UA is a recycled input, similar to using recycled plastic materials or recycled paper. Accounting for recycled inputs is a distinct allocation issue with a complicated and contested history in LCA (Frischknecht, 2010; Huppes and Curran, 2012; Toniolo et al., 2017; Weidema, 2000).

Examples:

A common practice to address recycling in LCA is the "simple cut off" method (Ekvall and Tillman, 1997). Here, the recycled product is cut off from the system that generated the waste, and enters the following system boundary when the waste material is transported to a recycling plant (Frischknecht, 2010). No impacts from the virgin material (for compost, this would be food or biomass production) are given to the system using the recycled product. Impacts of the recycling process and transport to the user are given to the system using the recycled material. This method can be refined by allocating some impacts from the recycling process to the upstream waste generator, considering the waste as a co-product that goes on to make a new good (Ekvall and Tillman, 1997). The ILCD Handbook (section 14.4.1.3) recommends this allocation method, considering that a valuable co-product is generated from the waste treatment process, and it is "inappropriate to attribute all preceding waste treatment processes to the eventually produced secondary good" (European Commission, 2010b). After allocating processes based on physical causality, an economic allocation is the preferred method to distribute impacts between the first system (i.e. that produced the waste) and the second system (i.e., the one that uses the compost) (European Commission, 2010b; Guinée et al., 2004). For compost, this has been done using the relative revenue at a composting plant between waste dumping fees and compost purchases (Christensen et al., 2018; Pepin, 2022).

258 For UA LCAs where off-farm compost was used, system modeling decisions have been mixed. In most cases, off-farm compost was included using the simple cut-off approach, 259 260 giving all impacts to the compost product, with no avoided burdens (Goldstein et al., 2016; Ledesma et al., 2020; Liang et al., 2019; Martin et al., 2019; Rothwell et al., 2016). 261 262 Recommendations: We recommend treating off-farm compost as a recycled input, using the refined cut off 263 method to give compost no impacts from the virgin material production and some impacts 264 265 from the composting process (Figure 1 and 2). Impacts from composting should be allocated 266 between organic waste treatment (assigned to the waste generator) and compost production (assigned to the compost user). Avoided burdens of fertilizer production should be credited to 267 the waste generator, and not the farm using compost, because the waste generator made the 268 decision that led to creation of the product displacing fertilizer (Schrijvers et al., 2016). 269 270 3.3.2 On-farm compost system modeling Challenge: 271 On-farm compost refers to the composting operations in a farm, mainly for composting 272 273 inedible plant biomass. There are several possible scenarios for on-farm compost and consequently several modeling options (Figure 1 and 2). On-farm compost may be: 274




Figure 1 A decision tree clarifies the different scenarios of composting for an urban farm and how to account for composting impacts. Blue circles represent impacts from composting emissions, and orange circles with dotted outlines represent substituted processes that can be subtracted from the farm's impacts, thanks to composting. Blue circles with gradients represent the fact that not all impacts from composting in that scenario will go to the farm: they should be allocated between the organic waste producer and the compost user. The numbered scenarios are detailed in section 3.3.2.

- 275 Scenario 1) made using on-farm biomass and used on the farm,
- 276 Scenario 2) made using on-farm biomass plus other green waste brought to the farm, and used
- 277 on the farm, or

282

- 278 Scenario 3) made using on-farm biomass and not used on the farm (i.e., for hydroponics
- 279 systems that generate biomass waste but do not use compost).
- 280 These possible scenarios, and the relevant system modeling decisions for LCA, have not been
- 281 explicitly examined before.

Examples and recommendations:

- Scenario 1 is a type of "closed loop" recycling system where the waste is generated and the
- recycled product is used within the same system. Examples of this are in Boneta et al. (2019)

and Sanyé-Mengual et al. (2015b). System modeling is straightforward with all impacts from composting given to the farm, with no avoided burdens or allocation (ISO, 2006b).




Figure 2 A process diagram shows the different composting scenarios as described in the text, from the perspective of the blue farm in the center. The numbers refer to the scenarios described in section 3.3.2, and scenario 4 refers to off-farm compost, described in section 3.3.1.

In scenario 2, composting is no longer a closed-loop system, because waste enters the system from elsewhere and is treated at the farm (i.e., food scraps from gardeners' homes or local restaurants). Here, the farm serves two functions: growing food, and treating waste. The additional function of avoided municipal waste treatment of biomass brought to the farm should be included. Allocation between these two functions is challenging because amounts of organic waste brought to the farm to be composted usually cannot be measured and tracked separately from organic waste generated on-farm. Then, the additional waste-treatment function should be accounted for using system expansion and substitution, by subtracting impacts of the alternate fate –incineration or landfilling–of organic waste from the UA system. This results in environmental credits to the UA system. This type of scenario is demonstrated in our case studies (Dorr et al., 2022b). Avoided fertilizer production should not be considered since the composted waste is used internally at the farm, so the benefit is

accounted for in the LCA results by showing smaller impacts than if the farm had used fertilizer.

Scenario 3 composting can be found at urban farms that create inedible biomass waste (all farms) but do not use compost, such as soilless hydroponics or aeroponics systems. This type of composting represents a multifunctional process: it treats the farm's waste and creates a compost product. Here the UA site is a waste generator, as discussed in the off-farm compost section (section 3.3.1). Farms should be credited with avoided environmental burdens from production of the fertilizer or potting soil that the produced compost substitutes (Corcelli et al., 2019; Goldstein et al., 2016). Vieira and Matheus (2019) provide a comprehensive review and recommendations on the matter. Composting for waste treatment of biomass can account for 10-15% of climate change impacts in UA (Corcelli et al., 2019; Sanjuan-Delmás et al., 2018), but avoided burdens of fertilizer production can generate net impact abatements (Corcelli et al., 2019).

312 3.3.3 Carbon sequestration

Challenge:

Compost is rich in organic carbon that is stabilized and stored after application to soil (Lal et al., 2015). Carbon sequestration through composting with low-carbon soil management can remove carbon from the atmosphere (Tiefenbacher et al., 2021). From an LCA perspective, this represents avoided climate change impacts, where farms using compost should receive environmental credits for sequestering CO₂ as organic carbon in their soils. However, the long-term fate of organic carbon is mostly unknown and highly context dependent because of complex soil ecology. This introduces high uncertainty to a process that can largely influence LCA results (Mclaren, 2010; Strohbach et al., 2012; Tidåker et al., 2017). Existing agriculture soil carbon models are highly time and data intensive, and are poorly adapted to UA where unique substrate and high composting rates predominate (Dorr et al., 2017).

Examples:

Several researchers argue for including carbon sequestration from compost in agricultural LCAs (Adewale et al., 2018; Martínez-Blanco et al., 2013) while others claim it is too poorly understood to be meaningfully considered (Joint Research Centre, Institute for Environment and Sustainability, 2012; Nordahl et al., 2022). Some compost LCAs (from a biowaste treatment perspective) have used carbon sequestration at rates of 10-14% of organic carbon (Boldrin et al., 2010; Tonini et al., 2020; Vaneeckhaute et al., 2018). Few UA LCAs have included soil carbon sequestration from compost. Dorr et al. (2017) used a soil model to estimate carbon sequestered from compost in substrate, potting soil, and amendments at an urban farm, , and concluded that sequestered carbon only offset 0.2-3% of GHG emissions of the farm. In a different UA LCA, Dorr et al. (2022b), applied standard carbon sequestration rates in a sensitivity analysis, which led to an offset of 3-23% of climate change impacts. LCAs of other urban green infrastructure, such as parks and golf courses, usually include carbon sequestration. This can largely affect results, sometimes even making the entire system a carbon sink (Bartlett and James, 2011; Nicese et al., 2021; Strohbach et al., 2012).

Recommendations:

We recommend excluding carbon sequestration from compost (or other organic inputs) in the main results of UA LCAs, due to the large uncertainties. It can be included in sensitivity analyses, or secondary results, to explore the extent to which it may be important, with care taken to highlight the uncertainty in those results.

3.3.4 Compost emission factors

Challenge:

The most impactful component of the compost life cycle is gaseous emissions of methane, nitrous oxide, ammonia, and volatile organic compounds during the composting process (Boldrin et al., 2009; Pergola et al., 2020). These emissions strongly affect climate change, acidification, eutrophication, and photochemical ozone formation impacts (Pergola et al., 2020). High variability in gaseous emissions from composting—due to differences in technical systems, feedstocks, and composting practices—result in high variability in

- 352 composting impacts (Joint Research Centre, Institute for Environment and Sustainability,
- 353 2012).

Reference	Type of composting system	N ₂ O emissions	CH ₄ emissions	GHG emissions	Notes (CO, NH ₃ , VOC emissions)		
	(kg/ton fresh waste)						
Andersen 2010 ^a	Home composting, closed unit	0.30-0.55	0.4-4.2	100-239	6 composting units		
Martínez-Blanco 2010 HC ^b	Home composting bin	0.676	0.158	205.4	$VOCs = 0.559$, $NH_3 = 0.842$.		
Martínez -Blanco 2010 IC ^b	Tunnel composting, with biofilters for fugitive gas	0.092	0.034	28.3	$VOCs = 1.21$, $NH_3 = 0.11$.		
Colón 2010 ^c	Fruit and vegetable scraps, yard waste, home composting	0.2	0.3	67.1	$VOCs = 0.32$, $NH_3 = 0.03$.		
Quirós 2014 HE ^d	Home composting, high-emission system	1.16	1.35	379.4	Leftover fruits and veg, yard waste. $NH_3 = 1.3$.		
Quirós 2014 LE ^d	Home composting, low-emission system	0.2	0.295	67.0	Leftover fruits and veg, yard waste. $NH_3 = 0.03$.		
Ecoinvent v3.5 ^e	Open windrow composting	0.025	1	32.5	Retrived from Ecoinvent.		
AgriBalyse- GW ^f	Green waste	0.48	0.21	148.3	Green waste. VOCs = 0.14 , $NH_3 = 1.87$		
AgriBalyse- BW ^f	Bio waste	0.13	1.15	67.5	Biowaste. VOCs = 0.21 , $NH_3 = 6.23$		
Nordahl 2022 YW ^g	Yard waste, average from review	0.043	2.31	70.6	Average of 9 values		
Nordahl 2022 OFMSW ^g	OFMSW, average, from review	0.068	0.879	42.2	Average of 21 and 19 values for CH_4 and N_2O		
Nordahl 2022 manure ^g	Manure, average, from review	0.354	2.82	176.0	Average of 41 and 45 values for CH_4 and N_2O		

Table 3 Emissions of N2O, CH4, and the sum of greenhouse gas (GHG) equivalents for N2O and CH4 are shown in kilograms of emission per ton of fresh waste composted, from some of the main sources of composting emission factors for urban agriculture life cycle assessments. GHG emissions are presented in kilograms of CO2 eq. OFMSW: organic fraction of municipal solid waste. a) Andersen et al., 2010, b) Martínez-Blanco et al., 2010, c) Colón et al., 2010, d) Quirós et al., 2014, e) Wernet et al., 2016, f) Asselin-Balençon et al., 2020, g) Nordahl et al., 2022.

Examples:

Many UA LCAs use composting emission factors from Andersen et al. (2012, 2011),

Martínez-Blanco et al. (2010), and Colón et al. (2010), because they measured inventory data specifically for home composting, which is representative of small scale, on-farm composting operations. The LCA database Ecoinvent (Wernet et al., 2016), which uses inventory data from Edelmann and Schleiss (1999), is also commonly used to model composting in UA and conventional agricultural LCAs. Table 3 shows the wide range in composting GHG emission factors from sources commonly used in agricultural LCAs. This selective list of emission values highlights the potential pitfalls from selecting composting inventories with such variability. Indeed, in our case study we found that climate change impacts were reduced by 2-14% when we used the inventory from Ecoinvent rather than from a meta-analysis by Nordahl et al. (2022). For more complete summaries of measured composting emission factors, see reviews papers by Nordahl (2022), Boldrin (2010), and Amlinger (2008), and discussion section reviews in Quiros (2015) and Avadi (2020).

Recommendations:

To address variability in composting emission factors for UA LCAs, we recommend modeling multiple scenarios with different emission factors when a farm applies large amounts of compost. Emission factors can be chosen from a specific source with a representative composting technology, or averages of multiple sources can be used. Monte Carlo simulations can be performed to include a distribution of composting emission factors to obtain a range of results and check for changes in directionality when comparing between systems (e.g., UA against conventional agriculture).

3.4 Substrate

Challenge:

A unique characteristic of UA compared to rural agriculture is that it is not necessarily carried out on soil. Soil, or top-soil, is defined as natural bodies made of organic and inorganic material that are formed at the surface as the result of complex biogeochemical and physical

processes (Brevik and Arnold, 2015; Hartemink, 2016). Using soil as a growing medium is often not an option in UA due to soil pollution in cities, or lack of greenfields. In these cases, soilless cultivation methods are used (such as hydroponics, aeroponics, or aquaponics), or a substrate/growing medium is imported. In an LCA, substrate can be considered infrastructure that requires material inputs of large quantities and variable types. Current practices around producing substrate in UA LCAs are unclear, because it often goes unmentioned, it seems to be inconsistently included, and system modeling decisions around the recycled materials often incorporated in substrate are variable (Dorr et al., 2021a). Yet, several UA LCAs have found that creating and replenishing substrate was the largest contributor for most impact categories (Dorr et al., 2017; Kim et al., 2018; Vacek et al., 2017). As a fixed input/infrastructure, substrate's lifetime directly affects its impacts, but very little information is available regarding the expected or actual lifetimes of substrate in UA. Since substrate is often amended, replenished, and used indefinitely, its lifetime is probably not limited by the material itself. Rather, substrate lifetime will likely be determined by the lifetime of the UA project or the building it is located on (Romanovska, 2019). There are few records of the lifetime of UA projects, but given UA's sometimes transient or uncertain economic nature, we suspect that such lifetimes may be shorter than anticipated (Demailly and Darly, 2017). Examples: Peat, coir, wood and compost are commonly used to produce substrate (Barrett et al., 2016). In UA, materials such as crushed brick, spent coffee grounds, spent brewer's grain, and shredded paper have also been observed (Dorr et al., 2021b; Grard et al., 2020; Martin et al., 2019). The numerous possible substrate inputs, mostly co- or by-products, lead to many options for modeling the materials. Limited details are available regarding lifetime and fate of permanent substrates in UA LCAs. Dorr et al. (2017) evaluated a research-oriented rooftop farm using substrate in raised beds,

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

403

404

405

406

407

408

409

and assumed a 10-year farm lifetime and that substrate had no end-of-life treatment (as it

would be donated and reused). Kim et al. (2018) evaluated a rooftop farm and green roof, and assumed a 40-year lifetime based on the durability of the roof membrane material. Vacek et al. (2017) did an LCA of green roofs and assumed a lifetime of 20 years, noting that they would require renovation after this point. They assumed that substrate would be landfilled, being too degraded for recycling/reuse.

Recommendations:

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

Regarding materials, the LCA guidelines published by Growing Media Europe (2021) detail how to model and what to include for numerous substrates found in UA. Peat and peat moss have been well studied, and the processes available in LCA databases should be used. Impacts for coconut and wood/bark-based materials should be allocated on an economic basis between the main coconut and forestry products and the substrate byproducts (European Commission, 2010b). Residual waste products that have negligible economic value should not incur impacts from the first use, according to economic allocation principles (ISO, 2006b). For both valuable byproducts and residual waste products, impacts from their transport after the original site of use, and energy and water needed for processing into substrate should be accounted for (Growing Media Europe, 2021). For permanent UA substrates (i.e. not disposable substrate in hydroponics and aeroponics), impacts from the initial installation of substrate should be allocated over the lifetime of the farm, similar to other pieces of infrastructure. This lifetime is usually highly uncertain, but a timeframe of 10-40 years can be considered, which can be refined based on the orientation and precarity of the case study. Results can be sensitive to this assumption so sensitivity analyses should evaluate scenarios with different farm/substrate lifetimes. Disposable substrate used in hydroponics and aeroponics do not have the same lifetime considerations and can be treated as a supply. Replenishing substrate helps maintain substrate volume and quality. Impacts of these

replenishments should be temporally allocated to the time between applications. For example,

if substrate is replenished every two years, then half of the amount applied can be allocated 436 the system in an LCA considering one year of production. 437 438 End-of-life for inorganic substrates will likely include municipal waste treatment or recycling. 439 Organic substrates are mostly composted or applied to fields as a soil improver (Growing Media Europe, 2021). For composting, the farm can be seen as the waste-generator described 440 in section 3.3.4, and impacts of composting should be allocated between the waste-generator 441 and the compost user. If substrate is applied as a soil improver by the next user, and no 442 443 treatment or processing are necessary, then no impacts for waste treatment should be given to 444 the farm. Increased transparency and improved reporting regarding substrates are necessary in UA 445 LCAs. The nature and the origins of substrate material should be clearly described, plus 446 447 physio-chemical characteristics, if available (Barrett et al., 2016). The amount of substrate 448 initially applied, the amount added in amendments, the lifetime, and end of life waste 449 treatment should be clearly stated.

3.5 Transport and delivery

451 *Challenge:*

450

452

453

454

455

456

457

458


459

460

461

462

A main supposed environmental benefit of UA is that it limits food-miles through proximity of producers and consumers (Kulak et al., 2013; Weidner et al., 2019). Yet, knowledge is scarce about how UA products are transported/delivered—yet alone their environmental performance. This benefit is sometimes dismissed, considering that on average, transportation accounts for 6-11% of climate change impacts from food systems (Poore and Nemecek, 2018; Weber and Matthews, 2008). However, fruits and vegetables can have larger contributions to climate change impacts from transport (often 10-25%, but as high as 54%), due to the potential relatively lower impacts at the farm-stage, long distances, refrigerated transport, and airplane travel (Barbier et al., 2019; Bell and Horvath, 2020; Poore and Nemecek, 2018; Weber and Matthews, 2008). The benefit of reduced transport is mostly tested through comparisons of UA LCA results to the supply chains of rural agriculture (Dorr et al., 2021a).



al., 2019; Pérez-Neira and Grollmus-Venegas, 2018). Other LCAs regarding urban food

consumption and food products have focused on the last-mile transport impacts (Bevilacqua et al., 2007; Melkonyan et al., 2020; Stelwagen et al., 2021).

Recommendations:

490

491

492

493

494

495

496

497

498

499

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

UA LCAs should include post-farm delivery processes to account for the unique urban position of farms (Weidner et al., 2019). Since there may be large uncertainties in delivery logistics and inconsistent system boundaries with rural systems, results should be presented with and without post-farm transport, giving cradle-to-farm-gate and cradle-to-consumer or market impacts (Sanyé-Mengual et al., 2015a). This is particularly relevant for comparisons to rural agriculture because proximity to the consumer is a core characteristic and environmental benefit of UA. The delivery scheme of a case study should be clearly described, including the transport distances, modes, and frequencies of deliveries. When comparing urban and rural agriculture, careful consideration must be taken to ensure that system boundaries are consistent. In particular, if the UA system has no impacts from transport, because it is done on foot or by bike, then the impacts are the same with a cradle-tofarm gate or cradle-to-consumer boundary. A cradle-to-consumer boundary is implied and should be considered, for comprehensiveness and to account for this environmental benefit of UA. Then, a scope including transport to the consumer should be included for the rural system. This stage is not represented in food products in LCA databases, and several additional transport steps are necessary for the product to reach the consumer. The feasibility of this is uncertain, however, given the lack of last mile transport data. Due to the difficulty of modeling these complex distribution networks, in-depth research on this topic may need to be done separately from production-focused UA LCAs (Coley et al., 2009; Stelwagen et al., 2021). This represents an opportunity for cross-disciplinary research on UA production and urban mobility. A city or foodshed scale may provide additional insight, as this topic quickly veers into the larger urban food logistics system rather than farm

systems (Benis and Ferrão, 2017; Melkonyan et al., 2020).

3.6 Variability and uncertainty of UA

Challenge:

516

517

518

519

520

521

522

523

524

525

526

527

528

529

530

531

532

533

534

535

536

537

538

539

540

541

542

Agricultural LCAs have particular issues with high variability because of diversity in controlled factors like farming practices and logistics, and in 'natural' factors like climate and soil characteristics (Lam et al., 2021; Mclaren, 2010; Notarnicola et al., 2017). We hypothesize that the controlled factors are even more variable in UA than in rural agriculture. Urban settings introduce physical limitations (i.e. shading from buildings, poor-quality anthropogenic soils, air pollution, and limited access to materials) which spur diverse growing practices and setups (Taylor, 2020; Wagstaff and Wortman, 2015). Human elements such as motivation for urban farming, years of experience, and access to agronomic information and training are highly variable, and likely affect growing practices (McClintock et al., 2016; Taylor, 2020). More broadly, the novel and semi-formal status of much of UA means that it has not converged towards optimized, standardized operations. In contrast, rural agriculture has been researched for decades, practiced for thousands of years, and is relatively consistent due to farmer trainings, university agricultural extension support, and technology such as tractors, crop varieties, and chemical inputs (Armanda et al., 2019; O'Sullivan et al., 2019). These factors lead to variability at a given farm (i.e. within systems). This can manifest as practices changing throughout the year, or spaces across the site being managed inconsistently. Uncertainty is also problematic, since many data are likely unavailable. This poses a problem for studying a system in its representative, average, 'steady' state. It also challenges the common LCA practice of substituting unavailable primary data with secondary data, based on the assumption that systems have somewhat standard and predictable practices. There is also high variability in UA overall (i.e. between systems). Indeed, in the review of UA LCAs (Dorr et al., 2021a), there were few actual replicates of systems due to diverse growing technology, motivation, climate, and others factors, making it difficult to compare results. Plus, many case studies were research-oriented or used innovative practices, suggesting that they may not have been representative systems. This poses a challenge to

understanding the general performance of UA, since there is not really a 'general' situation for UA.

Examples:

One of the most common ways of addressing variability and uncertainty in UA LCAs is presenting alternative scenarios in the form of sensitivity analyses. This was done in UA LCAs by modeling different infrastructure lifetimes (Dorr et al., 2017; Martin and Molin, 2019), crop yields (Romeo et al., 2018; Rufí-Salís et al., 2020b), or light efficiency for indoor systems (Pennisi et al., 2019; Shiina et al., 2011). Another strategy was to use ranges of inventory values, for example for delivery/distribution schemes, generating a range of results (Hu et al., 2019; Pérez-Neira and Grollmus-Venegas, 2018; Stelwagen et al., 2021). When parameters with high variability are identified, the goal of the LCA can shift to find tipping points where one system performs better/worse than another (usually UA vs rural). This was done for yield and distance from producer to consumer (Kulak et al., 2013; Sanyé-Mengual et al., 2015a). Alternatively, Monte Carlo simulations can be employed to quantify ranges of results based on distribution of parameters, such as composting emission factors and bulk density (Dorr et al., 2022b).

Recommendations:

Variability and uncertainty within systems can be reduced or accounted for with several strategies. Temporal variability, due to annual climate differences or changes in operations (for example due to farmer turnover), should be reduced by collecting data for multiple years and using an average of values, or selecting the most representative year (Loiseau et al., 2020). Specialized indicators can be used that quantify how important variability is for a system (Hauck et al., 2014). Variability of inventory items should be considered using distributions or ranges (Stelwagen et al., 2021), and probabilistic simulations, such as Monte Carlo simulations (Huijbregts, 1998).

Variability between systems is problematic when trying to compare or summarize results for similar systems. Such comparisons are necessary to draw trends and generalize LCA findings,

which is a feature of rather mature LCA research topics. Few technical recommendations can be made here, but we note that more complete, transparent descriptions of case studies would help readers interpret results and make more relevant, accurate comparisons between studies.

4 Research directions for UA LCAs

This section presents aspects of UA LCAs that should be the subject of future research. These topics should not necessarily be systematically included in UA LCAs, because more research and development are needed. Still, we present practical recommendations for including them in UA LCAs now. We discuss research directions that can improve UA LCAs, and how applying LCA to UA can lead to insights for LCA overall.

4.1 Align with urban land uses and green infrastructure LCAs

Presentation:

The UA LCA literature is dominated by a product-based perspective, which inherently places the focus on the food-production function of UA. UA distinguishes both the unique, non-rural position of agriculture, plus the non-conventional use of urban space (Neilson and Rickards, 2017). The latter perspective has not been widely studied with LCA, except for studies comparing different uses of rooftops for flower gardening, farming, or solar panels (Corcelli et al., 2019; Goldstein et al., 2016; Kim et al., 2018). UA is one option for urban green infrastructure among many others, and may be more comparable to a park or other social/recreational activity than it is to rural agriculture. There is a wealth of literature on environmental assessments of urban parks and forests (Strohbach et al., 2012), golf courses (Tidåker et al., 2017), urban wetlands (Duan et al., 2011), grassy areas (Smetana and Crittenden, 2014), and other green infrastructure (Nicese et al., 2021), and it would be useful to relate UA to these land uses. It could provide meaningful comparisons to similar systems, and illuminate shortcomings in UA LCAs that are obscured by a product-based perspective. For example, urban green infrastructure LCAs found that waste treatment of biomass can be highly impactful (Nicese et al., 2021; Tidåker et al., 2017), and results can be highly sensitive

to carbon sequestration (Strohbach et al., 2012; Tidåker et al., 2017), which has not emerged in UA LCAs.

Recommendations:

596

597

598

599

600

601

602

603

604

605

606

607

608

We call for increased attention to this unexplored research direction for UA LCAs: adopting an urban green infrastructure perspective of UA. Here, UA is seen as multifunctional with land use/green infrastructure as the main function, and food production is a secondary function that should be dealt with through allocation or system expansion. With system expansion, the impacts of producing an equivalent amount of food could be subtracted from the farm's impacts. With allocation, the repartition of revenue from food sales compared to grants or other sources of funding could be used for economic allocation.

UA is often presented as a tool for sustainable cities (Petit-Boix et al., 2017). Evaluating the

4.2 City-scale/Scaling up

Presentation:

effects of UA on resource consumption, food provisioning, and environmental impacts at the 609 city-scale is useful to determine the relative magnitude of findings from the farm-scale. It can 610 611 also identify emergent processes at the city-scale that are not evident at the farm-scale, such as effects on municipal organic waste treatment or urban transport logistics. 612 The effects on the city of "scaling up" or developing UA has been modeled under different 613 scenarios. Goldstein et al. (2017) evaluated the effect of installing UA in available land in 614 Boston, USA, and found that it could reduce food-related climate change impacts at the city 615 616 level by 1-3%, and increase land occupation by 1%. Mohareb et al. (2018) performed a 617 similar analysis for the USA and found food sector GHG emissions decreased by 1%. Other scaling-up analyses suggest that UA could 'absorb' and compost 9% of municipal organic 618 waste in Boston (Goldstein et al., 2017), 17% in Lyon, France, and 52% in Glasgow, Scotland 619 (Weidner and Yang, 2020). Extrapolating farm-level results to the city-scale helps provide 620 perspective, because if for example fruits and vegetables are substantially more or less 621 impactful than rural products, but at the city or individual diet scale the difference is meagre, 622

then UA LCAs should shift framing away from UA being a tool for reducing impacts of urban 623 food consumption. Such research requires estimates of the current diets of city inhabitants (to 624 625 evaluate substitution effects) (Dorr et al., 2022a), the available space for UA (Saha and Eckelman, 2017), and current city-scale flows of materials such as water and organic waste 626 (Weidner and Yang, 2020). 627 UA is embedded in the infrastructure and functioning of specific cities, which provide certain 628 629 environmental constraints or opportunities based on the city context (Martin et al., 2016). For UA LCAs, some characteristics of the specific city are inextricably included in the LCA 630 631 results. For example, a well-known factor at the country level is the electricity grid (Dorr et al., 2021a). Similar factors at the city-level may influence UA environmental performance, 632 such as city density, which may determine the proportion of rooftop vs ground-based UA, or 633 the transport mode for product delivery (Montealegre et al., 2021). The building stock in a 634 635 city may affect UA's form and impacts: for example, older buildings are more likely to need structural reinforcement for rooftop UA (Ledesma et al., 2020). The typical waste treatment 636 scheme for organic waste in a city would largely influence the potential for avoided burdens 637 638 related to compost—i.e. if organic waste is composted anyway through the city. Finally, the 639 benefits of reduced food miles for rural products are context-specific, and depend on the 640 actual source and distribution network of products to a city (Bell and Horvath, 2020; Edwards-Jones et al., 2008; Hospido et al., 2009). 641

Recommendations:

642

643

644

645

646

647

648

649

We recommend that researchers apply LCA to UA at the city scale, which can put farm-level impacts and benefits into perspective, and account for context-specific aspects of UA in a given city. As this scope veers away from on-farm production, and may focus on other aspects such as transport and delivery or external consequences of UA, primary data from farms may be less essential. Farm-level LCAs should include descriptions of the city to facilitate interpretation by others, such as the position of the farm in relation to the city center/boundary, city density, and the role of UA in the city (i.e. its history, orientation...).

4.3 Ecosystem services and positive impacts

650

Presentation: 651 LCA is designed to evaluate the negative (adverse) impacts of a system rather than its positive 652 653 impacts (benefits). The ecosystem service (ES) concept takes the opposite perspective, defined as the benefits that people obtain from ecosystems (Millennium Ecosystem 654 655 Assessment, 2005). ES assessments may better measure the benefits of UA than LCA, and combining the two ways of thinking would allow for more comprehensive assessments of 656 UA. There is no consensus on how best to measure ES, although there are many methods 657 658 available (Grêt-Regamey et al. (2017) evaluated 68 of them). Much work has been dedicated to the consideration of ES in LCA (Maia de Souza et al., 2018; Othoniel et al., 2016; Tang et 659 660 al., 2018; Zhang et al., 2010), although no method is consistently used. Some rural agriculture LCAs have performed allocation using ES (Boone et al., 2019) or with ES modeling 661 (Chaplin-Kramer et al., 2017), but no UA LCAs have incorporated ES. ES may be fully 662 integrated into the LCA methodology (i.e. with additional impact pathways for LCA,), or may 663 664 be more loosely integrated though qualitative or quantitative interpretation of results calculated separately from an LCA (De Luca Peña et al., 2022). 665 UA is a particularly rich topic through which to promote methodological development of ES 666 and LCA. It would offer useful case studies for future research because ES have been widely 667 measured as a benefit of UA, both qualitatively through interviews with stakeholders and 668 669 ranking of ES (Camps-Calvet et al., 2016; Sanyé-Mengual et al., 2020) or quantitatively with 670 indicators (Cabral et al., 2017; Grard et al., 2018). 671 There are four types of ES: provisioning, regulating, cultural and supporting (Millennium Ecosystem Assessment, 2005). Food production in UA is an obvious provisioning service. As 672 673 many UA LCAs use a FU based on food production, they essentially quantify the impact of 674 this ES. Boone et al. (2019) demonstrated a method to allocate between this provisioning ES of agriculture and other ES in an LCA, which highlighted that food was not the only ES (or 675 'output') of agriculture. 676

Regulating ES of UA that have been measured include water runoff regulation, organic waste 677 recycling, and microclimate regulation (Dennis and James, 2017; Grard et al., 2018). Benefits 678 679 of avoided stormwater runoff have been quantified with LCA, and offset 13-72% of several impact categories (Goldstein et al., 2016; Kim et al., 2018). Carbon sequestration can also be 680 evaluated using LCA or ES (Orsini et al., 2014), and its implication in LCA is described in 681 section 3.3.3. Reduction of the urban heat island effect is a frequently proposed regulation ES 682 of UA, and is generally excluded from all LCAs (Susca and Pomponi, 2020). 683 Cultural ES are sometimes perceived as the top benefit of UA, and include recreation, 684 beautification, cultural identity, social cohesion, community building, and education (Giacchè 685 et al., 2021; Sanyé-Mengual et al., 2018b). Indicators to measure cultural ES include the 686 687 volunteer hours, number of educational and recreational activities offered, and their number of participants (Dennis and James, 2017; Giacchè et al., 2021). Cultural ES may provide a 688 689 framework to include social benefits in UA LCA assessment (detailed more in section 4.4). 690 The role of biodiversity in ES is foundational, as it is defined as the source of ES (McDonald et al., 2013; Millennium Ecosystem Assessment, 2005), and is often used as a proxy indicator 691 for supporting ES (Cabral et al., 2017). Improved local biodiversity is perceived as an 692 important environmental benefit of UA (Sanyé-Mengual et al., 2018b) and is frequently 693 694 measured in the context of ES of UA (Dennis and James, 2017; Quistberg et al., 2016). This 695 benefit is not accounted for in LCA. Biodiversity impacts in LCA have been the subject of 696 methodological development for decades, and is usually framed as the impact *on* biodiversity 697 from land use (or other ecological damage, although most frequently land use) (Teixeira et al., 2016). LCA models the upstream and downstream impacts of materials and processes on 698 biodiversity around the world, and does not consider local biodiversity (Teixeira et al., 2016). 699 Other measures are more relevant for farm-scale biodiversity impacts like species richness, 700 701 habitat fragmentation, habitat vulnerability, or land use intensity indicators (Frischknecht et al., 2016; Pepin, 2022). 702

Recommendations:

703

- For practitioners looking to operationalize ES and LCA for UA, results from each method can
- be qualitatively assessed in parallel or quantitatively through composite indicators (De Luca
- Peña et al., 2022). For an integrated assessment, for example comparing types of UA within
- one study, results can be integrated in a multi-criteria decision analysis (Ledesma et al.,
- 708 2020).
- 709 Researchers looking to improve LCA methodology by integrating it with ES should consider
- 710 using UA as their application. UA represents a particularly relevant activity, due to its
- 711 multifunctionality and the fact that many ES have already been demonstrated.

712 4.4 Social benefits and life cycle sustainability assessment

- 713 *Presentation:*
- A main strength of UA is its multifunctionality, with important social functions (Gomez
- 715 Villarino et al., 2021; Orsini et al., 2020). This is rarely reflected in UA LCAs, but it should
- be, since core principles of LCA are evaluating the main function of a system (through
- selection of a FU), and accounting for multiple outputs (through allocation and system
- 718 expansion).
- 719 Accounting for social aspects of an activity is a main issue for LCA, and social LCA (S-LCA)
- 720 is a promising yet nascent strategy to overcome this (UNEP/SETAC, 2009; Zimek et al.,
- 721 2019). Using life-cycle thinking, S-LCA tracks the social impacts of a product's life cycle. S-
- 722 LCA quantifies negative impacts, and therefore may not be appropriate for evaluating the
- social benefits of UA. S-LCA databases offer data for social impacts embedded along the
- 724 supply chain, but the information necessary for UA is more relevant at the farm,
- neighborhood, or city scale (Romanovska, 2019). Plus, such databases are not as
- 726 generalizable as large LCA databases. A strength of S-LCA is its ability to account for the
- 727 perspectives of multiple stakeholders, such as workers, consumers, and the local community.
- 728 This is especially useful to evaluate the potential for UA to address social justice issues, by
- 729 highlighting not just which social benefits are brought, but who they are affecting. S-LCA
- currently lacks agreed upon social indicators, partly because they are situational and defined

through stakeholder engagement, making consistent methods and comparisons between 731 studies difficult (Fauzi et al., 2019). Peri et al. (2010) outlined indicators for S-LCA of green 732 733 roofs, including area of green roof made accessible to the public, fair salary, working hours, 734 air pollutant levels, and outside air temperature. Apart from S-LCA, an option to include social benefits of UA is to address its 735 multifunctionality with traditional LCA practices. For example, allocation can be used to 736 distribute impacts based on relative importance of food production vs. social benefits. This 737 738 allocation may be done based on the level of ES provided by each activity, as done in Boone 739 et al. (2019). Alternatively, it may be based on the relative sources of revenue from food sales vs. grants vs. other activities. If social goals are the main function of a farm, we can imagine 740 using a FU based on the social "output", such as volunteer hours or total number of new 741 people met by UA participants, which can be linked to cultural ES. 742 Social aspects of UA may be evaluated in parallel to environmental impacts from LCA rather 743 744 than being fully integrated into LCA. Indeed, many researchers acknowledge that LCA cannot capture everything, and it is useful to complement it with other methods (De Luca Peña et al., 745 2022; Fauzi et al., 2019). In practice, this would be most useful to compare different types of 746 UA within a study, where the same data can be collected from a set of urban farms. UA LCA 747 748 practitioners should strive to measure these indicators of social benefits and present them in 749 case studies, even when they are not based on a life-cycle approach. 750 The LCA community has promoted and strives for life cycle sustainability assessment, which combines environmental LCA, life cycle cost analysis (LCCA, which was reviewed for UA 751 by Peña and Rovera-Val (2020)), and S-LCA. Such holistic life cycle sustainability 752 assessments are still largely more aspirational than operational (Fauzi et al., 2019; Finkbeiner 753 et al., 2010). We urge UA LCA practitioners to consider measures of economic and social 754 sustainability even if they are not life-cycle based, which is indeed particularly data-755 demanding (Sanyé-Mengual et al., 2017). LCA results may even be included in broader 756 757 indicator-based sustainability assessments, which are operationalized in tools for rural

agriculture and are under development for UA (Clerino and Fargue-Lelièvre, 2020; Hély and Antoni, 2019).

Recommendations:

760

761

762

763

764

765

766

767

768

Researchers should work towards defining a set of S-LCA indicators relevant for UA. The concept and assessment of cultural ES may serve as a basis here, since they are both indicator-based, site-specific measures. New methods should be tested to use allocation or alternative, social-based FUs to account for social aspects of UA. Although we should ultimately strive for life cycle sustainability assessment, non-life cycle indicators and results, such as results from surveys and interviews, should be presented alongside LCA results to provide more holistic views of sustainability.

5 Conclusion

769 Since the first LCA of UA a decade ago, interest in and knowledge of the environmental performance of UA has increased. Still, large uncertainties remain regarding best practices for 770 771 these assessments, and even defining what questions we aim to address. In this article, we laid 772 out recommendations and research directions that are intended to improve LCAs of UA. 773 These improvements can lead to more thorough LCAs and more consistency between case studies. We also outlined the questions that UA LCAs may aim to answer, in the hopes of 774 bringing perspective and clarity to this field of research. Finally, this work highlights what 775 776 LCA can *learn* from UA through challenges in applying it to this complex and multifunctional 777 activity. To accurately support policy and decision-making around UA, LCAs must be more 778 comprehensive. To provide more meaningful support, UA LCA findings should be considered 779 alongside measurements of other sustainability dimensions, whether they are life-cycle based 780 or not. 781 By applying these guidelines and strengthening UA LCAs, this research topic can better support environmental sustainability of UA and cities. This research can better inform policy 782 783 makers about how UA implementation will affect environmental performance of cities, and

which types or characteristics of UA to leverage for specific goals. It can inform urban farmers on how to operate or design their farms to minimize environmental impacts. They can better understand which changes to implement, and which ones may not be worth the effort given small environmental gains. Finally, the research community can explore methods to enhance the use of LCA for multifunctional, complex activities, such as UA.

Acknowledgments

E.D, C.A, and B.G gratefully acknowledge financial support of lab recherche environnement
 VINCI ParisTech. A.H gratefully acknowledges the support of the National Science
 Foundation under Grant No. 1739676. Any opinions, findings, and conclusions or
 recommendations expressed in this material are those of the authors and do not necessarily
 reflect the views of the National Science Foundation.

Bibliography 796

804 805

806

807

809

810

811

812 813

814

815 816

- Acosta-Alba, I., Chia, E., Andrieu, N., 2019. The LCA4CSA framework: Using life cycle 797 798 assessment to strengthen environmental sustainability analysis of climate smart 799 agriculture options at farm and crop system levels. Agric. Syst. 171, 155–170. 800 https://doi.org/10.1016/j.agsy.2019.02.001
- Adewale, C., Reganold, J.P., Higgins, S., Evans, R.D., Carpenter-Boggs, L., 2018. Improving 801 802 carbon footprinting of agricultural systems: Boundaries, tiers, and organic farming. 803 Environ. Impact Assess. Rev. 71, 41–48. https://doi.org/10.1016/j.eiar.2018.04.004
 - Amlinger, F., Peyr, S., Cuhls, C., 2008. Green house gas emissions from composting and mechanical biological treatment. Waste Manag. Res. J. Int. Solid Wastes Public Clean. Assoc. ISWA 26, 47–60. https://doi.org/10.1177/0734242X07088432
- Andersen, J.K., Boldrin, A., Christensen, T.H., Scheutz, C., 2012. Home composting as an 808 alternative treatment option for organic household waste in Denmark: An environmental assessment using life cycle assessment-modelling. Waste Manag. 32, 31–40. https://doi.org/10.1016/j.wasman.2011.09.014
 - Andersen, J.K., Boldrin, A., Christensen, T.H., Scheutz, C., 2011. Mass balances and life cycle inventory of home composting of organic waste. Waste Manag. 31, 1934–1942. https://doi.org/10.1016/j.wasman.2011.05.004
 - Armanda, D.T., Guinée, J.B., Tukker, A., 2019. The second green revolution: Innovative urban agriculture's contribution to food security and sustainability – A review. Glob. Food Secur. 22, 13–24. https://doi.org/10.1016/j.gfs.2019.08.002
- 817 Audsley, A., Alber, S., Clift, R., Cowell, S., Crettaz, R., Gaillard, G., Hausheer, J., Jolliett, O., 818 Kleijn, R., Mortensen, B., Pearce, D., Roger, E., Teulon, H., Weidema, B., Zeijts, H. 819 van, 1997. Harmonisation of environmental Life Cycle Assessment for agriculture. 820 Final report (concerted action No. AIR3-CT94-2028). European commission DG VI, 821 Brussels, Belgium.
- Avadí, A., Aissani, L., Pradel, M., Wilfart, A., 2020. Life cycle inventory data on French 822 organic waste treatments yielding organic amendments and fertilisers. Data Brief 28, 823 105000. https://doi.org/10.1016/j.dib.2019.105000 824
- 825 Azunre, G.A., Amponsah, O., Peprah, C., Takyi, S.A., Braimah, I., 2019. A review of the role of urban agriculture in the sustainable city discourse. Cities 93, 104–119. 826 827 https://doi.org/10.1016/j.cities.2019.04.006
- Barbier, C., Couturier, C., Pourouchottamin, P., Cayla, J.-M., Sylvestre, M., Pharabod, I., 828 2019. L'empreinte énergétique et carbone de l'alimentation en France: de la 830 production à la consommation. Club Ingénierie Prospective Energie et Environnement, 831 Paris, France, IDDRI.
- Barrett, G.E., Alexander, P.D., Robinson, J.S., Bragg, N.C., 2016. Achieving environmentally 832 sustainable growing media for soilless plant cultivation systems – A review. Sci. 833 834 Hortic, 212, 220–234. https://doi.org/10.1016/j.scienta.2016.09.030
- Bartlett, M.D., James, I.T., 2011. A model of greenhouse gas emissions from the management 835 836 of turf on two golf courses. Sci. Total Environ. 409, 5137–5147. 837 https://doi.org/10.1016/j.scitotenv.2011.07.054
- Bartzas, G., Zaharaki, D., Komnitsas, K., 2015. Life cycle assessment of open field and 838 greenhouse cultivation of lettuce and barley. Inf. Process. Agric. 2, 191–207. 839 https://doi.org/10.1016/j.inpa.2015.10.001 840
- 841 Bell, E.M., Horvath, A., 2020. Modeling the carbon footprint of fresh produce: effects of 842 transportation, localness, and seasonality on US orange markets. Environ. Res. Lett. 15, 034040. https://doi.org/10.1088/1748-9326/ab6c2f 843
- 844 Benis, K., Ferrão, P., 2017. Potential mitigation of the environmental impacts of food systems through urban and peri-urban agriculture (UPA) – a life cycle assessment approach. J. 845 Clean. Prod. 140, 784–795. https://doi.org/10.1016/j.jclepro.2016.05.176 846

- Bevilacqua, M., Braglia, M., Carmignani, G., Zammori, F.A., 2007. Life Cycle Assessment of Pasta Production in Italy. J. Food Qual. 30, 932–952. https://doi.org/10.1111/j.1745-4557.2007.00170.x
- Bishop, G., Styles, D., Lens, P.N.L., 2021. Environmental performance comparison of bioplastics and petrochemical plastics: A review of life cycle assessment (LCA) methodological decisions. Resour. Conserv. Recycl. 168, 105451. https://doi.org/10.1016/j.resconrec.2021.105451
- Boldrin, A., Andersen, J.K., Møller, J., Christensen, T.H., Favoino, E., 2009. Composting and compost utilization: accounting of greenhouse gases and global warming contributions. Waste Manag. Res. 27, 800–812.

 https://doi.org/10.1177/0734242X09345275
- Boldrin, A., Hartling, K.R., Laugen, M., Christensen, T.H., 2010. Environmental inventory
 modelling of the use of compost and peat in growth media preparation. Resour.
 Conserv. Recycl. 54, 1250–1260. https://doi.org/10.1016/j.resconrec.2010.04.003
- Boneta, A., Rufí-Salís, M., Ercilla-Montserrat, M., Gabarrell, X., Rieradevall, J., 2019.
 Agronomic and Environmental Assessment of a Polyculture Rooftop Soilless Urban
 Home Garden in a Mediterranean City. Front. Plant Sci. 10, 12.
 https://doi.org/10.3389/fpls.2019.00341
- Boone, L., Roldán-Ruiz, I., Van linden, V., Muylle, H., Dewulf, J., 2019. Environmental
 sustainability of conventional and organic farming: Accounting for ecosystem services
 in life cycle assessment. Sci. Total Environ. 695, 133841.
 https://doi.org/10.1016/j.scitotenv.2019.133841
- Brankatschk, G., Finkbeiner, M., 2015. Modeling crop rotation in agricultural LCAs Challenges and potential solutions. Agric. Syst. 138, 66–76. https://doi.org/10.1016/j.agsy.2015.05.008
- 872 Brentrup, F., Küsters, J., Kuhlmann, H., Lammel, J., 2004. Environmental impact assessment 873 of agricultural production systems using the life cycle assessment methodology: I. 874 Theoretical concept of a LCA method tailored to crop production. Eur. J. Agron. 20, 875 247–264. https://doi.org/10.1016/S1161-0301(03)00024-8
- Brevik, E.C., Arnold, R.W., 2015. Is the Traditional Pedologic Definition of Soil Meaningful in the Modern Context? Soil Horiz. 56, sh15-01–0002. https://doi.org/10.2136/sh15-01-0002
- Cabral, I., Keim, J., Engelmann, R., Kraemer, R., Siebert, J., Bonn, A., 2017. Ecosystem services of allotment and community gardens: A Leipzig, Germany case study. Urban For. Urban Green. 23, 44–53. https://doi.org/10.1016/j.ufug.2017.02.008
 - Caffrey, K.R., Veal, M.W., 2013. Conducting an Agricultural Life Cycle Assessment: Challenges and Perspectives. Sci. World J. https://doi.org/10.1155/2013/472431

884

885

886 887

- Camps-Calvet, M., Langemeyer, J., Calvet-Mir, L., Gómez-Baggethun, E., 2016. Ecosystem services provided by urban gardens in Barcelona, Spain: Insights for policy and planning. Environ. Sci. Policy, Advancing urban environmental governance: Understanding theories, practices and processes shaping urban sustainability and resilience 62, 14–23. https://doi.org/10.1016/j.envsci.2016.01.007
- Caputo, P., Zagarella, F., Cusenza, M.A., Mistretta, M., Cellura, M., 2020. Energyenvironmental assessment of the UIA-OpenAgri case study as urban regeneration project through agriculture. Sci. Total Environ. 729, 138819. https://doi.org/10.1016/j.scitotenv.2020.138819
- Cerutti, A.K., Beccaro, G.L., Bruun, S., Bosco, S., Donno, D., Notarnicola, B., Bounous, G.,
 2014. Life cycle assessment application in the fruit sector: State of the art and
 recommendations for environmental declarations of fruit products. J. Clean. Prod.,
 Towards eco-efficient agriculture and food systems: Selected papers from the Life
 Cycle Assessment (LCA) Food Conference, 2012, in Saint Malo, France 73, 125–135.
 https://doi.org/10.1016/j.jclepro.2013.09.017
- Chaplin-Kramer, R., Sim, S., Hamel, P., Bryant, B., Noe, R., Mueller, C., Rigarlsford, G.,
 Kulak, M., Kowal, V., Sharp, R., Clavreul, J., Price, E., Polasky, S., Ruckelshaus, M.,
 Daily, G., 2017. Life cycle assessment needs predictive spatial modelling for

biodiversity and ecosystem services. Nat. Commun. 8, 15065.
 https://doi.org/10.1038/ncomms15065

907

908

909

914

915

916 917

924

925

926

927 928

929

930

931

932

933 934

935

936

937 938

939

940

941 942

943

944

945

- Christensen, L.O., Galt, R.E., Kendall, A., 2018. Life-cycle greenhouse gas assessment of
 Community Supported Agriculture in California's Central Valley. Renew. Agric. Food
 Syst. 33, 393–405. https://doi.org/10.1017/S1742170517000254
 - Clerino, P., Fargue-Lelièvre, A., 2020. Formalizing Objectives and Criteria for Urban Agriculture Sustainability with a Participatory Approach. Sustainability 12, 7503. https://doi.org/10.3390/su12187503
- 910 Cleveland, D., 1997. Are urban gardens an efficient use of resources? Arid Lands Newsl.
- CoDyre, M., Fraser, E.D.G., Landman, K., 2015. How does your garden grow? An empirical evaluation of the costs and potential of urban gardening. Urban For. Urban Green. 14, 72–79. https://doi.org/10.1016/j.ufug.2014.11.001
 - Cofie, O., Adam-Bradford, A., Drechsel, P., 2006. Recycling of Urban Organic Waste for Urban Agriculture, in: Cities Farming for TheFuture: Urban Agriculture for Green and Productive Cities. International Institute of Rural Reconstruction and ETC Urban Agriculture, Philippines, pp. 210–229.
- Coley, D., Howard, M., Winter, M., 2009. Local food, food miles and carbon emissions: A comparison of farm shop and mass distribution approaches. Food Policy 34, 150–155. https://doi.org/10.1016/j.foodpol.2008.11.001
- Colón, J., Martínez-Blanco, J., Gabarrell, X., Artola, A., Sánchez, A., Rieradevall, J., Font,
 X., 2010. Environmental assessment of home composting. Resour. Conserv. Recycl.
 54, 893–904. https://doi.org/10.1016/j.resconrec.2010.01.008
 - Corcelli, F., Fiorentino, G., Petit-Boix, A., Rieradevall, J., Gabarrell, X., 2019. Transforming rooftops into productive urban spaces in the Mediterranean. An LCA comparison of agri-urban production and photovoltaic energy generation. Resour. Conserv. Recycl. 144, 321–336. https://doi.org/10.1016/j.resconrec.2019.01.040
 - Cucurachi, S., Scherer, L., Guinée, J., Tukker, A., 2019. Life Cycle Assessment of Food Systems. One Earth 1, 292–297. https://doi.org/10.1016/j.oneear.2019.10.014
 - De Luca Peña, L.V., Taelman, S.E., Préat, N., Boone, L., Van der Biest, K., Custódio, M., Hernandez Lucas, S., Everaert, G., Dewulf, J., 2022. Towards a comprehensive sustainability methodology to assess anthropogenic impacts on ecosystems: Review of the integration of Life Cycle Assessment, Environmental Risk Assessment and Ecosystem Services Assessment. Sci. Total Environ. 808, 152125. https://doi.org/10.1016/j.scitotenv.2021.152125
 - Demailly, K.-E., Darly, S., 2017. Urban agriculture on the move in Paris: The routes of temporary gardening in the neoliberal city. ACME Int. E-J. Crit. Geogr.
 - Dennis, M., James, P., 2017. Ecosystem services of collectively managed urban gardens: Exploring factors affecting synergies and trade-offs at the site level. Ecosyst. Serv. 26, 17–26. https://doi.org/10.1016/j.ecoser.2017.05.009
 - Dijkman, T.J., Basset-Mens, C., Antón, A., Núñez, M., 2018. LCA of Food and Agriculture, in: Hauschild, M.Z., Rosenbaum, R.K., Olsen, S.I. (Eds.), Life Cycle Assessment: Theory and Practice. Springer International Publishing, Cham, pp. 723–754. https://doi.org/10.1007/978-3-319-56475-3_29
 - Dobson, M.C., Warren, P.H., Edmondson, J.L., 2021. Assessing the Direct Resource Requirements of Urban Horticulture in the United Kingdom: A Citizen Science Approach. Sustainability 13, 2628. https://doi.org/10.3390/su13052628
- Dorr, E., François, C., Poulhès, A., Wurtz, A., 2022a. A life cycle assessment method to support cities in their climate change mitigation strategies. Sustain. Cities Soc. 85, 104052. https://doi.org/10.1016/j.scs.2022.104052
- Dorr, E., Goldstein, B., Aubry, C., 2022b. Life cycle assessment of eight urban farms and community gardens in France and California. Resour. Conserv. Recycl.
- Dorr, E., Goldstein, B., Horvath, A., Aubry, C., Gabrielle, B., 2021a. Environmental impacts and resource use of urban agriculture: a systematic review and meta-analysis. Environ. Res. Lett. 16, 093002. https://doi.org/10.1088/1748-9326/ac1a39

- Dorr, E., Hawes, J.K., Goldstein, B., 2022c. Food production and resource use of urban farms and gardens: a five-country study. Agron. Sustain. Dev.
- Dorr, E., Koegler, M., Gabrielle, B., Aubry, C., 2021b. Life cycle assessment of a circular,
 urban mushroom farm. J. Clean. Prod. 288, 125668.
 https://doi.org/10.1016/j.jclepro.2020.125668
- Dorr, E., Sanyé-Mengual, E., Gabrielle, B., Grard, B.J.-P., Aubry, C., 2017. Proper selection
 of substrates and crops enhances the sustainability of Paris rooftop garden. Agron.
 Sustain. Dev. 37, 51. https://doi.org/10.1007/s13593-017-0459-1
- Duan, N., Liu, X.D., Dai, J., Lin, C., Xia, X.H., Gao, R.Y., Wang, Y., Chen, S.Q., Yang, J.,
 Qi, J., 2011. Evaluating the environmental impacts of an urban wetland park based on
 emergy accounting and life cycle assessment: A case study in Beijing. Ecol. Model.,
 Wetlands in China 222, 351–359. https://doi.org/10.1016/j.ecolmodel.2010.08.028
- Edelmann, W., Schleiss, K., 1999. Ökologischer, energetischer und ökonomischer Vergleich
 von Vergärung, Kompostierung und Verbrennung fester biogener Abfallstoffe.
 Bundesamt für Energie and Bundesamt für Umwelt, Wald und Landschaft, Bern,
 Switzerland.
- Edmondson, J.L., Davies, Z.G., Gaston, K.J., Leake, J.R., 2014. Urban cultivation in
 allotments maintains soil qualities adversely affected by conventional agriculture. J.
 Appl. Ecol. 51, 880–889. https://doi.org/10.1111/1365-2664.12254
- Edwards-Jones, G., Milà i Canals, L., Hounsome, N., Truninger, M., Koerber, G., Hounsome,
 B., Cross, P., York, E.H., Hospido, A., Plassmann, K., Harris, I.M., Edwards, R.T.,
 Day, G.A.S., Tomos, A.D., Cowell, S.J., Jones, D.L., 2008. Testing the assertion that
 'local food is best': the challenges of an evidence-based approach. Trends Food Sci.
 Technol. 19, 265–274. https://doi.org/10.1016/j.tifs.2008.01.008
- Egerer, M.H., Lin, B.B., Philpott, S.M., 2018. Water Use Behavior, Learning, and Adaptation to Future Change in Urban Gardens. Front. Sustain. Food Syst. https://doi.org/10.3389/fsufs.2018.00071
- Ekvall, T., Tillman, A.-M., 1997. Open-loop recycling: Criteria for allocation procedures. Int.
 J. Life Cycle Assess. 2, 155. https://doi.org/10.1007/BF02978810
 European Commission, 2017. PEFCR Guidance document, Guidance for the development of
 - European Commission, 2017. PEFCR Guidance document, Guidance for the development of Product Environmental Footprint Category Rules (PEFCRs) (No. version 6.3).
 - European Commission, 2010a. ILCD Handbook: General guide for Life Cycle Assessment: provisions and action steps, First. ed. Institute for Environment and Sustainability, Luxembourg. https://doi.org/10.2788/94987
- European Commission, 2010b. ILCD Handbook: General guide for Life Cycle Assessment:
 Detailed guidance, First. ed. Institute for Environment and Sustainability,
 Luxembourg. https://doi.org/10.2788/38479
 - Fauzi, R.T., Lavoie, P., Sorelli, L., Heidari, M.D., Amor, B., 2019. Exploring the Current Challenges and Opportunities of Life Cycle Sustainability Assessment. Sustainability 11, 636. https://doi.org/10.3390/su11030636
- Finkbeiner, M., Schau, E.M., Lehmann, A., Traverso, M., 2010. Towards Life Cycle
 Sustainability Assessment. Sustainability 2, 3309–3322.
 https://doi.org/10.3390/su2103309

987 988

989

993

994

- 999 Frischknecht, R., 2010. LCI modelling approaches applied on recycling of materials in view 1000 of environmental sustainability, risk perception and eco-efficiency. Int. J. Life Cycle 1001 Assess. 15, 666–671. https://doi.org/10.1007/s11367-010-0201-6
- Frischknecht, R., Fantke, P., Tschümperlin, L., Niero, M., Antón, A., Bare, J., Boulay, A.-M., Cherubini, F., Hauschild, M.Z., Henderson, A., Levasseur, A., McKone, T.E., Michelsen, O., i Canals, L.M., Pfister, S., Ridoutt, B., Rosenbaum, R.K., Verones, F., Vigon, B., Jolliet, O., 2016. Global guidance on environmental life cycle impact assessment indicators: progress and case study. Int. J. Life Cycle Assess. 21, 429–442. https://doi.org/10.1007/s11367-015-1025-1
- Giacchè, G., Consalès, J.-N., Grard, B.J.-P., Daniel, A.-C., Chenu, C., 2021. Toward an Evaluation of Cultural Ecosystem Services Delivered by Urban Micro-Farms. Sustainability 13, 1716. https://doi.org/10.3390/su13041716

- 1011 Gnielka, A.E., Menzel, C., 2021. The impact of the consumer's decision on the life cycle
 1012 assessment of organic pasta. SN Appl. Sci. 3, 839. https://doi.org/10.1007/s424521013 021-04822-y
- Goldstein, B., Hauschild, M., Fernández, J., Birkved, M., 2017. Contributions of Local
 Farming to Urban Sustainability in the Northeast United States. Environ. Sci. Technol.
 51, 7340–7349. https://doi.org/10.1021/acs.est.7b01011
- Goldstein, B., Hauschild, M., Fernández, J., Birkved, M., 2016. Testing the environmental performance of urban agriculture as a food supply in northern climates. J. Clean. Prod. 135, 984–994. https://doi.org/10.1016/j.jclepro.2016.07.004
- Gomez Villarino, M.T., urquijo, J., Gómez Villarino, M., García, A.I., 2021. Key insights of urban agriculture for sustainable urban development. Agroecol. Sustain. Food Syst. 0, 1–29. https://doi.org/10.1080/21683565.2021.1917471
- Gómez-Villarino, M.T., urquijo, J., Gómez Villarino, M., García, A.I., 2021. Key insights of urban agriculture for sustainable urban development. Agroecol. Sustain. Food Syst. 45, 1441–1469. https://doi.org/10.1080/21683565.2021.1917471
- Grard, B.J.-P., Chenu, C., Manouchehri, N., Houot, S., Frascaria-Lacoste, N., Aubry, C., 2018. Rooftop farming on urban waste provides many ecosystem services. Agron. Sustain. Dev. 38, 2. https://doi.org/10.1007/s13593-017-0474-2

1033

1034 1035

1036 1037

1038

1041

10421043

1044

1045

- Grard, B.J.-P., Manouchehri, N., Aubry, C., Frascaria-Lacoste, N., Chenu, C., 2020. Potential of Technosols Created with Urban By-Products for Rooftop Edible Production. Int. J. Environ. Res. Public. Health 17, 3210. https://doi.org/10.3390/ijerph17093210
 - Gregory, M.M., Leslie, T.W., Drinkwater, L.E., 2016. Agroecological and social characteristics of New York city community gardens: contributions to urban food security, ecosystem services, and environmental education. Urban Ecosyst. 19, 763–794. https://doi.org/10.1007/s11252-015-0505-1
 - Grêt-Regamey, A., Sirén, E., Brunner, S.H., Weibel, B., 2017. Review of decision support tools to operationalize the ecosystem services concept. Ecosyst. Serv., Putting ES into practice 26, 306–315. https://doi.org/10.1016/j.ecoser.2016.10.012
- Growing Media Europe, 2021. Growing Media Environmental Footprint Guideline V1.0. Guinée, J.B., Heijungs, R., Huppes, G., 2004. Economic allocation: Examples and derived
 - Guinée, J.B., Heijungs, R., Huppes, G., 2004. Economic allocation: Examples and derived decision tree. Int. J. Life Cycle Assess. 9, 23. https://doi.org/10.1007/BF02978533
 - Hall, G., Rothwell, A., Grant, T., Isaacs, B., Ford, L., Dixon, J., Kirk, M., Friel, S., 2014. Potential environmental and population health impacts of local urban food systems under climate change: a life cycle analysis case study of lettuce and chicken.
 - Hartemink, A.E., 2016. Chapter Two The definition of soil since the early 1800s, in: Sparks, D.L. (Ed.), Advances in Agronomy, Advances in Agronomy. Academic Press, pp. 73–126. https://doi.org/10.1016/bs.agron.2015.12.001
- Hauck, M., Steinmann, Z.J.N., Laurenzi, I.J., Karuppiah, R., Huijbregts, M.A.J., 2014. How to quantify uncertainty and variability in life cycle assessment: the case of greenhouse gas emissions of gas power generation in the US. Environ. Res. Lett. 9, 074005. https://doi.org/10.1088/1748-9326/9/7/074005
- Hély, V., Antoni, J.-P., 2019. Combining indicators for decision making in planning issues: A
 theoretical approach to perform sustainability assessment. Sustain. Cities Soc. 44,
 844–854. https://doi.org/10.1016/j.scs.2018.10.035
- Hospido, A., Milà i Canals, L., McLaren, S., Truninger, M., Edwards-Jones, G., Clift, R., 2009. The role of seasonality in lettuce consumption: a case study of environmental and social aspects. Int. J. Life Cycle Assess. 14, 381–391. https://doi.org/10.1007/s11367-009-0091-7
- Hu, Y., Zheng, J., Kong, X., Sun, J., Li, Y., 2019. Carbon footprint and economic efficiency of urban agriculture in Beijing—a comparative case study of conventional and home-delivery agriculture. J. Clean. Prod. 234, 615–625. https://doi.org/10.1016/j.jclepro.2019.06.122
- Huijbregts, M.A.J., 1998. Application of uncertainty and variability in LCA. Int. J. Life Cycle
 Assess. 3, 273. https://doi.org/10.1007/BF02979835

- Huppes, G., Curran, M.A., 2012. Environmental Life Cycle Assessment: Background and
 Perspective, in: Life Cycle Assessment Handbook. John Wiley & Sons, Ltd, pp. 1–14.
 https://doi.org/10.1002/9781118528372.ch1
- 1068 ISO, 2006a. ISO 14040, Environmental management Life cycle assessment Principles and framework.
- 1070 ISO, 2006b. ISO 14044, Environmental management Life cycle assessment Requirements and guidelines.
- Joint Research Centre, Institute for Environment and Sustainability, 2012. Supporting
 environmentally sound decisions for waste management: a technical guide to Life
 Cycle Thinking (LCT) and Life Cycle Assessment (LCA) for waste experts and LCA
 practitioners. Publications Office of the European Union, LU.
- Kim, E., Jung, J., Hapsari, G., Kang, S., Kim, K., Yoon, S., Lee, M., Han, M., Choi, Y., Choe,
 J.K., 2018. Economic and environmental sustainability and public perceptions of
 rooftop farm versus extensive garden. Build. Environ. 146, 206–215.
 https://doi.org/10.1016/j.buildenv.2018.09.046
- Kirkpatrick, J.B., Davison, A., 2018. Home-grown: Gardens, practices and motivations in urban domestic vegetable production. Landsc. Urban Plan. 170, 24–33. https://doi.org/10.1016/j.landurbplan.2017.09.023
- Kløverpris, J.H., Scheel, C.N., Schmidt, J., Grant, B., Smith, W., Bentham, M.J., 2020.
 Assessing life cycle impacts from changes in agricultural practices of crop production.
 Int. J. Life Cycle Assess. 25, 1991–2007. https://doi.org/10.1007/s11367-020-01767-z
- Kulak, M., Graves, A., Chatterton, J., 2013. Reducing greenhouse gas emissions with urban agriculture: A Life Cycle Assessment perspective. Landsc. Urban Plan. 111, 68–78. https://doi.org/10.1016/j.landurbplan.2012.11.007
- Lal, R., Negassa, W., Lorenz, K., 2015. Carbon sequestration in soil. Curr. Opin. Environ.
 Sustain., Environmental change issues 15, 79–86.
 https://doi.org/10.1016/j.cosust.2015.09.002
- Lam, W.Y., Sim, S., Kulak, M., van Zelm, R., Schipper, A.M., Huijbregts, M.A.J., 2021.
 Drivers of variability in greenhouse gas footprints of crop production. J. Clean. Prod.
 315, 128121. https://doi.org/10.1016/j.jclepro.2021.128121
- Laurent, A., Clavreul, J., Bernstad, A., Bakas, I., Niero, M., Gentil, E., Christensen, T.H.,
 Hauschild, M.Z., 2014. Review of LCA studies of solid waste management systems –
 Part II: Methodological guidance for a better practice. Waste Manag. 34, 589–606.
 https://doi.org/10.1016/j.wasman.2013.12.004
- Ledesma, G., Nikolic, J., Pons-Valladares, O., 2020. Bottom-up model for the sustainability
 assessment of rooftop-farming technologies potential in schools in Quito, Ecuador. J.
 Clean. Prod. 122993. https://doi.org/10.1016/j.jclepro.2020.122993
- Liang, L., Ridoutt, B.G., Wu, W., Lal, R., Wang, L., Wang, Y., Li, C., Zhao, G., 2019. A multi-indicator assessment of peri-urban agricultural production in Beijing, China. Ecol. Indic. 97, 350–362. https://doi.org/10.1016/j.ecolind.2018.10.040
- Loiseau, E., Colin, M., Alaphilippe, A., Coste, G., Roux, P., 2020. To what extent are short food supply chains (SFSCs) environmentally friendly? Application to French apple distribution using Life Cycle Assessment. J. Clean. Prod. 276, 124166.
 https://doi.org/10.1016/j.jclepro.2020.124166
- Maia de Souza, D., Lopes, G.R., Hansson, J., Hansen, K., 2018. Ecosystem services in life cycle assessment: A synthesis of knowledge and recommendations for biofuels.

 Ecosyst. Serv., SI: Human-Nature nexuses 30, 200–210.
- 1112 https://doi.org/10.1016/j.ecoser.2018.02.014
- Majewski, E., Komerska, A., Kwiatkowski, J., Malak-Rawlikowska, A., Wąs, A., Sulewski, P., Gołaś, M., Pogodzińska, K., Lecoeur, J.-L., Tocco, B., Török, Á., Donati, M.,
- 1115 Vittersø, G., 2020. Are Short Food Supply Chains More Environmentally Sustainable
- than Long Chains? A Life Cycle Assessment (LCA) of the Eco-Efficiency of Food
- 1117 Chains in Selected EU Countries. Energies 13, 4853.
- 1118 https://doi.org/10.3390/en13184853

- Martin, G., Clift, R., Christie, I., 2016. Urban Cultivation and Its Contributions to
 Sustainability: Nibbles of Food but Oodles of Social Capital. Sustainability 8, 409.
 https://doi.org/10.3390/su8050409
- Martin, M., Molin, E., 2019. Environmental Assessment of an Urban Vertical Hydroponic Farming System in Sweden. Sustainability 11, 4124. https://doi.org/10.3390/su11154124
- Martin, M., Poulikidou, S., Molin, E., 2019. Exploring the Environmental Performance of
 Urban Symbiosis for Vertical Hydroponic Farming. Sustainability 11, 6724.
 https://doi.org/10.3390/su11236724
- Martinez, S., del Mar Delgado, M., Marin, R.M., Alvarez, S., 2018. The environmental footprint of an organic peri-urban orchard network. Sci. Total Environ. 636, 569–579. https://doi.org/10.1016/j.scitotenv.2018.04.340
- Martínez-Blanco, J., Colón, J., Gabarrell, X., Font, X., Sánchez, A., Artola, A., Rieradevall,
 J., 2010. The use of life cycle assessment for the comparison of biowaste composting
 at home and full scale. Waste Manag. 30, 983–994.
 https://doi.org/10.1016/j.wasman.2010.02.023
- Martínez-Blanco, J., Lazcano, C., Christensen, T.H., Muñoz, P., Rieradevall, J., Møller, J.,
 Antón, A., Boldrin, A., 2013. Compost benefits for agriculture evaluated by life cycle
 assessment. A review. Agron. Sustain. Dev. 33, 721–732.
 https://doi.org/10.1007/s13593-013-0148-7
- Martínez-Blanco, J., Muñoz, P., Antón, A., Rieradevall, J., 2009. Life cycle assessment of the use of compost from municipal organic waste for fertilization of tomato crops. Resour. Conserv. Recycl. 53, 340–351. https://doi.org/10.1016/j.resconrec.2009.02.003
- McClintock, N., Mahmoudi, D., Simpson, M., Santos, J.P., 2016. Socio-spatial differentiation
 in the Sustainable City: A mixed-methods assessment of residential gardens in
 metropolitan Portland, Oregon, USA. Landsc. Urban Plan. 148, 1–16.
 https://doi.org/10.1016/j.landurbplan.2015.12.008
- McDonald, R.I., Marcotullio, P.J., Güneralp, B., 2013. Chapter 3. Urbanization and Global
 Trends in Biodiversity and Ecosystem Services, in: Urbanization, Biodiversity and
 Ecosystem Services: Challenges 31 and Opportunities: A Global Assessment.
 Springer, Dordrecht, pp. 31–52.
- McDougall, R., Kristiansen, P., Rader, R., 2019. Small-scale urban agriculture results in high
 yields but requires judicious management of inputs to achieve sustainability. Proc.
 Natl. Acad. Sci. 116, 129–134. https://doi.org/10.1073/pnas.1809707115
- Mclaren, S.J., 2010. Life Cycle Assessment (LCA) of food production and processing: An
 introduction, in: Environmental Assessment and Management in the Food Industry:
 Life Cycle Assessment and Related Approaches, Food Science, Technology and
 Nutrition. Woodhead Publishing Limited, Cambridge, United Kingdom, pp. 37–56.
- Meier, M.S., Stoessel, F., Jungbluth, N., Juraske, R., Schader, C., Stolze, M., 2015.
 Environmental impacts of organic and conventional agricultural products Are the differences captured by life cycle assessment? J. Environ. Manage. 149, 193–208.
 https://doi.org/10.1016/j.jenvman.2014.10.006
- Melkonyan, A., Gruchmann, T., Lohmar, F., Kamath, V., Spinler, S., 2020. Sustainability assessment of last-mile logistics and distribution strategies: The case of local food networks. Int. J. Prod. Econ. 228, 107746. https://doi.org/10.1016/j.ijpe.2020.107746
- Millennium Ecosystem Assessment, 2005. Ecosystems and Human Well-being: Synthesis.
 Island Press, Washington, DC.
- Mohareb, E., Heller, M., Novak, P., Goldstein, B., Fonoll, X., Raskin, L., 2017.
 Considerations for reducing food system energy demand while scaling up urban agriculture. Environ. Res. Lett. 12, 125004. https://doi.org/10.1088/1748-9326/aa889b
- Mohareb, E.A., Heller, M.C., Guthrie, P.M., 2018. Cities' Role in Mitigating United States Food System Greenhouse Gas Emissions. Environ. Sci. Technol. 52, 5545–5554. https://doi.org/10.1021/acs.est.7b02600
- Montealegre, A.L., García-Pérez, S., Guillén-Lambea, S., Monzón-Chavarrías, M., Sierra-1173 Pérez, J., 2021. GIS-based assessment for the potential of implementation of food-

- energy-water systems on building rooftops at the urban level. Sci. Total Environ. 149963. https://doi.org/10.1016/j.scitotenv.2021.149963
- Mougeot, L.J.A., 2000. Urban Agriculture: Definition, Presence, Potentials and Risks, and
 Policy Challenges (No. 31), Cities Feeding People Series. International Development
 Research Centre (IDRC).
- Neilson, C., Rickards, L., 2017. The relational character of urban agriculture: competing perspectives on land, food, people, agriculture and the city. Geogr. J. 183, 295–306. https://doi.org/10.1111/geoj.12188
- Nemecek, T., Gaillard, G., 2010. Challenges in assessing the environmental impacts of crop production and horticulture, in: Environmental Assessment and Management in the Food Industry: Life Cycle Assessment and Related Approaches, Food Science, Technology and Nutrition. Woodhead Publishing Limited, Cambridge, United Kingdom, pp. 98–116.
- Nicese, F.P., Colangelo, G., Comolli, R., Azzini, L., Lucchetti, S., Marziliano, P.A., Sanesi,
 G., 2021. Estimating CO2 balance through the Life Cycle Assessment prism: A case –
 Study in an urban park. Urban Green. 57, 126869.
 https://doi.org/10.1016/j.ufug.2020.126869
- Nicholls, É., Ely, A., Birkin, L., Basu, P., Goulson, D., 2020. The contribution of small-scale food production in urban areas to the sustainable development goals: a review and case study. Sustain. Sci. 15, 1585–1599. https://doi.org/10.1007/s11625-020-00792-z
- Nordahl, S., Preble, C.V., Kirchstetter, T.W., Scown, C.D., 2022. Gaseous emissions from composting. Environ. Sci. Technol.
- Notarnicola, B., Sala, S., Anton, A., McLaren, S.J., Saouter, E., Sonesson, U., 2017. The role of life cycle assessment in supporting sustainable agri-food systems: A review of the challenges. J. Clean. Prod., Towards eco-efficient agriculture and food systems: selected papers addressing the global challenges for food systems, including those presented at the Conference "LCA for Feeding the planet and energy for life" (6-8 October 2015, Stresa & Milan Expo, Italy) 140, 399–409. https://doi.org/10.1016/j.jclepro.2016.06.071
- Notarnicola, B., Tassielli, G., Renzulli, P.A., 2012. Modeling the Agri-Food Industry with Life Cycle Assessment, in: Curran, M.A. (Ed.), Life Cycle Assessment Handbook. John Wiley & Sons, Ltd, pp. 159–183. https://doi.org/10.1002/9781118528372.ch7
- Orsini, F., Gasperi, D., Marchetti, L., Piovene, C., Draghetti, S., Ramazzotti, S., Bazzocchi,
 G., Gianquinto, G., 2014. Exploring the production capacity of rooftop gardens
 (RTGs) in urban agriculture: the potential impact on food and nutrition security,
 biodiversity and other ecosystem services in the city of Bologna. Food Secur. 6, 781–
 792. https://doi.org/10.1007/s12571-014-0389-6
- Orsini, F., Pennisi, G., Michelon, N., Minelli, A., Bazzocchi, G., Sanyé-Mengual, E.,
 Gianquinto, G., 2020. Features and Functions of Multifunctional Urban Agriculture in
 the Global North: A Review. Front. Sustain. Food Syst. 4.
 https://doi.org/10.3389/fsufs.2020.562513
- O'Sullivan, C.A., Bonnett, G.D., McIntyre, C.L., Hochman, Z., Wasson, A.P., 2019.
 Strategies to improve the productivity, product diversity and profitability of urban agriculture. Agric. Syst. 174, 133–144. https://doi.org/10.1016/j.agsy.2019.05.007
- Othoniel, B., Rugani, B., Heijungs, R., Benetto, E., Withagen, C., 2016. Assessment of Life
 Cycle Impacts on Ecosystem Services: Promise, Problems, and Prospects. Environ.
 Sci. Technol. 50, 1077–1092. https://doi.org/10.1021/acs.est.5b03706
- Peña, A., Rovira-Val, M.R., 2020. A longitudinal literature review of life cycle costing applied to urban agriculture. Int. J. Life Cycle Assess. 25, 1418–1435. https://doi.org/10.1007/s11367-020-01768-y
- Pennisi, G., Sanyé-Mengual, E., Orsini, F., Crepaldi, A., Nicola, S., Ochoa, J., Fernandez,
 J.A., Gianquinto, G., 2019. Modelling Environmental Burdens of Indoor-Grown
 Vegetables and Herbs as Affected by Red and Blue LED Lighting. Sustainability 11,

1227 4063. https://doi.org/10.3390/su11154063

- Pepin, A., 2022. Performance environnementale de fermes maraîchères en agriculture biologique. Thesis submitted for defense.
- Pérez-Neira, D., Grollmus-Venegas, A., 2018. Life-cycle energy assessment and carbon
 footprint of peri-urban horticulture. A comparative case study of local food systems in
 Spain. Landsc. Urban Plan. 172, 60–68.
 https://doi.org/10.1016/j.landurbplan.2018.01.001
- Pergola, M., Persiani, A., Pastore, V., Palese, A.M., D'Adamo, C., De Falco, E., Celano, G.,
 2020. Sustainability Assessment of the Green Compost Production Chain from
 Agricultural Waste: A Case Study in Southern Italy. Agronomy 10, 230.
 https://doi.org/10.3390/agronomy10020230
- Peri, G., Traverso, M., Finkbeiner, M., Rizzo, G., 2010. Un possibile aproccio "Social LCA" per le coperture a verde. Presented at the Ecomondo 2010 "Ambiente-Economia Nel cuore delle azioni."
- Perrin, A., Basset-Mens, C., Gabrielle, B., 2014. Life cycle assessment of vegetable products: a review focusing on cropping systems diversity and the estimation of field emissions. Int. J. Life Cycle Assess. 19, 1247–1263. https://doi.org/10.1007/s11367-014-0724-3
- Petit-Boix, A., Llorach-Massana, P., Sanjuan-Delmás, D., Sierra-Pérez, J., Vinyes, E.,
 Gabarrell, X., Rieradevall, J., Sanyé-Mengual, E., 2017. Application of life cycle
 thinking towards sustainable cities: A review. J. Clean. Prod. 166, 939–951.
 https://doi.org/10.1016/j.jclepro.2017.08.030
- Pollard, G., Ward, J., Roetman, P., 2018. Water Use Efficiency in Urban Food Gardens: Insights from a Systematic Review and Case Study. Horticulturae 4, 27. https://doi.org/10.3390/horticulturae4030027
- Poore, J., Nemecek, T., 2018. Reducing food's environmental impacts through producers and consumers. Science 360, 987–992. https://doi.org/10.1126/science.aaq0216
- Pourias, J., Aubry, C., Duchemin, E., 2016. Is food a motivation for urban gardeners?

 Multifunctionality and the relative importance of the food function in urban collective gardens of Paris and Montreal. Agric. Hum. Values 33, 257–273.

 https://doi.org/10.1007/s10460-015-9606-y
- Quirós, R., Villalba, G., Gabarrell, X., Muñoz, P., 2015. Life cycle assessment of organic and mineral fertilizers in a crop sequence of cauliflower and tomato. Int. J. Environ. Sci. Technol. 12, 3299–3316. https://doi.org/10.1007/s13762-015-0756-7
 - Quistberg, R.D., Bichier, P., Philpott, S.M., 2016. Landscape and Local Correlates of Bee Abundance and Species Richness in Urban Gardens. Environ. Entomol. 45, 592–601. https://doi.org/10.1093/ee/nvw025
- Romanovska, L., 2019. Urban green infrastructure: perspectives on life-cycle thinking for holistic assessments. IOP Conf. Ser. Earth Environ. Sci. 294, 012011. https://doi.org/10.1088/1755-1315/294/1/012011
- Romeo, D., Vea, E.B., Thomsen, M., 2018. Environmental Impacts of Urban Hydroponics in Europe: A Case Study in Lyon. Procedia CIRP, 25th CIRP Life Cycle Engineering (LCE) Conference, 30 April 2 May 2018, Copenhagen, Denmark 69, 540–545. https://doi.org/10.1016/j.procir.2017.11.048
- Rothwell, A., Ridoutt, B., Page, G., Bellotti, W., 2016. Environmental performance of local food: trade-offs and implications for climate resilience in a developed city. J. Clean. Prod. 114, 420–430. https://doi.org/10.1016/j.jclepro.2015.04.096
- Rufí-Salís, M., Petit-Boix, A., Villalba, G., Ercilla-Montserrat, M., Sanjuan-Delmás, D.,
 Parada, F., Arcas, V., Muñoz-Liesa, J., Gabarrell, X., 2020a. Identifying eco-efficient
 year-round crop combinations for rooftop greenhouse agriculture. Int. J. Life Cycle
 Assess. https://doi.org/10.1007/s11367-019-01724-5
- Rufí-Salís, M., Petit-Boix, A., Villalba, G., Sanjuan-Delmás, D., Parada, F., Ercilla-Montserrat, M., Arcas-Pilz, V., Muñoz-Liesa, J., Rieradevall, J., Gabarrell, X., 2020b.
- 1279 Recirculating water and nutrients in urban agriculture: An opportunity towards
- environmental sustainability and water use efficiency? J. Clean. Prod. 261, 121213.
- 1281 https://doi.org/10.1016/j.jclepro.2020.121213

- Saha, M., Eckelman, M.J., 2017. Growing fresh fruits and vegetables in an urban landscape:

 A geospatial assessment of ground level and rooftop urban agriculture potential in

 Boston, USA. Landsc. Urban Plan. 165, 130–141.

 https://doi.org/10.1016/j.landurbplan.2017.04.015
- Sanjuan-Delmás, D., Llorach-Massana, P., Nadal, A., Ercilla-Montserrat, M., Muñoz, P.,
 Montero, J.I., Josa, A., Gabarrell, X., Rieradevall, J., 2018. Environmental assessment
 of an integrated rooftop greenhouse for food production in cities. J. Clean. Prod. 177,
 326–337. https://doi.org/10.1016/j.jclepro.2017.12.147
- Sanyé-Mengual, E., Cerón-Palma, I., Oliver-Solà, J., Montero, J.I., Rieradevall, J., 2013.
 Environmental analysis of the logistics of agricultural products from roof top
 greenhouses in Mediterranean urban areas: Life cycle assessment of the logistics of
 agricultural products. J. Sci. Food Agric. 93, 100–109.
 https://doi.org/10.1002/jsfa.5736
- Sanyé-Mengual, E., Gasperi, D., Michelon, N., Orsini, F., Ponchia, G., Gianquinto, G., 2018a.
 Eco-Efficiency Assessment and Food Security Potential of Home Gardening: A Case
 Study in Padua, Italy. Sustainability 10, 2124. https://doi.org/10.3390/su10072124
- Sanyé-Mengual, E., Oliver-Solà, J., Montero, J.I., Rieradevall, J., 2015a. An environmental and economic life cycle assessment of rooftop greenhouse (RTG) implementation in Barcelona, Spain. Assessing new forms of urban agriculture from the greenhouse structure to the final product level. Int. J. Life Cycle Assess. 20, 350–366. https://doi.org/10.1007/s11367-014-0836-9
- Sanyé-Mengual, E., Oliver-Solà, J., Montero, J.I., Riverdall, J., 2017. The role of interdisciplinarity in evaluating the sustainability of urban rooftop agriculture. Future Food J. Food Agric. Soc. 5, 13.
- Sanyé-Mengual, E., Orsini, F., Gianquinto, G., 2018b. Revisiting the Sustainability Concept of Urban Food Production from a Stakeholders' Perspective. Sustainability 10, 2175. https://doi.org/10.3390/su10072175
- Sanyé-Mengual, E., Orsini, F., Oliver-Solà, J., Rieradevall, J., Montero, J.I., Gianquinto, G.,
 2015b. Techniques and crops for efficient rooftop gardens in Bologna, Italy. Agron.
 Sustain. Dev. 35, 1477–1488. https://doi.org/10.1007/s13593-015-0331-0
- Sanyé-Mengual, E., Specht, K., Vávra, J., Artmann, M., Orsini, F., Gianquinto, G., 2020.
 Ecosystem Services of Urban Agriculture: Perceptions of Project Leaders,
 Stakeholders and the General Public. Sustainability 12, 10446.
 https://doi.org/10.3390/su122410446
- Schrijvers, D.L., Loubet, P., Sonnemann, G., 2016. Developing a systematic framework for consistent allocation in LCA. Int. J. Life Cycle Assess. 21, 976–993. https://doi.org/10.1007/s11367-016-1063-3
- Searchinger, T., Heimlich, R., Houghton, R.A., Dong, F., Elobeid, A., Fabiosa, J., Tokgoz, S.,
 Hayes, D., Yu, T.-H., 2008. Use of U.S. Croplands for Biofuels Increases Greenhouse
 Gases Through Emissions from Land-Use Change. Science 319, 1238–1240.
 https://doi.org/10.1126/science.1151861
- Shiina, T., Hosokawa, D., Roy, P., Nakamura, N., Thammawong, M., Orikasa, T., 2011. Life cycle inventory analysis of leafy vegetables grown in two types of plant factories.

 Acta Hortic. 919, 115–122. https://doi.org/10.17660/ActaHortic.2011.919.14
- Smetana, S.M., Crittenden, J.C., 2014. Sustainable plants in urban parks: A life cycle analysis of traditional and alternative lawns in Georgia, USA. Landsc. Urban Plan. 122, 140–1328 151. https://doi.org/10.1016/j.landurbplan.2013.11.011
- Specht, K., Siebert, R., Hartmann, I., Freisinger, U.B., Sawicka, M., Werner, A., Thomaier,
 S., Henckel, D., Walk, H., Dierich, A., 2014. Urban agriculture of the future: an
 overview of sustainability aspects of food production in and on buildings. Agric. Hum.
 Values 31, 33–51. https://doi.org/10.1007/s10460-013-9448-4
- Stelwagen, R.E., Slegers, P.M., de Schutter, L., van Leeuwen, E.S., 2021. A bottom-up approach to model the environmental impact of the last-mile in an urban food-system. Sustain. Prod. Consum. 26, 958–970. https://doi.org/10.1016/j.spc.2020.12.039

- Stillitano, T., Spada, E., Iofrida, N., Falcone, G., De Luca, A.I., 2021. Sustainable Agri-Food
 Processes and Circular Economy Pathways in a Life Cycle Perspective: State of the
 Art of Applicative Research. Sustainability 13, 2472.
 https://doi.org/10.3390/su13052472
- Strohbach, M.W., Arnold, E., Haase, D., 2012. The carbon footprint of urban green space—A
 life cycle approach. Landsc. Urban Plan. 104, 220–229.
 https://doi.org/10.1016/j.landurbplan.2011.10.013
- Susca, T., Pomponi, F., 2020. Heat island effects in urban life cycle assessment: Novel insights to include the effects of the urban heat island and UHI-mitigation measures in LCA for effective policy making. J. Ind. Ecol. 24, 410–423. https://doi.org/10.1111/jiec.12980
- Tang, L., Hayashi, K., Kohyama, K., Leon, A., 2018. Reconciling Life Cycle Environmental
 Impacts with Ecosystem Services: A Management Perspective on Agricultural Land
 Use. Sustainability 10, 630. https://doi.org/10.3390/su10030630
- Taylor, J.R., 2020. Modeling the Potential Productivity of Urban Agriculture and Its Impacts
 on Soil Quality Through Experimental Research on Scale-Appropriate Systems. Front.
 Sustain. Food Syst. 4.
- Teixeira, R.F.M., Maia de Souza, D., Curran, M.P., Antón, A., Michelsen, O., Milà i Canals,
 L., 2016. Towards consensus on land use impacts on biodiversity in LCA:
 UNEP/SETAC Life Cycle Initiative preliminary recommendations based on expert
 contributions. J. Clean. Prod. 112, 4283–4287.
 https://doi.org/10.1016/j.jclepro.2015.07.118
- Tidåker, P., Wesström, T., Kätterer, T., 2017. Energy use and greenhouse gas emissions from turf management of two Swedish golf courses. Urban For. Urban Green. 21, 80–87. https://doi.org/10.1016/j.ufug.2016.11.009
- Tiefenbacher, A., Sandén, T., Haslmayr, H.-P., Miloczki, J., Wenzel, W., Spiegel, H., 2021.
 Optimizing Carbon Sequestration in Croplands: A Synthesis. Agronomy 11, 882.
 https://doi.org/10.3390/agronomy11050882
- Tonini, D., Wandl, A., Meister, K., Unceta, P.M., Taelman, S.E., Sanjuan-Delmás, D.,
 Dewulf, J., Huygens, D., 2020. Quantitative sustainability assessment of household
 food waste management in the Amsterdam Metropolitan Area. Resour. Conserv.
 Recycl. 160, 104854. https://doi.org/10.1016/j.resconrec.2020.104854
- Toniolo, S., Mazzi, A., Pieretto, C., Scipioni, A., 2017. Allocation strategies in comparative
 life cycle assessment for recycling: Considerations from case studies. Resour.
 Conserv. Recycl. 117, 249–261. https://doi.org/10.1016/j.resconrec.2016.10.011
- Torres Pineda, I., Cho, J.H., Lee, D., Lee, S.M., Yu, S., Lee, Y.D., 2020. Environmental Impact of Fresh Tomato Production in an Urban Rooftop Greenhouse in a Humid Continental Climate in South Korea. Sustainability 12, 9029. https://doi.org/10.3390/su12219029
- 1375 UNEP/SETAC, 2009. Guidelines for social life cycle assessment of products. United Nations Environment Programme, Paris, France.
- Vacek, P., Struhala, K., Matějka, L., 2017. Life-cycle study on semi intensive green roofs. J.
 Clean. Prod. 154, 203–213. https://doi.org/10.1016/j.jclepro.2017.03.188
- van der Werf, H.M.G., Knudsen, M.T., Cederberg, C., 2020. Towards better representation of
 organic agriculture in life cycle assessment. Nat. Sustain. 3, 419–425.
 https://doi.org/10.1038/s41893-020-0489-6
- Vaneeckhaute, C., Styles, D., Prade, T., Adams, P., Thelin, G., Rodhe, L., Gunnarsson, I.,
 D'Hertefeldt, T., 2018. Closing nutrient loops through decentralized anaerobic
 digestion of organic residues in agricultural regions: A multi-dimensional
 sustainability assessment. Resour. Conserv. Recycl. 136, 110–117.
 https://doi.org/10.1016/j.resconrec.2018.03.027
- Vieira, V.H.A. de M., Matheus, D.R., 2019. Environmental assessments of biological treatments of biowaste in life cycle perspective: A critical review. Waste Manag. Res. 37, 1183–1198. https://doi.org/10.1177/0734242X19879222

- Wagstaff, R.K., Wortman, S.E., 2015. Crop physiological response across the Chicago
 metropolitan region: Developing recommendations for urban and peri-urban farmers
 in the North Central US. Renew. Agric. Food Syst. 30, 8–14.
 https://doi.org/10.1017/S174217051300046X
- Weber, C.L., Matthews, H.S., 2008. Food-Miles and the Relative Climate Impacts of Food Choices in the United States. Environ. Sci. Technol. 42, 3508–3513. https://doi.org/10.1021/es702969f
- Weidema, B., 2000. Avoiding Co-Product Allocation in Life-Cycle Assessment. J. Ind. Ecol.
 4, 11–33. https://doi.org/10.1162/108819800300106366
- Weidner, T., Yang, A., 2020. The potential of urban agriculture in combination with organic waste valorization: Assessment of resource flows and emissions for two european cities. J. Clean. Prod. 244, 118490. https://doi.org/10.1016/j.jclepro.2019.118490
- Weidner, T., Yang, A., Hamm, M.W., 2019. Consolidating the current knowledge on urban
 agriculture in productive urban food systems: Learnings, gaps and outlook. J. Clean.
 Prod. 209, 1637–1655. https://doi.org/10.1016/j.jclepro.2018.11.004
- Wernet, G., Bauer, C., Steubing, B., Reinhard, J., Moreno-Ruiz, E., Weidema, B., 2016. The
 ecoinvent database version 3 (part I): overview and methodology. Int. J. Life Cycle
 Assess. 21, 1218–1230. https://doi.org/10.1007/s11367-016-1087-8
- Whittinghill, L., Sarr, S., 2021. Practices and Barriers to Sustainable Urban Agriculture: A
 Case Study of Louisville, Kentucky. Urban Sci. 5, 92.
 https://doi.org/10.3390/urbansci5040092
- Zhang, Y., Baral, A., Bakshi, B.R., 2010. Accounting for Ecosystem Services in Life Cycle
 Assessment, Part II: Toward an Ecologically Based LCA. Environ. Sci. Technol. 44,
 2624–2631. https://doi.org/10.1021/es900548a
- Zimek, M., Schober, A., Mair, C., Baumgartner, R.J., Stern, T., Füllsack, M., 2019. The Third
 Wave of LCA as the "Decade of Consolidation." Sustainability 11, 3283.
 https://doi.org/10.3390/su11123283