



HAL
open science

Recent Advances in Enteric Methane Mitigation and the Long Road to Sustainable Ruminant Production

Simon Roques, Gonzalo Martinez-Fernandez, Yulixaxis Ramayo-Caldas, Milka Popova, Stuart Denman, Sarah J Meale, Diego Morgavi

► **To cite this version:**

Simon Roques, Gonzalo Martinez-Fernandez, Yulixaxis Ramayo-Caldas, Milka Popova, Stuart Denman, et al.. Recent Advances in Enteric Methane Mitigation and the Long Road to Sustainable Ruminant Production. Annual Review of Animal Biosciences, 2024, 12, pp.321-343. <10.1146/annurev-animal-021022-024931>. <hal-04342398>

HAL Id: hal-04342398

<https://hal.inrae.fr/hal-04342398v1>

Submitted on 13 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Recent Advances in Enteric Methane Mitigation and the Long Road to Sustainable Ruminant Production

Simon Roques,¹ Gonzalo Martinez-Fernandez,²
Yuliaxis Ramayo-Caldas,³ Milka Popova,¹
Stuart Denman,² Sarah J. Meale,⁴ and Diego P. Morgavi¹

¹ Université Clermont Auvergne, INRAE, VetAgro Sup, UMR Herbivores, Saint-Genes-Champanelle, France; email: simon.roques@inrae.fr, milka.popova@inrae.fr, diego.morgavi@inrae.fr

² Agriculture and Food, CSIRO, St. Lucia, Queensland, Australia; email: gonzalo.martinezfernandez@csiro.au, stuart.denman@csiro.au

³ Animal Breeding and Genetics Program, Institute of Agrifood Research and Technology (IRTA), Torre Marimon, Caldes de Montbui, Spain; email: yuliaxis.ramayo@irta.cat

⁴ School of Agriculture and Food Sustainability, Faculty of Science, University of Queensland, Gatton, Queensland, Australia; email: s.meale@uq.edu.au

ANNUAL
REVIEWS **CONNECT**

www.annualreviews.org

- Download figures
- Navigate cited references
- Keyword search
- Explore related articles
- Share via email or social media


Annu. Rev. Anim. Biosci. 2024. 12:321–43

First published as a Review in Advance on
December 11, 2023

The *Annual Review of Animal Biosciences* is online at
animal.annualreviews.org

<https://doi.org/10.1146/annurev-animal-021022-024931>

Copyright © 2024 by the author(s). This work is licensed under a Creative Commons Attribution 4.0 International License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See credit lines of images or other third-party material in this article for license information.



Keywords

methane emission, livestock, global warming, feed additives, breeding, methanogens

Abstract

Mitigation of methane emission, a potent greenhouse gas, is a world-wide priority to limit global warming. A substantial part of anthropogenic methane is emitted by the livestock sector, as methane is a normal product of ruminant digestion. We present the latest developments and challenges ahead of the main efficient mitigation strategies of enteric methane production in ruminants. Numerous mitigation strategies have been developed in the last decades, from dietary manipulation and breeding to targeting of methanogens, the microbes that produce methane. The most recent advances focus on specific inhibition of key enzymes involved in methanogenesis. But these inhibitors, although efficient, are not affordable and not adapted to the extensive farming systems prevalent in low- and middle-income countries. Effective global mitigation of methane emissions from

livestock should be based not only on scientific progress but also on the feasibility and accessibility of mitigation strategies.

1. INTRODUCTION

The production of food for human consumption challenges the sustainability of resources at a planetary level (1). Livestock production, particularly ruminant production, is at risk of exceeding some planetary boundaries beyond natural recovery (2). One boundary at risk associated with the ruminant sector is climate change, where enteric methane emissions are particularly relevant for global warming (3).

Methane is a greenhouse gas (GHG) that is second only to carbon dioxide in its importance for global warming. Methane concentrations in the atmosphere have accelerated over the past 15 years to the point where this increase could jeopardize the ongoing efforts to reduce carbon dioxide emissions (4). Besides, the radiative forcing of methane, the greenhouse effect, is now estimated to be 25% higher (5) than the values used in international agreements, including the UN Paris Agreement on climate change. Nevertheless, methane has a short perturbation lifetime in the atmosphere of approximately 12 years. This advocates methane as a preferred target for mitigating global warming in the near future, as proposed by the Global Methane Pledge initiative (<https://www.globalmethanepledge.org>).

The share of livestock emissions in anthropogenic GHG emissions (11%), and especially the share of methane in these emissions (45%) (6), highlights the need to consider the impact of livestock on the environment when considering global warming and the GHG mitigation targets set by the UN Framework Convention on Climate Change. Whether the recent increase in atmospheric methane concentration is due to increased emissions, reduced sinks, or both is unclear (4). This increase is accompanied by a shift in the $^{13}\text{C}/^{12}\text{C}$ isotopic ratio ($\delta^{13}\text{C}_{\text{CH}_4}$), with an increase in negative $\delta^{13}\text{C}_{\text{CH}_4}$ values that are characteristic of methane from biogenic sources, including ruminants but also wetlands and waste (4). Notwithstanding the uncertainties mentioned above, recent adjusted estimates suggest that enteric methane from ruminants is a major contributor to the drop in atmospheric $\delta^{13}\text{C}_{\text{CH}_4}$ and to the increase in overall methane emissions observed since the beginning of the twenty-first century (7).

Although ruminants contribute substantially to GHG emissions, ruminant-derived food plays a pillar role in food security and human nutrition, and that should not be underestimated (8, 9). Meat and milk from ruminants are major sources of high-quality protein and essential minerals and vitamins for many people, particularly in developing countries (10). The carbon footprint of meat and milk protein from ruminants is higher than that of plant proteins, but when quality aspects such as the digestible essential amino acids score are taken into account, the carbon footprint decreases by up to 40% compared to unadjusted values (11). Another comparative advantage of ruminant production systems is that up to 90% of feeds consumed are not human edible, such as forages and co-/by-products of the feed industry (12). Recent research shows that proteins from ruminants have on average a smaller GHG footprint than proteins from monogastric animals, which is particularly evident when feed production from crops is considered and the use of human-nonedible feeds is maximized for ruminants (13). If ruminants are “walking wetlands” (4) from a global warming perspective, they are also walking biological converters of cellulosic biomass into food and other useful products. Therefore, the urgent need to reduce the environmental impact of ruminants is often considered from the perspective of, and possible trade-offs with, providing people with an adequate supply of nutritious food.

Strategies to decrease enteric methane emissions have been explored for several decades already, with a steady increase since the mid-2000s following the publication of the UN Food and Agriculture Organization (14) report *Livestock's Long Shadow*. Enteric methane production is a complex trait driven by the rumen microbiome (all microbes present in the rumen, the microbiota, and their genes) and modulated by multiple factors, including the diet, the ruminant animal, and their interactions (15, 16). This review considers these animal-related approaches to mitigate methane emissions from farmed ruminants, focusing on the most recent and promising developments in the area. We critically assess present and future challenges the sector faces in adopting these solutions.

2. METHANOGENESIS AND EMISSIONS METRICS

2.1. How Enteric Methane Is Produced

Methanogenesis, the formation of methane, is a normal process that occurs during feed digestion in the ruminant gastrointestinal tract. Enteric methanogenesis takes place mostly in the rumen due to its size and anterior position in the ruminant gastrointestinal tract. It is a process of microbial origin that occurs in the absence of oxygen, when carbohydrates from the feeds are fermented by a collection of microbes to produce energetic substrates used by the animal (17). In normal conditions, fermentation produces hydrogen, which does not accumulate as it is assumed it would inhibit fermentation (18, 19), although animal studies in which methane inhibition increased hydrogen concentration without affecting fermentation of feeds challenge this assumption (20–22). Instead, most of the hydrogen is used by methanogenic microbes to produce methane (19), whereas a small part is emitted through belching.

In the rumen, as in other bioenvironments, methanogenesis is ensured by the methanogenic archaeal community. These microbes represent 0.3–3.3% of the rumen microbiota and are found in the rumen fluid, attached to feed particles, or associated to protozoa (23). As a reaction, methanogenesis is a reduction (the transfer of an electron) of carbon dioxide by dihydrogen to methane and water. Methane produced in this way, referred as the hydrogenotrophic pathway, represents a large proportion of the methane formed in normal conditions (24) and is performed mostly by archaea belonging to the *Methanobrevibacter* genus. However, other methanogenic archaea can synthesize methane using different pathways depending on available substrates. Two other pathways have been described, the acetate pathway, which produces methane from acetate, and methyl-based pathways (including methyl dismutase and methyl-reducing pathways) that use methylated compounds as substrates (24, 25). Other unusual pathways have been described in the literature (26) but are not yet demonstrated as active in ruminants. As mentioned above, the hydrogenotrophic pathway is dominant in the rumen, with the characteristic that methane produced from formate, once inside methanogens and transformed to carbon dioxide and dihydrogen, can represent up to 18% of total ruminal methane (27). The acetate pathway is not important in the rumen (28, 29), but the methyl-based pathway, once considered minor due to the low number of rumen methanogens using the methyl dismutation (methylotrophic) pathway (29), can contribute a significant proportion of the methane emitted by ruminants (30). The main methylotrophic pathway in the rumen, however, uses hydrogen as the electron donor to reduce methyl compounds. *Methanosphaera* spp. and methanogens from the *Methanomassiliicoccales* order are the main representatives carrying out this process and account for approximately 30% of the total number of methanogens (29) when adjusted by 16S ribosomal RNA copy numbers.

2.2. A Note on Expression Metrics

Several metrics are commonly used to account for methane emissions and to assess the efficacy of mitigation strategies. These are absolute emissions (g/day), yield [g/kg dry matter intake (DMI)],

and intensity (g/kg of milk, meat). We focus on absolute emissions. Methane intensity, convenient for calculating the carbon footprint of animal products, is linked to production efficiency. Improving efficiency, particularly in low-performing systems, will improve resource use and food security in line with the UN Sustainable Development Goals but not necessarily reduce overall enteric methane emissions. Another metric used is the methane conversion factor (Y_m), expressed as the proportion of the gross energy in the ingested feed that is lost as methane. It is often referred as the energy that could be used for productive purposes if it were saved by decreasing methane production in the rumen. Methane inhibition seems not to improve the energy balance of ruminants (31). Methane yield is, however, useful to decouple methane emissions from feed intake, which is the first dietary factor to modulate methane production.

3. DIETARY STRATEGIES

Enteric methanogenesis cannot be dissociated from feeds and the digestive processes that occur in the gastrointestinal tract (32). It is thus not surprising that mitigation strategies targeting diet composition and feed quality have been reported widely in the literature and continue to be explored. Extensive overviews of available dietary options have been published, some recently, covering the most successful strategies (16, 33, 34). The mitigation potential of these strategies has been quantitatively assessed in a recent meta-analysis (35). Of note, some successful dietary strategies decrease methane intensity but not absolute emissions (35). Here we present the main concepts by which dietary strategies, excluding additives that specifically affect methanogenesis (presented in Section 5), modulate methanogenesis, and we discuss knowledge gaps and shortcomings of current claims.

Dietary strategies modulate methanogenesis in two ways: by influencing the production of substrates used by methanogens or by affecting the methanogens themselves (**Figure 1b,c**). These two ways can, and often do, coexist. The main factors involved in the effect of diet on the rumen ecosystem can be summarized as those affecting feed digestibility, fermentation products, and pH. These factors are interrelated and interact mutually. Diet digestibility is positively correlated with methane emissions (36). It follows that rumen digestibility is related to intake and passage rate as determinants of methane production. In short, intake increases when animals are fed more digestible diets with a concomitant increase in emissions, up to the point where the residence time of feed in the rumen is shortened, i.e., less fermentation, causing a decrease in methane produced per kilogram of DMI. Similarly, including in the diet feeds that are not fermented in the rumen—those that are naturally inert, such as lipids, or technologically treated to bypass the rumen—decreases emissions. The use of lipids in ruminant diets as a carbohydrate substitute effectively reduces absolute methane emissions in a dose-dependent manner (35, 37). No to minor effects on productive traits have been reported when dietary lipids are not overfed, although lipid efficacy depends on the type of fatty acids. Medium-chain 12:0 and 14:0 and unsaturated long-chain (C18 and higher) fatty acids are the most effective (38). Another lipid-induced mechanism contributing to methanogenesis reduction, which is related to fatty acid composition, is toxicity to sensitive microbial groups, including microbes contributing to dihydrogen production, such as cellulolytic bacteria and protozoa (39, 40). Maia et al. (39) showed that microbial growth is particularly affected in cellulolytic, hydrogen-producing *Ruminococcus albus*, *Ruminococcus flavefaciens*, and some *Butyrivibrio* spp. This negative effect on microbial growth depends on carbon chain length and degree of saturation, i.e., from more to less toxic eicosapentaenoic acid (20:5), docosahexaenoic acid (22:6), α -linolenic acid (18:3), and linoleic acid (39). Hydrogenation of unsaturated fatty acids also acts as an electron sink, decreasing the amount of substrate available for methanogenesis, but the effect is minor (41).




Figure 1

Main strategies and their mechanisms for reducing methane emissions in ruminants. (a) At the herd level, methane emissions are reduced by breeding for low emitters, either by selecting for low emitters or by selecting individuals based on their genomic information related to methane emission. (b) Methane emissions are reduced by several dietary strategies, by increasing digestibility and passage rates or even bypassing the rumen, to limit nutrient availability to microbes. (c) In the rumen, methane emissions are mitigated by strategies that inhibit hydrogen-producing microbes; shift the substrate of methanogenesis; or inhibit methanogens, the microbes that produce methane. (d) Targeted strategies reduce methane emissions by inhibiting key enzymes in the methanogenesis pathway or by directly affecting methanogens' viability. Abbreviations: CNSL, cashew nut shell liquid; 3-NOP, 3-nitrooxypropanol.

Reducing the forage-to-concentrate ratio is another way to modulate the production of substrates used by methanogens. Concentrates are rapidly fermentable carbohydrates that shift fermentation pathways toward the production of propionate, a major electron sink under normal conditions (25, 42). In addition, these rapidly fermentable carbohydrates lower the pH of the rumen fluid and therefore inhibit methanogen growth. Methanogens found in the rumen have an optimum pH of approximately 7 and are sensitive to moderate changes, especially lower pH values (43–45). In mixed rumen cultures, methanogenesis was minimal at pH 6 and completely inhibited at pH 5.5 (46). Although excessive concentrate feeding leads to subacute acidosis impairing animal production, rumen pH values <6 are usually observed in lactating dairy cows after feeding (47). Low pH has a direct effect on methanogenesis that is independent of other factors, such as passage rate or substrate type (48), that usually co-occur in the rumen. Janssen (18) reviewed the effects of pH and other co-occurring effects on methanogens. Chemical composition

of the feed and specifically of the carbohydrate fraction also modulates methane emission, which has led to numerous dietary strategies to mitigate methane emissions. For instance, the hydrogenotrophic CO₂-reducing pathway is associated with structural plant carbohydrates cellulose and hemicellulose, whereas the hydrogenotrophic methyl-reducing pathway is associated with other dietary components, such as pectin, carnitine, and choline (49). This duality in substrate dependence influences the effect of dietary strategies on emissions and, more generally, on methanogenesis persistence.

Plants used to feed ruminants also contain secondary metabolites useful for methane mitigation. Particularly, tannins, and essential oils extracted from plants, have been shown to reduce absolute methane production (50–54), but not in all situations, as there are contrasting results. Meta-analyses show modest methane reduction effects of ~10% or less (53, 54), whereas single studies can show significant effects; e.g., a ~30% decrease in emissions was observed in sheep fed tannin-containing forages leucaena and glyricidia (51). These differences may arise because both tannins and essential oils are plant secondary metabolites comprising a diverse group of compounds with different chemical structures and, in many cases, specific biological activity. Contributing to this is the variation found in plants and extracts in the concentration and composition of the mixture of bioactive compounds. Plant secondary metabolites can affect overall feed digestibility and fermentation via mixed mechanisms involving the ones cited above and can also inhibit the microbiota. Tannins reduce the availability of substrates for fermentation in the rumen; affect the colonization of forages by microbes; and have a direct effect on microbes, particularly the fibrolytic microbial community and methanogens (55, 56). The mode of action of essential oils is attributed to their antimicrobial activities (57), and methanogens are more sensitive to some essential oils than fermentative microbes (57, 58). A recent meta-analysis shows that some, but not all, specific mixtures of essential oils can decrease methane production and increase milk yield, but reported changes were small, and more research is needed (53).

All these dietary strategies may, however, prove insufficient or irrelevant for the worldwide majority of farmed ruminants that are raised on extensive and mixed grazing systems (12). In addition, producers, particularly in low- and middle-income countries (LMICs), may not have the economic and technical means to implement these strategies. In this case, grassland management could be an easier solution to implement. Options include use of improved cultivars (e.g., brown midrib maize, high-sugar grasses), grazing at earlier stages of grass maturity for improved digestibility, and legumes containing tannins and other secondary plant metabolites. However, the efficacy of forage and fresh legumes on methane mitigation in ruminants is debated due to the multiple co-factors that might affect their efficacy. These include factors such as the temperature differences in regions of cultivation, where certain climates tend to favor the growth of plants with a lower methane mitigation potential, or animal-related traits such as increased feed intake (59, 60).

All of these dietary strategies, including promoting intake, could decrease methane intensity by up to 18% (35). However, they generally do not decrease absolute methane emissions unless the number of ruminants is reduced. For instance, DMI is closely correlated with absolute methane emissions and production; thus, when methane is expressed per unit of animal product, such as energy-corrected milk, intensity is reduced (34). As intake increases and energy requirements for maintenance are met, more nutrients are available for production (61). Reducing the carbon footprint intensity of animal products is particularly relevant in LMICs, where production efficiency is still low and consumption of ruminant-based foods is expected to grow in the coming years (62). This is acceptable only if increased production efficiency contributes to food security and nutrition for vulnerable populations, but clearly it does not contribute to the long-term sustainability perspective of the global livestock sector. Considering the looming prospect of GHGs leading to rising temperatures, we stress the negative impact of increasing productivity without reducing the

number of ruminants. Importantly, concerted efforts from all stakeholders and the right policies are needed to avoid negative impacts on vulnerable populations in LMICs.

4. ANIMAL BREEDING AND GENETICS

The individual variation in enteric methane emissions that is observed in animals under the same feeding and management conditions suggests the possibility of breeding for low-emissions cattle and sheep (63–66). To breed low-emitting ruminants, animals with the genetic potential for lower enteric methane emissions must be identified and selected (**Figure 1a**). The heritability (h^2) of methane production, which quantifies the impact of genetic background on methane variations (0: no impact, 1: extreme impact), ranges from 0.13 to 0.29 in sheep (65, 67) and 0.11 to 0.45 in dairy cattle (68–71). However, because direct and accurate individual measurements of methane in a large number of animals are not always possible, the use of proxies for indirect selection or prediction of methane has been recommended. These proxies include traits associated with methane production, such as rumen morphology, feed efficiency, milk mid-infrared spectra, milk volatile fatty acids, signatures of host-associated gastrointestinal microbes, or feeding behavior traits (72–75). Among these, milk mid-infrared spectra, which are readily available from milk recording agencies, and milk volatile fatty acids have been studied extensively. Diet influences milk composition, and these proxies show a wide range of accuracy depending on diet composition (76) or statistical model (77). Predicting methane levels from efficiency-related traits, such as intake or residual feed intake, has also been proposed, but considerable variation has been reported with conflicting results, including a lack of association with methane emissions (63, 78–80). Therefore, further studies are needed that consider the variability due to external factors such as diet, production system, recording period, and season. Alternatives based on the use of residual methane metrics estimated from variables such as DMI, body weight, and milk yield have also been recommended. Residual methane emission is the difference between the actual and expected methane output (81). For example, Manzanilla-Pech et al. (64) showed in dairy cattle that residual methane adjusted by metabolic body weight and energy-corrected milk appears to be a more suitable metric for inclusion in breeding selection indices than methane production, methane yield, or methane intensity.

Another critical limitation to the implementation of breeding programs to reduce methane is the economic value associated with methane emissions. To the best of our knowledge, there are currently no benefits to farmers associated with reducing methane emissions. In fact, several authors agreed that setting the price of methane production in the breeding target too low may make it impractical to implement (66, 82, 83). Therefore, government support, policies, and regulations should be put in place to support the adoption of breeding as a mitigation strategy to reduce methane emissions.

In addition to traditional breeding programs, the use of genomic information offers alternatives for selecting low-emitting ruminants. Today, hundreds of thousands of genetic markers covering the entire genome can be genotyped at reasonable cost. Having established the genetic basis of a trait of interest, a logical next step is to explore its genetic architecture using genome-wide association studies. In dairy cattle, the polygenic nature of individual variability in methane production has been demonstrated, and quantitative trait loci, single-nucleotide polymorphisms, and candidate genes associated with methane traits have been reported (84–87). However, current studies have ignored genotype \times environment interactions, and some have used relatively small numbers of animals (<300). Furthermore, independent replication between studies is lacking; interestingly, however, most of the studies report overlaps between genomic intervals associated with methane emissions and quantitative trait loci associated with feed efficiency, milk production, and conformation traits. Therefore, before considering the inclusion of associated single-nucleotide

polymorphisms in breeding programs, additional research must be conducted to investigate the potential pleiotropic effects of the observed associations, because they may have unintended negative effects on productivity or farm profitability.

The genomic information can also be used to improve the accuracy of breeding value prediction through genomic selection. However, implementing genomic selection requires a reference population of individuals with genotypes and accurate phenotypic records. For low-heritability traits such as methane emission, many genotyped animals with low levels of relatedness and accurate measurements of methane are essential to ensure accurate predictions (66, 88). However, accurate measurement of methane based on the gold standard of respiration chambers in thousands of samples is prohibitively expensive and impractical in extensive conditions. Although alternative technologies are available to measure methane emissions, many have been tested only in intensive dairy systems and are not suitable for pastoral production systems. Two options for quantifying methane under grazing conditions are the sulfur hexafluoride tracer technique and the GreenFeed system. However, these systems can be expensive to install and maintain and require Internet access for data collection, which can be an issue for LMICs with limited resources and infrastructure. In addition, performing genomic analyses requires access to computational infrastructure and expertise in data analysis, quantitative genetics, and modeling (89).

Importantly, breeding for low methane emissions has several advantages over other methane mitigation strategies, including a permanent and cumulative effect over generations. It is also a practical choice for extensive pastoral ruminant production systems. These advantages should be considered when assessing the suitability of genetic selection as a methane mitigation strategy. As a proof of concept, a pioneering initiative in New Zealand has demonstrated the potential of genetic selection to reduce methane emissions. After 10 years of divergent selection in ewes, Rowe et al. (65) reported a 12% difference in methane yield between selected lines. This demonstrates an effective response to selection and host genetic control. Integrative multi-omics approaches also offer many novel opportunities. Recent studies have confirmed that methane production results from a joint contribution of the host genome and the ruminal microbiome (70, 75, 84, 90, 91), opening up the possibility of developing innovative hologenomic methods for mitigation. Although still in its early stages, promising research results indicate that microbial data can significantly improve phenotype predictions, regardless of whether some microbes are under direct genetic control by the host (92). A recent study in sheep confirmed that holobiont models, incorporating host genome and metagenome information, provide better predictive accuracy compared to genome-centric approaches (91). The authors also showed that a reference-free metagenome profile performs better than a metagenome profile restricted to a reference database, explaining in combination with the host genotype more than 70% of the variation in methane emissions and residual feed intake. However, the reliability of these estimates is not yet well established, microbial profiling remains costly, and which statistical approach is most effective for implementing hologenomic predictions is unclear. Further research is therefore needed with larger sample sizes, during key life events, and covering different environments.

In summary, the genetic basis of methane emissions in ruminants is still not understood fully, and further research is needed, particularly regarding the use of proxies to predict methane emissions. Similarly, a comprehensive understanding of host–microbiome interactions is critical to the successful implementation of novel hologenomic strategies to reduce enteric methane emissions. Although we acknowledge the promise of breeding to reduce methane emissions, we believe it is insufficient as a stand-alone approach. Therefore, this challenging journey from classical breeding to hologenomic-based methane mitigation strategies requires additional multidisciplinary research and collaborative efforts to effectively reduce the environmental impact of ruminant production systems.

5. MODULATION OF THE RUMEN MICROBIOTA AND FERMENTATION

Dietary and management strategies described in previous sections have an average mitigation effect at the herd level estimated at 18% (35), but globally this figure is much lower (<10%) given the adoption rates and economics associated with the changes (93). In addition, most of the strategies would reduce the intensity of ruminant-based products, but emissions would be reduced only if ruminant numbers were reduced. In contrast, strategies that specifically target methanogens and the methanogenic pathway (**Figure 1d**) are effective in reducing emissions, often with limited negative side effects on the rumen microbial ecosystem. These strategies take the form of feed additives, some of which are now commercially available in some countries. In this section, we present the recent development of such targeted strategies and discuss the technical knowledge gaps for expanding their use.

5.1. Inhibition of Methanogenesis by 3-Nitrooxypropanol

The ability of 3-nitrooxypropanol (3-NOP) to inhibit methane production in ruminants results from its highly specific microbial target, the active Ni(I) site of the methyl-coenzyme M reductase that catalyzes the last step of methanogenesis (94). As a result, the relationship between 3-NOP dose and methane inhibition appears to be linear (95), with an average reduction in daily methane emissions of 32.5% (96). Despite its known mode of action, its effects on animal performance, rumen fermentation, and microbiota vary with dose, animal type, and basal diet (97, 98). Interestingly, inhibition of the methanogen population in the rumen is not always realized, even when reductions in methane emissions are reported (99, 100). When feeding beef cattle a 90% forage diet with 3-NOP at 200 mg/kg dry matter, Gruninger et al. (101) reported minimal impact on the relative abundance of predominant bacteria and archaeal communities but observed a reduction in the *Bacillota:Bacteroidota* (homotypic synonym *Firmicutes:Bacteroidetes*) ratio. Conversely, a reduction in the total methanogen population was observed when 3-NOP was included at doses of 2–2.5 g/animal in beef cattle fed a tropical forage diet (102), indicating that further research on the effects of 3-NOP on the microbiome is needed.

The anti-methanogenic effect of 3-NOP is linked directly to its presence in the rumen, yet as a result of its highly soluble nature, its ruminal residence time is short-lived (94). Consequently, current use of 3-NOP is limited to intensive feeding systems with the capability to offer at least one and preferably two or more feeding bouts per day. In attempts to overcome this, slow-releasing formulations of 3-NOP are being investigated to prolong the time that 3-NOP is active in the rumen (103). Further, a stacking of additives, such as that seen by Gruninger et al. (101) when 3-NOP was combined with canola oil, suggests an additive effect on methane inhibition with a reduction in methane observed for 21 h after feeding. A stacking of additives could provide productivity gains not commonly seen with 3-NOP supplementation alone. Despite an increase in propionate concentration in the rumen with 3-NOP supplementation (96), the expected increase in animal performance associated with an increase in energy-dense volatile fatty acids has not yet been realized. The ultimate fate of excess H₂ remains to be understood, although it does not appear to be fully used by the main fermentation pathways and is instead expelled as gaseous H₂ (104).

Early-life administration of 3-NOP as a mechanism to program the rumen for lower methane emissions has shown considerable promise. Meale et al. (100) successfully intervened with 3-NOP supplementation from birth to 11 weeks of age to imprint a methane reduction still observed at one year of life. An intervention targeting crucial periods of development, including microbial establishment in the newborn and the weaning transition, is necessary to overcome the highly resilient nature of the mature ruminal microbiota and the strong host–microbiota interaction, which often forces the microbial population to revert back to its preintervention state (105). However,

such an intervention is labor intensive, as the method of administration must not induce the reflexive closure of the esophageal groove in neonates, as would normally occur if the animal is suckling milk.

5.2. Inhibition of Methanogenesis by Halogenated Compounds

Halogenated compounds, synthetic (e.g., bromochloromethane, chloroform, and bromoform) and naturally occurring (e.g., bromoform produced by marine algae species), are very effective in inhibiting methane production in ruminants. Halogenated methane analogs have a direct inhibitory effect on methanogenesis, mainly by binding with coenzyme M methyltransferase, thus inhibiting methyl transfer in methanogenesis (106). Recently, Glasson et al. (106) suggested that halogenated compounds can inhibit both coenzyme M methyltransferase and methyl coenzyme M reductase in vivo, although the exact extent of the inhibition of both pathways requires confirmation. Bromochloromethane supplementation to cattle, sheep, and goats decreased methane production between 30% and 91%, with an increase in expelled H₂ levels (when measured), showing no detrimental effect on DMI or ruminal fermentation (21, 107, 108). Chloroform supplementation also decreases methane production in ruminants (38–89%), while increasing the H₂ expelled by the animal (20, 102, 109). These studies showed no apparent detrimental effects on rumen fermentation or feed intakes in cattle fed forage alone or supplemented with concentrate. Regarding naturally occurring halogens, marine algae are important producers of halogenated, low-molecular weight compounds (such as brominated and chlorinated haloforms) as a defense mechanism against predators or environmental stressors (110). Red seaweed species (*Asparagopsis taxiformis* and *Asparagopsis armata*) contain high concentrations of bromoform and inhibit methanogenesis in vivo. In recent years, *Asparagopsis* has been tested in sheep (111), dairy cattle (112, 113), and beef cattle (22, 114), showing a linear decrease in methane production (9–98% reduction) with increases in H₂ expelled. Although most studies reported nonapparent detrimental effects on DMI or ruminal fermentation, some studies found a decrease in DMI and milk yield with high doses above 1% of *Asparagopsis* per kilogram of DMI (112, 113, 115), which could be also due to a greater concentration of iodine or other minerals. In addition, some studies reported an increase in productivity and feed conversion efficiency (22, 114); however, these findings must be considered with caution due to the experimental designs used and the small number of animals.

The use of synthetic or naturally occurring halogenated compounds could also have collateral effects, as they might inhibit the methyl transferases in other microorganisms (116). For instance, these compounds may affect reductive acetogenesis, as B₁₂-dependent methyl transferases play a key role in one-carbon metabolism in this process (117). Other considerations for the use of these compounds in livestock are the environmental impact (ozone depletion), effects on animal health, and potential residues in products for human consumption. Some studies and reviews (22, 106, 112) suggested that animal health and product residues are not compromised with the *Asparagopsis* levels used for inhibiting methanogenesis in vivo, and that contribution to ozone depletion is minimal relative to the total sources of anthropogenic bromine (106). However, further research (e.g., longer-term studies using greater numbers of animals) must be carried out to confirm these claims.

5.3. Inhibition of Methanogenesis by Surfactant Anacardic Acid

Cashew nut shell liquid (CNSL) is an abundant coproduct of cashew production. It has been used for broad applications ranging from the chemical industry to biological applications such as antimicrobials (118). Indeed, CNSL contains phenolic compounds with antimicrobial activities, namely, anacardic acid, cardanol, and cardol, with anacardic acid being the main one. The antimicrobial activity of anacardic acid was already advocated a few decades ago to reduce methane

produced in the rumen (119). Watanabe et al. (120) later confirmed the anti-methanogenic potential of raw CNSL. They demonstrated a reduction of methane production along a shift of rumen microbes toward propionate producers in three independent in vitro experiments, namely, a batch culture and a semicontinuous culture of rumen fluid experiment and a pure culture of selected bacteria. Anacardic acid exhibits a surfactant activity that disrupts the cell wall of some microbes. Methanogens are particularly sensitive to anacardic acid, requiring lower inhibitory concentrations than most bacteria (121). Methanogens lacking an external proteinaceous surface layer in their cell envelope, such as the genus *Methanobrevibacter*, are susceptible, whereas other methanogens, such as *Methanosarcina barkeri* and *Methanomicrobium mobile*, do present such a surface and are not disrupted by CNSL. However, there are exceptions in both cases (121), and other membrane components seem to play a role in the resistance to CNSL that needs to be elucidated. Note that the presence of pseudomurein in the sensitive genera *Methanobrevibacter* and *Methanobacterium*, which are positive to Gram staining, is replaced by protein subunits and heteropolysaccharides, respectively, in the resistant genera *Methanomicrobium* and *Methanosarcina* (122). The surfactant property of anacardic acid also disrupts Gram-positive bacteria (118). For instance, CNSL disrupted the cell walls of *R. flavefaciens* and *Butyrivibrio fibrisolvens*, which are fibrolytic bacteria producers of H₂, in an in vitro assay (123).

CNSL's effects on methane emission were also observed in Holstein cows fed CNSL, with up to a 38% decrease in emissions (CH₄/kg of DMI) (124). Further in vivo trials confirmed the reduction of methane emission by CNSL (125–127). Together, these studies revealed the mode of action of CNSL. The reduction in the number of hydrogen producers is compensated, at least in relative abundance, by succinate and propionate producers, including *Succinivibrionaceae* and *Prevotellaceae*. Moreover, these studies highlighted a decrease in the relative abundance of *Methanobrevibacter*, the main methanogen genus in the rumen, characterized by both 16S rRNA and *mcrA* gene sequencing.

Although promising, CNSL use is limited by the stability of its main antimicrobial compound, anacardic acid. Indeed, heating of CNSL, a common process to enrich CNSL in cardanol for industrial uses, decarboxylates anacardic acid, thus converting it to cardanol (128). Cardanol itself has much lower antimicrobial activity, and heated CNSL does not reduce methane emission in vitro or in vivo (120, 129). Thus, the chemical industry's demand for cardanol reduces the availability of raw CNSL for the livestock sector. A second minor limitation is the nonadditive effect of CNSL with another inhibitor of methanogenesis, although only one study assessed a combined effect with encapsulated nitrate (130).

5.4. Inhibition of Methanogenesis by Nitrate

Nitrate has been evaluated in several methane mitigation studies and has shown consistent and persistent reductions in enteric methane emissions (53, 131). Nitrate supplementation is assumed to lower methane emissions by decreasing hydrogen availability in the rumen. Screening of the 501 reference genomes of cultured rumen bacteria showed that the *Selenomonadales* and *Campylobacterales* clades encode enzymes involved in the reduction of nitrate (NO₃⁻) to ammonia (NH₃) (132). Furthermore, in vitro studies with pure cultures of *Selenomonas ruminantium* ssp. *lactilytica*, *Wolinella succinogenes*, and *Veillonella parvula* have shown these species to be active for nitrate and nitrite reduction (133). In sheep with a low methane-emitting phenotype, the expression level of ammonia-forming nitrite reductases increased along with hydrogenotrophic acetogenesis and fumarate reduction, demonstrating that non-methanogenic pathways can be effective hydrogen sinks (132). New evidence suggests nitrate could reduce methane emissions via direct inhibition of methanogens. Dietary nitrate supplementation in dairy cows increases the concentration of rumen dissolved hydrogen and expelled hydrogen (134). These results suggest nitrate's

toxic effect on rumen methanogens as hydrogen consumption is reduced. Importantly, a significant decrease in the number of methanogens was reported in sheep receiving nitrate supplementation (135). In steers, changes in absolute numbers were not significant (136), but methanogen diversity was affected. Similarly, *mcrA* gene expression was reduced after nitrate feeding to dairy cows, but no change in the number of methanogens was observed (137). Lastly, a recent dynamic mechanistic modeling approach supported the claim that the methane mitigation effect of nitrate supplementation was due to methanogen inhibition by nitrites rather than reduced hydrogen availability (138).

At a low level, as it is in forage crops, nitrate is not a cause for concern for ruminants. However, the higher doses of nitrate required to decrease methane production in ruminants will result in absorption of nitrate and nitrite into the blood through the rumen wall due to incomplete reduction to ammonia. Nitrite in the blood binds to hemoglobin and converts it to methemoglobin, a metalloprotein incapable of carrying oxygen. High levels of methemoglobin are associated with a range of clinical symptoms, reviewed in detail by Lee & Beauchemin (131). Nitrate supplementation may contribute to increased N emissions from manure as it is further reduced to ammonia in the rumen and excreted as urea in the urine (139). However, as a methane mitigator, nitrate could also be a lever for dietary N reformulation, as nitrate increased microbial nitrogen synthesis *in vitro* (140), suggesting that it could enhance microbial protein flux from the rumen. Nitrate supplementation is a promising strategy for reducing methane production. However, animal health concerns may prevent its adoption in practice, and possible emissions trade-offs must be considered. Nitrate encapsulation is a commercially available technical solution that might circumvent potential health problems, as it induces a slow release of nitrate in the rumen.

Strategies that target the methanogens, either by inhibiting key enzymes in methanogenesis or by acting on the methanogens themselves, are highly effective, reducing enteric emissions by up to 80%. Nevertheless, questions remain about the fate of the breakdown products of some additives in the animal and the environment, as mentioned for halogenated compounds, or the possible adverse effects on animal health of nitrite and other compounds that are absorbed into the bloodstream. Although information on the long-term effects of these strategies appears promising, the results require confirmation with larger studies and under different conditions. Finally, the combination of strategies is promising but needs to be evaluated thoroughly both in terms of the real cumulative effect and in terms of market availability and economic constraints.

6. TECHNOLOGIES IN DEVELOPMENT

A much-hoped-for strategy in the field of enteric methane mitigation is the use of vaccines against rumen methanogens. The prospect of reducing emissions by stimulating the immunological response of the host ruminant is attractive because of its potential applicability under most farming conditions, including pastoral systems. The concept of modulating populations of specific rumen microbes by producing antibodies delivered via saliva after immunization was first demonstrated in the 1990s for ciliate protozoa and then for bacteria associated with rumen acidosis in cattle and sheep (141–143). Research into the approach's application to rumen methanogens soon followed (144, 145). However, the technology is challenging, and several hurdles to its application remain. Over the past 20 years, important advances have been made in several key fundamental areas needed to develop an effective vaccine. These include improved understanding and genomic information on rumen methanogens, which is essential for identifying potential antigens, as well as the demonstration of specific antibodies in blood, saliva, and rumen contents following immunization. Recently, the cross-reactivity of antibodies between various abundant species of the genus *Methanobrevibacter* has been shown (146). Cross-reactivity or the use of antigens

common to most rumen methanogens appears to be necessary to maintain the efficacy of produced antibodies in the long term and to avoid the replacement of targeted species by non-sensitive species (147). Studies on vaccines for methane mitigation were reviewed recently (148), summarizing advances in technology regarding antigen selection and adjuvant use, but also highlighting gaps in knowledge. Key findings of the systematic review (148) are the relatively limited number of publications and the lack of consistent evidence that enteric methane emissions can be reduced *in vivo*. Therefore, based on this publicly available information, the approach has not moved up the Technology Readiness Level ladder to be considered as a medium-term alternative for the ruminant sector. Potential emissions reductions from this technology have been assumed in some scenarios (149). However, based on published evidence, it is not yet possible to predict whether methane mitigation vaccines will be used in the field.

A novel approach being tested is the capture of exhaled methane from the animal. Different from other strategies that decrease enteric methane production by altering the ruminal environment or interfering with the methanogenic pathway, this proprietary technology uses a catalytic mechanism that converts exhaled methane into CO₂ and water. The portable catalyzer is positioned close to the nostrils via a halter. The developers' initial claims indicate a reduction potential of up to 50%, with no negative effect of the halter on animal behavior and welfare. Expected scientific publications in peer-reviewed journals should provide more information on the potential impact of this technology (150).

Novel slow-release technologies (such as rumen-bolus systems) are under development currently. These systems will allow release of anti-methanogenic compounds (e.g., bromoform, 3-NOP) at a constant rate in the rumen for prolonged periods of time. This is of particular interest in extensive grazing systems (such as the subtropical rangelands in Australia), where livestock consume less frequent and significantly lower amounts of supplements due to environmental conditions and availability of quality pastures during the wet season (3–5 months per year). Also, this technology could maximize the inhibition effect of the compounds, as they will be released continually in the rumen.

Various approaches at different stages of development are reported regularly in the scientific literature. It is beyond the scope of this review to list all these strategies (for comprehensive lists, see 33, 53, 54). However, some of these approaches have been studied for several years but have not progressed to higher Technology Readiness Levels and, if successful, may be available for field use only in the medium to long term. For instance, the use of homoacetogens as direct-fed microbials (DFMs) to decrease rumen methane emissions has been largely explored. The concept is attractive but has not been successful until now because the partial pressure of H₂ in the rumen is normally lower than the threshold for reductive acetogenesis, with homoacetogens having a lower H₂ affinity constant (K_s) compared to methanogens (15, 151). Other possible innovations in DFMs aimed at reducing methane production include the selection of lactic acid bacteria that produce methanogen-specific bacteriocins (152) and the use of sulfate-reducing bacteria that can oxidize hydrogen sulfide, thus avoiding the negative effects of this gas on animal performance and health (153). In the latter case, the DFMs must be combined with a sulfate-containing feed ingredient or additive. However, all these approaches have little chance of success in the field unless new DFMs are discovered or there is a breakthrough in the biology of these microbes, i.e., homoacetogens that compete with methanogens in the rumen. Expected emissions reductions from these innovations cannot be estimated at present.

A decade ago, Leng (154) reported that biochar, a charcoal product from organic matter pyrolysis, decreased enteric methane emissions and hypothesized that biochar's increased surface area would favor biofilm formation and methane oxidation. This has led to numerous research projects, but overall, the effects reported *in vivo* are variable, limited, and often contradictory,

although this might also be due to inaccurate ways to measure methane emission (155, 156). A recent report found that not all biochars can decrease enteric methane emissions in ruminants, and the effectiveness might depend on several factors, such as biomass type, pre- and post-pyrolysis manipulation, and compounds in the biochar (157). The *in vivo* methane reduction of the biochars and doses tested was less than 12% under controlled feeding conditions, and no significant effect was observed under grazing conditions (157). Further research is required to identify a biochar with much greater anti-methanogenic properties to be viable as feed supplement.

7. CHALLENGES AHEAD

The UN goal of achieving net zero emissions by 2050 to limit the rise in global temperatures will require a contribution from all sectors. Agriculture is no exception, and ruminant production, with its recognized high environmental impact, is under pressure to be sustainable. However, sustainability also rests on social and economic pillars that must be fulfilled for long-term solutions. Ruminant livestock also contribute substantially to UN Sustainable Development Goal 2, Zero Hunger, through their contribution to food security and improved nutrition and livelihoods, particularly for vulnerable populations in LMICs. Reducing emissions to net zero is a major challenge for the sector, compounded by global warming's increasing impact on animals and resources. Droughts and rising temperatures are already affecting forage and crop production, with tropical and subtropical regions with rain-fed pastoral and mixed systems most at risk (158). The estimated reduction in livestock production by the end of the century ranges from 4% to 10%, depending on the GHG emission pathway chosen (159). This figure, calculated from the reduction in DMI due to heat stress, is rather conservative, as the effects of heat stress on reproduction and health (158) were not considered (159). Breeding for heat-tolerant animals is one of the adaptation strategies the sector is exploring. These breeding programs should consider incorporating methane emission traits into selection indices to avoid unintended selection for high methane emitters.

Several aspects link enteric methane emissions and global warming. Higher ambient temperature decreases DMI and, consequently, enteric methane emissions. Concurrently, under heat stress, there is a decrease in the acetate-to-propionate ratio, along with an increase in lactate and a lower pH in the rumen (160). As described in previous sections of this review, all these factors contribute to lower methane emissions per animal, but for these poorly performing animals, the methane intensity of the meat and milk produced is much higher. However, temperature's effect on methane does not appear to be linear. In crossbred cattle adapted to tropical conditions, increases in temperature from 25°C to 35°C decreased methane emissions, but at 40°C, methane output was the same as at 25°C, whereas DMI continued to be affected (161). Why this happens is unclear, but if confirmed, it will have implications for overall emissions in hot regions and temperate regions now becoming warmer. A related aspect is that underfed ruminants, a common situation during the dry season in tropical areas, have higher emission yields (162).

Most farmed ruminants, 75% of large ruminants and 85% of small ruminants, come from grazing and mixed crop-livestock systems (163), with the highest concentration in tropical and subtropical regions (**Supplemental Figure 1**). The trend of increasing numbers of ruminants from LMICs in these regions has not abated since global data became available in the early 1960s (<https://www.fao.org/faostat/>). In contrast, ruminant numbers in high-income countries, mostly in temperate regions, have not increased or have even decreased (**Supplemental Figure 1**). Improvements in production efficiency have led to reductions in ruminant numbers and methane emissions, whereas milk and meat production has increased [e.g., for Germany and the United States (61, 164)]. Mitigation options (available and in development) can be applied to the intensified production system prevalent in temperate regions but are more difficult to apply in grazing

Supplemental Material >

and extensive systems prevalent in Africa, Australia, and South America. In addition, in some regions, such as dry tropics rangelands in northern Australia, animals graze in very extensive areas with little human interaction, in an environment dominated by a short wet season and a long dry season, making it difficult to deploy feed supplements or other methane abatement strategies consistently through the whole year. Other critical challenges, particularly for grazing systems, are that anti-methanogenic compounds must be continuously available in the rumen to be effective; thus, the timing, frequency, and quantity consumed are critical to maximize methane inhibition. Therefore, novel ways of delivering these strategies to cattle under these conditions must be developed and evaluated.

For global impact, novel strategies, including delivery methods, applicable to grazing and less-intensive mixed systems are needed. This is because the trajectory of intensification observed in high-income countries cannot be followed in LMICs due to the current and future constraints of global warming on production outlined above, as well as economic and infrastructure limitations. Intensification based on concentrate feeds also would not be recommended, given food competition with human feed production. Globally, increased energy costs render the use of cereals from arable land to feed ruminants less competitive, which naturally leads to more grazing and fewer animals (165).

8. CONCLUDING REMARKS

Recent years have seen significant progress in understanding and reducing enteric methane emissions in ruminants. Whereas dietary and breeding strategies have been shown to be effective, their impact on global emissions is limited. Research on and expansion of these strategies should continue, as no single solution can be applied to all production systems. The improved production efficiency achieved by these strategies decreases methane intensity and has allowed some regions of the world to reduce herd size and total methane emissions while increasing milk and meat production. However, the global increase in ruminant livestock numbers offsets these regional improvements.

Inhibition of enteric methane by targeting the microbiota is the most successful strategy with potential for high impact. The most promising and novel approaches are in this area, but technical hurdles for long-term delivery and efficacy must be overcome to have a global impact. Importantly, the cost of treatment is not offset by increased production and, if not offset by other mechanisms, will effectively prohibit its global adoption. Ruminant methane emission is a global problem. Part of the solution is to reduce enteric methane production, as highlighted in this review. Progress in this area is encouraging, but a massive research effort is needed to find solutions that can be applied globally and have a lasting positive effect. For this to happen, the right funding, policies, and government interventions must be in place to ensure access and improve equity.

DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

ACKNOWLEDGMENTS

We apologize for not being able to cite many excellent studies due to space limitations. We acknowledge the support by the EU Horizon 2020 research and innovation program under grant agreements 818368 (MASTER) and 101000213 (HoloRuminant). Y.R.-C. was financially supported by a Ramon y Cajal contract (RYC2019-027244-I) from the Spanish Ministry of Science and Innovation. Julien Marcetteau is credited for the graphic design of **Figure 1**.

LITERATURE CITED

1. Steffen W, Richardson K, Rockstrom J, Cornell SE, Fetzer I, et al. 2015. Planetary boundaries: guiding human development on a changing planet. *Science* 347:1259855
2. Bowles N, Alexander S, Hadjikakou M. 2019. The livestock sector and planetary boundaries: a 'limits to growth' perspective with dietary implications. *Ecol. Econ.* 160:128–36
3. Gerber PJ, Steinfeld H, Henderson B, Mottet A, Opio C, et al. 2013. *Tackling Climate Change Through Livestock – A Global Assessment of Emissions and Mitigation Opportunities*. Rome: Food Agric. Organ.
4. Nisbet EG, Manning MR, Dlugokencky EJ, Fisher RE, Lowry D, et al. 2019. Very strong atmospheric methane growth in the 4 years 2014–2017: implications for the Paris agreement. *Glob. Biogeochem. Cycles* 33:318–42
5. Etminan M, Myhre G, Highwood EJ, Shine KP. 2016. Radiative forcing of carbon dioxide, methane, and nitrous oxide: a significant revision of the methane radiative forcing. *Geophys. Res. Lett.* 43:12614–23
6. Food Agric. Organ. 2022. *Gleam v3 dashboard*. Shiny Apps, accessed Febr. 2023. https://foodandagricultureorganization.shinyapps.io/GLEAMV3_Public/
7. Chang J, Peng S, Ciaï P, Saunio M, Dungal SRS, et al. 2019. Revisiting enteric methane emissions from domestic ruminants and their $\delta^{13}\text{C}_{\text{CH}_4}$ source signature. *Nat. Commun.* 10:3420
8. Food Agric. Organ. 2020. *World Food and Agriculture: Statistical Yearbook 2020*. Rome: Food Agric. Organ.
9. Herrero M, Grace D, Njuki J, Johnson N, Enahoro D, et al. 2013. The roles of livestock in developing countries. *Animal* 7:3–18
10. Randolph TF, Schelling E, Grace D, Nicholson CF, Leroy JL, et al. 2007. Invited review: role of livestock in human nutrition and health for poverty reduction in developing countries. *J. Anim. Sci.* 85:2788–800
11. McAuliffe GA, Takahashi T, Beal T, Huppertz T, Leroy F, et al. 2023. Protein quality as a complementary functional unit in life cycle assessment (LCA). *Int. J. Life Cycle Assess.* 28:146–55
12. Mottet A, de Haan C, Falcucci A, Tempio G, Opio C, Gerber P. 2017. Livestock: On our plates or eating at our table? A new analysis of the feed/food debate. *Glob. Food Secur.* 14:1–8
13. Cheng L, Zhang X, Reis S, Ren C, Xu J, Gu B. 2022. A 12% switch from monogastric to ruminant livestock production can reduce emissions and boost crop production for 525 million people. *Nat. Food* 3:1040–51
14. Steinfeld H, Gerber P, Wassenaar T, Castel V, Rosales M, de Haan C. 2006. *Livestock's long shadow: environmental issues and options*. Rep., Food Agric. Organ., Rome
15. Morgavi DP, Forano E, Martin C, Newbold CJ. 2010. Microbial ecosystem and methanogenesis in ruminants. *Animal* 4:1024–36
16. Beauchemin KA, McAllister TA, McGinn SM. 2009. Dietary mitigation of enteric methane from cattle. *CAB Rev.* 4(035):1–18
17. Deusch S, Camarinha-Silva A, Conrad J, Beifuss U, Rodehutsord M, Seifert J. 2017. A structural and functional elucidation of the rumen microbiome influenced by various diets and microenvironments. *Front. Microbiol.* 8:1605
18. Janssen PH. 2010. Influence of hydrogen on rumen methane formation and fermentation balances through microbial growth kinetics and fermentation thermodynamics. *Anim. Feed Sci. Technol.* 160:1–22
19. Ungerfeld EM. 2020. Metabolic hydrogen flows in rumen fermentation: principles and possibilities of interventions. *Front. Microbiol.* 11:589
20. Martinez-Fernandez G, Denman SE, Yang C, Cheung J, Mitsumori M, McSweeney CS. 2016. Methane inhibition alters the microbial community, hydrogen flow and fermentation response in the rumen of cattle. *Front. Microbiol.* 7:1122
21. Mitsumori M, Shinkai T, Takenaka A, Enishi O, Higuchi K, et al. 2012. Responses in digestion, rumen fermentation and microbial populations to inhibition of methane formation by a halogenated methane analogue. *Br. J. Nutr.* 108:482–91
22. Kinley RD, Martinez-Fernandez G, Matthews MK, de Nys R, Magnusson M, Tomkins NW. 2020. Mitigating the carbon footprint and improving productivity of ruminant livestock agriculture using a red seaweed. *J. Clean. Prod.* 259:120836
23. Janssen PH, Kirs M. 2008. Structure of the archaeal community of the rumen. *Appl. Environ. Microbiol.* 74:3619–25

24. Tapio I, Snelling TJ, Strozzi F, Wallace RJ. 2017. The ruminal microbiome associated with methane emissions from ruminant livestock. *J. Anim. Sci. Biotechnol.* 8:7
25. Bueno de Mesquita CP, Wu D, Tringe SG. 2023. Methyl-based methanogenesis: an ecological and genomic review. *Microbiol. Mol. Biol. Rev.* 87(1):e00024-22
26. Kurth JM, Op den Camp HJM, Welte CU. 2020. Several ways one goal—methanogenesis from unconventional substrates. *Appl. Microbiol. Biotechnol.* 104:6839–54
27. Hungate RE, Smith W, Bauchop T, Yu I, Rabinowitz JC. 1970. Formate as an intermediate in the bovine rumen fermentation. *J. Bacteriol.* 102:389–97
28. Oppermann RA, Nelson WO, Brown RE. 1961. In vivo studies of methanogenesis in the bovine rumen: dissimilation of acetate. *J. Gen. Microbiol.* 25:103–11
29. Henderson G, Cox F, Ganesh S, Jonker A, Young W, et al. 2015. Rumen microbial community composition varies with diet and host, but a core microbiome is found across a wide geographical range. *Sci. Rep.* 5:14567
30. Poulsen M, Schwab C, Jensen BB, Engberg RM, Spang A, et al. 2013. Methylophilic methanogenic thermoplasmata implicated in reduced methane emissions from bovine rumen. *Nat. Commun.* 4:1428
31. Morgavi DP, Cantalapiedra-Hijar G, Eugène M, Martin C, Nozière P, et al. 2023. Review: Reducing enteric methane emissions improves energy metabolism in livestock: Is the tenet right? *Animal* 17:100830
32. Blaxter KL, Clapperton JL. 1965. Prediction of the amount of methane produced by ruminants. *Br. J. Nutr.* 19:511–22
33. Beauchemin KA, Ungerfeld EM, Abdalla AL, Alvarez C, Arndt C, et al. 2022. Invited review: current enteric methane mitigation options. *J. Dairy Sci.* 105:9297–326
34. Knapp JR, Laur GL, Vadas PA, Weiss WP, Tricarico JM. 2014. Invited review: enteric methane in dairy cattle production: quantifying the opportunities and impact of reducing emissions. *J. Dairy Sci.* 97:3231–61
35. Arndt C, Hristov AN, Price WJ, McClelland SC, Pelaez AM, et al. 2022. Full adoption of the most effective strategies to mitigate methane emissions by ruminants can help meet the 1.5°C target by 2030 but not 2050. *PNAS* 119:e2111294119
36. Ramin M, Huhtanen P. 2013. Development of equations for predicting methane emissions from ruminants. *J. Dairy Sci.* 96:2476–93
37. Eugène M, Masse D, Chiquette J, Benchaar C. 2008. Meta-analysis on the effects of lipid supplementation on methane production in lactating dairy cows. *Can. J. Anim. Sci.* 88:331–34
38. Patra AK. 2013. The effect of dietary fats on methane emissions, and its other effects on digestibility, rumen fermentation and lactation performance in cattle: a meta-analysis. *Livest. Sci.* 155:244–54
39. Maia MRG, Chaudhary LC, Figueres L, Wallace RJ. 2007. Metabolism of polyunsaturated fatty acids and their toxicity to the microflora of the rumen. *Antonie Van Leeuwenhoek* 91:303–14
40. Machmüller A, Soliva CR, Kreuzer M. 2003. Effect of coconut oil and defaunation treatment on methanogenesis in sheep. *Reprod. Nutr. Dev.* 43:41–55
41. Martin C, Morgavi DP, Doreau M. 2010. Methane mitigation in ruminants: from microbe to the farm scale. *Animal* 4:351–65
42. Wang K, Xiong B, Zhao X. 2023. Could propionate formation be used to reduce enteric methane emission in ruminants? *Sci. Total Environ.* 855:158867
43. Smith PH, Hungate RE. 1958. Isolation and characterization of *Methanobacterium-Ruminantium* N. Sp. *J. Bacteriol.* 75:713–18
44. Miller TL, Lin CZ. 2002. Description of *Methanobrevibacter gottschalkii* sp nov., *Methanobrevibacter thaueri* sp nov., *Methanobrevibacter woesei* sp nov. and *Methanobrevibacter wolinii* sp nov. *Int. J. Syst. Evol. Microbiol.* 52:819–22
45. Miller TL, Wolin MJ. 1985. *Methanosphaera stadtmaniae* gen. nov., sp. nov.: a species that forms methane by reducing methanol with hydrogen. *Arch. Microbiol.* 141:116–22
46. van Kessel JAS, Russell JB. 1996. The effect of pH on ruminal methanogenesis. *FEMS Microbiol. Ecol.* 20:205–10
47. Villot C, Meunier B, Bodin J, Martin C, Silberberg M. 2018. Relative reticulo-rumen pH indicators for subacute ruminal acidosis detection in dairy cows. *Animal* 12:481–90

48. Kim IS, Hwang MH, Jang NJ, Hyun SH, Lee ST. 2004. Effect of low pH on the activity of hydrogen utilizing methanogen in bio-hydrogen process. *Int. J. Hydrogen Energy* 29:1133–40
49. Thomas CM, Desmond-Le Quéméner E, Gribaldo S, Borrel G. 2022. Factors shaping the abundance and diversity of the gut archaeome across the animal kingdom. *Nat. Commun.* 13:3358
50. Belanche A, Newbold CJ, Morgavi DP, Bach A, Zweifel B, Yáñez-Ruiz DR. 2020. A meta-analysis describing the effects of the essential oils blend agolin ruminant on performance, rumen fermentation and methane emissions in dairy cows. *Animals* 10:620
51. Jayanegara A, Leiber F, Kreuzer M. 2012. Meta-analysis of the relationship between dietary tannin level and methane formation in ruminants from *in vivo* and *in vitro* experiments. *J. Anim. Physiol. Anim. Nutr.* 96:365–75
52. Archimède H, Rira M, Barde DJ, Labirin F, Marie-Magdeleine C, et al. 2016. Potential of tannin-rich plants, *Leucaena leucocephala*, *Glyricidia sepium* and *Manihot esculenta*, to reduce enteric methane emissions in sheep. *J. Anim. Physiol. Anim. Nutr.* 100:1149–58
53. Hegarty R, Cortez Passetti R, Dittmer K, Wang Y, Shelton S, et al. 2021. *An evaluation of emerging feed additives to reduce methane emissions from livestock*. Rep., Clim. Change Agric. Food Secur., N.Z. Agric. Greenh. Gas Res. Cent., Glob. Res. Alliance. <https://hdl.handle.net/10568/116489>
54. Almeida AK, Hegarty RS, Cowie A. 2021. Meta-analysis quantifying the potential of dietary additives and rumen modifiers for methane mitigation in ruminant production systems. *Anim. Nutr.* 7:1219–30
55. Frutos P, Hervás G, Giráldez FJ, Mantecón AR. 2004. Review. Tannins and ruminant nutrition. *Span. J. Agric. Res.* 2:191–202
56. Rira M, Morgavi DP, Popova M, Maxin G, Doreau M. 2022. Microbial colonisation of tannin-rich tropical plants: interplay between degradability, methane production and tannin disappearance in the rumen. *Animal* 16:100589
57. Benchaar C, Greathead H. 2011. Essential oils and opportunities to mitigate enteric methane emissions from ruminants. *Anim. Feed Sci. Technol.* 166–67:338–55
58. Macheboeuf D, Morgavi DP, Papon Y, Mousset JL, Arturo-Schaan M. 2008. Dose-response effects of essential oils on *in vitro* fermentation activity of the rumen microbial population. *Anim. Feed Sci. Technol.* 145:335–50
59. Lagrange S, Beauchemin KA, MacAdam J, Villalba JJ. 2020. Grazing diverse combinations of tanniferous and non-tanniferous legumes: implications for beef cattle performance and environmental impact. *Sci. Total Environ.* 746:140788
60. Archimède H, Eugène M, Marie Magdeleine C, Boval M, Martin C, et al. 2011. Comparison of methane production between C3 and C4 grasses and legumes. *Anim. Feed Sci. Technol.* 166–67:59–64
61. Capper JL, Cady RA, Bauman DE. 2009. The environmental impact of dairy production: 1944 compared with 2007. *J. Anim. Sci.* 87:2160–67
62. OECD. 2022. *OECD-FAO Agricultural Outlook 2022–2031*. Paris: OECD Publ. <https://doi.org/10.1787/f1b0b29c-en>
63. Lassen J, Difford GF. 2020. Review: genetic and genomic selection as a methane mitigation strategy in dairy cattle. *Animal* 14:s473–s83
64. Manzanilla-Pech CIV, Løvendahl P, Mansan Gordo D, Difford GF, Pryce JE, et al. 2021. Breeding for reduced methane emission and feed-efficient Holstein cows: an international response. *J. Dairy Sci.* 104:8983–9001
65. Rowe SJ, Hickey SM, Jonker A, Hess MK, Janssen P, et al. 2019. Selection for divergent methane yield in New Zealand sheep—a ten-year perspective. In *Proceedings of the 23rd Conference of the Association for the Advancement of Animal Breeding and Genetics, Armidale, NSW, Aust., Oct. 27–Novemb. 1*, pp. 306–9. Armidale, Aust.: Assoc. Adv. Anim. Breed. Genet.
66. de Haas Y, Veerkamp RF, de Jong G, Aldridge MN. 2021. Selective breeding as a mitigation tool for methane emissions from dairy cattle. *Animal* 15(Suppl. 1):100294
67. Johnson PL, Hickey S, Knowler K, Wing J, Bryson B, et al. 2022. Genetic parameters for residual feed intake, methane emissions, and body composition in New Zealand maternal sheep. *Front. Genet.* 13:911639

68. Breider IS, Wall E, Garnsworthy PC. 2019. Short communication: heritability of methane production and genetic correlations with milk yield and body weight in Holstein-Friesian dairy cows. *J. Dairy Sci.* 102:7277–81
69. Lassen J, Lovendahl P, Madsen J. 2012. Accuracy of noninvasive breath methane measurements using Fourier transform infrared methods on individual cows. *J. Dairy Sci.* 95:890–98
70. Saborío-Montero A, Gutiérrez-Rivas M, García-Rodríguez A, Atxaerandio R, Goiri I, et al. 2020. Structural equation models to disentangle the biological relationship between microbiota and complex traits: methane production in dairy cattle as a case of study. *J. Anim. Breed. Genet.* 137:36–48
71. Difford GF, Olijhoek DW, Hellwing ALF, Lund P, Bjerring MA, et al. 2018. Ranking cows' methane emissions under commercial conditions with sniffers versus respiration chambers. *Acta Agric. Scand. A* 68:25–32
72. de Haas Y, Pszczola M, Soyeurt H, Wall E, Lassen J. 2017. Invited review: phenotypes to genetically reduce greenhouse gas emissions in dairying. *J. Dairy Sci.* 100:855–70
73. Shadpour S, Chud TCS, Hailemariam D, Plastow G, Oliveira HR, et al. 2022. Predicting methane emission in Canadian Holstein dairy cattle using milk mid-infrared reflectance spectroscopy and other commonly available predictors via artificial neural networks. *J. Dairy Sci.* 105:8272–85
74. Sepulveda BJ, Muir SK, Bolormaa S, Knight MI, Behrendt R, et al. 2022. Eating time as a genetic indicator of methane emissions and feed efficiency in Australian maternal composite sheep. *Front. Genet.* 13:883520
75. Ramayo-Caldas Y, Zingaretti L, Popova M, Estellé J, Bernard A, et al. 2020. Identification of rumen microbial biomarkers linked to methane emission in Holstein dairy cows. *J. Anim. Breed. Genet.* 137:49–59
76. Engelke SW, Das G, Derno M, Tuchscherer A, Wimmers K, et al. 2019. Methane prediction based on individual or groups of milk fatty acids for dairy cows fed rations with or without linseed. *J. Dairy Sci.* 102:1788–802
77. Vanlierde A, Vanrobays ML, Dehareng F, Froidmont E, Soyeurt H, et al. 2015. Hot topic: innovative lactation-stage-dependent prediction of methane emissions from milk mid-infrared spectra. *J. Dairy Sci.* 98:5740–47
78. de Haas Y, Windig JJ, Calus MPL, Dijkstra J, de Haan M, et al. 2011. Genetic parameters for predicted methane production and potential for reducing enteric emissions through genomic selection. *J. Dairy Sci.* 94:6122–34
79. Renand G, Vinet A, Decruyenaere V, Maupetit D, Dozias D. 2019. Methane and carbon dioxide emission of beef heifers in relation with growth and feed efficiency. *Animals* 9:1136
80. Flay HE, Kuhn-Sherlock B, Macdonald KA, Camara M, Lopez-Villalobos N, et al. 2019. Hot topic: Selecting cattle for low residual feed intake did not affect daily methane production but increased methane yield. *J. Dairy Sci.* 102:2708–13
81. Donoghue KA, Bird-Gardiner T, Arthur PF, Herd RM, Hegarty RF. 2016. Genetic and phenotypic variance and covariance components for methane emission and postweaning traits in angus cattle. *J. Anim. Sci.* 94:1438–45
82. Richardson CM, Amer PR, Quinton C, Crowley J, Hely FS, et al. 2022. Reducing greenhouse gas emissions through genetic selection in the Australian dairy industry. *J. Dairy Sci.* 105:4272–88
83. Beauchemin KA, Ungerfeld EM, Eckard RJ, Wang M. 2020. Review: fifty years of research on rumen methanogenesis: lessons learned and future challenges for mitigation. *Animal* 14:s2–s16
84. Difford GF, Plichta DR, Lovendahl P, Lassen J, Noel SJ, et al. 2018. Host genetics and the rumen microbiome jointly associate with methane emissions in dairy cows. *PLOS Genet.* 14:e1007580
85. Manzanilla-Pech CIV, Difford GF, Sahana G, Romé H, Lovendahl P, Lassen J. 2022. Genome-wide association study for methane emission traits in Danish Holstein cattle. *J. Dairy Sci.* 105:1357–68
86. Jalil Sarghale A, Moradi Shahrehabak M, Moradi Shahrehabak H, Nejati Javaremi A, Saatchi M, et al. 2020. Genome-wide association studies for methane emission and ruminal volatile fatty acids using Holstein cattle sequence data. *BMC Genet.* 21:129
87. Pszczola M, Strabel T, Mucha S, Sell-Kubiak E. 2018. Genome-wide association identifies methane production level relation to genetic control of digestive tract development in dairy cows. *Sci. Rep.* 8:15164

88. Lee SH, Clark S, van der Werf JHJ. 2017. Estimation of genomic prediction accuracy from reference populations with varying degrees of relationship. *PLOS ONE* 12:e0189775
89. Ducrocq V, Laloe D, Swaminathan M, Rognon X, Tixier-Boichard M, Zerjal T. 2018. Genomics for ruminants in developing countries: from principles to practice. *Front. Genet.* 9:251
90. Martinez-Alvaro M, Auffret MD, Duthie CA, Dewhurst RJ, Cleveland MA, et al. 2022. Bovine host genome acts on rumen microbiome function linked to methane emissions. *Commun. Biol.* 5:16
91. Hess MK, Zetouni L, Hess AS, Budel J, Dodds KG, et al. 2023. Combining host and rumen metagenome profiling for selection in sheep: prediction of methane, feed efficiency, production, and health traits. *Genet. Sel. Evol.* 55:53
92. Pérez-Enciso M, Zingaretti LM, Ramayo-Caldas Y, de los Campos G. 2021. Opportunities and limits of combining microbiome and genome data for complex trait prediction. *Genet. Sel. Evol.* 53:65
93. Thornton PK, Herrero M. 2010. Potential for reduced methane and carbon dioxide emissions from livestock and pasture management in the tropics. *PNAS* 107:19667–72
94. Duin EC, Wagner T, Shima S, Prakash D, Cronin B, et al. 2016. Mode of action uncovered for the specific reduction of methane emissions from ruminants by the small molecule 3-nitrooxypropanol. *PNAS* 113:6172–77
95. Jayanegara A, Sarwono KA, Kondo M, Matsui H, Ridla M, et al. 2017. Use of 3-nitrooxypropanol as feed additive for mitigating enteric methane emissions from ruminants: a meta-analysis. *Ital. J. Anim. Sci.* 17:650–56
96. Dijkstra J, Bannink A, France J, Kebreab E, van Gastelen S. 2018. Short communication: antimethanogenic effects of 3-nitrooxypropanol depend on supplementation dose, dietary fiber content, and cattle type. *J. Dairy Sci.* 101:9041–47
97. Kim H, Lee HG, Baek YC, Lee S, Seo J. 2020. The effects of dietary supplementation with 3-nitrooxypropanol on enteric methane emissions, rumen fermentation, and production performance in ruminants: a meta-analysis. *J. Anim. Sci. Technol.* 62:31–42
98. Yu GH, Beauchemin KA, Dong RL. 2021. A review of 3-nitrooxypropanol for enteric methane mitigation from ruminant livestock. *Animals* 11:3540
99. Martínez-Fernández G, Abecia L, Arco A, Cantalapiedra-Hijar G, Martín-García A, et al. 2014. Effects of ethyl-3-nitrooxy propionate and 3-nitrooxypropanol on ruminal fermentation, microbial abundance, and methane emissions in sheep. *J. Dairy Sci.* 97:3790–99
100. Meale SJ, Popova M, Saro C, Martin C, Bernard A, et al. 2021. Early life dietary intervention in dairy calves results in a long-term reduction in methane emissions. *Sci. Rep.* 11:3003
101. Gruninger RJ, Zhang XM, Smith ML, Kung L Jr., Vyas D, et al. 2022. Application of 3-nitrooxypropanol and canola oil to mitigate enteric methane emissions of beef cattle results in distinctly different effects on the rumen microbial community. *Anim. Microbiome* 4:35
102. Martínez-Fernández G, Duval S, Kindermann M, Schirra HJ, Denman SE, McSweeney CS. 2018. 3-NOP vs. halogenated compound: methane production, ruminal fermentation and microbial community response in forage fed cattle. *Front. Microbiol.* 9:1582
103. Muetzel S, Lowe K, Janssen P, Pacheco D, Bird N, et al. 2019. *Towards the application of 3-nitrooxypropanol in pastoral farming systems*. Poster presented at the New Zealand Agricultural Climate Change Conference, Palmerston North, N.Z., April 8–9
104. Guyader J, Ungerfeld EM, Beauchemin KA. 2017. Redirection of metabolic hydrogen by inhibiting methanogenesis in the rumen simulation technique (RUSTEC). *Front. Microbiol.* 8:393
105. Weimer PJ. 2015. Redundancy, resilience and host specificity of the ruminal microbiota: implications for engineering improved ruminal fermentations. *Front. Microbiol.* 6:296
106. Glasson CRK, Kinley RD, de Nys R, King N, Adams SL, et al. 2022. Benefits and risks of including the bromoform containing seaweed *Asparagopsis* in feed for the reduction of methane production from ruminants. *Algal Res.* 64:102673
107. Sawyer MS, Hoover WH, Sniffen CJ. 1974. Effects of a ruminal methane inhibitor on growth and energy metabolism in the ovine. *J. Anim. Sci.* 38:908–14
108. Tomkins NW, Colegate SM, Hunter RA. 2009. A bromochloromethane formulation reduces enteric methanogenesis in cattle fed grain-based diets. *Anim. Prod. Sci.* 49:1053–58

109. Knight T, Ronimus RS, Dey D, Tootill C, Naylor G, et al. 2011. Chloroform decreases rumen methanogenesis and methanogen populations without altering rumen function in cattle. *Anim. Feed Sci. Technol.* 166–67:101–12
110. Paul NA, de Nys R, Steinberg PD. 2006. Chemical defence against bacteria in the red alga *Asparagopsis armata*: linking structure with function. *Mar. Ecol. Prog. Ser.* 306:87–101
111. Li X, Norman HC, Kinley RD, Laurence M, Wilmot M, et al. 2016. *Asparagopsis taxiformis* decreases enteric methane production from sheep. *Anim. Prod. Sci.* 58:681–88
112. Roque BM, Salwen JK, Kinley R, Kebreab E. 2019. Inclusion of *Asparagopsis armata* in lactating dairy cows' diet reduces enteric methane emission by over 50 percent. *J. Clean. Prod.* 234:132–38
113. Stefenoni HA, Räisänen SE, Cueva SF, Wasson DE, Lage CFA, et al. 2021. Effects of the macroalga *Asparagopsis taxiformis* and oregano leaves on methane emission, rumen fermentation, and lactational performance of dairy cows. *J. Dairy Sci.* 104:4157–73
114. Roque BM, Venegas M, Kinley RD, de Nys R, Duarte TL, et al. 2021. Red seaweed (*Asparagopsis taxiformis*) supplementation reduces enteric methane by over 80 percent in beef steers. *PLOS ONE* 16:e0247820
115. Muizelaar W, Groot M, van Duinkerken G, Peters R, Dijkstra J. 2021. Safety and transfer study: transfer of bromoform present in *Asparagopsis taxiformis* to milk and urine of lactating dairy cows. *Foods* 10:584
116. Graham DE, White RH. 2002. Elucidation of methanogenic coenzyme biosyntheses: from spectroscopy to genomics. *Nat. Prod. Rep.* 19:133–47
117. Banerjee R, Ragsdale SW. 2003. The many faces of vitamin B₁₂: catalysis by cobalamin-dependent enzymes. *Annu. Rev. Biochem.* 72:209–47
118. Kubo I, Muroi H, Himejima M, Yamagiwa Y, Mera H, et al. 1993. Structure-antibacterial activity relationships of anacardic acids. *J. Agric. Food Chem.* 41:1016–19
119. Van Nevel CJ, Demeyer DI, Henderickx HK. 1971. Effect of fatty acid derivatives on rumen methane and propionate in vitro. *Appl. Microbiol.* 21:365–66
120. Watanabe Y, Suzuki R, Koike S, Nagashima K, Mochizuki M, et al. 2010. In vitro evaluation of cashew nut shell liquid as a methane-inhibiting and propionate-enhancing agent for ruminants. *J. Dairy Sci.* 93:5258–67
121. Wakai M, Hayashi S, Chiba Y, Koike S, Nagashima K, Kobayashi Y. 2021. Growth and morphologic response of rumen methanogenic archaea and bacteria to cashew nut shell liquid and its alkylphenol components. *Anim. Sci. J.* 92:e13598
122. Balch WE, Fox GE, Magrum LJ, Woese CR, Wolfe RS. 1979. Methanogens: reevaluation of a unique biological group. *Microbiol. Rev.* 43:260–96
123. Oh S, Suzuki Y, Hayashi S, Suzuki Y, Koike S, Kobayashi Y. 2017. Potency of cashew nut shell liquid in rumen modulation under different dietary conditions and indication of its surfactant action against rumen bacteria. *J. Anim. Sci. Technol.* 59:27
124. Shinkai T, Enishi O, Mitsumori M, Higuchi K, Kobayashi Y, et al. 2012. Mitigation of methane production from cattle by feeding cashew nut shell liquid. *J. Dairy Sci.* 95:5308–16
125. Mitsumori M, Enishi O, Shinkai T, Higuchi K, Kobayashi Y, et al. 2014. Effect of cashew nut shell liquid on metabolic hydrogen flow on bovine rumen fermentation: methane and metabolic hydrogen in rumen. *Anim. Sci. J.* 85:227–32
126. Su C, Shinkai T, Miyazawa N, Mitsumori M, Enishi O, et al. 2021. Microbial community structure of the bovine rumen as affected by feeding cashew nut shell liquid, a methane-inhibiting and propionate-enhancing agent. *Anim. Sci. J.* 92:e13503
127. Maeda K, Nguyen VT, Suzuki T, Yamada K, Kudo K, et al. 2021. Network analysis and functional estimation of the microbiome reveal the effects of cashew nut shell liquid feeding on methanogen behaviour in the rumen. *Microb. Biotechnol.* 14:277–90
128. Tyman JHP, Wilczynski D, Kashani MA. 1978. Compositional studies on technical cashew nutshell liquid (CNSL) by chromatography and mass spectroscopy. *J. Am. Oil Chem. Soc.* 55:663–68
129. Branco AF, Giallongo F, Frederick T, Weeks H, Oh J, Hristov AN. 2015. Effect of technical cashew nut shell liquid on rumen methane emission and lactation performance of dairy cows. *J. Dairy Sci.* 98:4030–40

130. El-Zaiat HM, Araujo RC, Soltan YA, Morsy AS, Louvandini H, et al. 2014. Encapsulated nitrate and cashew nut shell liquid on blood and rumen constituents, methane emission, and growth performance of lambs. *J. Anim. Sci.* 92:2214–24
131. Lee C, Beauchemin KA. 2014. A review of feeding supplementary nitrate to ruminant animals: nitrate toxicity, methane emissions, and production performance. *Can. J. Anim. Sci.* 94:557–70
132. Greening C, Geier R, Wang C, Woods LC, Morales SE, et al. 2019. Diverse hydrogen production and consumption pathways influence methane production in ruminants. *ISME J.* 13:2617–32
133. Iwamoto M, Asanuma N, Hino T. 2002. Ability of *Selenomonas ruminantium*, *Veillonella parvula*, and *Wolinella succinogenes* to reduce nitrate and nitrite with special reference to the suppression of ruminal methanogenesis. *Anaerobe* 8:209–15
134. Guyader J, Eugène M, Meunier B, Doreau M, Morgavi DP, et al. 2015. Additive methane-mitigating effect between linseed oil and nitrate fed to cattle. *J. Anim. Sci.* 93:3564–77
135. van Zijderveld SM, Gerrits WJJ, Apajalahti JA, Newbold JR, Dijkstra J, et al. 2010. Nitrate and sulfate: effective alternative hydrogen sinks for mitigation of ruminal methane production in sheep. *J. Dairy Sci.* 93:5856–66
136. Zhao LP, Meng QX, Li Y, Wu H, Huo YL, et al. 2018. Nitrate decreases ruminal methane production with slight changes to ruminal methanogen composition of nitrate-adapted steers. *BMC Microbiol.* 18:21
137. Popova M, Guyader J, Silberberg M, Seradj AR, Saro C, et al. 2019. Changes in the rumen microbiota of cows in response to dietary supplementation with nitrate, linseed, and saponin alone or in combination. *Appl. Environ. Microbiol.* 85:e02657-18
138. van Lingen HJ, Fadel JG, Yáñez-Ruiz DR, Kindermann M, Kebreab E. 2021. Inhibited methanogenesis in the rumen of cattle: microbial metabolism in response to supplemental 3-nitrooxypropanol and nitrate. *Front. Microbiol.* 12:705613
139. Villar ML, Godwin IR, Hegarty RS, Erler DV, Farid HT, Nolan JV. 2021. Nitrate and nitrite absorption, recycling and retention in tissues of sheep. *Small Rumin. Res.* 200:106392
140. Guo WS, Schaefer DM, Guo XX, Ren LP, Meng QX. 2009. Use of nitrate-nitrogen as a sole dietary nitrogen source to inhibit ruminal methanogenesis and to improve microbial nitrogen synthesis in vitro. *Asian-Australas. J. Anim. Sci.* 22:542–49
141. Shu Q, Gill HS, Hennessy DW, Leng RA, Bird SH, Rowe JB. 1999. Immunisation against lactic acidosis in cattle. *Res. Vet. Sci.* 67:65–71
142. Gill HS, Shu Q, Leng RA. 2000. Immunization with *Streptococcus bovis* protects against lactic acidosis in sheep. *Vaccine* 18:2541–48
143. Gnanasampanthan G. 1993. *Immune responses of sheep to rumen ciliates and the survival and activity of antibodies in the rumen fluid*. PhD Thesis, Univ. Adelaide, Adelaide, Aust.
144. Baker SK. 2000. *Method for improving utilisation of nutrients by ruminant or ruminant-like animals*. USA Patent No. 6,036,950
145. Wright AD, Kennedy P, O'Neill CJ, Toovey AF, Popovski S, et al. 2004. Reducing methane emissions in sheep by immunization against rumen methanogens. *Vaccine* 22:3976–85
146. Khanum S, Roberts JM, Heathcott RW, Bagley S, Wilson T, et al. 2022. Cross-reactivity of antibodies to different rumen methanogens demonstrated using immunomagnetic capture technology. *Front. Microbiol.* 13:14
147. Williams YJ, Popovski S, Rea SM, Skillman LC, Toovey AF, et al. 2009. A vaccine against rumen methanogens can alter the composition of archaeal populations. *Appl. Environ. Microbiol.* 75:1860–66
148. Baca-González V, Asensio-Calavia P, González-Acosta S, Pérez de la Lastra JM, Morales de la Nuez A. 2020. Are vaccines the solution for methane emissions from ruminants? A systematic review. *Vaccines* 8:460
149. Reisinger A, Clark H, Cowie AL, Emmet-Booth J, Gonzalez Fischer C, et al. 2021. How necessary and feasible are reductions of methane emissions from livestock to support stringent temperature goals? *Philos. Trans. A* 379:20200452
150. Budan A, Roman Garcia Y, Piantoni P, Humphries D, Sun Y. 2022. Potential of ZELP to improve the Cargill holistic approach to mitigate enteric methane emissions. In *Proceedings of the 8th International Greenhouse Gas and Animal Agriculture Conference, Orlando, FL*, p. 36. Gainesville: Univ. Fla.

151. Jeyanathan J, Martin C, Morgavi DP. 2014. The use of direct-fed microbials for mitigation of ruminant methane emissions: a review. *Animal* 8:250–61
152. Doyle N, Mbandlwa P, Kelly WJ, Attwood G, Li Y, et al. 2019. Use of lactic acid bacteria to reduce methane production in ruminants, a critical review. *Front. Microbiol.* 10:13
153. Zhao Y, Zhao G. 2022. Decreasing ruminal methane production through enhancing the sulfate reduction pathway. *Anim. Nutr.* 9:320–26
154. Leng RA. 2014. Interactions between microbial consortia in biofilms: a paradigm shift in rumen microbial ecology and enteric methane mitigation. *Anim. Prod. Sci.* 54:519–43
155. Qomariyah N, Ella A, Nurdin Ahmad S, Yusriani Y, Miftakhus Sholikin M, et al. 2023. Dietary biochar as a feed additive for increasing livestock performance: a meta-analysis of *in vitro* and *in vivo* experiment. *Czech J. Anim. Sci.* 68:72–86
156. Schmidt HP, Hagemann N, Draper K, Kammann C. 2019. The use of biochar in animal feeding. *PeerJ* 7:e7373
157. Fernandez GM, Durmic Z, Vercoe P, Joseph S. 2022. *Fit-for-purpose biochar to improve efficiency in ruminants*. Rep., Meat Livest. Aust., North Sydney, NSW, Aust.
158. Nardone A, Ronchi B, Lacetera N, Ranieri MS, Bernabucci U. 2010. Effects of climate changes on animal production and sustainability of livestock systems. *Livest. Sci.* 130:57–69
159. Thornton P, Nelson G, Mayberry D, Herrero M. 2022. Impacts of heat stress on global cattle production during the 21st century: a modelling study. *Lancet Planet. Health* 6:E192–201
160. Uyeno Y, Sekiguchi Y, Tajima K, Takenaka A, Kurihara M, Kamagata Y. 2010. An rRNA-based analysis for evaluating the effect of heat stress on the rumen microbial composition of Holstein heifers. *Anaerobe* 16:27–33
161. Yadav B, Singh G, Wankar A, Dutta N, Chaturvedi VB, Verma MR. 2016. Effect of simulated heat stress on digestibility, methane emission and metabolic adaptability in crossbred cattle. *Asian Australas. J. Anim. Sci.* 29:1585–92
162. Goopy JP, Korir D, Pelster D, Ali AIM, Wassie SE, et al. 2020. Severe below-maintenance feed intake increases methane yield from enteric fermentation in cattle. *Br. J. Nutr.* 123:1239–46
163. Herrero M, Havlík P, Valin H, Notenbaert A, Rufino M, et al. 2013. Biomass use, production, feed efficiencies, and greenhouse gas emissions from global livestock systems. *PNAS* 110:20888–93
164. Kuhla B, Viereck G. 2022. Enteric methane emission factors, total emissions and intensities from Germany's livestock in the late 19th century: a comparison with the today's emission rates and intensities. *Sci. Total Environ.* 848:157754
165. Benoit M, Mottet A. 2023. Energy scarcity and rising cost: towards a paradigm shift for livestock. *Agric. Syst.* 205:103585