



Effect of the microbiota of children with autism spectrum disorder(ASD) on behavior and various biological parameters in germ-free mice

Léa Roussin

► To cite this version:

Léa Roussin. Effect of the microbiota of children with autism spectrum disorder(ASD) on behavior and various biological parameters in germ-free mice. Optinutribrain Symposia, Optinutribrain, Food4BrainHealth, Université de Laval, Oct 2021, Online, France. hal-04343162

HAL Id: hal-04343162


<https://hal.inrae.fr/hal-04343162v1>

Submitted on 13 Dec 2023


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

➤ **Effect of the microbiota of children with autism spectrum disorder(ASD) on behavior and various biological parameters in germ-free mice**




➤ Autism spectrum disorder



embracingasd.com

tacanow.org



 **TACA**
THE AUTISM COMMUNITY IN ACTION

- Studies show a rate of heritability lower than 50%, highlighting an important impact of environment on the disorder

- It is important to characterize and better understand these environmental factors



- Notably: Prevalence of GI symptoms 4X higher in ASD patients → implication of the Gut-microbiota-brain axis



➤ Microbiota-gut-brain axis in ASD

Difference in microbiota composition

- Many studies in the past decade show a difference in microbiota composition in ASD patients compared to neurotypical individuals
- However, there is great variability in the results that could be in part caused by the impact of environment and alimentation on microbiota, or differences in the choice of patients and control groups:



From Dr. Ruth Ann Luna's presentation at the 74th Annual Meeting Society of Biological Psychiatry, Chicago (May 2019)

→ Controls groups used: difference in microbiota composition in unrelated neurotypical controls compared to neurotypical siblings of ASD individuals.

→ GI symptoms: difference in microbiota composition depending on presence or absence of GI symptoms

> Our project




Stool sampling
and preservation



Stool sampling
and preservation



Stool sampling
and preservation



Stool sampling
and preservation



FMT of pooled samples from each group to mice



Germ-free mice
(two strains)




Analysis of **behavior**
(8 weeks after FMT)
and various biological
parameters




3 mouse models of ASD:
-CMA model
-VPA model
- *PCDH9* KO



➤ Behavioral tests



Anxiety: Open field



Cognition: spatial memory



Social interaction and novelty



Repetitive behaviors
(grooming)

Hypothesis:

- Altered behavior in the ASD groups compared to their respective sibling groups.
- Potential differences between A and AG groups


Our project

Biological parameters

5-HT neurons



KYN/TRP ratio



Brain


Neuroinflammation (microglia)

T cell populations

Gut permeability

Gut
inflammation

Microbiota composition
and activity




Microbiota


Images: biorender.com

First results

Behavior



C57BL/6JCrI




Cognition: spatial memory

Recognition index (%) = $DO/(DO+NDO) \times 100$

DO=mean time interacting with **displaced** objets

NDO=mean time interacting with **non displaced** objets




- Decreased recognition index in the AG group compared to S-AG and A.

→ **Microbiota from children with ASD and GI symptoms can impact spatial memory in germ-free mice**

First results

Analysis of microbiota composition

Principal coordinate Analysis (PcoA):



→ Microbiota from each group forms a distinct cluster in PcoA analysis, highlighting that our 4 groups form distinct microbiota populations

➤ Conclusions and perspectives

Behavior:

- Altered spacial memory in germ-free C57BL/6 mice that received microbiota from the AG group
- No differences between groups for the other behavioral tests.
- Another study is planned to study behavior of Germ-free mice (2022)

Other parameters:

- Some differences in SFCAs profiles and T cell populations (analysis still ongoing)
- A lot left to analyse (gut inflammation/permeability, neuroinflammation, 5-HT etc).

1 month stay at
Utrecht University for
some of the
experiments!



Thanks for your attention!

PhD supervisors:

Laurent Naudon
Sylvie Rabot



The Amipem team:



Utrecht Team:

Aletta Kraneveld
Paula-Perez Pardo
Naika Prince



Thanks to the
Anaxem animal
facility





› Extras

➤ Preclinical studies

FMT from ASD patients to germ-free mice

Xiao et al. 2021:

ASD children or NT controls → pool of stool samples → transplantation to mice



In ASD-FMT mice compared to TD-FMT:

- Increased anxiety (time in center of open-field)
- Altered social behavior


Microbiota from ASD patients can transfer altered behaviors to mice

► T cell analysis in sleep (flow cytometry)

Expressed in % of
CD4+ cells



C57BL/6 (2nd cohort)



- Th1/Th2 ratio is lower in the AG group, highlighting a less inflammatory profile in those mice compared to S-AG and AG groups.