



HAL
open science

Conservation agriculture reduces climate change impact of a popcorn and wheat crop rotation

Maria Vittoria Guidoboni, Annie Duparque, Joachim Boissy, Jean-Christophe Mouny, Julie Auberger, Hayo Mg van der Werf

► **To cite this version:**

Maria Vittoria Guidoboni, Annie Duparque, Joachim Boissy, Jean-Christophe Mouny, Julie Auberger, et al.. Conservation agriculture reduces climate change impact of a popcorn and wheat crop rotation. PLoS ONE, 2023, 18 (5), pp.e0285586. 10.1371/journal.pone.0285586 . hal-04356702

HAL Id: hal-04356702

<https://hal.inrae.fr/hal-04356702>

Submitted on 15 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RESEARCH ARTICLE

Conservation agriculture reduces climate change impact of a popcorn and wheat crop rotation

Maria Vittoria Guidoboni¹, Annie Duparque², Joachim Boissy², Jean-Christophe Mouny², Julie Auberger¹, Hayo MG van der Werf^{1*}

1 UMR SAS, INRAE, Institut Agro, Rennes, France, **2** Agro-Transfert Ressources et Territoires, Estrées-Mons, France

* hayo.van-der-werf@inrae.fr



OPEN ACCESS

Citation: Guidoboni MV, Duparque A, Boissy J, Mouny J-C, Auberger J, van der Werf HMG (2023) Conservation agriculture reduces climate change impact of a popcorn and wheat crop rotation. PLOS ONE 18(5): e0285586. <https://doi.org/10.1371/journal.pone.0285586>

Editor: Farhan Hafeez, COMSATS University Islamabad Abbottabad Campus, PAKISTAN

Received: November 25, 2022

Accepted: April 27, 2023

Published: May 23, 2023

Copyright: © 2023 Guidoboni et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its [Supporting information files](#).

Funding: Hayo van der Werf received funding from the Carbon Farming research and innovation project (www.climate-kic.org/innovation-spotlight/carbon-farming-is-innovating-land-use/) under grant agreement no. TCL 210065, this funding paid the salary of Maria Vittoria Guidoboni. The other authors did not receive specific funding for the project. The funders encouraged publication of the

Abstract

Urgent action is needed to ensure humanity's future under climate change. Agriculture faces major challenges as it is both influenced by and contributes to climate change. Conservation agriculture sequesters carbon (C) in the soil due to practices such as reduced tillage and planting of cover crops. This study assessed effects of an innovative conservation agriculture popcorn (*Zea mays*) and wheat (*Triticum aestivum*) crop rotation in south-western France on soil C sequestration, GHG emissions and several environmental impacts. Two complementary approaches were used: i) a comparison based on field data and expert judgement to assess short-term effects and ii) modelling of three scenarios to quantify long-term outcomes. In both approaches Life cycle assessment (LCA) was used to compare popcorn and wheat rotations. The conventional rotation used ploughing, and its soil was bare between wheat harvest and popcorn sowing. Conservation agriculture used reduced tillage, cover crops, and compost of green waste. Impacts of compost production were allocated mainly to its waste treatment function, based on waste treatment cost and compost price. Simulation modelling of soil C was used to estimate the amount of C sequestered by the conservation and conventional crop rotations. LCA was combined with soil C modelling over 100 years to assess the long-term climate change impact of three scenarios for the popcorn and wheat rotation. These scenarios were 1) Conventional agriculture, 2) Conservation agriculture with cover crops only, 3) Conservation agriculture with cover crops + compost. Mean annual C sequestration and net climate change impact were -0.24 t/ha and 3867 kg CO₂-eq./ha, respectively, for the conventional rotation and 0.91 t/ha and 434 kg CO₂-eq./ha, respectively, for the conservation rotation. The climate change impact of the conservation rotation depended strongly on the allocation of composting impacts between the waste treatment and compost production functions. Compared to the conventional rotation, the conservation rotation had a lower marine eutrophication impact (-7%) but higher impacts for terrestrial acidification (+9%), land competition (+3%), and cumulative energy demand (+2%). Modelling over 100 years revealed that, at near soil C equilibrium, a conventional scenario lost 9% of soil C, whereas conservation agriculture scenarios gained 14% (only

study results, but had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Abbreviations: °C, degree Celsius; C, carbon; C_{comp}, organic C input from compost; C_{cover}, organic C input from cover crops; C_{crop}, organic C input from cash-crop residues; Cd, cadmium; C_{min}, mineralisation of soil C; CO₂, carbon dioxide; CO₂-eq., carbon dioxide-equivalent; Cons_{CC}, conservation agriculture with cover crops scenario; Cons_{full}, conservation agriculture with cover crops and compost scenario; Conv, conventional agriculture scenario; Cr, chromium; C_{seq}, organic C input from humified C sequestered; Cu, copper; GHG, greenhouse gas; Ha, hectare; Hg, mercury; IPCC, Intergovernmental Panel on Climate Change; ISO, international organization for standardization; LCA, life cycle assessment; LCI, life cycle inventory; N, nitrogen; N₂O, nitrous oxide; NH₃, ammonia; Ni, nickel; NO₃, nitrate; NO_x, nitrogen oxides; P, phosphorous; Pb, lead; PO₄, phosphate; SOC, soil organic carbon; t, ton; Zn, zinc.

cover crop) and 26% of soil C (cover crop + compost). Conservation agriculture resulted in soil C sequestration over several decades, until a new soil C equilibrium was reached.

1. Introduction

The Intergovernmental Panel on Climate Change [1] has reiterated the need for urgent action to ensure the future of humanity under climate change. To remain within the target of no more than 1.5°C of warming, urgent action at unprecedented scales and across all sectors is required within the next 10 years. According to IPCC [2], 24% of total greenhouse gas (GHG) emissions worldwide are caused by the agricultural sector. Consequently, agriculture has a major role in meeting the 1.5°C target through lowering its GHG emissions and sequestering carbon dioxide (CO₂) in the soil. At the same time, agricultural systems need to adapt to climate change, and produce sufficient food for a growing world population, while preserving biodiversity, water and soil.

Soil carbon (C) sequestration can contribute to reducing net anthropogenic GHG emissions, while preserving and restoring soil health by maintaining and increasing soil organic matter, thus helping to increase production resilience and food security [3]. Conservation agriculture is based on three crop management principles: no tillage, permanent organic soil cover with crop residues and/or cover crops, and crop rotation [4].

Conservation agriculture addresses several environmental and management issues. Planting of cover crops can improve soil quality by providing a source of substrate for microbial activity, which enhances soil C and nitrogen (N) cycles [5]. Soil erosion decreases when soils are less disturbed by tillage and are enriched with biomass from cover crops and organic fertilisation [6]. Cover crops also increase N-use efficiency: they take up soil N left by the previous crop, thus decreasing nitrate (NO₃) leaching and providing N from cover crop residues to the next crop [7]. Finally, conservation agriculture usually has lower production costs than conventional agriculture, due to reduced use of inputs (e.g. fuel) and less time spent in the field [8, 9].

Conservation agriculture is not common in popcorn (*Zea mays*) and wheat (*Triticum aestivum*) rotations in France. In current conventional popcorn and wheat rotations, the soil is ploughed in autumn, remains bare in winter, and only mineral fertiliser is applied. This type of management is prone to soil erosion and NO₃ leaching, and results in low stocks of soil organic matter, which negatively influences soil structure and soil water retention [10]. The Natais company, Europe's main popcorn producer, started transitioning towards conservation agriculture approximately two decades ago [11], with reduced tillage, inclusion of cover crops, and application of organic fertiliser.

Several studies have highlighted the importance of soil organic C (SOC) dynamics when estimating the C footprint of agricultural systems, particularly when conservation agriculture practices are used [12, 13]. These studies have shown that planting cover crops and using reduced or no tillage are good strategies to increase soil C content. These practices also tend to increase in-field N₂O emissions, due to the incorporation of cover crop residues, and may cause additional CO₂ emissions related to fuel combustion. Overall, conservation agriculture has lower net GHG emissions than conventional agriculture [13, 14] because the effect of soil sequestration on net GHG emissions is superior to that of increased N₂O and CO₂ emissions. Similarly, Prechsl et al. [14] found that using organic fertiliser (cattle slurry) rather than mineral N fertiliser in a 6-year rotation with cover crops in Switzerland emitted more in-field N₂O. However, the GHG emissions associated with production of mineral N fertiliser tipped the net GHG emissions in favour of organic fertilisation.

Life cycle assessment (LCA) has been used in environmental assessment of agricultural products and systems for several decades. LCA is a methodological framework that estimates potential environmental impacts of a product by quantifying the resources consumed and pollutant emissions to the environment in the course of its life cycle [15, 16]. It allows a multi-criteria assessment, considering a range of environmental impacts. The carbon footprint method assesses the total amount of GHG emissions that are directly and indirectly caused by an activity or are accumulated over the life stages of a product [17]. It quantifies a single criterion, the LCA climate change impact.

Conservation agriculture practices have been studied using LCA, e.g. [14] compared conventional tillage and no tillage and [18] assessed the effect of the introduction of cover crops. Yang et al. [19] used carbon footprint methodology to assess the effect of diversification of crop rotation. Surprisingly, we found no LCA or carbon footprint studies that compared conservation agriculture, i.e. the combination of no tillage, organic soil cover and crop rotation, to conventional agriculture.

The objectives of this study were to (i) assess short-term effects of an innovative conservation agriculture popcorn and wheat crop rotation on SOC sequestration, GHG emissions, and the environmental impacts climate change, terrestrial acidification, marine eutrophication, land competition and cumulative energy demand, and (ii) to quantify the potential mid- to long-term benefits of conservation agriculture for SOC sequestration and these environmental impacts.

2. Materials and methods

2.1 Life cycle assessment

Conservation agriculture differs from conventional agriculture both with respect to input use (e.g. tractors, seed for cover crops) and with respect to its sequestration of SOC. LCA allows a whole system comparison of conventional and conservation agriculture by considering simultaneously off-farm resource use and emissions of GHG and other pollutants (associated with inputs), on-farm resource use and emissions of GHG and other pollutants (e.g. from diesel combustion) and SOC changes corresponding to CO₂ emissions or sequestration.

2.1.1 Goal and scope definition. The goal of the study was to assess and compare potential environmental impacts of conventional and conservation agriculture in a popcorn and wheat crop rotation using attributional LCA. The system boundaries of the crop rotations began with the extraction of resources and ended at the farm gate. Processes included the production of all inputs and all processes related to crop production: (1) soil cultivation, sowing, weed control, fertilisation, pest and pathogen control, harvest; (2) machines and the buildings or areas used to park them; (3) seeds for cash and cover crops, fertilisers including compost, pesticides, water for irrigation and fuel, as well as their transport to the farm; (4) direct emissions of fuel combustion, tire abrasion and pollutant emissions in the field. Post-harvest processes, such as drying, sorting and storage were not included. The function considered was “land management”, defined as the occupation of agricultural and non-agricultural land for a given amount of time. The land occupied included both “direct” land (on-farm) and “indirect” land (off-farm, e.g. to produce seeds for sowing). The functional unit was 1 ha of land occupied for 1 year (i.e. ha.year).

To assess conventional and conservation agriculture rotations we used two complementary approaches: i) a comparison based on field data (for conservation agriculture) and expert judgement (for conventional agriculture) to assess short-term effects (Table 1) and ii) modelling three agricultural scenarios to quantify long-term outcomes (Table 2).

Table 1. Yields and inputs for conventional and conservation wheat and popcorn crop rotations in 2017–2019 and 2018–2020. WW = Winter Wheat; PC = Popcorn. The year indicates the harvest year. Unless indicated otherwise, inputs of cover crops were attributed to conservation popcorn. For conservation agriculture yields and inputs were based on field data, for conventional agriculture they were based on expert judgement.

	Unit	Conventional				Conservation			
		WW_2018	PC_2019	WW_2019	PC_2020	WW_2018	PC_2019	WW_2019	PC_2020
Yield, cash crop	t/ha	5.9	6.5	9.5	6.2	5.9	6.5	9.5	6.2
Input/Process									
Seeds, cash crop	kg/ha	127	15	122	16	127	15	122	16
Seeds, cover crop ¹	kg/ha	-	-	-	-	-	120	-	154
Fava bean/phacelia cover crop dry matter	t/ha	-	-	-	-	-	3.1	-	3.9
Sorghum cover crop dry matter	t/ha	-	-	-	-	-	-	-	3
Compost of green waste	t/ha	-	-	-	-	-	6.5	-	9.9
N compost of green waste	kg/ha	-	-	-	-	-	52	-	79
N mineral fertiliser	kg/ha	179	190	202	190	179	188	202	182
N total	kg/ha	179	190	202	190	179	240	202	261
P ₂ O ₅ total	kg/ha	24	50	24	50	24	53	24	66
K ₂ O total	kg/ha	0	50	0	50	0	89	0	122
Diesel	kg/ha	40	73	40	75	32	68	32	78
Irrigation water	m ³ /ha	-	1 659	-	2 396	-	1 659	-	2 396
Pesticide active ingredient	kg/ha	5.2	2.8	6.6	2.8	5.2	3.7	6.6	3.7

¹ in 2019 only fava bean and phacelia seeds, in 2020 both fava bean, phacelia and sorghum seeds

<https://doi.org/10.1371/journal.pone.0285586.t001>

2.1.1.1 Comparison to assess short-term effects. For the conservation agriculture crop rotation field data (i.e. farmer practices and yields) was collected on the Natais commercial farm (43°52' North, 0°89' East) in the Gers administrative department (south-western France) during two rotations: 2017–2019 and 2018–2020 (Table 1).

Conservation agriculture results for each crop were calculated by averaging results for five wheat or popcorn fields. With slopes of 4–12%, the Natais farm's soils are prone to erosion, even though they contain 25–30% clay. The study site has a temperate oceanic climate according to the Köppen-Geiger climate classification [20], with mean monthly precipitation of 54 mm and mean annual air temperature of 15.1 °C from 2018–2020. The 2018–2020 period was relatively warm, and the 30-year mean air temperature was 13.5 °C. Soil texture was mainly clay-limestone. The initial soil organic matter content (0–30 cm) at the experimental sites was 1.8–2.1%, and the initial SOC stock was 10.3–12.0 g/kg.

In the conservation agriculture rotation, popcorn received compost of green waste along with mineral fertiliser, while winter wheat received only mineral fertiliser. Green waste consisted mainly of grass clippings, leaves and branch cuttings from private and public gardens. Popcorn was sown according to a technique called “green-tillage” as described hereafter. For the 2017–2019 rotation, winter wheat was sown in October 2017. In July 2018, after the wheat harvest, stubble was tilled twice, at 5 and 10 cm. In mid-September 2018, the soil was tilled

Table 2. Scenarios of conventional and conservation popcorn and wheat crop rotations.

Scenario	Agricultural system	Cover crops	Compost applied (t/ha)	N applied (kg N/(ha))
Conv	Conventional	Absent	0	190
Cons_CC	Conservation	sorghum and fava bean + phacelia	0	190
Cons_full	Conservation	sorghum and fava bean + phacelia	6.5	220

<https://doi.org/10.1371/journal.pone.0285586.t002>

with a spring-toothed harrow, to a depth of 20 cm on the future popcorn row with one set of teeth, and to a depth of 8 cm with a second set of teeth. Some of the soil of the future inter row was moved to the future row to form a ridge. In the same operation, winter fava bean (98.6%) and phacelia (1.4%) was sown to establish a cover crop in the future inter-row. During winter, the soil of the ridge settled, which created a seedbed for the popcorn. In April 2019, herbicide application killed the cover crop. Subsequently, popcorn was sown on the ridges using a machine that flattened the cover crop and simultaneously applied fertiliser (diammonium phosphate, 58 kg/ha), insecticide (cypermethrin, 92 g/ha), and molluscicide (methaldehyde, 79 g/ha) on the row. Popcorn was sown on ridges because this favours germination due to higher soil temperature. After harvesting the popcorn in October 2019, winter wheat was directly sown and mineral fertiliser was applied. For the 2018–2020 rotation, the operations were identical, but two cover crops were grown: a sorghum cover crop was sown at the beginning of July, and chopped and left in the field in mid-September before sowing the fava bean-phacelia cover crop.

For the conventional agriculture rotation farmer practices reflected dominant practices for popcorn and winter wheat rotations in the Gers department according to expert judgment (S. Hypolite, Agro d'Oc, pers. comm.) (Table 1). Popcorn and wheat yields were assumed to be the same as those in the conservation agriculture rotation based on expert judgement. Both crops received only mineral fertilisers. For popcorn, the soil was ploughed in autumn and superficially tilled twice in spring. After the popcorn harvest, stubble was tilled at a depth of 10 cm before sowing the winter wheat. No cover crops were grown. All other farming practices were the same as in the conservation agriculture system.

Wheat yielded 5.9 and 9.5 t/ha in 2018 and 2019 respectively, whereas popcorn yielded 6.5 and 6.2 t/ha in 2019 and 2020, respectively (Table 1). The fava bean/phacelia cover crop produced 3.1 and 3.9 t dry matter/ha in 2019 and 2020, respectively, whereas the sorghum cover crop produced 3 t dry matter/ha in 2020.

Conservation popcorn received 6.5 and 9.9 t/ha of compost in 2019 and 2020, respectively, corresponding to 22% and 30% of total N applied, respectively. Compared to conventional popcorn, conservation popcorn (including the cover crop) used 26% (2019) and 37% (2020) more fertiliser N, 6% less (2019) and 5% more (2020) diesel, and 33% (2019) and 32% (2020) more pesticide active ingredient. Both conservation wheat crops (2018 and 2019) used 20% less diesel than the conventional ones. Yields and amounts of cash-crop seed, irrigation water, mineral fertiliser, and pesticide applied were identical for conventional and conservation wheat crops.

2.1.1.2 Comparison of three scenarios to assess long-term outcomes. In the scenario-modelling approach, a conventional scenario (Conv) and two conservation scenarios (Cons_CC and Cons_full) were compared (Table 2). The Conv scenario corresponded to the conventional popcorn and wheat rotation described previously. The two conservation scenarios were based on Nataï's conservation agriculture rotation. The Cons_CC scenario used sorghum and fava bean + phacelia as cover crops (as in the 2018–2020 cropping system), whereas the Cons_full scenario used these cover crops as well as 6.5 t/ha of compost of green waste (as in the 2017–2019 cropping system). For all scenarios, yields were set at 6.8 t/ha for winter wheat and 6.3 t/ha for popcorn, based on expert judgement for long-term average yield level of these crops in the region.

2.1.2 Life cycle inventory. Life cycle inventories (LCIs) were calculated using the MEAN-S-InOut web application v3.0, which is a customized agricultural LCA tool that generates LCIs of agricultural production systems [21]. It contains forms to guide data entry, a reference data set for the pollutant emissions and resource use of main inputs of agri-food systems, analytical models to estimate direct pollutant emissions and resource use, and an export function that

generates LCI files ready to be imported into LCA software (here we used Simapro, see 2.1.3) to calculate impact indicators. Databases used for background processes (e.g. production of fertilizer, irrigation infrastructure), were AGRIBALYSE v3.0.1 and ecoinvent v3.5.

Emissions to the air (ammonia (NH₃), nitrous oxide (N₂O), N oxides (NO_x), CO₂), water (NO₃, phosphorus (P), phosphate (PO₄), cadmium (Cd), chromium (Cr), copper (Cu), mercury (Hg), nickel (Ni), lead (Pb), zinc (Zn)) and soil (Cd, Cr, Cu, Hg, Ni, Pb, Zn, pesticides) were calculated using models recommended by the AGRIBALYSE methodology [22]. NH₃ emissions from organic fertiliser application were modelled using EMEP/EEA 2016 Tier 2 [23]. N₂O emissions were modelled using IPCC 2019 Tier 1 [24]. NO_x emissions were modelled using EMEP/EEA 2009 Tier 1 [25]. NO₃ leaching was modelled according to Tailleur et al. [26]. CO₂ emissions from fuel combustion and emissions of active substances of pesticides were modelled according to ecoinvent[®] v2 [27].

Composting of green waste consists of the following processes: shredding, stabilisation, mixing, composting and screening [28]. The compost of green waste LCI available in the AGRIBALYSE database [28] attributes all polluting emissions of the composting process to the compost, and thus to the crop on which the compost is applied. From a methodological point of view, this is questionable, as the composting process has two functions: waste treatment and compost production. According to the LCI for compost in the AGRIBALYSE database, producing 1 t of compost requires 3 t of green waste. The cost of treating green waste by composting was assumed to be 60 €/t [29] (i.e. 180 € per t of compost), and the price of the green waste compost delivered to the Nataïs farm was 14 €/t. Given a total cost of 1 t of compost of 194 €/t, 7.2% of it (i.e. 14/194×100) was for the compost and the rest for the waste treatment process. Consequently, following an approach proposed by Christensen et al. [30], we used economic allocation to allocate pollutant emissions and resource use between the two functions: 7.2% to compost production and 92.8% to waste treatment.

2.1.3 Impact assessment: LCA impact categories. Impacts were assessed using SimaPro software v9.0.0.35. We calculated climate change, terrestrial acidification, and marine eutrophication impacts according to the ReCiPe 2016 Midpoint (H) characterisation method [31]. We also calculated land competition impact according to the CML-IA non-baseline method [32] and cumulative energy demand [33].

2.2 Estimating changes in soil organic carbon stock

Most LCA studies of agricultural production systems assume that farming practices remain constant over time, and thus that the SOC stock remains constant. This assumption is often reasonable, but not when considering a change in farming practices that increases the input of organic C to the soil, such as the introduction of cover crops and organic fertiliser. When such practices are implemented, the increase in SOC stock needs to be included in the LCI, as it represents an absorption of CO₂ from the atmosphere (i.e. a negative emission).

The change in SOC stock for the comparison of the two rotations was estimated using the AMG model [34], which considers three compartments: fresh exogenous organic C, active SOC, and stable SOC. The sources of fresh organic C are organic fertilisers and crop residues. The active SOC compartment is fed by fresh organic C inputs and influenced by annual mineralisation, while stable SOC is considered to be completely inert over the short and mid-terms (i.e. its turnover time is millenary). The AMG model represents the generally accepted fact that the SOC pool is heterogeneous. The AMG model has three main parameters: the humification coefficient, which is the conversion factor from fresh organic C inputs into humified SOC; the annual rate of SOC mineralisation; and the initial proportion of stable C in the initial SOC stock. The humification coefficient depends only on the nature of fresh organic C inputs,

whereas the annual rate of SOC mineralisation depends on the soil (i.e. clay and lime contents, pH, C:N ratio of soil organic matter), soil tillage (type and depth), and meteorological conditions (i.e. mean annual air temperature and water balance). The model runs on a yearly basis at the field level to estimate SOC stock at a depth of 0–30 cm.

To estimate the amount of crop biomass left on the field after harvest, crop yield is converted into above-ground biomass using a crop-specific harvest index. Below-ground biomass is estimated using an allometric relationship with aerial biomass [35]. Total (above- and below-ground) biomass minus crop yield equals the amount of biomass that provides fresh organic C inputs. In the comparison of the conservation and the conventional agriculture crop rotation field data for crop yields were available for conservation agriculture but not for conventional agriculture. We assumed that crop yields of the latter were identical to those of the former. This is not necessarily true, we therefore conducted a sensitivity analysis assessing the effects of a 10% increase or decrease of the yield of conventional crops on carbon storage and subsequent environmental impacts.

For the comparison of three scenarios, changes in the SOC stock over 100 years were estimated using the SIMEOS-AMG tool [36, 37], which is a decision-support tool based on the AMG model designed to facilitate simulation of long-term changes in SOC. Model description and model inputs and outputs for AMG and SIMEOS-AMG are presented in the S1 and S2 Tables in [S1 File](#).

3. Results

3.1 Characteristics of crop rotations

Emissions of NH₃, NO, and N₂O were higher for the conservation popcorn crops (including cover crop) than for the conventional ones by 11%, 15%, and 32%, respectively, in 2019 and by 17%, 22%, and 58%, respectively, in 2020, whereas CO₂ emissions were similar (Table 3). NH₃ and NO emissions were higher for conservation wheat crops than for conventional wheat crops by 3% and 4%, respectively, in 2019 and for both gases by 6% in 2019. NO₃ leaching was 17% lower for conservation wheat crops than for conventional wheat crops (Table 3).

According to the AMG model, conventional wheat and popcorn sequestered -0.24 t C/ha for both the 2017–2019 and the 2018–2020 crop rotation (Table 4). A 10% decrease or increase of assumed yields of the crops in the conventional rotation resulted in a moderate (-0.33/-0.16 and -0.34/-0.15) variation of these values (Table 4). C sequestration of conservation wheat was similar to that of conventional wheat, while conservation popcorn, due to the cover crop and compost application, sequestered 1.6 (2019) and 2.7 (2020) t C/ha. Mean annual C sequestration was 0.6 and 1.2 t C/ha for the 2017–2019 and 2018–2020 conservation crop rotations, respectively. The difference in annual C sequestration between the two systems was thus 0.83 and 1.47 t C/ha, with C input from compost of 0.46 and 0.69 t/ha, for 2017–2019 and 2018–2020, respectively. For conservation popcorn, percentages of C input from crop residues, cover crop, and compost were 36%, 29%, and 35%, respectively, in 2019 and 23%, 43%, and 34%, respectively, in 2020.

3.2 Climate change impact of crop rotations

The conservation crop rotation had higher GHG emissions than the conventional crop rotation (15% higher in 2017–2019; 25% in 2018–2020) and also sequestered much more C (-347% as much in 2017–2019; -608% in 2018–2020) (Fig 1). Consequently, conventional rotations had positive climate change impacts (7440 kg CO₂-eq./ha in 2017–2019 and 8027 kg CO₂-eq./ha in 2018–2020), as well as the conservation rotation in 2017–2019 (2422 kg CO₂-eq./ha). On the other hand, the conservation rotation in 2018–2020 had a negative climate change impact

Table 3. Main direct emissions for conventional and conservation wheat and popcorn crop rotations in 2017–2019 and 2018–2020. WW = Winter Wheat; PC = Popcorn. The year indicates the harvest year. Unless indicated otherwise, direct emissions of cover crops were attributed to conservation popcorn.

	Unit	Conventional				Conservation			
		WW_2018	PC_2019	WW_2019	PC_2020	WW_2018	PC_2019	WW_2019	PC_2020
Direct emissions									
To air									
NH ₃	kg/ha	18.0	24.5	19.1	22.3	18.6	27.3	20.2	26.1
NO	kg/ha	4.4	4.7	4.9	4.5	4.6	5.4	5.1	5.4
N ₂ O	kg/ha	5.7	6.3	6.6	6.2	5.7	8.3	6.6	9.7
CO ₂	kg/ha	129	209	143	414	129	208	143	410
To water									
NO ₃	kg/ha	133	111	133	111	111	111	111	111

¹ in 2019 only fava bean and phacelia seeds, in 2020 both fava bean, phacelia and sorghum seeds

<https://doi.org/10.1371/journal.pone.0285586.t003>

(-688 kg CO₂-eq./ha), because C sequestration was superior to GHG emissions. For the conventional crop rotations, direct emissions contributed most to the climate change impact. For the conservation crop rotation, direct emissions had a similar magnitude as those of conventional crop rotations, but C sequestered contributed most to the climate change impact.

When averaged over the two crop rotations, the climate change impact considering only emissions was 1% lower for wheat and 8% higher for popcorn for conservation agriculture than for conventional agriculture (Fig 2). Expressed in kg CO₂-eq./ha, impacts were 2629 (conventional) and 2598 (conservation) for wheat, 3407 (conventional) and 3680 (conservation) for popcorn, and 966 for cover crops.

For conservation popcorn, climate change impacts for other field operations and mineral fertilisers were 45% and 10% lower, respectively, than those for conventional popcorn. Transport of inputs and pesticides, which contributed little (ca. 1%) to impacts for all crops, were 78% and 16% higher respectively, for conservation popcorn than for conventional popcorn. The processes that contributed most were direct emissions (64% conventional and 59% conservation), mineral fertiliser (16% conventional and 13% conservation), irrigation (10% conventional and 9% conservation), other field operations (9% conventional and 4% conservation), and compost 11% (only for conservation).

For conservation wheat, climate change impacts for direct emissions and other field operations were similar to and 22% lower, respectively, than those for conventional wheat. Impacts associated with mineral fertilisers, pesticides, and transport of inputs did not differ for

Table 4. Carbon (C) balance per crop and its components for wheat and popcorn estimated by the AMG model: Organic C input from cash-crop residues (Ccrop), mineralisation of soil C (Cmin), organic C input from cover crops (Ccover), organic C input from compost (Ccomp), and humified C sequestered (Cseq) for the conventional and conservation rotations. Values are expressed as t C/ha/year at a depth of 0–30 cm. Values in brackets for Cseq of the conventional crops correspond to the effects of a 10% increase or decrease of their yields.

Year	Crop	Conventional			Conservation				
		Ccrop	Cmin	Cseq	Ccrop	Ccover	Ccomp	Cmin	Cseq
2017–2018	Wheat	0.78	-1.15	-0.37 (-0.45/-0.29)	0.78	0.00	0.00	-1.15	-0.37
2019	Popcorn	0.95	-1.06	-0.11 (-0.20/-0.02)	0.95	0.76	0.91	-1.06	1.56
Mean		0.87	-1.11	-0.24 (-0.33/-0.16)	0.87	0.38	0.46	-1.11	0.59
2018–2019	Wheat	1.25	-1.38	-0.13 (-0.25/-0.01)	1.25	0.00	0.00	-1.47	-0.22
2020	Popcorn	0.92	-1.28	-0.36 (-0.43/-0.28)	0.92	1.71	1.39	-1.35	2.67
Mean		1.09	-1.33	-0.24 (-0.34/-0.15)	1.09	0.86	0.69	-1.41	1.23

<https://doi.org/10.1371/journal.pone.0285586.t004>




Fig 1. Contribution of soil organic carbon, direct emissions, seed, compost, mineral fertilisers, pesticides, field operations (including irrigation), and transport of inputs to the climate change impact for the two-year popcorn and wheat crop rotation in 2017–2019 and 2018–2020, for conventional and conservation rotations. Dots indicate net impact, which is equal to C sequestration minus greenhouse gas emissions. The inputs and direct emissions of the cover crops were attributed to the popcorn crops.

<https://doi.org/10.1371/journal.pone.0285586.g001>

conservation and conventional wheat because they used identical amounts. The processes that contributed most were direct emissions (75% conventional and 76% conservation), mineral fertiliser (15% for both conventional and conservation), and other field operations (6% conventional and 4% conservation). For the cover crops, the processes that contributed most were direct emissions (84%), other field operations (13%), and seeds (3%).

3.3 Multi-impact comparison of crop rotations

Impacts of the conventional and conservation rotations differed (Table 5). The conservation rotation had lower impacts than the conventional rotation for climate change (-89%) and marine eutrophication (-7%). A 10% decrease or increase of assumed yields of the crops in the conventional rotation resulted in a moderate (-97%/-82%) variation of the difference for the climate change impact (Table 5). The conservation rotation had somewhat higher impacts than the conventional rotation for terrestrial acidification (+9%), land competition (+3%), and cumulative energy demand (+2%).

When comparing the 2017–2019 and 2018–2020 crop rotations, differences were small, except for climate change for the conservation rotation, which had much lower impacts for




Fig 2. Contribution of direct emissions, seed, compost, mineral fertilisers, pesticides, irrigation, other field operations, and transport of inputs to the climate change impact for individual crops. Values are means for 2017–2018 and 2018–2019 for wheat, for 2019 and 2020 for popcorn, and for 2018–2019 and 2019–2020 for cover crops.

<https://doi.org/10.1371/journal.pone.0285586.g002>

2018–2020 than for 2017–2019 (-169%). Cumulative energy demand was 17% higher for 2018–2020 than for 2017–2019 for both the conservation and conventional systems.

3.4 Change in soil organic C stock over time of scenarios

According to the SIMEOS-AMG simulation tool, initial SOC stock was 38.4 t/ha for the three scenarios (Fig 3). Over time, SOC stock decreased in the Conv scenario but increased in the Cons_CC and Cons_full scenarios. After 100 years, SOC had not completely reached

Table 5. Impacts of conventional and conservation crop rotations (2017–2019 and 2019–2020) for the land management (ha.year) functional unit. Difference expresses impacts of the conservation rotation relative to those of the conventional rotation. Values in brackets for Difference correspond to the effects of a 10% increase or decrease of the conventional crop yields.

Impact category	Unit	Crop rotation						Difference (%)
		Conventional			Conservation			
		2017–2019	2018–2020	Mean	2017–2019	2018–2020	Mean	
Climate change	kg CO ₂ eq	3720	4014	3867	1211	-344	434	-89 (-97/-82)
Terrestrial acidification	kg SO ₂ eq	48.5	47.9	48.2	51.9	53.3	52.6	+9
Marine eutrophication	kg N eq	8.38	8.43	8.41	7.75	7.85	7.80	-7
Land competition	m ² a	10 293	10 436	10 365	10 531	10 774	10 653	+3
Cumulative energy demand	MJ	21 549	24 626	23 087	21 777	25 513	23 645	+2

<https://doi.org/10.1371/journal.pone.0285586.t005>




Fig 3. Soil organic carbon stock (0–30 cm) over 100 years according to three scenarios of popcorn and wheat crop rotations as estimated by the SIMEOS-AMG simulation tool. See Table 2 for a description of the scenarios.

<https://doi.org/10.1371/journal.pone.0285586.g003>

equilibrium. When 95% of the change in SOC stock at 100 years was reached, the subsequent annual change was near zero for each scenario: -0.0005% (Conv), 0.0005% (Cons_CC), and 0.0009% (Cons_full). The Conv scenario reached this 95% threshold after 31 years, losing 9% of the initial SOC stock, whereas Cons_CC and Cons_full reached it after 34 years, gaining 14% and 26% of the initial SOC stock, respectively. After 100 years, estimated SOC stocks were 35.7 (Conv), 43.9 (Cons_CC), and 48.8 (Cons_full) t C/ha.

In the south west of France approximately 7500 ha of popcorn are grown in a conventional crop rotation. Let us assume that all of this popcorn were grown in a popcorn-wheat rotation which would transition from conventional to conservation agriculture according to the Cons_CC scenario. Over a 34-year period this would allow the sequestration of $43.9 - 38.4 = 5.5$ t/ha on 15000 ha (twice the popcorn area, because popcorn is present one year out of two in the rotation). This corresponds to an average annual C sequestration of $(5.5 * 15000) / 34 = 2426$ t. The annual C footprint of a French citizen being 9 t CO₂-eq. [38], this corresponds to the C footprint of 270 citizens.

Initial annual climate change impacts for the scenarios were 0.19 (Cons_full), 1.64 (Cons_CC), and 4.31 (Conv) t CO₂-eq./ha. As mentioned, the Cons_full scenario included a sorghum cover crop followed by a fava bean + phacelia cover crop and 6.5 t/ha of compost (Table 1). The Cons_full scenario resembled the 2017–2019 conservation system (Table 3), but differed in that it had a single cover crop (fava bean + phacelia) and slightly different crop yields (6.8 and 6.3 t/ha for wheat and popcorn, respectively vs. 5.9 and 6.5 t/ha, respectively in the 2017–2019 system). Assuming constant management practices, the climate change impact decreased over time for Conv but increased for Cons_CC and Cons_full. Climate change

impacts for the three systems intersected after 23–28 years, when they became larger for Cons_CC and Cons_full than for Conv.

4. Discussion

4.1 C inputs and sequestration

According to field data for the Natais conservation agriculture system, C input from cover crops was 0.8 and 1.7 t/ha in 2019 (fava bean) and 2020 (sorghum followed by fava bean), respectively (Table 3). The 2020 value is close to the mean C input by cover crops (1.9 t/ha) estimated from the meta-analysis of Poeplau and Don [39]. Considering both C inputs by crop residues and cover crops as well as mineralisation of soil C, mean annual C sequestration for the cropping system was 0.6 and 1.2 t/ha for 2017–2019 and 2018–2020, respectively.

According to the meta-analysis of Sun et al. [40], the difference in mean annual SOC sequestration of conservation vs. conventional agriculture was 0.35 t C/ha. Conservation agriculture practices do not include organic fertilisation, so for comparison with our results, the contribution of compost to SOC sequestration (0.46 (2019) and 0.69 (2020) t C/ha) should be subtracted. This yields differences in annual C sequestration of conservation vs. conventional system of 0.37 (i.e. $0.83 - 0.46$) and 0.78 (i.e. $1.47 - 0.69$) t/ha for 2019 and 2020, respectively. When a single cover crop was present, our results thus agreed with those from the meta-analysis, and when two consecutive cover crops were grown, they were twice as high.

In the Natais conservation agriculture system, compost contributed 34% (2019) and 35% (2020) to the flow of C to the soil, and was thus a major contributor to the system's climate change impact (Table 3). Based on composting cost and compost price, we allocated 7.2% of impacts of the composting process to the compost. As composting cost and compost price can vary, a sensitivity analysis was conducted. A 50% decrease or 50% increase in the allocation percentage had a small effect (-5% and +5%, respectively) on the climate change impact of conservation popcorn (Fig 4). However, when 100% of the impact of the composting process was allocated to the compost, the climate change impact increased by 140%.

The French low-C label [41] developed a method to quantify SOC sequestration and reductions in GHG emissions. This method uses the AGRIBALYSE LCI of compost of green waste, which allocates 100% of the impact of the composting process to the compost. The climate change impact of this compost is 694 kg CO₂-eq./t. In our study, compost of green waste contained 172 kg C/t, 41% of which (i.e. 70.5 kg, corresponding to 259 kg CO₂-eq.) was estimated to be sequestered [42]. Consequently, using this method, applying 1 t of compost of green waste increases the climate change impact of a given crop by 435 kg CO₂-eq. (i.e. $694 - 259$), not counting emissions associated with applying compost. This example illustrates the importance of allocation choices on potential climate change impacts, and the need to establish a meaningful consensus on this.

4.2 SOC dynamics and climate change impact

4.2.1 SOC dynamics. In their recent review of 106 mainly short-term (2–3 year) experiments on the impact of cover crops, Abdalla et al. [43] found mean annual C sequestration of 0.54 t/ha. For our Cons_CC scenario, the short-term (2-year) annual C sequestration was 0.57 t/ha, close to their value.

4.2.2 Climate change impact. No tillage, permanent organic soil cover and crop rotation are the three crop management principles at the heart of conservation agriculture. The environmental impacts of each of these principles separately have been assessed using LCA or carbon footprint studies. However, we found no LCA or carbon footprint studies of conservation agriculture systems, implementing these three principles simultaneously. This study is thus a




Fig 4. Contribution of direct emissions, seed, compost, other field operations, irrigation, mineral fertilisers, pesticides and transport of inputs to the climate change impact of conservation agriculture popcorn as a function of the percentage of impact allocated to compost: 7.2% (this study), 3.6% (50% decrease), 10.8% (50% increase), and 100% (original life cycle inventory from AGRIBALYSE; Avadi (2020)).

<https://doi.org/10.1371/journal.pone.0285586.g004>

first of its kind. One of its main findings was that conservation agriculture had higher GHG gas emissions than conventional agriculture, due mainly to the cover crop and, to a lesser extent, to compost production and application. However, conservation agriculture sequestered much more SOC, resulting in a much smaller net climate change impact compared to conventional agriculture. For both systems GHG emissions consisted for approximately two-thirds of direct emissions, the remaining GHG emissions were due to fertiliser production (including compost), irrigation and field operations.

Over time, the soil was a C sink in the conservation scenarios but a C source in the conventional scenario (Fig 3). Considering both GHG emissions and changes in SOC stock, the net climate change impact of the conservation scenarios increased over time, eventually overtaking that of the conventional scenario (Fig 5). This was due to the slightly higher GHG emissions of the conservation scenarios and the gradual decrease in the amount of SOC sequestered by the conservation scenarios and released by the conventional scenario. This illustrates that conservation agriculture can sequester a large amount of C until the SOC stock reaches a new equilibrium after a few decades.

It would be possible to increase C sequestration of the conservation scenario further by increasing C inputs, for instance by shifting towards more complex systems such as agroforestry [44]. In their review, Cardinael et al. [45] indicate that converting an arable system to a silvo-arable system yielded mean annual C sequestration rates of 0.21, 0.28, and 1.12 t/ha in SOC, below-ground biomass, and above-ground biomass, respectively.

4.3 Effects of the conservation system on other impact categories

Conventional and conservation agriculture systems differed strongly in their climate change impact, but differences for other impact categories were relatively small. Terrestrial acidification was 9% higher for conservation agriculture than for conventional agriculture. This




Fig 5. Net climate change impact over 50 years considering greenhouse gas emissions and soil organic carbon sequestration according to three scenarios of popcorn and wheat crop rotations. See Table 2 for a description of the scenarios.

<https://doi.org/10.1371/journal.pone.0285586.g005>

difference was due to NH_3 emissions, which were higher for the conservation agriculture system, which used compost in addition to mineral fertiliser, whereas the conventional agriculture system used only mineral fertiliser. Marine eutrophication was 7% lower for the conservation system than for the conventional system due to lower NO_3 leaching because of the cover crop in the former. Land competition was 3% higher for the conservation system than for the conventional system because more land was needed to produce cover crop seeds. Cumulative energy demand was 2% higher for the conservation system than for the conventional system because the cover crops required more inputs (i.e. seeds, diesel, and transport of inputs).

4.4 Other beneficial effects of conservation agriculture

SOC sequestration due to conservation agriculture ceases when a new equilibrium is reached, but this does not mean that its benefits cease. Under climate change, droughts are becoming increasingly frequent and severe, with negative effects on crop productivity [46]. Increase in soil organic matter can increase plant available water capacity in the root zone, and thus improve climate-resilience of soils and agroecosystems and reduce the need for irrigation [47]. More generally, an increase of SOC has been shown to increase yields of wheat and maize and reduce the need for fertilizer N [48].

Furthermore, cover crops have beneficial effects beyond their effect on SOC [10]. Due to their albedo effect, they mitigate climate change by reflecting more sunlight than most bare soils [49, 50]. The annual albedo effect of cover crops on climate change has been estimated to

be equivalent to 120–460 kg CO₂-eq./ha [49], smaller than the average effect of cover crops on SOC, but not insignificant. Other continued beneficial effects of cover crops include reduction of soil erosion, fixation of atmospheric N, reduction of NO₃ leaching, weed suppression, and improvement of the structure of soil microbial communities [10].

4.5 Study limitations

The amount of dry matter (crop residues and compost) received by the conservation agriculture system was 9.6 t/ha in 2019, and 16.8 t/ha in 2020. Application of crop residues and compost increases N₂O emissions. This was taken into consideration in our estimation of N₂O emissions. However, N₂O emissions are particularly increased when residues are left on the soil as a mulch rather than incorporated in the soil due to inversion tillage [51, 52]. This effect was not taken into account in the modelling of N₂O emissions according to the IPCC 2019 Tier 1 method we used. Consequently, we may have somewhat underestimated N₂O emissions of the conservation agriculture system.

The assessment of short-term effects of conservation agriculture was based on real-life farm data on yield and farming practices for the conservation rotation, while farming practices of the conventional rotation were based on expert judgement and its crop yields were assumed to be identical to those of the conservation rotation. Although the assumption regarding yield level is supported by the literature and we consider the expert judgement on farming practices to be of high quality, the lack of field data on the reference conventional system constitutes a limitation of this study. A 10% decrease or increase of assumed conventional crop yields had a modest effect on C sequestration and climate change impact, suggesting that our results are nevertheless robust.

Furthermore effects of the conservation and conventional agriculture rotations on soil carbon dynamics are based on modelling results. This also constitutes another limitation of this study. However, an evaluation of the AMG model by comparing it to SOC results from 60 long-term field trials in France revealed that the model accurately simulated the changes in SOC stocks over time [34]. Thus, in spite of the absence of field data, we consider these model-based results to be sufficiently solid.

5. Conclusions

This study revealed that implementing an innovative conservation agriculture system initially decreased the climate change impact of a popcorn and wheat rotation strongly, while moderately influencing marine eutrophication, terrestrial acidification, land competition, and cumulative energy demand. Scenario modelling revealed that a new SOC equilibrium was nearly reached after ca. 30 years, when C sequestration virtually ceased, and the climate change impact of the conservation scenarios was slightly higher than that of a conventional scenario. Modelling over 100 years revealed that at near soil C equilibrium, the SOC stock of the conventional scenario had decreased by 9%, whereas that of the conservation scenario had increased by 14% and, when compost was used, by 26%.

The climate change impact of the system using compost of green waste depended strongly on how impacts of the composting process were allocated between its waste treatment and compost production functions. This finding illustrates the need to establish a consensus on the allocation of composting impacts.

Conservation agriculture can sequester a large amount of C until the SOC stock reaches a new equilibrium after a few decades. Shifting towards more complex systems such as agroforestry could enhance C sequestration further.

Supporting information

S1 File.
(XLSX)

Acknowledgments

We thank the Natais team for its support, in particular Anne-Marie Joliet, who answered our many questions during data collection. We thank Eric Ceschia of CESBIO/INRAE for his reactivity and help, as well as Sylvain Hypolite (Agro d'Oc) for his expert knowledge. We thank our colleagues of the Carbon Farming project, especially Gécica Yogo for her coordination role.

Author Contributions

Conceptualization: Hayo MG van der Werf.

Data curation: Julie Auberger.

Formal analysis: Maria Vittoria Guidoboni, Annie Duparque, Joachim Boissy, Jean-Christophe Mouny.

Methodology: Maria Vittoria Guidoboni, Annie Duparque, Joachim Boissy, Julie Auberger, Hayo MG van der Werf.

Software: Jean-Christophe Mouny, Julie Auberger.

Supervision: Hayo MG van der Werf.

Writing – original draft: Maria Vittoria Guidoboni.

Writing – review & editing: Annie Duparque, Julie Auberger, Hayo MG van der Werf.

References

1. IPCC. Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte. V. et al., (eds.)]. Cambridge University Press, 2021.
2. IPCC. Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team. R.K. Pachauri and L.A. Meyer (eds.)]. IPCC. Geneva, Switzerland, 2014. 151 pp.
3. Smith P, Bustamante M, Ahammad H, Clark H, Dong H, Elsiddig EA, et al. Agriculture Forestry and Other Land Use (AFOLU) in: Edenhofer O et al, (Eds) Climate Change 2014: Mitigation of Climate Change Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change Cambridge University Press Cambridge United Kingdom and New York, 2014, NY USA.
4. Pittelkow CM, Liang X, Linquist BA, van Groenigen KJ, Lee J, Lundy ME, et al. Productivity limits and potentials of the principles of conservation agriculture *Nature* 2015; 517: 365–368.
5. Mbuthia LW, Acosta-Martinez V, DeBruyn J, Schaeffer S, Tyler D, Odoi E, et al. Long term tillage, cover crop, and fertilization effects on microbial community structure. activity: Implications for soil quality. *Soil Biol. Biochem.* 2015; 89: 24–34.
6. Lal R. Restoring Soil Quality to Mitigate Soil Degradation. *Sustainability* 2015; 7: 5875–5895.
7. Hartwig NL, Ammon HU. 50th Anniversary—Invited article—Cover crops and living mulches. *Weed Sci.* 2002; 50: 688–699.
8. Giller KE, Andersson JA, Corbeels M, Kirkegaard J, Mortensen D, Erenstein O, et al. Beyond conservation agriculture. *Front. Plant Sci.* 2015; 6: 870. <https://doi.org/10.3389/fpls.2015.00870> PMID: 26579139

9. Hobbs PR, Sayre K, Gupta R. The role of conservation agriculture in sustainable agriculture. *Philos. Trans. R. Soc. B-Biol. Sci.* 2008; 363: 543–555. <https://doi.org/10.1098/rstb.2007.2169> PMID: 17720669
10. Blanco-Canqui H, Shaver TM, Lindquist JL, Shapiro CA, Elmore RW, Francis CA, et al. Cover crops and ecosystem services: insights from studies in temperate soils. *Agron. J.* 2015; 107: 2449–2474.
11. Bodoville G. Le Green Tillage: Une technique innovante pour l'implantation du maïs pop-corn dans le Sud-Ouest. *Agronomie, écologie et innovation* 2014; 79: 4–8.
12. Plaza-Bonilla D, Nogue-Serra I, Raffailac D, Cantero-Martinez C, Justes E. Carbon footprint of cropping systems with grain legumes and cover crops: A case-study in SW France. *Agric. Syst.* 2018; 167: 92–102.
13. Yao ZY, Zhang DB, Yao PW, Zhao N, Liu N, Zhai BN, et al. Coupling life-cycle assessment and the RothC model to estimate the carbon footprint of green manure-based wheat production in China. *Sci. Total Environ.* 2017; 607: 433–442. <https://doi.org/10.1016/j.scitotenv.2017.07.028> PMID: 28704669
14. Prechsl UE, Wittwer R, van der Heijden MGA, Luscher G, Jeanneret P, Nemecek T. Assessing the environmental impacts of cropping systems and cover crops: Life cycle assessment of FAST a long-term arable farming field experiment. *Agric. Syst.* 2017; 157: 39–50.
15. ISO. ISO 14040—Environmental management—life cycle assessment—principles and framework. ISO, Geneva, 2006a.
16. ISO. ISO 14044—Environmental management—life cycle assessment—requirements and guidelines. ISO, Geneva, 2006b.
17. Galli A, Wiedmann T, Ercinc E, Knoblauch D, Ewing B, Giljum S. Integrating Ecological, Carbon and Water footprint into a “Footprint Family” of indicators: Definition and role in tracking human pressure on the planet. *Ecol. Indic.* 2012; 16: 100–112.
18. Cecchin A, Pourhashem G, Gesch RW, Mohammed YA, Patel S, Lenssen AW, et al. The environmental impact of ecological intensification in soybean cropping systems in the U.S. upper Midwest. *Sustainability* 2021, 13: 1696.
19. Yang X, Gao W, Zhang M, Chen Y, Sui P. Reducing agricultural carbon footprint through diversified crop rotation systems in the North China Plain. *J. Clean. Prod.* 2014, 76: 131–139.
20. Peel MC, Finlayson BL, McMahon TA. Updated world map of the Köppen-Geiger climate classification. *Hydrol. Earth Syst. Sci.* 2007; 11: 1633–1644.
21. Auberger J, Biard Y, Colomb V, Grasselly D, Martin E, van der Werf H, et al. MEANS- InOut: user-friendly software to generate LCIs of farming systems. 11th International Conference on Life Cycle Assessment of Food 2018 (LCA Food), Oct 2018, Bangkok, Thailand. <https://hal.archives-ouvertes.fr/hal-02014939>.
22. Koch P, Salou T. AGRIBALYSE®: Methodology. Agricultural stage—Version 3.0. ADEME. Angers. France, 2020. p. 303.
23. EMEP/EEA. EMEP/EEA air pollutant emission inventory guidebook 2016. Technical guidance to prepare national emission inventories. EEA. Copenhagen, Denmark, 2016. p. 28.
24. IPCC. Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Calvo Buedia, E. et al., (eds). Published: IPCC, Switzerland, 2019.
25. EMEP/EEA. EMEP/EEA air pollutant emission inventory guidebook 2009. Technical guidance to prepare national emission inventories. Copenhagen, Denmark, 2009. p. 16.
26. Tailleur A, Cohan J, Laurent F, Lellahi A. A simple model to assess nitrate leaching from annual crops for life cycle assessment at different spatial scales in: Corson, MS, van der Werf, HMG (Eds) LCA Food 2012 Proceedings of the 8th International Conference on Life Cycle Assessment in the Agri-Food Sector Saint-Malo France pp 903–904.
27. Nemecek T, Kägi T. Life Cycle Inventories of Swiss and European Agricultural Production Systems—Data v2.0. ecoinvent® Report No. 15a. Life Cycle Inventories of Agricultural Production Systems Data v2.0, 2007. Zürich and Dübendorf. Switzerland. p. 360.
28. Avadi A. Screening LCA of French organic amendments and fertilisers. *Int. J. Life Cycle Assess.* 2020; 25: 698–718.
29. ADEME. Etude des coûts de collecte et de compostage de biodéchets de quatre sites Qualorg. Etude par AWIPLAN, 2002, 19 p.
30. Christensen LO, Galt RE, Kendall A. Life-cycle greenhouse gas assessment of Community Supported Agriculture in California's Central Valley. *Renewable Agriculture and Food Systems* 2017; 33(5): 393–405.

31. Huijbregts MAJ, Steinmann ZJN, Elshout PMF, Stam G, Verones F, Vieira M, et al. ReCiPe2016: a harmonised life cycle impact assessment method at midpoint and endpoint level (vol 22. pg 138. 2017). *Int. J. Life Cycle Assess.* 2020; 25: 1635–1635.
32. Guinée J. Handbook on life cycle assessment—Operational guide to the ISO standards. *Int. J. Life Cycle Assess.* 2002; 6: 255–255.
33. Frischknecht R., Jungbluth N, Althaus HJ, Bauer C, Doka G, Dones R, et al. Implementation of Life Cycle Impact Assessment Methods. Swiss Centre for Life Cycle Inventories 2007. Dübendorf.
34. Clivot H, Mouny JC, Duparque A., Dinh JL, Denoroy P., Houot S., et al. Modeling soil organic carbon evolution in long-term arable experiments with AMG model. *Environmental Modelling and Software* 2019; 118: 99–113.
35. Baret F, Olioso A, Luciani JL. Root biomass fraction as a function of growth degree days in wheat. *Plant Soil* 1992; 140: 137–144.
36. Duparque A, Tomis V, Mary B, Boizard H, Damay N. Le bilan Humique AMG, pour une démarche de conseil fondée sur des cas-types régionaux. In: 10th Rencontres de la fertilisation raisonnée et de l'analyse de terre. GEMAS-COMIFER 2011 Reims: 16p.
37. Duparque A, Dinh JL, Mary B. AMG: a simple SOC balance model used in France for decision support. International workshop SOMpatic Rauischholtzhausen (Germany), November 20–22, 2013.
38. SDES Datalab—Chiffres clés du climat France, Europe et Monde Le service des données et études statistiques, France, 2022, 93p.
39. Poeplau C, Don A Carbon sequestration in agricultural soils via cultivation of cover crops—A meta-analysis. *Agric. Ecosyst. Environ.* 2015; 200: 33–41.
40. Sun W, Canadell JG, Yu L, Yu L, Zhang W, Smith P, et al. Climate drives global soil carbon sequestration and crop yield changes under conservation agriculture *Glob. Change. Biol.* 2020; 26: 3325–3335.
41. Soenen B, Henaff M, Lagrange H, Lanckriet E, Schneider A, Duval R, et al. Méthode Label Bas-Carbone Grandes Cultures (version 10), 2021 133p www.ecologiegouv.fr/label-bas-carbone.
42. Pellerin S, Bamière L, Launay C, Martin R, Schiavo M, Angers D, et al. Stocker du carbone dans les sols français Quel potentiel au regard de l'objectif 4 pour 1000 et à quel coût? Synthèse du rapport d'étude INRA France, 2019 p 114.
43. Abdalla M, et al. A critical review of the impacts of cover crops on nitrogen leaching, net greenhouse gas balance and crop productivity. *Glob. Change Biol.* 2019; 25: 2530–2543. <https://doi.org/10.1111/gcb.14644> PMID: 30955227
44. Lorenz K, Lal R. Soil organic carbon sequestration in agroforestry systems. A review. *Agron. Sustain. Dev.* 2014; 34: 443–454.
45. Cardinael R, Umulisa V, Toudert A, Olivier A, Bockel L, Bernoux M. Revisiting IPCC Tier 1 coefficients for soil organic and biomass carbon storage in agroforestry systems. *Environ. Res. Lett.* 2018; 13: 124020.
46. Kim W, Iizumi T, Nishimori M. Global patterns of crop production losses associated with droughts from 1983 to 2009, *Journal of Applied Meteorology and Climatology* 2019; 58(6): 1233–1244.
47. Lal R. Soil organic matter and water retention. *Agron. J.* 2020; 112: 3265–3277.
48. Oldfield EE, Bradford MA, Wood SA. Global meta-analysis of the relationship between soil organic matter and crop yields. *Soil* 2019; 5: 15–32.
49. Kaye JP, Quemada M. Using cover crops to mitigate and adapt to climate change. A review. *Agronomy for Sustainable Development* 2017; 3: 17.
50. Carrer D, Pique G., Ferlicoq M., Ceamanos X., Ceschia E., What is the potential of cropland albedo management in the fight against global warming? A case study based on the use of cover crops. *Environ. Res. Lett.* 2018; 13: 11.
51. Hu N., Chen Q., Zhu L. The Responses of Soil N₂O Emissions to Residue Returning Systems: A Meta-Analysis. *Sustainability* 2019; 11: 748.
52. Shan J, Yan X, 2013. Effects of crop residue returning on nitrous oxide emissions in agricultural soils. *Atmospheric Environment* 2013; 71: 170–175.