



HAL
open science

Probiotics impact on *Tenebrio molitor* growth, microbial composition and pathogen infection

Christina Nielsen-Leroux, Carlotta Savio

► To cite this version:

Christina Nielsen-Leroux, Carlotta Savio. Probiotics impact on *Tenebrio molitor* growth, microbial composition and pathogen infection. International Congress on Invertebrate Pathology and Microbial Control and the 55th Annual Meeting of the Society for Invertebrate Pathology, Robert Harrison, Jul 2023, Park City, United States. hal-04370193

HAL Id: hal-04370193

<https://hal.inrae.fr/hal-04370193>

Submitted on 2 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Oral student presentation

The 2023 International Congress on Invertebrate Pathology and Microbial Control and the 55th Annual Meeting of the Society for Invertebrate Pathology (SIP 2023)

July 30 to August 3 , 2023 , College Park, Maryland, USA.

Probiotics impact on *Tenebrio molitor* growth, microbial composition and pathogen infection

Savio Carlotta^{1, 2, 5}, Herren Pascal^{2,3,4}, Agnès Rejasse¹, Annette Bruun Jensen², Joop J.A. van Loon⁵, Christina Nielsen-Leroux¹

¹Université Paris-Saclay, INRAE, AgroParisTech, Micalis Institute, Domaine de Vilvert, 78350, Jouy-en-Josas, France; ²University of Copenhagen, Department of Plant and Environmental Sciences, Thorvaldsensvej 40, 1871, Frederiksberg, Denmark; ³UK Centre for Ecology & Hydrology, Maclean Building, Benson Lane, Crowmarsh Gifford, Wallingford, Oxfordshire OX10 8BB, United Kingdom; ⁴University of Leeds, Faculty of Biology, Leeds LS2 9JT, United Kingdom; ⁵Wageningen University, Department of Plant Sciences, Laboratory of Entomology, 6708 PB Wageningen, The Netherlands

Address for correspondence: carlotta.savio@inrae.fr

The yellow mealworm *Tenebrio molitor* is an insect model for infection and immunity studies and is mass-produced as feed and food. The industrial rearing of *T. molitor* on agricultural by-products may expose larvae and adults to entomopathogens used as biocontrol agents, like bacterial spores/toxins of *Bacillus thuringiensis* or conidia of the fungus *Metarhizium brunneum*, that could impact *T. molitor* survival and growth. Therefore, as for other animal livestock, the possible benefits of addition of living or tantalized probiotic bacteria to the insect feed are investigated. Two probiotic stains, *Pediococcus pentosaceus* KVL B09-1 and *Lactobacillus plantarum* WJB were added to wheat bran (WB) or wheat bran added with dried egg white (WE) during development from the egg to larvae weighing 20 mg, after which larvae were exposed during 72 h to *B. thuringiensis*, *M.brunneum* or their combination in simulated optimal rearing conditions. Larval survival and weight gain were recorded for 14 days and the bacterial microbiota composition was analyzed by 16S rDNA sequencing at day 0, 3 and 14 after contact with the pathogens, to define the impact of feed composition and probiotics on *T. molitor* survival, growth and microbial composition. Results show evidence of increased weight gain and differences in microbial community in larvae fed with WE diet. Positive effects on growth and increased survival rate on *T. molitor* larvae reared on feed supplemented with probiotics, conferring increased survivability and weigh gain case of co-infections and suggesting a role of probiotic-host-microbiota interactions in maintenance of insect health. At the end of the experience, all individuals presented a comparable microbial community composition, highlight the inability of the probiotic strains to persist in the larvae after probiotic removal from the diet.