



HAL
open science

Mirroring Amino Acids Hold Sway to Scare Pathobionts Away

Jeremy Denizot, Benoit Chassaing

► **To cite this version:**

Jeremy Denizot, Benoit Chassaing. Mirroring Amino Acids Hold Sway to Scare Pathobionts Away. Cellular and Molecular Gastroenterology and Hepatology, 2023, 16 (6), pp.1038-1039. 10.1016/j.jcmgh.2023.09.009 . hal-04372753

HAL Id: hal-04372753

<https://hal.inrae.fr/hal-04372753v1>

Submitted on 4 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0
International License

Mirroring Amino Acids Hold Sway to Scare Pathobionts Away



All amino acids (aa), except glycine, are stereoisomers found in 2 different configurations in living organisms: the L-configuration and its chiral counterparts, the D-configuration. In mammals, L-aa are used to synthesize proteins, while microorganisms mostly use D-aa to synthesize their cell wall via a racemase enzyme converting L-aa into D-aa.¹ Accordingly, one might expect that D-aa supplementation will fuel intestinal bacteria overgrowth. In this issue of *Cellular and Molecular Gastroenterology and Hepatology*, Umeda et al² unexpectedly reveal the beneficial role played by D-aa in tempering intestinal inflammation through microbiota modulation.

In inflammatory bowel diseases (IBDs), accumulating evidence highlights that the associated intestinal dysbiosis is characterized frequently by an increase in *Escherichia coli* pathobiont bacteria, together with a decrease in *Firmicutes* members, which could drive or foster chronic intestinal inflammation.³ Moreover, various dietary factors are suspected to play a critical role in IBD pathogenesis through their impact on the intestinal microbiota.⁴ Here, Umeda et al² first observed a decreased D-aa/L-aa ratio in fecal samples collected from UC patients compared with healthy individuals. The negative correlation between fecal D-aa abundance and microbial families such as *Enterobacteriaceae* further suggested that depletion of fecal D-aa could play a role in UC patients' dysbiosis. Strikingly, oral administration of 5 major D-aa (Ala, Ser, Glu, Trp, and Asn), either in their L or D configuration, revealed that D-aa supplementation could efficiently alleviate inflammation in mouse models of dextran sulfate sodium-induced colitis and of primary sclerosing cholangitis. Moreover, fecal microbiota transplantation from D-aa-treated mice into germ-free mice was sufficient to confer protection against dextran sulfate sodium-induced colitis, suggesting that the protective effect provided by D-aa supplementation is mediated, at least in part, by modulation of the intestinal microbiota. Various D-aa next were tested independently for their protective potential against colitis, identifying D-Ala, D-Trp, and D-Glu as central actors in microbiota composition modulation, especially by decreasing Proteobacteria relative abundance, and prevention of chronic intestinal inflammation.

Umeda et al² next focused their study on *Klebsiella pneumoniae*, *E coli*, and *Proteus mirabilis*, 3 Proteobacteria selectively decreased in mice treated with D-aa. They reported that D-aa supplementation could prevent colonic mucosal invasion by bacteria in the mouse model of cholangitis, as well as limit pore formation induced by *K pneumoniae* in murine organoid cultures. Bacterial growth experiments indicated that D-Ala supplementation inhibited the growth of the various Proteobacteria tested. Finally, Umeda et al² identified the bacterial *FtsZ* gene, essential for the last step of cell division, as strongly down-regulated in

E coli and *K pneumoniae* upon D-ala supplementation. Overall, this study by Umeda et al² provides compelling evidence that select D-aa hold a potential therapeutic avenue, notably in IBD patients carrying an excess of Proteobacteria.

This work leaves open several questions. Although the microbiota was sufficient to transfer the protective effect conferred by D-aa supplementation, it cannot be excluded that D-aa also directly impacted the host, notably by modulating gene expression. The effect of D-aa supplementation on microbiota composition might indirectly restore butyrate-producing microbiota members, which frequently are reduced in IBD patients. Because butyrate is a potent inhibitor of histone deacetylases in intestinal epithelial cells,⁵ it can be envisioned that intestinal D-aa impacts the epigenetic landscape in host epithelial and/or immune cells, and thereby modulates expression of antimicrobial peptides and genes involved in intestinal barrier function. Another point warranting further investigation relates to the depletion of fecal D-aa observed in IBD patients, which might be a cause, but also a consequence, of chronic intestinal inflammation. Similarly, the cause-consequence relationship between the fecal D-aa load and the relative abundance of various microbiota members remains to be elucidated.

The authors nicely reported how D-aa can inhibit *E coli* expansion in vitro and in vivo in mice, but further studies appear needed to confirm these observations in human beings. Among pathobionts, adherent-invasive *E coli* (AIEC) bacteria are thought to play an important role in the etiology of IBD.⁶ AIECs are found encroached within the ileal mucosa of Crohn's disease (CD) patients, where they hold the potential to induce a chronic inflammatory response.⁷ To date, no specific therapeutic approach targeting AIEC bacteria in CD patients have been developed, but clinical trials are ongoing to assess the efficiency of AIEC eradication in CD patients through antibiotics, phages therapy, or FimH blockage. The important role suggested here for D-aa in regulating the colibiome opens another innovative therapeutic approach to specifically inhibit AIEC intestinal colonization. Hence, these data are of great interest for future targeted modulation of the intestinal microbiota in IBD or individuals at risk of developing IBD.

JEREMY DENIZOT

Microbes, Intestin, Inflammation et Susceptibilité de l'Hôte
Université Clermont Auvergne/Inserm U1071 Unité sous
contrat Institut National de recherche pour l'agriculture
l'alimentation et l'environnement 1382
Clermont-Ferrand, France

BENOIT CHASSAING

Mucosal Microbiota in Chronic Inflammatory Diseases Team
INSERM U1016
CNRS UMR 8104, Université de Paris
Paris, France

References

1. Bastings JAJ, van Eijk HM, Olde Damink SW, et al. D-amino Acids in health and disease: a focus on cancer. *Nutrients* 2019;11:2205.
2. Umeda S, Sujino T, Miyamoto K, et al. D-amino acids ameliorate experimental colitis and cholangitis by inhibiting growth of Proteobacteria: potential therapeutic role in inflammatory bowel disease. *Cell Mol Gastroenterol Hepatol* 2023;16:1011–1031.
3. Santana PT, Rosas SLB, Ribeiro BE, et al. Dysbiosis in inflammatory bowel disease: pathogenic role and potential therapeutic targets. *Int J Mol Sci* 2022;23:3464.
4. Srour B, Kordahi MC, Bonazzi E, et al. Ultra-processed foods and human health: from epidemiological evidence to mechanistic insights. *Lancet Gastroenterol Hepatol* 2022;7:1128–1140.
5. Davie JR. Inhibition of histone deacetylase activity by butyrate. *J Nutr* 2003;133:2485S–2493S.
6. Chervy M, Barnich N, Denizot J. Adherent-Invasive *E. coli*: update on the lifestyle of a troublemaker in Crohn's disease. *Int J Mol Sci* 2020;21:3734.
7. Darfeuille-Michaud A, Boudeau J, Bulois P, et al. High prevalence of adherent-invasive *Escherichia coli* associated with ileal mucosa in Crohn's disease. *Gastroenterology* 2004;127:412–421.

Correspondence

Address correspondence to: Benoit Chassaing, PhD, INSERM, U1016, Mucosal Microbiota in Chronic Inflammatory Diseases Team, CNRS UMR 8104, Université de Paris, 24 Rue du Faubourg Saint-Jacques, 75014 Paris, France. e-mail: benoit.chassaing@inserm.fr.

Conflicts of interest

The authors disclose no conflicts.

Funding

Supported by the Ministère de la Recherche et de la Technologie, Inserm (UMR 1071), INRAE (USC-1382), the French government's IDEX-ISITE initiative 16-IDEX-0001 (CAP 20-25), Agence Nationale de la Recherche (ANR-21-CE14-0014-01), the National Program "Microbiote" Inserm, and Association François Aupetit (J.D.); and by a Starting Grant from the European Research Council under the European Union's Horizon 2020 research and innovation program (ERC-2018-StG-804135), a Chaire d'Excellence from IdEx Université de Paris (ANR-18-IDEX-0001), an Innovator Award from the Kenneth Rainin Foundation, an award from the Fondation de l'Avenir (AP-RM-21-032), Agence Nationale de la Recherche grants EMULBIONT (ANR-21-CE15-0042-01) and DREAM (ANR-20-PAMR-0002), and the national program "Microbiote" from INSERM (B.C.). The funders had no role in writing the manuscript.



Most current article

© 2023 The Authors. Published by Elsevier Inc. on behalf of the AGA Institute. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).
2352-345X
<https://doi.org/10.1016/j.jcmgh.2023.09.009>