



**HAL**  
open science

# GENOME-WIDE ASSOCIATION STUDY OF INFECTIOUS PANCREATIC NECROSIS IN TWO SUCCESSIVE GENERATIONS OF RAINBOW TROUT

Jonathan D'ambrosio, Pierre Patrice, Yoannah François, Thierry Morin, Joëlle Cabon, Jean Ruche, Alexandre Desgranges, Pierrick Haffray, Florence Phocas

► **To cite this version:**

Jonathan D'ambrosio, Pierre Patrice, Yoannah François, Thierry Morin, Joëlle Cabon, et al.. GENOME-WIDE ASSOCIATION STUDY OF INFECTIOUS PANCREATIC NECROSIS IN TWO SUCCESSIVE GENERATIONS OF RAINBOW TROUT. *Aquaculture Europe* 2021, Oct 2021, Madeira, Spain. hal-04385905

**HAL Id: hal-04385905**

**<https://hal.inrae.fr/hal-04385905>**

Submitted on 10 Jan 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# GENOME-WIDE ASSOCIATION STUDY OF INFECTIOUS PANCREATIC NECROSIS IN TWO SUCCESSIVE GENERATIONS OF RAINBOW TROUT



J. D'Ambrosio<sup>1\*</sup>, P. Patrice<sup>1</sup>, Y. François<sup>1</sup>, T. Morin<sup>3</sup>, J. Cabon<sup>3</sup>, J. Ruche<sup>4</sup>, A. Desgranges<sup>4</sup>, P. Haffray<sup>1</sup>, F. Phocas<sup>2</sup>

<sup>1</sup>SYSAAF, French Poultry and Aquaculture Breeders Association, 35042 Rennes, France

<sup>2</sup>Université Paris-Saclay, INRAE, AgroParisTech, GABI, 78350 Jouy-en-Josas, France

<sup>3</sup>ANSES, Ploufragan-Plouzané-Niort Laboratory, Unit Virology, immunology and ecotoxicology of fish (VIMEP), 29280 Plouzané, France

<sup>4</sup>SARL Milin Nevez, 29610 Plouigneau, France

## Aims of the study


- Moderate heritability was estimated for Infectious Pancreatic Necrosis (IPN) resistance. However, there is little work to precisely characterize the architecture of IPN resistance in rainbow trout and **previous analyses only included a single cohort of phenotyped fish.**
- Therefore, the two objectives of this study were to **estimate heritability** and to **detect quantitative trait loci (QTLs)** associated with IPN resistance **in two successive generations of a commercial French rainbow trout line.**

## Material & methods

- Challenge : 4,000 fish were derived sibs from the Bretagne Truite multitraits breeding program challenged by immersion with IPN virus at the ANSES-SYSAAF Fortior Genetics platform.
- 3,885 fish and their 372 parents were genotyped with Axiom™ Trout Genotyping array (57K SNP).
- After quality control : 1,799 and 1,978 phenotyped and genotyped (for 25,512 SNPs) fish of two successive cohorts were retained.
- IPN resistance was calculated as a survival trait (dead/alive at 37 days).
- BLUPf90 package was used for the analysis.
- First cohort (C1)** is represented in **green color** and **second cohort (C2)** in **blue color** in following figures.

## Genetic architecture

- Moderate heritability for survival :  $h^2 = 0,20$




Manhattan plot showing associations between SNPs and IPN resistance. Lines correspond respectively to 1% chromosome-wide and 1% genome-wide significance thresholds (after Bonferroni correction).

- Two main QTLs for survival on chromosomes 1 and 16
- These QTLs were also observed in the two analyses of each fish cohort

## Survival of fish according to status in QTLs

### Chromosome 1




- For C2, survivals of  $Q_1Q_1$  and  $Q_1q_1$  fish are respectively **+27%** and **+11%** greater than  $q_1q_1$  trout survival.

Percentage of survival of fish according to their genotyping. Horizontal lines represent the average survival rates for each challenge.


### Chromosome 16

- For C2, survivals of  $Q_1Q_1$  and  $Q_1q_1$  fish are respectively **+23%** and **+10%** greater than  $q_1q_1$  trout survival.



## Survival of fish with combination of QTLs

### Chromosome 1 & chromosome 16



Percentage of survival of fish according to their genotyping. Horizontal lines represent the average survival rates for each challenge.

- The association of favorable alleles across QTLs allows a small additional increase in survival.

## Conclusion

- We confirm two significant and additive QTLs located on chromosomes 1 and 16. They were already identified in the literature as playing a role on IPN resistance (Santi et al. 2019; Ozaki et al. 2007).
- The two QTLs are observed in each of the two studied cohorts and are therefore consistently segregating QTLs in the population.
- These QTLs offer the possibility of marker-assisted selection for rapid dissemination of genetic improvement for IPN resistance of rainbow trout.