



HAL
open science

Deep Gaussian Process Metamodeling with Mixed Variables for Vegetative Filter Strip Design in Agricultural Context

Guerlain Lambert, Céline Helbert, Claire Lauvernet

► **To cite this version:**

Guerlain Lambert, Céline Helbert, Claire Lauvernet. Deep Gaussian Process Metamodeling with Mixed Variables for Vegetative Filter Strip Design in Agricultural Context. Journées Scientifiques du Consortium en mathématiques appliquées CIROQUO, Nov 2023, Palaiseau (France), France. hal-04389102

HAL Id: hal-04389102

<https://hal.inrae.fr/hal-04389102v1>

Submitted on 11 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Deep Gaussian Process Meta-Modeling with Mixed Variables for Vegetative Filter Strip Design in Agricultural Context

Guerlain Lambert¹, Céline Helbert¹, Claire Lauvernet²

¹ Univ. Lyon, UMR CNRS 5208 Ecole Centrale de Lyon ICJ, France
 guerlain.lambert@master.ec-lyon.fr

celine.helbert@ec-lyon.fr

² INRAE, RIVERLY, Villeurbanne, France
 claire.lauvernet@inrae.fr

Abstract

Conventional farming practices often result in soil erosion, loss of biodiversity, and pollution of soil and water resources. To mitigate pollution risks, European regulations require the implementation of vegetated buffer strips along watercourses. However, these buffer zones may not always be sufficient, necessitating additional buffer zones with dimensions tailored to each case. The Riverly lab at INRAE has developed the BUVARD tool to determine optimal buffer zone widths based on field data. The initial BUVARD model, which includes the VFSMOD numerical model [7, 5], is complex and requires numerous input variables with both quantitative and qualitative uncertainty, making it challenging for risk management [1].

Previous work has employed a metamodeling approach to simplify the use of BUVARD and aid users in designing buffer strips adapted to specific contexts [6, 4]. Bayesian and non-Bayesian surrogates, such as Polynomial Chaos Expansions (PCE) and Mixed-kriging, have been compared, addressing the mixed variables issue for GP and PCE, but not for Deep Gaussian Processes (DGP). In this work, we explore new surrogate methods for BUVARD with quantitative and qualitative variables: Deep Gaussian Processes (DGP) [2] and Deep Sigma Point Processes (DSPP) [3], which are capable of handling mixed variables. These methods leverage their deep structure to efficiently handle zero value plateaus, given the non-stationary and parsimonious structure of BUVARD data.




Figure 1: Depiction of the model chain integrated in BUVARD and its meta modelling as a function of only a few input variables, quantitative (black) and qualitative (green).

We examine various approaches for incorporating categorical variables into the DGP and DSPP models, including processing data by pairs of modalities, converting modalities to quantitative values, and using appropriate kernel combinations. Our findings contribute to the development of more accessible and effective buffer zone management tools for the agricultural community, addressing the challenges posed by mixed variables and the complex model chain in BUVARD, while also tackling non-stationarity and zero value plateaus. Furthermore, the metamodel serves as a relevant first guess for pollution reduction device design and provides an effective uncertainty tool for visualizing the impact of parameter uncertainty on filter efficiency in risk analysis and management.

References

- [1] C. Catalogne, Claire Lauvernet, and Nadia Carlier. Guide d'utilisation de l'outil buvard pour le dimensionnement des bandes tampons végétalisées destinées à limiter les transferts de pesticides par ruissellement. Technical report, irstea, 2018.
- [2] Andreas Damianou and Neil D. Lawrence. Deep Gaussian processes. In Carlos M. Carvalho and Pradeep Ravikumar, editors, *Proceedings of the Sixteenth International Conference on Artificial Intelligence and Statistics*, volume 31 of *Proceedings of Machine Learning Research*, pages 207–215, Scottsdale, Arizona, USA, 29 Apr–01 May 2013. PMLR.
- [3] Martin Jankowiak, Geoff Pleiss, and Jacob R. Gardner. Deep Sigma Point processes, 2020.
- [4] C. Lauvernet, C. Helbert, Z. Xujia, and B. Sudret. Metamodeling approaches to help designing vegetative filter strips and improve the water quality. In *EGU General Assembly 2022*, Vienna, Austria, May 23–27 2022.
- [5] C. Lauvernet and R. Muñoz Carpena. Shallow water table effects on water, sediment, and pesticide transport in vegetative filter strips - part 2: model coupling, application, factor importance, and uncertainty. *Hydrology and Earth System Sciences*, 22(1):71–87, 2018.
- [6] Claire Lauvernet and Céline Helbert. Metamodeling methods that incorporate qualitative variables for improved design of vegetative filter strips. *Reliability Engineering System Safety*, 204:107083, 2020.
- [7] Rafael Muñoz-Carpena, John E. Parsons, and J. Wendell Gilliam. Modeling hydrology and sediment transport in vegetative filter strips. *Journal of Hydrology*, 214(1-4):111–129, 1999.