



**HAL**  
open science

## Characterization of a uranium-tolerant green microalga of the genus *Coelastrella* with high potential for the remediation of metal-polluted waters

Camille Beaulier, Marie Dannay, Fabienne Devime, Adrien Galeone, Célia Baggio, Nabila El Sakkout, Camille Raillon, Olivier Courson, Jacques Bourguignon, Claude Alban, et al.

### ► To cite this version:

Camille Beaulier, Marie Dannay, Fabienne Devime, Adrien Galeone, Célia Baggio, et al.. Characterization of a uranium-tolerant green microalga of the genus *Coelastrella* with high potential for the remediation of metal-polluted waters. *Science of the Total Environment*, 2023, 908, pp.168195. 10.1016/j.scitotenv.2023.168195 . hal-04390380

**HAL Id: hal-04390380**

**<https://hal.inrae.fr/hal-04390380>**

Submitted on 31 Jan 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Characterization of a uranium-tolerant green microalga of the genus *Coelastrella* with high potential for the remediation of metal-polluted waters

Camille Beaulier<sup>a</sup>, Marie Dannay<sup>a</sup>, Fabienne Devime<sup>a</sup>, Adrien Galeone<sup>a</sup>, Célia Baggio<sup>a</sup>, Nabila El Sakkout<sup>a</sup>, Camille Raillon<sup>a</sup>, Olivier Courson<sup>b</sup>, Jacques Bourguignon<sup>a</sup>, Claude Alban<sup>a</sup>, Stéphane Ravanel<sup>a\*</sup>

<sup>a</sup>Univ. Grenoble Alpes, INRAE, CEA, CNRS, IRIG, LPCV, F-38000 Grenoble, France

<sup>b</sup>Univ. Strasbourg, UMR 7178, CNRS, IPHC, F-67000 Strasbourg, France

## \*Correspondence

stephane.ravanel@cea.fr

Univ. Grenoble Alpes, INRAE, CEA, CNRS, IRIG, LPCV, 38000 Grenoble, France

## Highlights


- A *Coelastrella* microalgal species was isolated from U-contaminated liquid wastes
- *Coelastrella* is highly tolerant to U compared to model green microalgae
- *Coelastrella* can accumulate remarkably high amounts of U from contaminated media
- *Coelastrella* is able to efficiently remediate U from metal-polluted natural waters
- *Coelastrella* accumulates lipids when growing in metal-polluted natural waters

**Keywords:** microalga, uranium, tolerance, accumulation, polluted waters, remediation

## Abstract

Uranium (U) contamination of terrestrial and aquatic ecosystems poses a significant threat to the environment and human health due to the chemotoxicity of this actinide. The characterization of organisms that tolerate and accumulate U is crucial to decipher the mechanisms evolved to cope with the radionuclide and to propose new effective strategies for the bioremediation of U-contaminated environments. Here, we isolated a unicellular green microalga of the genus *Coelastrella* from U-contaminated wastewater. We showed that *Coelastrella* sp. PCV is much more tolerant to U than *Chlamydomonas reinhardtii* and *Chlorella vulgaris*. *Coelastrella* sp. PCV is able to accumulate U very rapidly and then gradually release it into the medium, behaving as an excluder to limit the toxic effects of U. The ability of *Coelastrella* sp. PCV to accumulate U is remarkably high, with up to 240 mg of tightly bound U per g of dry biomass. *Coelastrella* sp. PCV is able to grow and maintain high photosynthesis in natural metal-contaminated waters from a wetland near a reclaimed U mine. In a single one-week growth cycle, *Coelastrella* sp. PCV is able to capture 25-55% of the U from the contaminated waters and shows lipid droplet accumulation. *Coelastrella* sp. PCV is a very promising microalga for the remediation of polluted waters with valorization of algal biomass that accumulates lipids.

Graphical abstract



## 1. Introduction

Over the last several decades, the anthropogenic pollution of aquatic ecosystems has emerged as a major issue with global implications (Hader et al, 2020). Freshwater ecosystems cover less than 1% of the Earth's surface but are important reservoirs of biodiversity. Consequently, their pollution not only compromises the supply of drinking water but also poses a threat to many organisms. One of the main source of ecosystem pollution is trace metal elements, also known as heavy metals. These elements are of particular concern because they are not degradable, persistent, and hazardous to all living organisms. Conventional physical and chemical technologies for metal remediation are costly, poorly effective in treating large metal-contaminated water bodies, and can have adverse effects on aquatic ecosystems (Gandhi et al, 2022). Phytoremediation is an emerging, ecofriendly, sustainable and cost-effective green technology using metal-accumulating land or aquatic plants, macro- or micro-algae, to remove toxic metals from polluted soils and waters (Plohn et, 2021a; Bhat et al, 2022). Microalgae are also promising organisms to combine wastewater reclamation and biomass valorization, as they accumulate high-value compounds such as neutral lipids and carotenoids in response to stress (Nayana et al, 2022). In this context, the characterization of organisms with high metal tolerance and accumulation capacity is particularly relevant and timely.

Uranium (U) is a ubiquitous element in the Earth's crust, with average concentrations of 3 ppm in soils and aquatic sediments, 3 ppb in sea water, and 0.01-30 ppb in freshwater, with a strong influence of the geochemical environment (Gandhi et al, 2022; Markich 2002). This radionuclide is primarily redistributed in the environment by anthropogenic activities related to U mining and milling industries, civil and military nuclear activities, and agricultural practices, mainly through phosphate fertilizers that are significantly contaminated with U (Anke et al, 2009; Vandenhove, 2002). Natural U is composed of >99%  $^{238}\text{U}$ , an isotope that is poorly radioactive but chemically toxic to all living organisms (Gao et al, 2019). In recent years, our knowledge of U toxicity and the molecular mechanisms used by land plants to respond to U stress has increased significantly (Chen et al, 2021). Recent studies shed light on the uptake of U through calcium channels in root cells (Sarhou et al, 2022) or by endocytosis in cultured

plant cells (John et al, 2022), and identified uranium-binding proteins in *Arabidopsis thaliana* (Vallet et al, 2023). The fate of U is less well understood in eukaryotic microalgae than in land plants.

Uranium toxicity has been studied in various microalgae grown in synthetic media, natural waters, or synthetic media mimicking natural waters (Franklin et al, 2000; Hogan et al, 2005; Gunter et al, 2008). Uranium uptake and toxicity in microalgae depend on its speciation, as in other organisms (Markich 2002; Gao et al, 2019; Chen et al, 2021). Thus, U toxicity is dependent on pH (Franklin et al, 2000; Lavoie et al, 2014), water hardness (Charles et al, 2002; Fortin et al, 2007), and its bioavailability is significantly decreased in the presence of dissolved organic carbon (Trenfield et al, 2011; Trenfield et al, 2012; Hogan et al, 2015). The consequences of U intoxication on microalgal cells have been mainly reported in *Chlamydomonas* and *Chlorella* species. They include a significant impairment of cell division (Franklin et al, 2000; Charles et al, 2002; Pradines et al, 2005; Hogan et al, 2005; Lavoie et al, 2014; Trenfield et al, 2011) and an inhibition of photosynthesis (Pradines et al, 2005; Herlory et al, 2013).

Most microalgae have the ability to bind high amounts of metals, and thus the potential to remove pollutants from wastewater (Plohn et al, 2021a). The rapid, reversible, and metabolism-independent binding of metals onto living or dead algal cells is referred to as adsorption or biosorption. The incorporation of metals inside cells through active or passive transport mechanisms is referred to as absorption or uptake. Several studies have described the biosorption of U onto microalgae. This process is dependent on U speciation, pH and cell viability (dead or living cells), but is independent on temperature, light or metabolic activity (Horikoshi et al, 1979; Zhang et al, 1997; Gunter et al, 2008; Vogel et al, 2010). Cell wall composition and architecture are critical determinants of metal biosorption capacity in microalgae (Spain et al, 2021). The combination of spectroscopic and imaging methods has indicated that carboxyl, amino and organic/inorganic phosphate groups are involved in U complexation to algal cell walls (Gunter et al, 2008; Vogel et al, 2010; Zhang and Luo, 2022). The maximum biosorption capacity for U has been investigated in various microalgae by using

very diverse experimental conditions (e.g. cell density, U concentration, incubation medium). The capacity measured for living cells ranged from 4 to 75 mg of U per g of dry biomass (Horikoshi et al, 1979; Zhang et al, 1997; Vogel et al, 2010; Baselga-Cervera et al, 2018; Zhang and Luo, 2022). Only a few reports described the accumulation of U inside microalgal cells. However, these results might be interpreted with caution because rigorous discrimination between adsorption and absorption is missing or cell intactness is not established (Fortin et al, 2004; Fortin et al, 2007; Lavoie et al, 2014; Cheng et al, 2023). In an extremotolerant *Chlamydomonas* sp., biosorption onto the cell wall accounted for >90% U accumulated by algal cells, with the rest irreversibly sequestered (Baselga-Cervera et al, 2018). Electron microscopy dispersive X-ray spectroscopy was in agreement with the presence of U deposits in both the cell wall and cytoplasm (Garcia-Balboa et al, 2013).

Most of the information on U accumulation and toxicity in microalgae was obtained from strains that had not been previously exposed to the radionuclide. However, examination of phytoplankton diversity in former U mining and milling sites in Salamanca, Spain, indicated that a few microalgal species were able to rapidly adapt and colonize extremely U-polluted water bodies (Garcia-Balboa et al, 2013; Baselga-Cervera et al, 2020). The extremotolerant *Chlamydomonas* sp. isolated from this extreme environment is a very promising strain for sustainable management of U tailings water and for deciphering the genetic basis of adaptation to U stress (Baselga-Cervera et al, 2018).

In this study, we report the isolation and characterization of a new green microalga of the genus *Coelastrella* from laboratory liquid waste contaminated with U. We show that this species, named *Coelastrella* sp. PCV, is much more tolerant to U than the reference green microalgae *Chlamydomonas reinhardtii* and *Chlorella vulgaris*. In addition, it is able to capture remarkably high amounts of this radionuclide from artificially contaminated media, accounting for up to 60% of its dry biomass. *Coelastrella* sp. PCV was also challenged with natural waters from a reclaimed U mine, deprived of essential nutrients and contaminated with toxic elements. *Coelastrella* sp. PCV was able to grow in natural waters and to capture 25 to 55% of the U initially present. The production of lipid droplets by stressed algal cells indicated that

*Coelastrella* sp. PCV is a good candidate for both the remediation of polluted waters and valorization of the algal biomass accumulating lipids.

## 2. Materials and methods

### 2.1. Algal strains and culture conditions

*Coelastrella* sp. PCV was isolated from wastes of culture media used to study U toxicity in plants (Berthet et al, 2018). Green algae blooms were observed in liquid wastes containing approx. 5  $\mu\text{M}$  uranyl nitrate, after a few weeks of storage in daylight at room temperature. To get rid of bacterial and fungal contamination, samples were plated onto Tris-acetate-phosphate (TAP) agar solidified medium (Harris, 2009) supplemented with a cocktail of antibiotics (ampicillin 500  $\mu\text{g.mL}^{-1}$ , cefotaxime 100  $\mu\text{g.mL}^{-1}$ ) and a broad-spectrum fungicide (carbendazim 40  $\mu\text{g.mL}^{-1}$ ) (Kan and Pan, 2010) Two rounds of antibiotic selection were sufficient to obtain axenic cultures of *Coelastrella* sp. PCV. The strains *Chlorella vulgaris* CCAP 211/11B and *Chlamydomonas reinhardtii* CC-125 have been used in this study.

Algal strains were maintained by weekly sub-culturing in TAP medium and growth at 21°C in continuous light at 40  $\mu\text{mol photons s}^{-1}.\text{m}^{-2}$ . For testing U toxicity, the amount of phosphate was decreased to limit the formation of precipitates (Markich, 2002). The resulting TAP LoP (low phosphate; pH 7.0) medium contained 50  $\mu\text{M}$  phosphate instead of 1 mM in TAP. TAP LoP was supplemented with uranyl nitrate and inoculated with algal cells from an exponentially growing culture in TAP. The growth of *Coelastrella* sp. PCV was also analyzed in photoautotrophic conditions using the Sueoka's high salt medium (HSM) medium (Harris 2009) and natural waters from the Rophin wetland (see below). Growth rates were assessed by cell counting using a Luna automated cell counter (Logos biosystems) for both *Coelastrella* sp. PCV and *Chlorella vulgaris*, or a Countess automated cell counter (ThermoFisher) for *Chlamydomonas reinhardtii*.

All experiments based on microalgal cultures in the presence of U or in natural waters from Rophin were repeated several times independently. An independent experiment is defined as



a repetition performed at a different period with n=1-3 replicates corresponding to independent cultures (flasks).

## **2.2. Photosynthetic parameters**

Algal cells (200  $\mu\text{L}$ ) were placed in a black-bottom 96-well plate and dark-adapted for 30 min at room temperature (21°C) before measurements. Chlorophyll fluorescence was measured using a Maxi-Imaging PAM fluorometer (Heinz Walz). The maximum quantum efficiency of photosystem II photochemistry ( $F_v/F_m$ ) was calculated as  $F_v/F_m = (F_m - F_0)/F_m$ , where  $F_0$  is the minimal fluorescence in the dark-adapted state and  $F_m$  the maximum fluorescence in the dark-adapted state (Maxwell and Johnson, 2000). The electron transport rate (ETR) was calculated as  $\text{ETR} = Y(\text{II}) \times 0.5 \times 0.84 \times \text{PAR}$ , where  $Y(\text{II})$  is the effective quantum yield  $(F_m' - F)/F_m'$  and PAR is the incident photosynthetically active radiation (pulsing from 0 to 700  $\mu\text{mol photons} \cdot \text{s}^{-1} \cdot \text{m}^{-2}$ ).

## **2.3. Water sampling at the Rophin mining site**

Water samples were collected close to the former U mining site of Rophin (Puy-de-Dome, France) (Martin et al, 2020). The Rophin site is a pilot area for a French multidisciplinary scientific network studying territories concerned by the presence of natural and anthropogenically-enhanced radioactivity (ZATU, Zone Atelier Territoires Uranifères; <https://zatu.org/>). Water samples were collected in June 2019 at three points of the Rophin area. Dose rates were measured using a FH 40G-L10 survey radiometer (Thermo Scientific™). Sample E1 was from circulating water collected at the southern boundary of the mine (geographic coordinates 46.00846,3.55247; measured gamma dose rate 240 nSv/h). Water samples H1 and H2 were collected in the wetland located 200 meters downstream of the Rophin site (South-West direction). Sample H2 was collected from a stream (46.0069,3.54995; 600-800 nSv/h) while sample H1 was pore water from water-saturated soil (46.00696,3.55017; 1000-1300 nSv/h). Water samples were filtrated using fritted glass within

8 hours after sampling, then stored at 4°C. They were filtrated using 0.2 µm Nylon filters before use for chemical analysis or algal cultivation assays.

#### **2.4. Inductively coupled plasma mass spectrometry analyses**

*Coelastrella* cells challenged with U in TAP LoP medium or in natural waters were sampled every day (2 mL of culture) and centrifuged for 4 min at 9,400 *g*. Supernatants were collected and acidified with 0.65 % (w/v) nitric acid before ICP-MS analysis. Cell pellets were washed three times with 500 µL sodium carbonate (Na<sub>2</sub>CO<sub>3</sub>) 10 mM, then dried overnight at 80°C. Dried samples were resuspended in 100 µL of 65% (w/v) HNO<sub>3</sub> and mineralized for 2 hours at 90°C. Samples were diluted to the appropriate concentration and analyzed using an iCAP RQ quadrupole mass instrument (Thermo Fisher Scientific GmbH, Germany). <sup>238</sup>U concentration (in ppb) was determined using a standard curve and corrected using the internal standard <sup>172</sup>Yb (standard acquisition mode).

For elemental analysis of waters from the Rophin area, samples were filtrated through 0.2 µm nylon filters and acidified with 0.65 % (w/v) HNO<sub>3</sub>. Elements were analyzed using the standard mode (<sup>7</sup>Li, <sup>11</sup>B, <sup>23</sup>Na, <sup>24</sup>Mg, <sup>25</sup>Mg, <sup>27</sup>Al, <sup>31</sup>P, <sup>39</sup>K, <sup>95</sup>Mo, <sup>98</sup>Mo, <sup>107</sup>Ag, <sup>109</sup>Ag, <sup>111</sup>Cd, <sup>206</sup>Pb, <sup>208</sup>Pb, <sup>238</sup>U) and the collision mode with helium as a cell gas (<sup>31</sup>P, <sup>39</sup>K, <sup>44</sup>Ca, <sup>52</sup>Cr, <sup>53</sup>Cr, <sup>55</sup>Mn, <sup>56</sup>Fe, <sup>57</sup>Fe, <sup>59</sup>Co, <sup>63</sup>Cu, <sup>65</sup>Cu, <sup>64</sup>Zn, <sup>66</sup>Zn, <sup>69</sup>Ga). Quantification was as described above using the internal standards <sup>103</sup>Rh and <sup>172</sup>Yb.

#### **2.5. Determination of total dissolved organic carbon**

Water samples filtrated using 0.2 µm Nylon filters were used for carbon analysis. Organic carbon concentration was determined by catalytic combustion (680°C) using a TOC-VCPH analyzer (Shimadzu). Total dissolved organic carbon (TOC) was determined by two methods. First, TOC was calculated from inorganic carbon (IC) and total carbon (TC) measurements using the formula TOC=TC-IC. Second, non-purgeable organic carbon (NPOC) measurements were carried out to verify TOC values obtained by the TC-IC determination.

## 2.6. Phylogenetic studies

Single colonies of *Coelastrella* sp. PCV picked from TAP-agar were used for PCR amplification of the 18S rDNA gene. Colonies were dispersed in sterile water, sonicated for 1 min, and the suspension was used as a DNA template in PCR reactions. Primers 5'-ACCTGGTTGATCCTGCAG-3' and 5'-TGATCCTCCYGCAGGTTTAC-3' were used to amplify the 18S rDNA region (Khaw et al, 2020). PCR products were cloned in the pSC101 vector and sequenced. The partial 18S rRNA gene sequence from *Coelastrella* sp. PCV was deposited to the GenBank database with accession number OQ202154. This sequence was compared to the Chlorophyceae sequence dataset using BLASTn from NCBI (<https://blast.ncbi.nlm.nih.gov/>). Thirty-one 18S rDNA sequences from the top ranking list were retrieved and analyzed using the Mega X software (Kumar et al, 2018): *Coelastrella tenuithea* FACHB-2315 (GenBank accession numbers MH176106), *Coelastrella tenuithea* FACHB-2313 (MH176107), *Coelastrella tenuithea* FACHB-2314 (MH176108), *Coelastrella vacuolata* (X56104), *Coelastrella vacuolata* MACC-549 (MW979811), *Coelastrella* sp. KNUA037 (KT883907), *Coelastrella thermophila* FACHB-2304 (MH176098), *Coelastrella thermophila* FACHB-2303 (MH176094), *Coelastrella* sp. YACCYB176 (MH636637), *Coelastrella thermophila* FACHB-2309 (MH176099), *Coelastrella* sp. YACCYB207 (MH636662), *Coelastrella* sp. YACCYB208 (MH636663), *Coelastrella* sp. SAG2471 (KM020087), *Coelastrella* sp. SAG2470 (KM020171), *Coelastrella* sp. SAG2123 (JX513883), *Coelastrella saipanensis* CCAP217/9 (JX316760), *Coelastrella saipanensis* (AB055800), *Coelastrella saipanensis* FACHB-2294 (MH176093), *Coelastrella saipanensis* LY31-2 (MF407353), *Coelastrella* sp. IN-2019 SYKOA Ch-047-11 (MK478814), *Coelastrella* sp. ACSSI083 (MG696570), *Coelastrella oocystiformis* FACHB-2312 (MH176105), *Enallax costatus* SAG46.88 (AB773883), *Tetradesmus arenicola* WD-1-6 (MH703774), *Tetradesmus arenicola* SAG2564 (MH703775), *Tetradesmus obliquus* CCAP276/1A (MW471023), *Tetradesmus obliquus* BEA D01\_12 (MW065553), *Scenedesmus rubescens* IPPAS D-292 (KU057946), *Scenedesmus rubescens* KMMCC263 (JQ315585), *Scenedesmus pectinatus* MIC-G8 (JF834707), *Scenedesmus pectinatus* var. *distendus* (AB037093). Sequences were aligned

using Muscle (default settings). The phylogenetic analysis was done by using the Maximum Likelihood method and Kimura 2-parameter mode. A discrete Gamma distribution was used to model evolutionary rate differences among sites (5 categories (+G, parameter = 0.0500)). The rate variation model allowed for some sites to be evolutionarily invariable ([+I], 48.22% sites). The bootstrap consensus tree was inferred from 1000 replicates.

### **2.7. Light microscopy and Nile Red staining of lipid droplets**

The morphological analysis of *Coelastrella* cells was done using a Zeiss Axioplan 2 microscope. Images were acquired in bright field with magnification x63/x100 and post-processed using Fiji/ImageJ. The accumulation of lipid droplets was monitored using the Nile Red fluorescent staining procedure (Jaussaud et al, 2020). *Coelastrella* cells grown in HSM and waters from the Rophin area were incubated for 20 min in darkness with Nile Red (Sigma Aldrich) at a concentration of 0.5  $\mu\text{g}\cdot\text{mL}^{-1}$ . Cells were imaged using an inverted confocal microscope (LSM880 Zeiss) equipped with a 63x/1.4NA oil objective. Laser beams at 561 nm and 633 nm were used to excite Nile Red and chlorophyll, respectively. The software control was done using Zen Black and image post-processing was done using Fiji/ImageJ.

## **3. Results**

### **3.1. Isolation, genetic identification, growth and morphological characteristics of *Coelastrella* sp. PCV**

The development of a green algal bloom was observed in laboratory liquid wastes from culture media used for the study of U stress in plants (Berthet et al, 2018). In order to identify the green alga(e) able to grow in this medium contaminated by uranyl nitrate ( $\text{UO}_2(\text{NO}_3)_2$ ; about 5  $\mu\text{M}$ , 1.2 ppm), we performed two rounds of selection on antibiotics/fungicide and obtained an axenic algal culture (**Figure S1**). Colony PCR with 18S rDNA primers was used for the genetic identification of selected microalgae. A unique partial 18S rDNA sequence was obtained from all tested colonies. It showed >99% sequence identity with 18S rDNA sequences of microalgae

from the Scenedesmaceae family (within the Chlorophyceae class and Sphaeropleales order), including members of the genera *Coelastrella*, *Scenedesmus*, *Tetradesmus* or *Asterarcys*. A phylogenetic tree inferred from 31 related 18S rDNA sequences showed that the isolated microalga was clustered with *Coelastrella* species (**Figure 1A**). The genus *Coelastrella* was first reported by Chodat (Chodat, 1922) and 24 species isolated from aero-terrestrial or freshwater habitats are currently described in the AlgaeBase (algaebase.org) (Guiry, 2021). The strain selected from liquid waste is likely to have originated from airborne contamination of the laboratory environment. Since the isolate does not correspond to any described species, it was named *Coelastrella* sp. PCV (with reference to the laboratory acronym).

*Coelastrella* sp. PCV was maintained under mixotrophic growth conditions in Tris-acetate-phosphate (TAP) medium in continuous light, either in liquid cultures with weekly subcultures or on TAP-agar plates. The microalga was also able to grow under photoautotrophic conditions in media such as the Sueoka's high salt medium (HSM) or Bold's basal medium (BBM), in continuous light or light/dark cycles. *Coelastrella* sp. PCV is a fast-growing microalga in mixotrophic or autotrophic conditions and batch cultures reach a maximal cell density of 5-6 million cells.mL<sup>-1</sup> at the stationary phase. The maximal cell density is quite low and can be explained by the large size of algal cells (Agusti and Kalff, 1989). Indeed, the morphological analysis of *Coelastrella* sp. PCV by light microscopy showed green unicellular and spherical cells with an average diameter of  $9 \pm 3 \mu\text{m}$  (**Figure 1B**). Cell size distribution in liquid cultures is quite variable, with young cells of  $7 \pm 1 \mu\text{m}$  in diameter and mature cells up to  $16 \pm 2 \mu\text{m}$  in diameter. Vegetative reproduction occurs by autospore formation, with sporangia containing 2 to 16 daughter cells (**Figure 1B**). Autospores are released after the rupture of the parental cell wall. Sexual reproduction was not observed in standard growth conditions.

### **3.2. Tolerance of *Coelastrella* sp. PCV to uranium in synthetic media**

The isolation of *Coelastrella* sp. PCV from liquid wastes contaminated by U prompted us to analyze its tolerance to this radionuclide, which is chemically toxic to photosynthetic organisms (Chen et al, 2021). The speciation of U in natural or artificial environments is very complex and

determines its toxicity to living organisms (Gao 2019). In particular, the presence of phosphate in plant or algal culture media is known to limit U bioavailability due to U-phosphate complex formation and precipitation (Markich, 2002; Fortin et al, 2004; Pradines et al, 2005). We found that phosphate limitation in TAP medium did not impaired significantly the growth of *Coelastrella* sp. PCV (**Figure 2**). In contrast, phosphate limitation in phototrophic media was accompanied with a drastic reduction in cell division, which was not compatible for measuring U toxicity. Thus, the tolerance of *Coelastrella* sp. PCV to U stress was analyzed in a TAP medium containing 50  $\mu\text{M}$  of phosphate (TAP LoP).

*Coelastrella* sp. PCV cells were challenged with 50 to 400  $\mu\text{M}$  uranyl nitrate in TAP LoP for 4 days. Under these conditions, U concentrations were stable over time in control experiments without algal cells. Growth was measured by cell counting and photosynthetic activity was used as a proxy to assess U toxicity (Herlory et al, 2013). As shown in **Figure 2A**, algal growth was inhibited by U in a dose-dependent manner. At the highest U concentration tested (400  $\mu\text{M}$ ), the growth of *Coelastrella* cells was severely impaired. Determination of the doubling time during the exponential growth phase allowed calculation of a half-maximal inhibitory concentration ( $\text{IC}_{50}$ ) of 80  $\mu\text{M}$  (**Figure S2A**). Chlorophyll fluorescence was used to measure the maximum quantum yield of photosystem II in dark-adapted cells ( $F_v/F_m$ ) and the electron transport rate (ETR) at various light intensities (Maxwell et al, 2000). We did not observed significant effect of U on photosynthetic parameters for concentrations up to 200  $\mu\text{M}$ . However, a significant decrease in  $F_v/F_m$  was observed at 400  $\mu\text{M}$  uranyl nitrate, starting from 1 day of treatment and reaching 65% reduction after 4 days of U exposure (**Figure 2B**). In addition, the ETR measured during the exponential growth phase was decreased by 40% at the highest light intensity (700  $\mu\text{mol photons}\cdot\text{s}^{-1}\cdot\text{m}^{-2}$ ) compared to the control TAP LoP condition (**Figure 2C**). Dose-effect relationships gave  $\text{IC}_{50}$  values of >400  $\mu\text{M}$  for ETR and >500  $\mu\text{M}$  for  $F_v/F_m$  (**Figure S2B-C**), indicating that the photosynthetic machinery of *Coelastrella* cells was moderately affected by U stress. In line with these results, when cells challenged for 4 days with 400  $\mu\text{M}$  uranyl nitrate were reintroduced into fresh TAP medium, growth and photosynthetic rates were comparable to unstressed cells, demonstrating that *Coelastrella* sp.

PCV was able to survive severe U stress. Together, these experiments showed that *Coelastrella* sp. PCV is tolerant to high U concentrations. Noteworthy, tolerance experiments have been conducted over a 3-year period since the isolation of the microalga. Growth curves and photosynthetic properties were similar throughout the period, indicating that the ability of *Coelastrella* sp. PCV to tolerate U is genetically fixed.

The physiological consequences of U stress in green microalgae have been reported in the models *Chlamydomonas* and *Chlorella* (Franklin et al, 2000; Charles et al, 2002; Pradines et al, 2005; Hogan et al, 2005; Trenfield et al, 2011; Herlory et al, 2013; Lavoie et al, 2014). Comparing U tolerance of *Coelastrella* sp. PCV (this work) with data from the literature is not relevant as U speciation, bioavailability and therefore toxicity differ significantly due to the variety of stress conditions. Therefore, we included *Chlamydomonas reinhardtii* and *Chlorella vulgaris* in our study for comparison. As previously, cell growth and photosynthetic parameters were measured to assess tolerance to U stress in TAP LoP containing uranyl nitrate. The inhibitory effect of U in *C. reinhardtii* and *C. vulgaris* was observed from 50  $\mu\text{M}$  uranyl nitrate (**Figure 3; Figure S3**), and growth was fully abolished and  $F_v/F_m$  values were null at 100  $\mu\text{M}$ , indicating that the cells were dead. Under these conditions, the growth of *Coelastrella* sp. PCV was only reduced by 50% and  $F_v/F_m$  was not affected. These data show that *Coelastrella* sp. PCV is much more tolerant to U than freshwater model microalgae.

Next, we addressed the fate of U in algal cultures over a 4-day incubation period. The amount of U remaining in the medium and tightly bound to algal cells was determined by ICP-MS. To this end, algal cells were extensively washed with sodium carbonate to remove U that is loosely adsorbed to the cell wall (Revel et al, 2022; Sarthou et al, 2022). For uranyl nitrate concentrations ranging from 50 to 200  $\mu\text{M}$ , the kinetics of U accumulation in the medium and cells clearly showed two main phases (**Figure 4A-C**). During the first 24 hours, U was markedly depleted from the medium and reached its highest accumulation level in the cells. Eighty to 95% of U initially present in the medium was captured by algal cells, of which a large amount was tightly bound and could not be released with carbonate. From 48 hours, a second phase corresponding to a gradual release of U from the cells to the medium was observed (**Figure**

**4A-C).** As a result of this release process, the amount of U in the medium returned to 50 to 80% of its initial concentration after 4 days of culture. When cells were challenged with 400  $\mu\text{M}$  uranyl nitrate, the first phase was similar to that observed for lower metal concentrations, with a strong decrease (90%) of U in the medium and the accumulation of tightly-bound U in the cells (**Figure 4D**). The second phase was different, however, because we did not observe U release from cells but, instead, constant levels of U in the cells and medium until the end of the stress period (**Figure 4D**). These data suggest that U release from *Coelastrella* cells is a defense process whose efficiency is dependent on stress intensity and algal physiology (**Figure 2**).

### **3.3. Accumulation of uranium in *Coelastrella* sp. PCV**

To gain further insight into the accumulation of U by *Coelastrella* sp. PCV, we analyzed the fate of the radionuclide during the first 24-hour period. Algal cells were incubated with different concentrations of uranyl nitrate (from 50 to 400  $\mu\text{M}$ ) in the TAP LoP medium and aliquot fractions were withdrawn at regular time intervals to measure U in the medium, in whole cells, and in cells washed with sodium carbonate. As shown in **Figure 5A**, 80 to 98% of U initially present in the medium was depleted within 1.5 hours and the amount of U was maintained at a very low level up to 24 hours. A very rapid accumulation of U in *Coelastrella* cells was concomitant with these kinetics (**Figure 5B**). Because the formation of U precipitates can interfere to some extent with whole-cell measurements, we further investigated metal accumulation in cells that had been thoroughly washed with carbonate. At 24 hours of incubation, the amount of U tightly bound to algal cells ranged from 30 to 50% of total U (**Figure 5C**). To estimate the maximum capacity of *Coelastrella* cells to capture U, we plotted the amount of U accumulated in cells (values from **Figure 5BC**, expressed on a dry mass basis) as function of the initial U concentration in the medium. As shown in **Figure 5D**, both the total amount of U and the tightly-bound fraction increased linearly with U concentration in the medium. These data indicate that the accumulation (adsorption and absorption) of U by *Coelastrella* cells was not saturated at 400  $\mu\text{M}$  uranyl nitrate in TAP LoP medium. In these



conditions, *Coelastrella* sp. PCV was able to capture up to 600 mg U per gram dry biomass, of which 240 mg.g<sup>-1</sup> DW was tightly bound and/or incorporated within algal cells. Together, these data show that *Coelastrella* sp. PCV has a remarkably high capacity to accumulate U.

#### 3.4. Remediation of natural polluted waters by *Coelastrella* sp. PCV

To analyze the ability of *Coelastrella* sp. PCV to remediate natural waters polluted by U, we collected water samples at the vicinity of the former U mining site of Rophin (Puy-de-Dome, France). Ore bodies from the Rophin site are rich in parsonsite, a lead-uranium-phosphate mineral ( $\text{Pb}_2(\text{UO}_2)(\text{PO}_4)_2 \cdot 2\text{H}_2\text{O}$ ), and were exploited between 1949 and 1957 (Martin et al, 2020). During this period and before complete rehabilitation of the site, the wetland located a few hundred meters downstream from the mine accumulated U of anthropogenic origin. Water samples were collected from three locations of the Rophin site with measured gamma dose rates of 240, 600-800 and 1000-1300 nS.h<sup>-1</sup> for samples E1, H2, and H1, respectively. The local geological background of the area was 200-250 nS.h<sup>-1</sup> (Martin et al, 2020). The elemental composition of waters from Rophin was analyzed by ICP-MS (**Table S1**). Waters were found to be quite poor in elements that are essential for microalgal growth, with magnesium, potassium, phosphorus, calcium or iron in the micromolar concentration range, and manganese, copper, zinc or molybdenum in the nanomolar concentration range. Also, Rophin waters contain non-essential and potentially toxic elements, including U at a concentration of  $15 \pm 1$ ,  $62 \pm 12$  and  $228 \pm 40$  nM in the E1, H2, and H1 waters, respectively (**Table S1**). Total dissolved organic carbon (TOC) was also determined using the differential (total carbon minus dissolved inorganic carbon) and NPOC methods. Both methods gave similar results and indicated that the H1 pore water contained 3 to 4 times more TOC ( $18.7 \pm 3.9$  mg.L<sup>-1</sup>) than the E1 ( $4.2 \pm 0.6$  mg.L<sup>-1</sup>) and H2 ( $4.8 \pm 1.1$  mg.L<sup>-1</sup>) flowing waters (**Table S1**).

We analyzed the ability of *Coelastrella* sp. PCV to proliferate in Rophin waters. As shown in **Figure 6A**, *Coelastrella* sp. PCV was able to grow exponentially in all three waters to reach a maximal cell density of 2-2.5 million cells.mL<sup>-1</sup>. The stationary phase was reached 4 days after inoculation while growth continued in the control HSM medium, suggesting growth impairment

due to nutrient limitation and/or toxicity of contaminating metals (e.g. U, lead, aluminum) (**Table S1**). Monitoring of Fv/Fm and ETR indicated that *Coelastrella* was able to maintain significant photosynthetic activity for at least 7 days of growth in Rophin waters (**Figure 6B-C**).

The distribution of U between polluted waters and *Coelastrella* cells was analyzed during a 7-day growing period. As the goal of this experiment was to evaluate the remediation potential of *Coelastrella* sp. PCV, we first determined the total amount of U associated with the algal biomass, including loosely (carbonate washable) and tightly (unaffected by carbonate treatment) bound U. Data in **Figure 7A-B** show that growth of *Coelastrella* sp. PCV in the E1 and H2 waters resulted in the removal of 50 to 55% of U after 2 to 3 days of growth. In the H1 water, the removal of U was slower and less efficient with a 25% decrease of U concentration in 7 days (**Figure 7C**). In contrast with accumulation profiles observed in synthetic TAP LoP medium containing up to 200  $\mu$ M uranyl nitrate (**Figure 4**), there was no release of U captured by algal cells during prolonged incubation periods. After 7 days of growth in Rophin waters, 40 to 50% of U was tightly bound to algal cells and the rest could be released by washing with sodium carbonate (**Figure 7D**). In addition to U, Rophin waters are significantly contaminated by lead (**Table S1**), mostly originating from the decay chain of U as demonstrated by the analysis of lead isotopy (Martin et al, 2020). The ability of *Coelastrella* sp. PCV cells to bind lead was analyzed. After 7 day of growth, *Coelastrella* cells captured 40 to 45% of lead present in E1 and H2 waters, and 5% from the H1 water (**Figure S4**). Thus, *Coelastrella* sp. PCV was able to reclaim waters from at least two heavy metals (U and lead) resulting from the anthropogenic contamination of the Rophin wetland.

The growth of *Coelastrella* sp. PCV in waters from the Rophin wetland was significantly reduced as compared to a nutrient-sufficient and contaminant-free synthetic medium (**Figure 6A**). The impairment of algal cell growth due to nutrient deprivation or another stressful situation is known to trigger neutral lipid production (Sun et al, 2018). Under adverse conditions, several *Coelastrella* species have been shown to accumulate high amount of lipids with fatty acid profiles of interest for the biofuel industry (Nayana et al, 2022). Lipid production in *Coelastrella* sp. PCV grown in Rophin waters was investigated using the Nile red staining

procedure. As shown in **Figure 8**, several large lipid droplets were observed in the cytoplasm of *Coelastrella* cells grown in either E1, H1 or H2 waters. As a comparison, cells grown in HSM displayed very small lipid droplets. We also analyzed lipid production in *Coelastrella* cells challenged with uranyl nitrate in TAP LoP medium. As shown in **Figure S5**, *Coelastrella* cells accumulated neutral lipid droplets in response to U stress. These data indicate that the growth of *Coelastrella* sp. PCV in nutrient-deprived and metal-polluted waters, or in a synthetic medium supplemented with U, triggers a reorientation of cellular metabolism towards the production and accumulation of neutral lipids, which is of potential interest for industrial and technological applications.

#### 4. Discussion

The green unicellular microalga isolated from U-contaminated liquid wastes belongs to the genus *Coelastrella*. Genetically, 18S rDNA sequence analysis indicated that *Coelastrella* sp. PCV is related to at least one *Coelastrella saipanensis* strain, but branch support values are not enough robust to allow accurate determination at the species level (**Figure 1A**). The morphological characteristics of *Coelastrella* sp. PCV (**Figure 1B**) did not allow to refine this classification but are in agreement with this relationship with *C. saipanensis* (Smith-Baedorf et al, 2013; Wang et al, 2019). The genus *Coelastrella* was first described a century ago (Chodat, 1922) and is currently represented by dozen of species (Guiry, 2021; Nayana et al, 2022). These microalgae have been found worldwide in a variety of aero-terrestrial and aquatic environments. Some *Coelastrella* species are characterized by the harshness of their natural habitat, including warm water from Roman baths (Smith-Baedorf et al, 2013), municipal wastewater in Nordic climate (Ferro et al, 2018), or dry terrestrial surfaces exposed to high light and elevated temperature (Narayanan et al, 2018; Kawasaki et al, 2020).

The ability of *Coelastrella* species to tolerate metal intoxication or to live in habitats contaminated with metals has been reported. Toxicity evaluation of zinc and iron oxide nanoparticles in *Coelastrella terrestris* indicated that this species has the potential for the remediation of metallic nanoparticles (Saxena and Harish, 2019; Saxena et al, 2020).

*Coelastrella* sp. (3-4) and *Coelastrella* sp. BGV displayed high absorption capacity for cadmium, copper or lead, and can remove metals from contaminated environments very efficiently (Plohn et al, 2021b; Karcheva et al, 2022). Also, a *Coelastrella* sp. was isolated from copper mine tailings sand (Wu et al, 2022). Co-inoculation of this microalgal species and two native fungi onto the contaminated tailings sand resulted in a 15% decrease of copper content. Here, we describe for the first time the physiological response and accumulation capacity of a *Coelastrella* species exposed to U. The tolerance to U has been previously investigated in *Chlorella* sp. (Franklin et al, 2000; Charles et al, 2002), *C. reinhardtii* (Lavoie et al, 2014), or *Ankistrodesmus* sp. (Cheng et al, 2023). Since U toxicity has been studied in media with different U speciation and bioavailability, a comparison of U tolerance between these microalgae is not relevant. In this study, we compared the tolerance of *Coelastrella* sp. PCV, *C. reinhardtii*, and *C. vulgaris* in similar conditions and could determine that *Coelastrella* sp. PCV is the most U-tolerant species. In the TAP LoP medium, *C. reinhardtii* and *C. vulgaris* cells died at 100  $\mu$ M uranyl nitrate while *Coelastrella* sp. PCV was still able to grow (albeit at a reduced rate) and maintain efficient photosynthesis up to 200  $\mu$ M uranyl nitrate (**Figure 3**).

A U-tolerant microalgal species has been formerly isolated from open pits and tailings ponds at Spanish mining sites extensively contaminated with metals, primarily U and zinc. (Baselga-Cervera et al, 2018). This *Chlamydomonas* sp. has the ability to grow rapidly in natural contaminated waters from U mine and in synthetic media containing U. A comparison of U tolerance between this *Chlamydomonas* sp. and *Coelastrella* sp. PCV is not possible for the above mentioned reasons. However, it can be hypothesized that these two U-tolerant algae are derived from different processes. On the one hand, the capacity of *Coelastrella* sp. PCV to tolerate elevated U concentrations is probably due to its genetic background (prior to colonization of U-containing liquid waste) and some physiological and molecular acclimation to the U-contaminated environment from which it was isolated. Indeed, the short duration (a few weeks) and low intensity (1.2 ppm U) of the stress cannot be considered an extreme condition leading to genetic adaptation through the selection of spontaneous mutations (Garcia-Balboa et al, 2013). On the other hand, the U-tolerant *Chlamydomonas* sp. is resulting

from an adaptive process that took place over several years to ensure survival under very hostile conditions (U ranging from 25 to 48 ppm) (Baselga-Cervera et al, 2018).

Several mechanisms of tolerance to U have been described in microorganisms (Kolhe et al, 2018). They include metal biosorption onto the cell wall, metal uptake and then sequestration by compartmentalization, metal detoxification through binding to chelators (metabolites or proteins/polypeptides), metal exclusion by efflux transport, metal biomineralization (formation of an insoluble mineral precipitate inside or outside the cell), or metal bioreduction (reduction of soluble U(VI) into insoluble U(IV)). When challenged with U, *Coelastrella* cells very rapidly capture the metal from the medium, demonstrating a high capacity for U adsorption (binding to the cell wall) and absorption (uptake). In a second stage, much slower and dose-dependent, the alga is able to progressively release U into the environment, behaving as an excluder to limit the toxic outcomes of U. This process of desorption (from the cell wall) and efflux (from the cytoplasm of algal cells) was observed for U concentrations ranging from 50 to 200  $\mu$ M in TAP LoP (**Figure 4A-C**). It was not detected when *Coelastrella* cells were cultivated in natural waters from Rophin (**Figure 7**), in which U is much less abundant and bioavailable (**Table S1**), or in TAP LoP containing 400  $\mu$ M uranyl nitrate (**Figure 4D**), in which algal physiology and metabolism are significantly reduced (**Figure 2**). The mechanisms behind U desorption and efflux by *Coelastrella* cells are still unknown. However, they are only active when the amount of U accumulated in cells is sufficient to trigger stress signaling, and can be overwhelmed when cells are not physiologically and metabolically competent. Vogel *et al.* (2010) previously observed desorption of U from *C. vulgaris* living cells during prolonged exposure (2-3 days) to the metal. The authors suggested, but did not demonstrate, that the release of metabolites such as organic acids could explain this behavior (Vogel et al, 2010). Recently, an unusual U biomineralization process was described in a microalga of the genus *Ankistrodesmus* (Cheng et al, 2023). While the mechanisms of U biomineralization described in some bacteria and fungi involve the formation of U-phosphate complexes (Kolhe et al, 2018), intra- and extracellular crystals of a mineral containing U and potassium (compreignacite) were observed in *Ankistrodesmus* sp., suggesting an important role of potassium in response to U stress.

Further analyses will be required to specifically address whether U biomineralization is part of the defense mechanisms evolved by *Coelastrella* sp. PCV to tolerate U.

The rapid, reversible and energy-independent biosorption of U onto various microalgae has been described (Horikoshi et al, 1979; Zhang et al, 1997; Fortin et al, 2004; Vogel et al, 2010; Baselga-Cervera et al, 2018). It is difficult to compare U biosorption capacity in different species because experimental parameters such as algal biomass, medium composition, pH, contact time or U concentration that influence this process are not similar. However, our data showed that the ability of *Coelastrella* sp. PCV to capture U is remarkably high. Indeed, titration of U accumulation indicated that up to 600 mg U per g DW was sorbed to *Coelastrella* cells, of which 240 mg.g<sup>-1</sup> DW was tightly bound or internalized (i.e. not extractable with carbonate). These values do not reflect the maximum accumulation capacity of *Coelastrella* sp. PCV because titration of this process did not show saturation, even at 400 µM U (**Figure 5**). For comparison, the original and artificially modified U-tolerant *Chlamydomonas* sp. isolated from U mine tailings accumulated 4 to 6 mg U per g DW (Baselga-Cervera et al, 2018) and the highest adsorption capacity reported to date for a green microalga was 75 mg.g<sup>-1</sup> DW for *Scenedesmus obliquus* (Zhang et al, 1997), a species belonging to the Scenedesmaceae family, as does *Coelastrella* sp. PCV. It is well known that the capacity of algae to bind cationic metals is due to the anionic nature of functional groups of cell wall components (Spain et al, 2021). The involvement of carboxylic and organic/inorganic phosphate groups in the coordination of U by *C. vulgaris* cells was demonstrated (Gunther et al, 2008; Vogel et al, 2010). The very high capacity of *Coelastrella* sp. PCV to accumulate U suggests i/ a unique composition of its cell wall as compared with other microalgae, and/or ii/ a high potential to absorb the radionuclide inside algal cells. Cell wall analysis of *Coelastrella* sp. (3-4) revealed a high quantity of polysaccharides, and thus a highly hydrophilic cell surface, and pointed out important changes in cell wall composition and thickness in relation with growth conditions (Spain and Funk, 2022). The U uptake mechanisms have been recently characterized in *Saccharomyces cerevisiae* (Revel et al, 2022) but are still unknown in green microalgae. Deciphering the composition and dynamics of the cell wall during U stress, together with U

uptake mechanisms, will deserve special attention to understand the unique ability of *Coelastrella* sp. PCV to capture U.

Besides its ability to tolerate a high concentration of U in a synthetic culture medium, *Coelastrella* sp. PCV was able to grow and maintain high photosynthetic activity in natural waters from a wetland near a reclaimed U mine (**Figure 6**). The amount of U found in waters from Rophin is similar to that found in natural waters from U mines in Germany (Bernhard et al, 1998), but lower than that from a U mine tailings pond where a U-tolerant *Chlamydomonas* species was isolated (Baselga-Cervera et al, 2018). The concentration of essential elements in waters from Rophin is two to three orders of magnitude lower than in artificial algal culture media and the concentration of TOC ranges from 4 to 19 mg.mL<sup>-1</sup> (**Table S1**). Thus, the speciation, bioavailability and toxicity of U are different in natural or synthetic media used in this study (Markich, 2002). In a single one-week growth cycle, *Coelastrella* was able to capture 25 to 55% of U from natural contaminated waters (**Figure 7**). The efficiency of U remediation was not related to the initial U level but was inversely correlated with the concentration of TOC (**Table S1**), as suggested by the decreased bioavailability of U with increasing TOC level (Hogan et al, 2005, Trenfield et al, 2011; Trenfield et al, 2012). The efficiency of U removal could also be related to differences in the elemental composition of natural waters (**Table S1**). Compared to H2 and E1, H1 water is enriched in several microelements (e.g. manganese, 130 times, iron >10 times), in magnesium (2-3 times) and in calcium (>2 times). These elements can compete with U for adsorption to cell walls or uptake (Charles et al., 2002) and thus, together with TOC, contribute to less efficient sequestration of the radionuclide in H1 than in E1 and H2. The fact that *Coelastrella* sp. PCV was able to remediate waters containing nanomolar amount of U is another illustration of the very high intrinsic capacity of the alga to capture U, even when metal speciation is complex (Markich, 2002). In addition to U, *Coelastrella* sp. PCV was able to remediate part of lead contaminating waters from Rophin (**Figure S4**). This suggests that *Coelastrella* sp. PCV has the potential to successfully address remediation of different contaminating metals, as previously described for the *Coelastrella* sp. (3-4) and *Coelastrella* sp. BGV strains (Plohn et al, 2021; Karcheva et al, 2022).

Due to nutritional deficiency and/or U toxicity (**Figure S5**), *Coelastrella* sp. PCV accumulated lipid droplets when grown in natural waters from Rophin (**Figure 8**). *Coelastrella* sp. of diverse origins have been shown to accumulate large amounts of neutral lipids (17-22% of total biomass) when cultivated in stress conditions (nutrient deprivation, high salinity, light or temperature) (Nayana et al, 2022). This response is shared with many photosynthetic microalgae, which can adjust lipid metabolism to stress conditions to produce substantial amounts of neutral lipids, mainly in the form of triacylglycerols (Zienkiewicz et al, 2016). As an example, several genes involved in lipid biosynthesis pathways are up-regulated under osmotic stress conditions in *Coelastrella* sp. M-60 (Karpagam et al, 2018). Several *Coelastrella* species have also been reported to produce carotenoids in stress conditions, including *C. astaxantica*, *C. rubescens* or *C. aeroterrestica* (Nayana et al, 2022). *Coelastrella* sp. PCV has been cultivated in different stress conditions, including nutrient deprivation or metal stress, but none of these treatments resulted in carotenoid accumulation. This suggests that metabolic changes induced by nutrient or metal stress are not similar in all *Coelastrella* species (Karpagam et al, 2018).

## 5. Conclusions

In this study, we characterized a resilient microalgal species demonstrating both tolerance to high U concentration in artificial culture media and the ability to survive in nutrient-depleted natural waters containing substantial amounts of toxic elements (including U, lead or aluminum). As it is much more tolerant to U than *C. reinhardtii* or *C. vulgaris*, *Coelastrella* sp. PCV is a very promising model for identifying mechanisms of response and adaptation to U stress, in particular the processes of desorption and efflux, which are likely to protect algal cells from the toxic effects of U during a prolonged period of stress. To support this assumption, the characterization of (hyper)tolerant microbial species has allowed the identification of novel mechanisms of U tolerance, including biomineralization processes or a high-affinity U-binding protein (e.g. Choudhary and Sar, 2011, Pinel-Cabello et al, 2021, Gallois et al, 2022). *Coelastrella* sp. PCV also demonstrated very interesting properties for treating natural or



wastewater with low to high U contamination. On the one hand, this fast growing and high biomass-producing microalga could be used to reclaim heavily contaminated environments, taking advantage of its remarkable ability to capture U. On the other hand, *Coelastrella* sp. PCV could be grown in natural soft water contaminated with moderate levels of toxic metals, allowing both habitat remediation and valorization of the lipid accumulating algal biomass.

### **Declaration of Competing Interest**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

### **CRedit authorship contribution statement**

Camille Beaulier: conceptualization, investigation, writing – original draft, review and editing. Marie Dannay: conceptualization, investigation. Fabienne Devime: investigation. Adrien Galeone: investigation. Célia Baggio: investigation. Nabila El Sakkout: investigation. Camille Raillon: investigation. Olivier Courson: investigation. Jacques Bourguignon: funding acquisition, investigation, writing - review and editing. Claude Alban: conceptualization, investigation, writing - review and editing. Stéphane Ravanel: supervision, funding acquisition, conceptualization, investigation, writing - original draft, review and editing.

### **Acknowledgments**

This work has received funding from the University Grenoble Alpes (PhD fellowship to Camille Beaulier), the Plant Biology and Breeding division from INRAE (DemoniaCo project), the “Nucléaire, énergie, environnement, déchets, société - NEEDS” program from the CNRS (INSPECT project), the “Diversity of biological mechanisms” program from the Institute for Biological Sciences at CNRS (SilverCoela project), and the Agence Nationale de la Recherche (ANR-21-CE34-0004, DemoniaCo project; ANR-17-EURE-0003, CBH-EUR-GS). We gratefully acknowledge Dr. Virginie Chapon and Dr. Jonathan Przybyla-Toscano for helpful discussions.

## Legend to figures

**Figure 1: Phylogenetic and morphological analysis of *Coelastrella* sp. PCV.** A- Phylogenetic analysis inferred from 18S rDNA sequences. The 18S rDNA sequence from *Coelastrella* sp. PCV was obtained by PCR and used for a NCBI BLASTn search against the Chlorophyceae sequence dataset. Thirty-one 18S rDNA sequences from the top ranking list were retrieved and aligned using Muscle (MEGA X software, default settings). The phylogenetic analysis was done by using the Maximum Likelihood method and Kimura 2-parameter mode. The consensus bootstrap tree is shown and branch support values (in % for 1000 replicates) are indicated. The GenBank accession numbers are provided in the 'Materials and methods', section 2.3. B- Morphological analysis of *Coelastrella* sp. PCV at different stages of growth. Bright field (left) and chlorophyll fluorescence (right) images of the five cell samples were captured using a Zeiss Axioplan 2 microscope and post-processed using ImageJ/Fiji. Scale bar = 5  $\mu$ m.

**Figure 2: Effect of uranium on the growth and photosynthesis of *Coelastrella* sp. PCV.** A- Growth curves were established for cells growing in TAP (○) or TAP LoP (50  $\mu$ M of phosphate) supplemented with uranyl nitrate (0 ●, 50 ▲, 100 ▼, 200 ◆, and 400  $\mu$ M ■). Cells were grown at 21°C in continuous white light (40  $\mu$ E) and counted daily. B,C- Photosynthetic efficiency of *Coelastrella* cells challenged with U. Fv/Fm in dark-adapted cells (B) and electron transfer rates (ETR) under increasing light irradiance were determined after 2 days of U stress. Data from n=5 to 8 independent experiments (with n=2-3 replicates) analyzed over a 3-year period have been gathered to produce growth and Fv/Fm kinetics (mean  $\pm$  SD). Light response curves are representative of n=2 independent experiments with n=3 replicates (mean  $\pm$  SD).

**Figure 3: Comparison of uranium tolerance in *Coelastrella* sp. PCV, *Chlamydomonas reinhardtii* and *Chlorella vulgaris*.** A- Growth curves were established for *Coelastrella* sp.

PCV (●), *Chlamydomonas reinhardtii* (◆) and *Chlorella vulgaris* (▼) grown in TAP LoP containing 0-200  $\mu\text{M}$  of uranyl nitrate. Cells were grown at 21°C in continuous white light (40  $\mu\text{E}$ ) and cell density after 2 or 3 days of U stress was used to calculate relative growth with reference to TAP LoP (U0). B- Fv/Fm was determined in dark-adapted cells after 2 or 3 days of U stress. Data from n=3 to 5 independent experiments (and n=2-3 replicates) are displayed (mean  $\pm$  SD).

**Figure 4: Time course of uranium distribution between the medium and algal cells during growth under stress conditions.** *Coelastrella* sp. PCV was grown at 21°C in continuous white light (40  $\mu\text{E}$ ) in TAP LoP containing 50 (A), 100 (B), 200 (C) and 400 (D)  $\mu\text{M}$  of uranyl nitrate (see Figure 2). Aliquots of the medium (■) and cells (●) were withdrawn every day and U was determined by ICP-MS. Before U quantification, cells were washed with sodium carbonate to remove loosely-bound metals. The amount of U (in  $\mu\text{g}$ ) is indicated for a standard 20-mL culture. Data from n=5 independent experiments (and n=2-3 replicates) are shown (mean  $\pm$  SD).

**Figure 5: Biosorption of uranium by *Coelastrella* sp. PCV.** Time course of U distribution between the medium (A) and algal cells (B) during short-term incubation periods. *Coelastrella* sp. PCV was challenged with 50 (▲), 100 (▼), 200 (◆) and 400 (■)  $\mu\text{M}$  uranyl nitrate in TAP LoP at 21°C in continuous white light (40  $\mu\text{E}$ ). Uranium was determined by ICP-MS in the medium and whole cells (not washed with sodium carbonate) at different time intervals. C- Quantification of loosely and tightly-bound U to *Coelastrella* cells. Cells were washed with sodium carbonate to remove loosely-bound U. Uranium was determined in both the washed cells and the carbonate wash solution. Data show the distribution of U at time point 24 hours. In panels A,B,C, the amount of U (in  $\mu\text{g}$ ) is indicated for a standard 20-mL culture. D- Titration of U biosorption capacity of *Coelastrella* cells. The maximal amount of metal bound to whole

cells (■) or carbonate-washed cells (●) is derived from data in Fig 5BC and is expressed in mg U per gram dry weight. Data from n=5 independent experiments (and n=1-3 replicates) are shown (mean ± SD).

**Figure 6: Growth and photosynthesis of *Coelastrella* sp. PCV in natural metal-polluted**

**waters.** A- Growth curves were established for cells growing in HSM (●) and natural waters from the Rophin area (E1 ▼, H1 ■, H2 ▲). Cells were grown at 21°C in continuous white light (40 μE) and counted daily. Picture in the insert shows cultures on day 7. Data from n=5 independent experiments with n=2 replicates are shown (mean ± SD). B,C- Photosynthetic efficiency of *Coelastrella* cells grown in natural waters. Fv/Fm was determined in dark-adapted cells. Data from n=5 independent experiments with n=2 replicates are shown (mean ± SD). The ETR was determined under increasing light irradiance after 3 days of growth. Data from n=2 independent experiments with n=2 replicates are shown (mean ± SD).

**Figure 7: Bioremediation of uranium from natural polluted waters by *Coelastrella* sp.**

**PCV.** A,B,C- Time courses of U distribution between the medium (■) and algal cells (●) during growth of *Coelastrella* in waters from the locations E1 (A), H2 (B), and H1 (C) of the Rophin area. Growth was conducted at 21°C in continuous white light (40 μE). Uranium was determined by ICP-MS in the medium and whole cells (not washed with sodium carbonate) at regular time intervals. The amount of U (in ng) is indicated for a standard 20-mL culture. D- Distribution of U remaining in waters and loosely or tightly-bound to *Coelastrella* cells. Cells were washed with sodium carbonate to remove loosely-bound U. Data show the distribution of U after 7 days of growth in contaminated waters. Data from n=3 to 6 independent experiments and n=2 replicates are shown (mean ± SD).

**Figure 8: Lipid droplet accumulation in *Coelastrella* sp. PCV grown in natural polluted waters.** *Coelastrella* sp. PCV was cultivated in waters from Rophin for 3 days at 21°C in continuous white light (40  $\mu$ E) and then stained with Nile Red. Two representative cells from each condition were imaged using an inverted confocal microscope (LSM880 Zeiss). Chlorophyll autofluorescence is in red; lipid droplets stained with Nile Red are in yellow. Scale bar = 5  $\mu$ m. The images have been selected to highlight lipid droplets; they are not representative of cell size because the z-depth does not necessarily correspond to the median plane of cells. Additional experiments showed that the average diameter of *Coelastrella* cells grown in HSM and H1 water was  $7.3 \pm 2.0$  and  $6.0 \pm 0.8$   $\mu$ m, respectively (mean  $\pm$  SD, n=200 cells per condition).

## References

- Agusti S, Kalff J (1989) The influence of growth conditions on the size dependence of maximal algal density and biomass. *Limnology and Oceanography* 34: 1104-1108
- Anke M, Seeber O, Mueller R, Schaefer U, Zerull J (2009) Uranium transfer in the food chain from soil to plants, animals and man. *Chemie Der Erde-Geochemistry* 69: 75-90
- Baselga-Cervera B, Garcia-Balboa C, Diaz-Alejo HM, Costas E, Lopez-Rodas V (2020) Rapid Colonization of Uranium Mining-Impacted Waters, the Biodiversity of Successful Lineages of Phytoplankton Extremophiles. *Microb Ecol* 79: 576-587
- Baselga-Cervera B, Romero-Lopez J, Garcia-Balboa C, Costas E, Lopez-Rodas V (2018) Improvement of the Uranium Sequestration Ability of a *Chlamydomonas* sp. (ChISP Strain) Isolated From Extreme Uranium Mine Tailings Through Selection for Potential Bioremediation Application. *Front Microbiol* 9: 523
- Bernhard G, Geipel G, Brendler V, Nitsche H (1998) Uranium speciation in waters of different uranium mining areas. *Journal of Alloys and Compounds* 271-273: 201-205
- Berthet S, Villiers F, Alban C, Serre NBC, Martin-Laffon J, Figuet S, Boisson AM, Bligny R, Kuntz M, Finazzi G, Ravanel S, Bourguignon J (2018) *Arabidopsis thaliana* plants challenged with uranium reveal new insights into iron and phosphate homeostasis. *New Phytol* 217: 657-670
- Bhat SA, Bashir O, Ul Haq SA, Amin T, Rafiq A, Ali M, Américo-Pinheiro JHP, Sher F (2022) Phytoremediation of heavy metals in soil and water: An eco-friendly, sustainable and multidisciplinary approach. *Chemosphere* 303: 134788
- Charles AL, Markich SJ, Stauber JL, De Filippis LF (2002) The effect of water hardness on the toxicity of uranium to a tropical freshwater alga (*Chlorella* sp.). *Aquatic Toxicology* 60: 61-73
- Chen L, Liu JR, Zhang WX, Zhou JQ, Luo DQ, Li ZM (2021) Uranium (U) source, speciation, uptake, toxicity and bioremediation strategies in soil-plant system: A review. *Journal of Hazardous Materials* 413

- Cheng Y, Zhang T, Chen S, Li F, Qing R, Lan T, Yang Y, Liao J, Liu N (2023) Unusual uranium biomineralization induced by green algae: Behavior investigation and mechanism probe. *Journal of Environmental Sciences* 124: 915-922
- Chodat R (1922) Matériaux pour l'histoire des algues de la Suisse. *In* Bulletin de la Société botanique de Genève, Vol ser. 2, v. 13-14 (1921-22). Société botanique de Genève, Genève, [Switzerland], pp 66-114
- Choudhary S, Sar P (2011) Uranium biomineralization by a metal resistant *Pseudomonas aeruginosa* strain isolated from contaminated mine waste. *Journal of Hazardous Materials* 186: 336-343
- Ferro L, Gentili FG, Funk C (2018) Isolation and characterization of microalgal strains for biomass production and wastewater reclamation in Northern Sweden. *Algal Research-Biomass Biofuels and Bioproducts* 32: 44-53
- Fortin C, Denison FH, Garnier-Laplace J (2007) Metal-phytoplankton interactions: Modeling the effect of competing ions ( $H^+$ ,  $Ca^{2+}$ , and  $Mg^{2+}$ ) on uranium uptake. *Environ Toxicol Chem* 26: 242-248
- Fortin C, Dutel L, Garnier-Laplace J (2004) Uranium complexation and uptake by a green alga in relation to chemical speciation: The importance of the free uranyl ion. *Environmental Toxicology and Chemistry* 23: 974-981
- Franklin NM, Stauber JL, Markich SJ, Lim RP (2000) pH-dependent toxicity of copper and uranium to a tropical freshwater alga (*Chlorella* sp.). *Aquatic Toxicology* 48: 275-289
- Gallois N, Alpha-Bazin B, Bremond N, Ortet P, Barakat M, Piette L, Mohamad Ali A, Lemaire D, Legrand P, Theodorakopoulos N, Floriani M, Février L, Den Auwer C, Arnoux P, Berthomieu C, Armengaud J, Chapon V (2022) Discovery and characterization of UipA, a uranium- and iron-binding PepSY protein involved in uranium tolerance by soil bacteria. *ISME Journal* 16: 705-716
- Gandhi TP, Sampath PV, Maliyekkal SM (2022) A critical review of uranium contamination in groundwater: Treatment and sludge disposal. *Sci Total Environ* 825: 153947

- Gao N, Huang ZH, Liu HQ, Hou J, Liu XH (2019) Advances on the toxicity of uranium to different organisms. *Chemosphere* 237
- Garcia-Balboa C, Baselga-Cervera B, Garcia-Sanchez A, Igual JM, Lopez-Rodas V, Costas E (2013) Rapid adaptation of microalgae to bodies of water with extreme pollution from uranium mining: An explanation of how mesophilic organisms can rapidly colonise extremely toxic environments. *Aquatic Toxicology* 144: 116-123
- Guiry MD (2021) AlgaeBase: a global database for algae. *Current Science* 121: 10-11
- Gunther A, Raff J, Geipel G, Bernhard G (2008) Spectroscopic investigations of U(VI) species sorbed by the green algae *Chlorella vulgaris*. *Biometals* 21: 333-341
- Häder D-P, Banaszak AT, Villafañe VE, Narvarte MA, González RA, Helbling EW (2020) Anthropogenic pollution of aquatic ecosystems: Emerging problems with global implications. *Sci Total Environ* 713: 136586
- Harris EH (2009) *The Chlamydomonas Sourcebook (Second Edition)*. Academic Press, London
- Herlory O, Bonzom JM, Gilbin R (2013) Sensitivity evaluation of the green alga *Chlamydomonas reinhardtii* to uranium by pulse amplitude modulated (PAM) fluorometry. *Aquatic Toxicology* 140: 288-294
- Hogan AC, van Dam RA, Markich SJ, Camilleri C (2005) Chronic toxicity of uranium to a tropical green alga (*Chlorella* sp.) in natural waters and the influence of dissolved organic carbon. *Aquatic Toxicology* 75: 343-353
- Horikoshi T, Nakajima A, Sakaguchi T (1979) Studies on the Accumulation of Heavy-Metal Elements in Biological-Systems .4. Uptake of Uranium by *Chlorella-Regularis*. *Agricultural and Biological Chemistry* 43: 617-623
- Jaussaud A, Lupette J, Salvaing J, Jouhet J, Bastien O, Gromova M, Marechal E (2020) Stepwise Biogenesis of Subpopulations of Lipid Droplets in Nitrogen Starved *Phaeodactylum tricornutum* Cells. *Frontiers in Plant Science* 11



- John WA, Luckel B, Matschiavelli N, Hubner R, Matschi S, Hoehenwarter W, Sachs S (2022) Endocytosis is a significant contributor to uranium(VI) uptake in tobacco (*Nicotiana tabacum*) BY-2 cells in phosphate-deficient culture. *Sci Total Environ* 823: 153700
- Kan Y, Pan J (2010) A one-shot solution to bacterial and fungal contamination in the green alga *Chlamydomonas reinhardtii* culture by using an antibiotic cocktail. *Journal of Phycology* 46: 1356-1358
- Karcheva Z, Georgieva Z, Tomov A, Petrova D, Zhiponova M, Vasileva I, Chaneva G (2022) Heavy metal stress response of microalgal strains *Arthrocnemum africanum* and *Coelastrella* sp. *BGV. BioRisk* 17: 83-94
- Karpagam R, Jawaharraj K, Ashokkumar B, Sridhar J, Varalakshmi P (2018) Unraveling the lipid and pigment biosynthesis in *Coelastrella* sp M-60: Genomics-enabled transcript profiling. *Algal Research-Biomass Biofuels and Bioproducts* 29: 277-289
- Kawasaki S, Yoshida R, Ohkoshi K, Toyoshima H (2020) *Coelastrella astaxanthina* sp. nov. (Sphaeropleales, Chlorophyceae), a novel microalga isolated from an asphalt surface in midsummer in Japan. *Phycological Research* 68: 107-114
- Khaw YS, Khong NMH, Shaharuddin NA, Yusoff FM (2020) A simple 18S rDNA approach for the identification of cultured eukaryotic microalgae with an emphasis on primers. *Journal of Microbiological Methods* 172: 105890
- Kolhe N, Zinjarde S, Acharya C (2018) Responses exhibited by various microbial groups relevant to uranium exposure. *Biotechnol Adv* 36: 1828-1846
- Kumar S, Stecher G, Li M, Knyaz C, Tamura K (2018) MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. *Mol Biol Evol*
- Lavoie M, Sabatier S, Garnier-Laplace J, Fortin C (2014) Uranium accumulation and toxicity in the green alga *Chlamydomonas reinhardtii* is modulated by pH. *Environmental Toxicology and Chemistry* 33: 1372-1379
- Markich SJ (2002) Uranium speciation and bioavailability in aquatic systems: an overview. *ScientificWorldJournal* 2: 707-729


- Martin A, Hassan-Loni Y, Fichtner A, Peron O, David K, Chardon P, Larrue S, Gourgiotis A, Sachs S, Arnold T, Grambow B, Stumpf T, Montavon G (2020) An integrated approach combining soil profile, records and tree ring analysis to identify the origin of environmental contamination in a former uranium mine (Rophin, France). *Sci Total Environ* 747
- Maxwell K, Johnson GN (2000) Chlorophyll fluorescence - a practical guide. *Journal of Experimental Botany* 51: 659-668
- Narayanan GS, Kumar G, Seepana S, Elankovan R, Arumugan S, Premalatha M (2018) Isolation, identification and outdoor cultivation of thermophilic freshwater microalgae *Coelastrella* sp. Fl69 in bubble column reactor for the application of biofuel production. *Biocatalysis and Agricultural Biotechnology* 14: 357-365
- Nayana K, Sudhakar MP, Arunkumar K (2022) Biorefinery potential of *Coelastrella* biomass for fuel and bioproducts-a review. *Biomass Conversion and Biorefinery*
- Pinel-Cabello M, Jroundi F, López-Fernández M, Geffers R, Jarek M, Jauregui R, Link A, Vilchez-Vargas R, Merroun ML (2021) Multisystem combined uranium resistance mechanisms and bioremediation potential of *Stenotrophomonas bentonitica* BII-R7: Transcriptomics and microscopic study. *Journal of Hazardous Materials* 403: 123858
- Plohn M, Escudero-Onate C, Funk C (2021b) Biosorption of Cd(II) by Nordic microalgae: Tolerance, kinetics and equilibrium studies. *Algal Research-Biomass Biofuels and Bioproducts* 59
- Plohn M, Spain O, Sirin S, Silva M, Escudero-Onate C, Ferrando-Climent L, Allahverdiyeva Y, Funk C (2021a) Wastewater treatment by microalgae. *Physiologia Plantarum* 173: 568-578
- Pradines C, Wiktor V, Camilleri V, Gilbin R (2005) Development of biochemical methods to estimate the subcellular impact of uranium exposure on *Chlamydomonas reinhardtii*. *Radioprotection* 40: S163-S168
- Revel B, Catty P, Ravanel S, Bourguignon J, Alban C (2022) High-affinity iron and calcium transport pathways are involved in U(VI) uptake in the budding yeast *Saccharomyces cerevisiae*. *J Hazard Mater* 422: 126894

- Sarthou MCM, Devime F, Baggio C, Figuet S, Alban C, Bourguignon J, Ravanel S (2022) Calcium-permeable cation channels are involved in uranium uptake in *Arabidopsis thaliana*. *Journal of Hazardous Materials* 424: 127436
- Saxena P, Harish (2019) Toxicity assessment of ZnO nanoparticles to freshwater microalgae *Coelastrella terrestris*. *Environ Sci Pollut Res Int* 26: 26991-27001
- Saxena P, Sangela V, Harish (2020) Toxicity evaluation of iron oxide nanoparticles and accumulation by microalgae *Coelastrella terrestris*. *Environ Sci Pollut Res Int* 27: 19650-19660
- Schmidt-Kloiber A, Bremerich V, De Wever A, Jähnig SC, Martens K, Strackbein J, Tockner K, Hering D (2019) The Freshwater Information Platform: a global online network providing data, tools and resources for science and policy support. *Hydrobiologia* 838: 1-11
- Smith-Baedorf HD, Chuck CJ, Mokebo KR, MacDonald H, Davidson MG, Scott RJ (2013) Bioprospecting the thermal waters of the Roman baths: isolation of oleaginous species and analysis of the FAME profile for biodiesel production. *AMB Express* 3: 1-14
- Spain O, Funk C (2022) Detailed Characterization of the Cell Wall Structure and Composition of Nordic Green Microalgae. *Journal of Agricultural and Food Chemistry* 70: 9711-9721
- Spain O, Plohn M, Funk C (2021) The cell wall of green microalgae and its role in heavy metal removal. *Physiologia Plantarum* 173: 526-535
- Sun XM, Ren LJ, Zhao QY, Ji XJ, Huang H (2018) Microalgae for the production of lipid and carotenoids: a review with focus on stress regulation and adaptation. *Biotechnol Biofuels* 11: 272
- Trenfield MA, Ng JC, Noller B, Markich SJ, van Dam RA (2012) Dissolved organic carbon reduces uranium toxicity to the unicellular eukaryote *Euglena gracilis*. *Ecotoxicology* 21: 1013-1023
- Trenfield MA, Ng JC, Noller BN, Markich SJ, van Dam RA (2011) Dissolved Organic Carbon Reduces Uranium Bioavailability and Toxicity. 2. Uranium[VI] Speciation and Toxicity to Three Tropical Freshwater Organisms. *Environmental Science & Technology* 45: 3082-3089


- Vallet A, Martin-Laffon J, Favier A, Revel B, Bonnot T, Vidaud C, Armengaud J, Gaillard JC, Delangle P, Devime F, Figuet S, Serre NBC, Erba EB, Brutscher B, Ravanel S, Bourguignon J, Alban C (2023) The plasma membrane-associated cation-binding protein PCaP1 of *Arabidopsis thaliana* is a uranyl-binding protein. *J Hazard Mater* 446: 130668
- Vandenhove H (2002) European sites contaminated by residues from the ore-extracting and -processing industries. *International Congress Series* 1225: 307-315
- Vogel M, Gunther A, Rossberg A, Li B, Bernhard G, Raff J (2010) Biosorption of U(VI) by the green algae *Chlorella vulgaris* in dependence of pH value and cell activity. *Sci Total Environ* 409: 384-395
- Wang QH, Song HY, Liu XD, Zhu H, Hu ZY, Liu GX (2019) Deep genomic analysis of *Coelastrella saipanensis* (Scenedesmaceae, Chlorophyta): comparative chloroplast genomics of Scenedesmaceae. *European Journal of Phycology* 54: 52-65
- Wu WY, Ke T, Zhou XJ, Li QQ, Tao Y, Zhang YR, Zeng YY, Cao J, Chen LZ (2022) Synergistic remediation of copper mine tailing sand by microalgae and fungi. *Applied Soil Ecology* 175
- Zhang JG, Luo XG (2022) Bioaccumulation characteristics and acute toxicity of uranium in *Hydrodictyon reticulatum*: An algae with potential for wastewater remediation. *Chemosphere* 289
- Zhang X, Luo S, Yang Q, Zhang H, Li J (1997) Accumulation of uranium at low concentration by the green alga *Scenedesmus obliquus* 34. *Journal of Applied Phycology* 9: 65-71
- Zienkiewicz K et al., Stress-induced neutral lipid biosynthesis in microalgae — Molecular, cellular and physiological insights. *Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids*, 2016. 1861(9, Part B): p 1269-1281.

**FIGURE 1**


**A**




**B**




**FIGURE 2**



**FIGURE 3**




**FIGURE 4**







**FIGURE 5**




**FIGURE 6**



**FIGURE 7**



**FIGURE 8**



## **Characterization of a uranium-tolerant green microalga of the genus *Coelastrella* with high potential for the remediation of metal-polluted waters**

Camille Beaulier<sup>a</sup>, Marie Dannay<sup>a</sup>, Fabienne Devime<sup>a</sup>, Adrien Galeone<sup>a</sup>, Célia Baggio<sup>a</sup>, Nabila El Sakkout<sup>a</sup>, Camille Raillon<sup>a</sup>, Olivier Courson<sup>b</sup>, Jacques Bourguignon<sup>a</sup>, Claude Alban<sup>a</sup>, Stéphane Ravanel<sup>a\*</sup>


### **Supporting information**

**Figure S1:** Isolation of *Coelastrella* sp. PCV from liquid wastes contaminated by uranium.


**Figure S2:** Inhibition of growth and photosynthesis by uranium in *Coelastrella* sp. PCV.

**Figure S3:** Lipid droplet accumulation in *Coelastrella* sp. PCV challenged with uranium.

**Table S1:** Physico-chemical properties of natural waters from the Rophin area.




**Figure S1: Isolation of *Coelastrella* sp. PCV from liquid wastes contaminated by uranium.** The algal bloom developed in laboratory wastes from plant culture medium contaminated with uranyl nitrate (about 5  $\mu$ M). The original sampling streaked on TAP-agar was contaminated with bacteria and fungi. Single algal colonies (1 and 2) could be cleared of contaminants by two successive rounds of selection on TAP-agar supplemented with antibiotics (ampicillin and cefotaxime) and a broad-spectrum fungicide (carbendazim). Plates were incubated at 21°C in continuous light (20  $\mu$ E) for 7 days.



**Figure S2: Inhibition of growth and photosynthesis by uranium in *Coelastrella sp. PCV*.**

A- Effect of U on growth. Doubling times were determined at day 2 from growth curves displayed in Figure 2, plus additional experiments with a similar design (for 25 and 75  $\mu\text{M}$  uranyl nitrate). B,C- Effect of U on photosynthesis. Fv/Fm in dark-adapted cells (B) and electron transfer rates at a light irradiance of 336  $\mu\text{E}$  (C) were measured at day 2 (see Figure 2). Dose-response curves have been drawn from  $n=5$  to 8 independent experiments (mean  $\pm$  SD) and fitted using sigmoidal equations, indicating IC50 values  $>80$   $\mu\text{M}$  for doubling time,  $>500$   $\mu\text{M}$  for Fv/Fm, and  $>400$   $\mu\text{M}$  for ETR.



**Figure S3: Lipid droplet accumulation in *Coelastrella* sp. PCV challenged with uranium.** *Coelastrella* sp. PCV was cultivated for 1 day at 21°C in continuous white light (40  $\mu$ E) in TAP LoP containing 50 to 400  $\mu$ M of uranyl nitrate. Cells were stained with Nile Red and imaged using a Zeiss Axioplan 2 microscope. Chlorophyll autofluorescence is in red; lipid droplets stained with Nile Red are in yellow. Scale bar = 5  $\mu$ m.



**Table S1: Physico-chemical properties of natural waters from the Rophin area.**

	E1	H1	H2	Unit
<b>pH</b>	7.0-7.2	6.9-7.1	7.0-7.3	
<b>Na</b>	223 ± 46	214 ± 27	214 ± 24	
<b>Mg</b>	31 ± 0.8	91 ± 7.8	47 ± 2.9	
<b>Al</b>	1.1 ± 0.4	2.2 ± 0.4	1.3 ± 0.4	
<b>P</b>	0.73 ± 0.1	1.23 ± 0.5	0.80 ± 0.2	x10 <sup>-6</sup> M
<b>K</b>	24 ± 1.2	37 ± 1.9	36 ± 2.1	
<b>Ca</b>	66 ± 12	151 ± 41	97 ± 24	
<b>Fe</b>	4.4 ± 0.8	53.9 ± 5.6	4.5 ± 1.0	
<b>Li</b>	79 ± 22	106 ± 30	95 ± 32	
<b>B</b>	206 ± 79	261 ± 102	510 ± 159	
<b>Cr</b>	2.0 ± 0.3	11.8 ± 2.8	3.3 ± 0.7	
<b>Mn</b>	70 ± 29	9260 ± 2909	69 ± 25	
<b>Co</b>	0.8 ± 0.3	38.0 ± 17.8	1.0 ± 0.2	
<b>Cu</b>	32 ± 8	327 ± 74	135 ± 22	
<b>Zn</b>	89 ± 26	305 ± 66	90 ± 13	x10 <sup>-9</sup> M
<b>Ga</b>	53 ± 6	199 ± 28	81 ± 8	
<b>Mo</b>	1.5 ± 0.4	10.3 ± 2.8	1.6 ± 0.3	
<b>Ag</b>	0.12 ± 0.06	0.21 ± 0.14	0.36 ± 0.11	
<b>Cd</b>	0.20 ± 0.03	0.60 ± 0.14	0.40 ± 0.06	
<b>Pb</b>	3.8 ± 0.5	277 ± 71	21 ± 8	
<b>U</b>	15 ± 1	228 ± 40	62 ± 12	
<b>Total carbon (TC)</b>	8.2 ± 0.5	24.5 ± 1.2	8.9 ± 0.4	
<b>Inorganic carbon (IC)</b>	4.0 ± 0.2	5.8 ± 2.9	4.1 ± 1.1	mg C.l <sup>-1</sup>
<b>Total organic carbon (TC – IC)</b>	4.2 ± 0.6	18.7 ± 3.9	4.8 ± 1.1	
<b>Total organic carbon (NPOC)</b>	3.8 ± 0.3	14.8 ± 2.8	4.4 ± 0.7	

**Physico-chemical properties of natural waters from the Rophin area.** Water samples from the Rophin area were collected at the E1, H1 and H2 locations and filtered using fritted glass and then 0.2 µm nylon filters before analysis. Elemental composition was determined by ICPMS. Data are mean ± SD of n=3-5 measurements. Total carbon (TC), dissolved inorganic carbon (DIC), and non-purgeable organic carbon (NPOC) were determined by the high temperature catalytic oxidation methods and non-dispersive infrared detection. Data are mean ± SD of n=3-6 measurements with n=2 technical replicates.