



The effect of contaminants on taxonomic composition of micromeiofauna present in biofilms

Jacky Vedrenne, Gardebled Eugénie, Antoine Tourret, Maelle Baillergeau,
Nicolas Creusot, Chloé Bonnineau

► To cite this version:

Jacky Vedrenne, Gardebled Eugénie, Antoine Tourret, Maelle Baillergeau, Nicolas Creusot, et al.. The effect of contaminants on taxonomic composition of micromeiofauna present in biofilms. 49th Annual Canadian Ecotoxicity Workshop, Oct 2023, Ottawa, Canada. 2023. hal-04403941

HAL Id: hal-04403941

<https://hal.inrae.fr/hal-04403941>

Submitted on 18 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The effect of contaminants on taxonomic composition of micromeiofauna present in biofilms

Eugénie Gardebled^{1,2}, Maelle Baillergeau¹, Antoine Tourret^{1,2}, Chloé Bonnneau¹, Nicolas Creusot¹, Jacky Vedrenne¹

¹ Institut national de recherche pour l'agriculture, l'alimentation et l'environnement, 50 Avenue de Verdun, 33612 Cestas, France

² Institut national de la recherche scientifique, Centre Eau Terre Environnement, 490 rue de la Couronne, Québec, QC, G1K 9A9

1 Introduction

Micro-metazoans play an important role in freshwater, estuarine and marine ecosystems. Their complex interactions with their environments, including other organisms, create an intricate web of relationships that collectively affect ecosystem processes.


Considering that micro-metazoans are widespread, easy to sample, and highly responsive to environmental changes, they may serve as bio-indicators of pollution or disturbances, yet their full potential in this regard remains largely untapped.

In this preliminary work, we investigated the effects of a mixture of contaminants at environmental concentrations on the biodiversity of biofilm micro-metazoans: the optical brightener (**DAST**); the insecticide fenoxy carb (**FNX**); the herbicide glyphosate; and its main metabolite AMPA (**GA**).

2 Methods

CHANNEL EXPERIENCE

131-Day Maturation
For 28 days




GA : Glyphosate (1 $\mu\text{g.L}^{-1}$) + AMPA (100 $\mu\text{g.L}^{-1}$)

DAST_GA : DAST (30 $\mu\text{g.L}^{-1}$) + Gly (1 $\mu\text{g.L}^{-1}$) + AMPA (100 $\mu\text{g.L}^{-1}$)

FNX_GA : FNX (30 mg.L⁻¹) + Gly (1 $\mu\text{g.L}^{-1}$) + AMPA (100 $\mu\text{g.L}^{-1}$)

Sampling days during exposure : 1, 14 et 28 days

3 Results



4 Discussion

- * **Ciliates** are more resistant to the cocktail with FNX with DAST
- * **Rotifers** are resistant to GA and DAST_GA up to 14 and 28 days, respectively, whereas they are sensitive to FNX_GA.
- * **Zooflagellates** are sensitive to GA, DAST_GA and FNX_GA cocktails or to experimental conditions.

The results revealed that these contaminants modified the taxonomic composition of the micro-metazoans. The study highlights the sensitivity of micro-metazoans to contaminants in biofilms.


- * Rotifers feed on diatoms

Colurella uncinata is phytophagous (Obertegger and Wallace, 2023)

- * Ciliates feed on cyanobacteria

Aspidisca cicada feed on the order of 15 μm , corresponding to the diameter of the cyanobacteria observed (Neury-Ormanni et al., 2020)

5 Pictures



6 Acknowledgments