



Bioconversion of H₂ to Single Cell Protein by Purple Bacteria consortia: Influence of environmental conditions on microbial kinetics

María del Rosario Rodero, Jose Antonio Magdalena, Jean-Philippe Steyer, Nicolas Bernet, Gabriel Capson-Tojo

► To cite this version:

María del Rosario Rodero, Jose Antonio Magdalena, Jean-Philippe Steyer, Nicolas Bernet, Gabriel Capson-Tojo. Bioconversion of H₂ to Single Cell Protein by Purple Bacteria consortia: Influence of environmental conditions on microbial kinetics. 6th IWA International Conference on eco-Technologies for Wastewater Treatment, Jun 2023, Girona, Spain. hal-04466592

HAL Id: hal-04466592

<https://hal.inrae.fr/hal-04466592>

Submitted on 19 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Bioconversion of H₂ to Single Cell Protein by Purple Bacteria consortia: Influence of environmental conditions on microbial kinetics

M. del R. Rodero*, **, J.A. Magdalena*, ***, J. P. Steyer*, N. Bernet* and G. Capson-Tojo*

* INRAE, Univ Montpellier, LBE, 102 avenue des Étangs, 11100 Narbonne, France (E-mail: maria-del-rosario.rodero-raya@inrae.fr; jose-antonio.magdalena-cadelo@inrae.fr, jean-philippe.steyer@inrae.fr; nicolas.bernet@inrae.fr; gabriel.capson-tojo@inrae.fr)

** Institute of Sustainable Processes, University of Valladolid, 47011, Valladolid, Spain

*** Vicerrectorado de Investigación y Transferencia de la Universidad Complutense de Madrid, 28040 Madrid, Spain

Abstract

Single cell proteins (SCP) have emerged as an alternative protein source to partially alleviate the current problem of food scarcity. However, there is a need for the production of SCP from renewable and pathogen-free sources, such as gaseous streams. This study evaluated, for the first time, the use of enriched phototrophic purple bacteria (PPB) consortia for the production of SCP using H₂ and CO₂. The influence of pH (6-8.5), temperature (15-50 °C) and light intensity (0-50 W m⁻²) on the growth kinetics and biomass yields was investigated. Optimal conditions were found at a pH of 7, temperature of 25 °C and light intensities over 30 W·m⁻². High biomass yields (~ 1 g COD_{biomass}·g⁻¹ COD_{consumed}) and protein yields of ~4 g protein·g⁻¹ H₂ were achieved, regardless of the environmental conditions. The biomass had high protein contents (51–64 %w/w), potentially suitable as SCP. The biomass yields are amongst the highest ones reported from gaseous streams.

Keywords

Autotrophic growth; Environmental conditions; Hydrogen; Phototrophic Purple Bacteria; Single Cell Protein

INTRODUCTION

The sustainable production of food and feed is nowadays a serious global concern. Ever-growing population, climate change and limited natural resources call for the development of new sustainable food/feed alternatives (Zha et al., 2021). Proteins play a key role in human and animal diet as a source of nitrogen and essential amino acids (Ritala et al., 2017). The use of microorganisms as a protein-rich feedstock, the so-called single cell protein (SCP), is a promising alternative to plant or animal-based proteins, since microorganisms provide higher nitrogen recovery during protein synthesis (Puyol et al., 2017). The production of SCP has many environmental benefits compared to plant/animal proteins, such as lower land requirements and GHGs emissions, or a reduced water footprint (Alloul et al., 2022). Despite these advantages, SCP is currently produced from costly agricultural feedstocks or from unsustainable raw materials such as molasses, sucrose, starch, n-alkanes, methanol or natural gas.

Phototrophic purple bacteria (PPB) grown on pathogen-free sources are a potential SCP source, thanks to their high yields and their high contents on amino acids, pigments and vitamins. PPB exhibit a highly versatile metabolism, capable of performing anoxygenic photosynthesis, using solar light as energy source, and a wide range of electron/carbon donors such as organic compounds (photoheterotrophic growth), or H₂/H₂S with CO₂ as carbon source (litoautotrophic and photoautotrophic growth) (Capson-Tojo et al., 2020). H₂ from fermentation of organic waste, syngas, or generated via water electrolysis using surplus of electricity from renewable sources, and CO₂ from off-gases, represent promising electron and carbon sources for the sustainable and pathogen-free production of SCP. Spanoghe et al. (2021) proved, for the first time, the feasibility of the production of SCP from H₂ using pure PPB cultures. However, to the best of our knowledge, the potential of an enriched PPB community (without axenic conditions) growing photoautotrophically with H₂ as SCP source remains unexplored.

Here, the use of an enriched PPB consortia for the bioconversion of H_2 into SCP has been tested for the first time. The influence of environmental conditions (temperature, pH, light intensity) on microbial growth kinetics, protein contents, and microbial population, has been evaluated.

MATERIALS AND METHODS

An enriched PPB community grown photoheterotrophically from continuous photobioreactors used for wastewater treatment in Madrid (Spain) was used as a pre-inoculum. A series of batch enrichments (over 14) in Schott flasks of 500 mL were performed to obtain a PPB consortium able to achieve an efficient photoautotrophic growth. During each enrichment, aliquots of 20 mL from the previous culture were added to the Schott flasks, together with 180 mL of fresh medium. The medium was based on the mineral synthetic medium (MSM) proposed by Ormerod et al. (1961), modified to ensure photoautotrophic growth. The bottles were closed and the headspace was flushed with N_2 prior to H_2 . The initial headspace pressure was adjusted to 1.3 bar. The flasks were covered with UV/VIS filters (Capson-Tojo et al., 2021) and incubated under continuous illumination at $\sim 50 \text{ W}\cdot\text{m}^{-2}$ using infrared LED lights (850 nm). The flasks pressure was monitored daily. Once the pressure dropped below 1.0 bar, hydrogen was added. Each enrichment lasted for around one week.

Once the enrichment was stable and showed a constant performance, batch tests at different environmental conditions were tested. Assays at initial pH values of 6, 7 and 8.5, temperatures of 15, 25, 38 and 50 °C, and infrared light intensities of 0, 5, 15, 30 and 50 $\text{W}\cdot\text{m}^{-2}$ were carried out in triplicate, as described above for the enrichments (Figure 1). The gas composition and pressure were monitored 4-5 times per day. These data were used to calculate H_2 consumption rates. Liquid samples were also drawn at the beginning and at the end of each assay in order to determine biomass yields and productivities, crude protein contents, amino acids profiles and microbial communities.




Figure 1. Simplified experimental procedure followed for the evaluation of the influence of environmental conditions on the bioconversion of H_2 into SCP using an enriched PPB culture.

RESULTS

After 2-3 enrichments, H_2 consumption was observed, confirming the ability of the mixed PPB culture for growing photoautotrophically. Population analyses (16S r-RNA sequencing) of two enrichment samples after seven cycles confirmed that the consortium was dominated by PPB (> 80%), with *Rhodobacter* sp. and *Rhodopseudomonas* sp. as dominant genera. Consistent biomass yields of 0.9-1.0 g COD_{biomass}·g⁻¹ COD_{consumed} were obtained during the last six enrichments.

Environmental conditions had a noticeable impact on H_2 consumption, impacting biomass growth, kinetics, and protein production (Figure 2, Table 1). The highest H_2 consumption was observed at initial pH of 7, (with a final value of 8.4 due to CO_2 consumption). The most unfavourable initial pH was 8.5. The highest H_2 consumption was observed at 25°C, with 38°C decreasing the growth kinetics and 50°C inhibiting growth. Temperatures down to 15°C also reduced H_2 consumption. Light limitation was observed at intensities lower than 30 $W \cdot m^{-2}$ but similar results were obtained for both 30 and 50 $W \cdot m^{-2}$. Biomass yields close to 1 g $COD_{biomass} \cdot g^{-1} COD_{consumed}$ and protein yields over 4 g protein $\cdot g^{-1} H_2$ were achieved in almost all the conditions tested (Table 1). These proteins yields were higher than those obtained using pure cultures (Spanoghe et al., 2021). High protein contents ($> 50\%$ w/w) were achieved, in agreement with PPB grown photoheterotrophically (Hülßen et al., 2022a, 2022b). Similar C and N contents in the biomass (42 and 9 %w/w) were obtained (Table 1).




Figure 2. Time course of H_2 consumption by the enriched PPB culture at a) different pH values, b) different temperatures and c) different light intensities. Each data point shows the average and standard deviation ($n=3$).

Table 1. Main results obtained along with the corresponding standard deviation ($n=3$) during the batch tests.

	pH		Temperature (°C)		Intensity ($W \cdot m^{-2}$)	
	6	8.5	15	38	30	50*
Biomass yield (g $COD_{biomass} \cdot g^{-1} COD_{consumed}$)	1.00±0.07	0.93±0.01	0.85±0.07	0.99±0.07	1.05±0.01	1.04±0.02

Biomass production rate (mgCOD·L⁻¹·d⁻¹)	202±37	77±26	56±44	182±59	378±72	351±60
Protein content of biomass (%w/w)	-	56±5	54±2	51±3	64±2	51±2
Protein yield (g protein·g⁻¹ H₂)	-	3.9±0.3	4.4±0.6	4.3±0.5	4.9±0.8	4.2±0.3
C content of biomass (%w/w)	42±1	41±2	41±1	41±1	43±3	41±2
N content of biomass (%w/w)	8±1	10±0	8±0	9±0	10±1	10±0

*These values also corresponded to the pH 7 and temperature 25°C tests.

CONCLUSIONS

This study demonstrated, that PPB consortia can be efficiently used for production of SCP from H₂. Neutral pH, temperatures of ~25°C and light intensities higher than 30 W·m⁻² are the best conditions for biomass growth. High biomass yields and protein contents were achieved. The high yields and rates achieved confirm the great potential of enriched PPB cultures for H₂ valorisation.

ACKNOWLEDGEMENTS

The authors acknowledge NextGenerationEU Margarita Salas programme from the European Union for the research contract of María del Rosario Rodero.

REFERENCES

- Alloul, A., Spanoghe, J., Machado, D., Vlaeminck, S.E. 2022. Unlocking the genomic potential of aerobes and phototrophs for the production of nutritious and palatable microbial food without arable land or fossil fuels. *Microb. Biotechnol.* **15**, 6–12.
- Capson-Tojo, G., Batstone, D.J., Grassino, M., Vlaeminck, S.E., Puyol, D., Verstraete, W., Kleerebezem, R., Oehmen, A., Ghimire, A., Pikaar, I., Lema, J.M., Hülsen, T. 2020. Purple phototrophic bacteria for resource recovery: Challenges and opportunities. *Biotechnol. Adv.* **43**, 107567.
- Capson-Tojo, G., Lin, S., Batstone, D. J., Hülsen, T. (2021). Purple phototrophic bacteria are outcompeted by aerobic heterotrophs in the presence of oxygen. *Water Res.* **194**, 116941.
- Hülsen, T., Stegman, S., Batstone, D.J., Capson-Tojo, G. 2022a. Naturally illuminated photobioreactors for resource recovery from piggery and chicken-processing wastewaters utilising purple phototrophic bacteria. *Water Res.* **214**, 118194.
- Hülsen, T., Züger, C., Gan, Z.M., Batstone, D.J., Solley, D., Ochre, P., Porter, B., Capson-Tojo, G. 2022b. Outdoor demonstration-scale flat plate photobioreactor for resource recovery with purple phototrophic bacteria. *Water Res.* **216**, 118327.
- Ormerod, J.G., Ormerod, K.S., Gest, H. 1961. Light-Dependent Utilization of Organic Compounds Photoproduction of Molecular Hydrogen by Photosynthetic Bacteria ; Relationships with Nitrogen Metabolism. *Arch. Biochem. Biophys.* **94**, 449–463.
- Puyol, D., Batstone, D.J., Hülsen, T., Astals, S., Peces, M., Krömer, J.O. 2017. Resource recovery from wastewater by biological technologies: Opportunities, challenges, and prospects. *Front. Microbiol.* **7**, 1–23.
- Ritala, A., Häkkinen, S.T., Toivari, M., Wiebe, M.G. 2017. Single cell protein-state-of-the-art, industrial landscape and patents 2001-2016. *Front. Microbiol.* **8**, 2009.
- Spanoghe, J., Vermeir, P., Vlaeminck, S.E. 2021. Microbial food from light, carbon dioxide and hydrogen gas: Kinetic, stoichiometric and nutritional potential of three purple bacteria. *Bioresour. Technol.* **337**, 125364.
- Zha, X., Tsapekos, P., Zhu, X., Khoshnevisan, B., Lu, X., Angelidaki, I. 2021. Bioconversion of wastewater to single cell protein by methanotrophic bacteria. *Bioresour. Technol.* **320**, 124351.