



HAL
open science

Genetic control of abiotic stress-related specialized metabolites in sunflower

Marco Moroldo, Nicolas Blanchet, Harold Duruflé, Stéphane Bernillon, Thierry Berton, Olivier Fernandez, Yves Gibon, Annick Moing, Nicolas Langlade

► To cite this version:

Marco Moroldo, Nicolas Blanchet, Harold Duruflé, Stéphane Bernillon, Thierry Berton, et al.. Genetic control of abiotic stress-related specialized metabolites in sunflower. *BMC Genomics*, 2024, 25 (1), pp.199. 10.1186/s12864-024-10104-9 . hal-04468891

HAL Id: hal-04468891

<https://hal.inrae.fr/hal-04468891>

Submitted on 21 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

RESEARCH

Open Access



Genetic control of abiotic stress-related specialized metabolites in sunflower

Marco Moroldo^{1*}, Nicolas Blanchet¹, Harold Duruflé^{1,4}, Stéphane Bernillon^{2,3,5}, Thierry Berton², Olivier Fernandez^{2,6}, Yves Gibon^{2,3}, Annick Moing^{2,3} and Nicolas B. Langlade¹

Abstract

Background Abiotic stresses in plants include all the environmental conditions that significantly reduce yields, like drought and heat. One of the most significant effects they exert at the cellular level is the accumulation of reactive oxygen species, which cause extensive damage. Plants possess two mechanisms to counter these molecules, i.e. detoxifying enzymes and non-enzymatic antioxidants, which include many classes of specialized metabolites. Sunflower, the fourth global oilseed, is considered moderately drought resistant. Abiotic stress tolerance in this crop has been studied using many approaches, but the control of specialized metabolites in this context remains poorly understood. Here, we performed the first genome-wide association study using abiotic stress-related specialized metabolites as molecular phenotypes in sunflower. After analyzing leaf specialized metabolites of 450 hybrids using liquid chromatography-mass spectrometry, we selected a subset of these compounds based on their association with previously known abiotic stress-related quantitative trait loci. Eventually, we characterized these molecules and their associated genes.

Results We putatively annotated 30 compounds which co-localized with abiotic stress-related quantitative trait loci and which were associated to seven most likely candidate genes. A large proportion of these compounds were potential antioxidants, which was in agreement with the role of specialized metabolites in abiotic stresses. The seven associated most likely candidate genes, instead, mainly belonged to cytochromes P450 and glycosyltransferases, two large superfamilies which catalyze greatly diverse reactions and create a wide variety of chemical modifications. This was consistent with the high plasticity of specialized metabolism in plants.

Conclusions This is the first characterization of the genetic control of abiotic stress-related specialized metabolites in sunflower. By providing hints concerning the importance of antioxidant molecules in this biological context, and by highlighting some of the potential molecular mechanisms underlying their biosynthesis, it could pave the way for novel applications in breeding. Although further analyses will be required to better understand this topic, studying how antioxidants contribute to the tolerance to abiotic stresses in sunflower appears as a promising area of research.

Keywords Sunflower (*Helianthus annuus*), Abiotic stresses, Metabolome, Liquid chromatography-mass spectrometry, Genome-wide association study, Antioxidants, Specialized metabolites, Reactive oxygen species

*Correspondence:

Marco Moroldo

marco.moroldo@inrae.fr

Full list of author information is available at the end of the article



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

Abiotic stresses in plants can be defined as all the environmental conditions that decrease growth and yield below optimum levels [1]. They include, among others, drought, salinity, low and high temperatures, nutrient deficiencies, and ultraviolet radiation [2]. The impact of most of these stresses is becoming more severe because of climate change [1]. Abiotic stresses exert their effects in many complex and diverse ways, and plants have evolved a vast array of mechanisms to cope with them. Some of these mechanisms include the accumulation of wax and cutin on leaf surfaces, the desaturation of membrane lipids, and the accumulation of compatible solutes [2].

At the molecular level, one of the most significant effects of abiotic stresses is the accumulation of reactive oxygen species (ROS) [3], which arises from an imbalance between ROS production and scavenging [4, 5]. ROS are strong oxidizers and cause extensive damage to many biological molecules, like for instance proteins, lipids, and DNA [3, 5].

Plants use two mechanisms to counterbalance oxidative stress. The first is represented by detoxifying enzymes, such as superoxide dismutase, catalase, ascorbate peroxidase, and glutathione reductase [4]. The second corresponds to non-enzymatic antioxidants, i.e. ascorbic acid, reduced glutathione, α -tocopherol, and several classes of secondary or specialized metabolites such as carotenoids, flavonoids, and phenolic acids, whose ROS-scavenging activity has been demonstrated across many plant species [5, 6].

Terpenes are another class of specialized metabolites with antioxidant properties. Although better known as constituents of essential oils, allelopathic agents, and attractants or repellants in plant–herbivore interactions [7], there is increasing evidence of their implication in ROS scavenging [8–10]. Taken together, it can be stated that most specialized metabolites induced by abiotic stresses show antioxidative activity [11], which makes these molecules key players for plant adaptation to more stressful environments and for breeding tolerant varieties.

Sunflower (*Helianthus annuus* L.) is the fourth most important oilseed worldwide. It can maintain stable yields across many conditions and, largely thanks to its well-developed tap roots, it is adapted to low water-input regimes in warm to semi-arid zones [12]. Although this crop is usually considered moderately drought tolerant, the challenges posed by climate change will require major efforts in terms of breeding and crop management.

To cope more efficiently with hydric stress, tolerant varieties will have to be developed [13], and tolerance to heat will have to be jointly prioritized, because high

temperatures dramatically affect pollination, fertilization, and seed set [13]. Another way to avoid drought is early sowing. This strategy allows to anticipate the timing of flowering, thus avoiding the summer periods in which evaporative demand is higher [13, 14]. However, this practice presents side-effects, because the crop is more exposed to cold stress at germination [15]. From this perspective, developing hybrids with improved tolerance to cold will be another relevant goal.

The molecular mechanisms underlying tolerance to abiotic stresses in sunflower have been studied using different approaches over the last years. Non-targeted metabolomics and proteomics have been used to profile a set of inbred lines and hybrid genotypes [16, 17] and to find biomarkers for drought tolerance [18], while transcriptome and metabolome have been integrated to identify transcription factors regulated under the same condition [19].

Transcriptome profiling has been used to describe the impact of drought using co-expression networks [20], differential analysis [21], differential analysis coupled to association genetics [22] and gene-phenotype networks [23]. It has also been chosen to characterize low-nutrient stress and three water-related stresses [24]. Eventually, tolerance to salt stress and its link with vigor have been studied through association genetics [25]. Nevertheless, to date no information is available concerning the genetic control of specialized metabolome in sunflower under abiotic stress.

In this work, we present the results of the first GWAS performed in sunflower using specialized metabolites related to abiotic stresses as molecular phenotypes. Our approach consisted of three main steps. First, we analyzed the semi-polar fraction of leaf extracts of a panel of sunflower hybrids using untargeted liquid chromatography-mass spectrometry (LC-MS), which allowed us to focus our analysis on specialized metabolites. Second, we selected a subset of these compounds based on their genetic association with some previously known quantitative trait loci (QTLs) related to yield and abiotic stress tolerance. Third, we characterized *in silico* these compounds and the genes associated with them.

It has also to be noticed that, in addition to our own work, GWAS using molecular phenotypes in sunflower has been used so far only in another case, i.e. to disentangle the genetic basis of oil fatty acid content [26]. Our results can therefore be considered original by a methodological point of view.

Results

Association mapping

To study the genetic control of specialized metabolites in sunflower and how this relates to abiotic stress

tolerance, we obtained the metabolomic profiles of the leaves of 450 hybrids originating from crosses among 36 restorer and 36 cmsPET1 sterile lines grown in agronomical conditions. After the partial removal of redundancy due to isotopes and adducts, the final metabolome dataset consisted of 2557 LC-MS features (Table S01) characterized by a retention time (RT) and a mass over charge ratio (m/z) (Table S02). A total of 21 features already had an annotation based on previous works (Table S02) [16, 18, 27].

A PCA performed using these data showed that the first two principal components accounted for 11% and 5% of total variability, which was consistent with the results obtained in similar contexts [28]. A clustering of most of the hybrids according to their male parental line was observed (Fig. 1). This was especially evident for SF295, SF324, SF330, SF342 and SF281 male lines, but could be observed in other cases as well. On the contrary, no clustering was observed according to female parental lines.

The 2557 LC-MS features were then used as an input for the first step of the association analysis, which consisted in performing GWAS using reference SNPs. A visual overview of this analysis step, as well as all the other ones included in our workflow, is found in Fig. 2. Similarly to what observed in other plant species [29], 955 LC-MS features (i.e. 37.3% of the total number used) were associated to at least one SNP. This corresponded to 2560 associations (Table S03). On average, an LC-MS feature was therefore associated to 2.7 SNPs, with 472 features associated to only one SNP and 483 features associated to two to 19 SNPs.

Among the 955 significantly associated LC-MS features, 798 (i.e. 83.6%) presented genomic heritability (h^2_g) values higher than 0.50, the average being 0.66 (Fig. S01), which was in line with what reported in the literature [29, 30]. The 2560 detected associations corresponded to 1716 unique reference SNPs, with 378 SNPs associated to two to 43 features. This suggested that many LC-MS features were under pleiotropic




Fig. 1 Individual plot of the first two components of the PCA based on the normalized intensities of 2557 LC-MS features measured in the leaves of 450 sunflower hybrids. The different hybrids are colored according to their male parental lines, which are indicated in the box on the right




Fig. 2 Graphical abstract illustrating the main steps of our analysis workflow. (1) First step of GWAS: detection of the associations among LC-MS features (orange boxes) and reference SNPs (red vertical bars); (2) Second step of GWAS: reference SNPs are linked to co-inherited SNPs sets (blue vertical bars); (3) Co-localization among co-inherited SNPs and SNPs belonging to abiotic stress-related QTLs (green vertical bar). Co-inherited SNPs falling in a 50 kb interval downstream or upstream of a SNP belonging to a QTL are considered co-localizing with the same QTL. All the SNPs in complete linkage disequilibrium (LD) with co-localizing SNPs and mapping to exons (yellow transparent boxes) are then used to identify putative candidate genes; (4) LC-MS features associated to reference SNPs in complete LD with co-localizing SNPs are selected and tentatively annotated

control, as already described in other species [31, 32], or that some biochemical information redundancy was still present in our data set after LC-MS data filtering. Five features already possessed an annotation based on previous works (Table S03).

The second step of association analysis consisted in linking reference SNPs to their corresponding sets of co-inherited SNPs, which included all the SNPs in complete linkage disequilibrium (LD) with them (see Fig. 2 and the Methods section). After this step, 62,134 associations were found. The number of associations per LC-MS feature ranged from one to 3666, the average being of 65.1 (Table S04). Overall, these associations corresponded to 27,246 unique SNPs.

To gather a first functional understanding of the genetic control of sunflower specialized metabolome, we then investigated all the possible associations among LC-MS features and genes in an unsupervised way, i.e. without any further biological information. A gene was considered associated to a feature if at least one SNP of a co-inherited set mapped to one of its exons (Fig. 2). Our analysis highlighted that 1768 SNPs belonging to

co-inherited sets out of 27,246 (i.e. 6.5%) were found in exons.

Exonic SNPs corresponded to 533 genes, with an average of 3.3 associations per gene (Table S05). These genes appeared to be involved in several pathways, with a slight over-representation of those related to glutathione, lipids and specialized metabolite biosynthesis, like for instance flavonoids. Anyway, it must be considered that the levels of enrichment, especially in terms of numbers of genes associated to each ontology, were rather low (Table S06).

As observed in other species [29, 30], metabolite-associated SNPs and genes were not randomly distributed across the genome, but appeared to be especially concentrated in some specific 'hot spots' (Fig. S02). To explore this pattern of distribution, we first used a sliding window approach and subsequently, by applying a threshold based on the proportion and on the absolute number of metabolite-associated genes in each window, we defined six hot spots on chromosomes 5, 6, 7, 9, 12 and 16. These regions contained 88 metabolite-related genes and spanned 48 Mb, which corresponds to a sixth of all metabolite-related genes on slightly less than 1.5% of the

sunflower genome (Table S07). In some instances, these genes were arranged in small families, like in the case of the hot spot on chromosome 5, which contained six putative quinate O-hydroxycinnamoyltransferase, and in the case of the hot spot on chromosome 6, where seven putative glutathione transferases were detected (Table S07). Anyway, despite the presence of these specific patterns, we could not find any evidence of functional metabolic clusters as defined by Nützmann and coworkers [33].

Co-localization of SNPs associated to LC-MS features with QTLs related to abiotic stresses

As illustrated in Fig. 2, to study the genetic control of specialized metabolites linked to abiotic stresses we tested the co-localization of co-inherited SNPs associated to LC-MS features with previously identified QTL regions for drought, cold, and nutrient stress tolerance and for productivity and development-related traits [15, 34]. These last two groups of traits were added because they were considered as indirectly related to abiotic stress tolerance.

We detected a total of 638 SNPs that co-localized with 20 QTLs, among which seven were related to drought stress tolerance, five to cold stress tolerance, four to nutrient stress tolerance and four to productivity and development-related traits, with the QTLs for abiotic stress tolerances showing some overlap among them (Table S08). We then took these 638 co-localizing SNPs and searched for all the other SNPs which were in complete LD with them, and which were 10,793. On the one hand, these co-inherited SNPs were associated to 137 LC-MS features (Table S09), of which 16 were related to drought stress tolerance, 92 to cold stress tolerance, four to nutrient stress tolerance and 17 to productivity and development-related traits. A further eight LC-MS features were associated to two QTLs at the same time (Table S09). On the other hand, the same co-inherited SNPs were also associated to 155 putative candidate genes (Table S09).

Annotation of LC-MS features of interest and identification of the most likely candidate genes

Because the LC-MS protocol used in this work did not involve data dependent MS/MS (i.e. tandem mass spectrometry; see the [Methods](#) section), the only way to perform the annotation of the 137 previously identified LC-MS features of interest was by relying on an in silico workflow. This procedure allowed to tentatively annotate 30 features, among which one was related to drought stress tolerance, 21 to cold stress tolerance, one to nutrient stress tolerance and four to productivity and development-related traits. A further three features were associated to two QTLs at the same time. Most of the

annotated molecules (Table 1) belonged to the biochemical classes of terpenes (30%), flavonoids (17%), polyacetylenes (17%) and cinnamic acids (10%).

Subsequently, the results obtained from GWAS showed that 13 out of the 30 aforementioned metabolites were associated to at least one gene, corresponding to a total of 80 'initial' genes (Table S10). Two metabolites, namely 1,4-tridecadiene-7,9-diyne (a polyacetylene) and 4,5,9,10-dehydroisolongifolene (a terpene), were associated to the same genes, potentially suggesting pleiotropy.

After the process of functional characterization of all of these associations, we focused on a final set of four metabolites that could be related to seven most likely candidate genes (Table 2). Three associations involved metabolites and enzyme-encoding genes, namely (i) the flavonoid pentahydroxychalcone and a member of the P450 cytochrome family; (ii) the sesquiterpene heliannuol F and a uridine diphosphate (UDP) glucosyltransferase (UGT); (iii) the flavonoid hexahydroxydimethylflavanone and three UGTs. The sesquiterpene 4,5,9,10-dehydroisolongifolene, instead, was associated to two transcription factors (TFs) of the AP2/ERF family (Table 2).

Overall, these genes were linked to 13 SNPs belonging to co-inherited sets, 10 of which caused missense mutations (Table 2). Eventually, the genotypic boxplots corresponding to the four characterized metabolites showed in all of the cases a good correlation among the different allelic states and the phenotypic values of the LC-MS features (Fig. 3).

Discussion

Potential antioxidants represent a large proportion of sunflower leaf metabolome under abiotic stress

Oxidative damage is one of the most important modifications induced by abiotic stresses in plant cells. It is caused by the accumulation of ROS and has been reported, for instance, in the cases of drought, cold and salinity [5, 11]. Plants use many strategies to mitigate the impact of ROS, one of which is the biosynthesis of non-enzymatic antioxidants. These compounds include, among others, several important groups of semi-polar specialized metabolites like carotenoids, flavonoids, and phenolic acids [6]. Indeed, the majority of specialized metabolites produced by plants under abiotic stress show antioxidative activity in vitro, even if an in vivo experimental confirmation of their function is still lacking in many cases [11].

In this work, we studied the metabolome of 450 sunflower hybrids by performing LC-MS on the semi-polar fraction of leaf extracts, thus specifically targeting specialized metabolites. We then selected a subset of these compounds based on their genetic association to some previously known QTLs which were mainly related to

Table 1 Putative annotation of the 30 LC-MS features measured in the leaves of sunflower hybrids and co-localizing with QTLs of interest. All the metabolites were assigned an MSI level 3 (see Methods)

LC-MS feature	[M+H] ⁺ m/z	RT (min)	isotopic pattern ion	putative chemical formula for [M+H] ⁺	putative metabolite name	metabolite ID	metabolite class
M201T581	201.1637	9.69	M0	C15H21	4,5,9,10-dehydroisolongifolene isomer A	CID: 588771	sesquiterpenes
M203T379	203.1066	6.32	M0	C13H15O2	demethoxyencecalin	CID: 177040	chromenes
M287T548	287.0550	9.14	M0	C15H11O6	cyanidin	CID: 128861	flavonoids
M289T534	289.0706	8.89	M0	C15H13O6	pentahydroxychalcone	CID: 129636553	flavonoids
M307T374	307.1288	6.23	M0	C15H19O5N2	brachystemidine A	CID: 10892118	pyrroles
M349T519	349.0917	8.64	M0	C17H17O8	hexahydroxydimethylflavanone	–	flavonoids
M131T321	131.0490	5.36	M0	C9H7O	2-nonene-4,6,8-triyn-1-ol	CID: 57449462	polyacetylenes
M294T138	294.1547	2.31	M0	C12H24O7N	deoxyfructosyl-leucine	CID: 131752244	amino acid derivatives
M145T650	145.1011	10.83	M0	C11H13	1,9-undecadiene-5,7-diyne	CID: 15736690	polyacetylenes
M151T299	151.0754	4.99	M0	C9H11O2	coumaryl-alcohol isomer A	CID: 5280535	cinnamic acids
M151T342	151.0754	5.71	M0	C9H11O2	coumaryl-alcohol isomer B	CID: 5280535	cinnamic acids
M165T301	165.0545	5.02	M0	C9H9O3	coumaric acid	CID: 637542	cinnamic acids
M173T648	173.1324	10.80	M0	C13H17	1,4-tridecadiene-7,9-diyne	CID: 101410972	polyacetylenes
M175T609	175.0753	10.14	M0	C11H11O2	(2E,4E)-5-phenylpenta-2,4-dienoic acid	CID: 1549512	styrenes
M201T687	201.1637	11.45	M0	C15H21	4,5,9,10-dehydroisolongifolene isomer B	CID: 588771	sesquiterpenes
M202T649	202.1671	10.81	M1	C15H21	4,5,9,10-dehydroisolongifolene isomer C	CID: 588771	sesquiterpenes
M217T336	217.1585	5.60	M0	C15H21O	(3S,9Z)-pentadeca-9,14-dien-4,6-diy-3-ol	CID: 163193273	polyacetylenes
M231T299	231.1378	4.99	M0	C15H19O2	dehydrocostus lactone isomer A	CID: 73174	sesquiterpenes
M231T362	231.1378	6.03	M0	C15H19O2	dehydrocostus lactone isomer B	CID: 73174	sesquiterpenes
M231T604	231.1378	10.06	M0	C15H19O2	dehydrocostus lactone isomer C	CID: 73174	sesquiterpenes
M243T272	243.0881	4.54	M0	C12H11O2N4	lumichrome	CID: 5326566	alloxazines
M247T334	247.1327	5.57	M0	C15H19O3	annuolide A	CID: 44583825	sesquiterpenes
M249T273	249.1484	4.55	M0	C15H21O3	annuolide E	CID: 131752335	sesquiterpenes
M266T572	266.1467	9.53	M1	C15H21O4	heliannuol F	CID: 10730325	sesquiterpenes
M273T341	273.1485	5.69	M0	C17H21O3	8-acetoxy-1,9,14-pentadecatriene-4,6-diy-3-ol	CID: 14037439	polyacetylenes
M273T635_2	273.2213	10.59	M0	C19H29O	androst-5-en-4-one	CID: 22213540	steroids
M299T342	299.1278	5.70	M0	C18H19O4	3-phenyl-1-(2,3,4-trimethoxyphenyl)prop-2-en-1-one	CID: 67199147	flavonoids
M346T682	346.1001	11.37	M1	C18H17O7	tambulin	CID: 5281700	flavonoids
M372T338	372.1288	5.64	M0	C16H22O9N	4-(2-amino-3-hydroxyphenyl)-4-oxobutanoic acid glucoside	CID: 10948689	phenolic compounds
M515T343	515.2122	5.72	M0	C24H35O12	eugenol acetylramnosylglucoside	–	phenolic compounds

abiotic stresses and, to a lesser extent, development-related traits. Eventually, we annotated this subset of molecules. However, it must be considered that because tandem mass spectrometry had not been performed and because MS commercial standards were not available for the large majority of these metabolites, only putative annotations could be assigned.

Most of the tentatively annotated metabolites belonged to four biochemical classes, namely terpenes,

flavonoids, polyacetylenes and cinnamic acids (Table 1). These findings were in line with our previous research on sunflower, and specifically with the works focusing on drought stress [16, 18]. However, the detection of polyacetylenes was unique to this study.

Flavonoids, represented by five molecules in our data, are among the best characterized ROS scavengers in plants [7, 35]. Their activity is known to provide

Table 2 List of the four tentatively annotated metabolites that were associated to seven most likely candidate genes. Column seven reports the results of the tBlastN analysis when they appear to be helpful in clarifying the already available gene descriptions

LC-MS feature	putative metabolite name	metabolite class	v2.1 gene	v2.1 gene description	tBLASTn result	co-inherited v1.0 SNP	type of mutation	base mutation	amino acid change	reference v1.0 SNP
M201T581	4,5,9,10-dehydroisolongifolene	sesquiterpenes	HanXRQr2_Chr07g0314841	putative transcription factor AP2-EREBP family	AP2/ERF transcription factor	HanXRQChr07_98876273	synonymous			HanXRQChr07_97029247
M266T572	heliannuol F	sesquiterpenes	HanXRQr2_Chr07g0314871	putative transcription factor AP2-EREBP family	AP2/ERF transcription factor	HanXRQChr07_98748814	missense	T > G	ASP>GLU	HanXRQChr07_97029247
M289T534	pentahydroxy-chalcone	flavonoids	HanXRQr2_Chr09g0363371	putative UDP-glucuronosyl/UDP-glucosyltransferase	Glycosyltransferase 74G1, likely family 1	HanXRQChr09_10756950 HanXRQChr09_10757103	missense missense	G > A G > T	ALA>THR ALA>SER	HanXRQChr09_10756950 HanXRQChr09_10756950
M349T519	hexahydroxy-dimethylflavanone	flavonoids	HanXRQr2_Chr11g0515571	putative UDP-glucuronosyl/UDP-glucosyltransferase	Glycosyltransferase 73E1, likely family 1	HanXRQChr03_139760168	synonymous			HanXRQChr03_139760112
			HanXRQr2_Chr11g0515581	hypothetical protein	Glycosyltransferase 73E1, likely family 1	HanXRQChr11_157852754 HanXRQChr11_157852773	missense missense	G > A C > T	ASP>ASN SER>LEU	HanXRQChr11_157849826 HanXRQChr11_157849826
			HanXRQr2_Chr11g0515591	Putative trans-zeatin O-beta-D-glucosyltransferase	Glycosyltransferase 73E1, likely family 1	HanXRQChr11_157874063 HanXRQChr11_157873968 HanXRQChr11_157874469	synonymous missense missense	C > T A > G	TYR > HIS ARG > GLY	HanXRQChr11_157849826 HanXRQChr11_157849826 HanXRQChr11_157849826

tolerance towards many abiotic stresses, like drought [36], cold [37], and nutrient depletion [38].

Even if better known as constituents of essential oils or allelopathic agents [7], there is increasing evidence that terpenes are also involved in ROS scavenging, as described for instance in tea plant [10], sage, and rosemary [8, 9]. In other instances, the involvement of terpenes in abiotic stress tolerance has been demonstrated, even if the underlying antioxidant mechanism has not been proven yet [39].

The class of cinnamic acids, here intended as including all the derivatives of cinnamic acid, is found in most plant families, including Asteraceae [18, 27, 40]. Chlorogenic acid shows antioxidant properties, although this could be true also in the case of other cinnamic derivatives [41]. However, no ROS-scavenging activity has been demonstrated in the case of coumaric acid and coumaryl alcohol. This latter molecule, instead, is one of the precursors of lignin [42] whose accumulation, in turn, is increased for instance under drought and cold [43].

Polyacetylenes are found in a few botanical families, Apiaceae and Asteraceae being among the most relevant [44, 45]. They exhibit a wide range of antibacterial, antifungal, and insecticidal activities [45, 46].

The remaining annotated compounds belonged to several different classes. Phenolic acids, represented by 4-(2-amino-3-hydroxyphenyl)-4-oxobutanoic acid glucoside and eugenol acetylramnosylglucoside, are well characterized as antioxidants in plants [6, 47].

The mammal steroid androstenone is found in many plant species [48]. Mammal steroids in plants are known to be involved in processes such as root and shoot growth [49], but no information is specifically available for androstenone.

Demethoxyencecalin is a chromene. The compounds in this class have been described in plants such as mulberry [50] and *Hypericum polyanthemum* [51], but they are especially frequent in Asteraceae [52]. They are repellent towards herbivorous insects.

(2E,4E)-5-phenylpenta-2,4-dienoic acid belongs to styrenes, which are naturally synthesized for instance by mulberry [50] and styrax [53]. Some styrenes show

antifeedant activity against insects in pear [54], but no evidence is available in the case of (2E,4E)-5-phenylpenta-2,4-dienoic acid.

The last three molecules highlighted by our study are lumichrome (i.e. an alloxazine), brachystemidine A, (i.e. a pyrrole), and deoxyfructosyl-leucine (i.e. an amino acid derivative). Their role is not clear and therefore difficult to discuss in our biological context.

Altogether, it is possible to affirm that a relevant fraction of the metabolites that we have tentatively annotated, i.e. from seven (considering flavonoids and phenolic acids) to 16 (including also terpenes) out of 30, fit in biochemical classes with oxygen scavenging properties.

Although the number of characterized molecules in our work is relatively small, and even if their specific *in vivo* activity has not been proven yet, our findings can be considered in agreement with the high proportion of antioxidants observed by many authors in abiotic stress-related specialized metabolome [11, 31, 55, 56].

The most likely candidate genes mainly belong to two highly diverse superfamilies

Interestingly, the five enzyme-encoding most likely candidate genes found in our study belonged to only two families, namely cytochromes P450 (CYPs) and glycosyltransferases (GTs). Although very different in their functionalities, both of them are large and catalyze greatly diverse reactions that create a vast array of chemical modifications. This is consistent with the high level of plasticity of specialized metabolism pathways in plants [57].

The reactions catalyzed by CYPs play a basic role in defining the skeletal structure of many metabolites, such as flavonoids and terpenes [58, 59]. These reactions include hydroxylations, reductive activations, ring couplings, ring formations, ring expansions and oxidative aryl migrations [60, 61].

Today, a large number of CYPs involved in flavonoid biosynthesis are known in many plant families, including Asteraceae [62, 63]. It is therefore possible to hypothesize that the enzyme encoded by the

(See figure on next page.)

Fig. 3 Manhattan plots and genotypic boxplots of the functionally characterized metabolites, here indicated using the codes of LC-MS features. For each metabolite, the corresponding Manhattan plot (left) and genotypic boxplot (right) are shown. Manhattan plots show the reference SNPs obtained from GWAS on the X-axis and the corresponding *p*-values on the Y-axis. SNPs filtered according to the eBIC criterion are shown as asterisks (Table S03). The reference SNP associated to a most likely candidate gene is highlighted by a grey circle. Genotypic boxplots show the normalized intensity values of the corresponding LC-MS feature (Y-axis) grouped according to the three possible allelic states (i.e. 00, 01|10, and 11). The classes identified with the Tukey's test are indicated using colored squares. It is to note that the SNPs used to produce the genotypic boxplots are, by definition, reference SNPs, and therefore they are not the same ones found associated to the corresponding most likely candidate gene, which belong instead to co-inherited sets. The correspondences among the reference SNPs used for genotypic plots and the co-inherited used to identify most likely candidate genes are given in Table 2




Fig. 3 (See legend on previous page.)

HanXRQr2_Chr03g0130651 gene performs one of the molecular reactions that lead to the biosynthesis of the flavonoid pentahydroxychalcone.

Unlike cytochromes P450, all glycosyltransferases catalyze the same type of reaction, i.e. the transfer of a sugar moiety to an acceptor. Anyway, they act on a broad range of compounds such as lipids, proteins, nucleic acids and other molecules [64].

GTs are classified in many families according to the CAZy database (<http://www.cazy.org/>). Family 1 is defined by the presence of a specific domain [64, 65] and is usually referred to as UDP-glycosyltransferases (UGTs). Plant UGTs are especially involved in the glycosylation of specialized metabolites such as flavonoids and terpenes [66].

Glycosylation increases the activity and the availability of both these categories of compounds and hence the ROS-scavenging capacity of the plant, which confers tolerance to several abiotic stresses. In *Arabidopsis*, for instance, the enzymes encoded by *UGT79B2* and *UGT79B3* add a UDP-rhamnose to the flavonoids cyanidin and cyanidin 3-O-glucoside. An increased concentration of these two molecules provides tolerance to cold, salinity and drought [67]. In tea plant, instead, *CsUGT78A14-1* and *CsUGT78A14-2* are involved in the biosynthesis of the flavonoids kaempferol 3-O-glucoside and kaempferol diglucoside, which reduces oxidative damage and increases tolerance to cold stress [68]. Another similar example is provided by the action of sesquiterpene nerolidol in tea plant. Again, this metabolite is glycosylated by the protein encoded by *CsUGT91Q2*, which causes an enhanced level of tolerance against cold stress [10].

In light of this, it could be speculated that the three UGTs associated to the flavonoid hexahydroxydimethylflavanone, i.e. *HanXRQr2_Chr11g0515571*, *HanXRQr2_Chr11g0515581* and *HanXRQr2_Chr11g0515591*, could be involved in the glycosylation of this molecule, thus contributing to reduce the impact of oxidative damage in sunflower. Likewise, the UGT *HanXRQr2_Chr09g0363371* could be involved in ROS scavenging through the glycosylation of the terpene heliannol F.

Besides enzyme-encoding genes, we also found that two transcription factors of the AP2/ERF family, i.e. *HanXRQr2_Chr07g0314841* and *HanXRQr2_Chr07g0314871* were associated to the terpene 4,5,9,10-dehydroisolongifolene. Because some transcription factors of this family are implicated in the biosynthesis of terpenes, as in the cases of orange [69] and *Litsea cubeba* [70], it is possible to imagine a potential link with the aforementioned metabolite.

As already stated, 4,5,9,10-dehydroisolongifolene was associated to the same co-inherited SNPs sets that were

linked to the polyacetylene 1,4-tridecadiene-7,9-diyne. This could suggest a potential case of pleiotropic control of metabolite biosynthesis, which has already been described in other plants [31, 32]. Because the biochemical pathways of terpenes and polyacetylenes are completely different, the only genes that could explain this case of pleiotropy are indeed the previously indicated AP2/ERF transcription factors. Anyway, to date information about which TFs could be involved in the biosynthesis of polyacetylenes is lacking, thus making it difficult to draw conclusions in this respect.

Despite the limited number of most likely candidate genes identified, our results appear globally in agreement with those obtained from similar metabolic GWAS analyses performed under abiotic stress in *Arabidopsis thaliana* [71] and maize [72].

It has also to be considered that our capability to identify the most likely candidate genes was reduced by some technical limitations and specific features of our study. On the one hand, gene functions are largely unknown in plants, particularly in a species such as sunflower. Indeed, 12 genes out of the 80 that were initially found associated with co-inherited SNPs sets, i.e. previous to the process of functional characterization, were annotated as 'hypothetical' or 'putative' proteins. On the other hand, our approach to link SNPs to genes was rather stringent, because it required the SNPs to directly land on exons in order to identify a potential candidate.

Conclusions

Our work represents the first characterization of the genetic control of abiotic stress-related specialized metabolites in sunflower. It provides hints concerning the importance of antioxidant compounds in this biological context, and it highlights some of the potential molecular mechanisms underlying their biosynthesis, thus paving the way for novel applications in sunflower breeding. Even if our capability to identify candidate genes was diminished by some technical limitations, our results were consistent with those obtained from similar metabolic GWAS performed under abiotic stress in plants such as *Arabidopsis thaliana* and maize. Although further analyses will be needed to obtain a deeper understanding of the topic, studying how antioxidants contribute to the tolerance to abiotic stresses in sunflower appears as a promising area of research.

Methods

Plant material and sampling

A panel of 475 sunflower hybrids, corresponding to an incomplete factorial design, was obtained by crossing 36 male and 36 female inbred lines as previously described [73, 74]. Each hybrid was named using its respective

female and male parental lines and adding an underscore to separate them.

As already described [75], each hybrid was grown in a single 13 m² plot, and all the plots were cultivated on the same field trial in Anais (Charente-Maritime, France) from 2 May 2015 to 29 September 2015. Four control hybrids, corresponding to 65 plots, were included in the field trial to allow for the subsequent adjustment of spatial biases. Therefore, the trial included a total number of 540 plots.

For each single plot, *n*-4 topmost leaves without petioles were sampled from four different plants on July 22 2015, i.e. 7 days (\pm 3 days according to genotypes) after blooming, between 11:00 to 12:30 (CET time), and then pooled. Each pool was immediately frozen in dry ice and stored at -80°C until grinding.

Metabolome profiling

Leaf samples were cryoground using a Retsch Mill MM 400 ball mixer (Thermo Scientific, Waltham, MA, USA) and lyophilized. Aliquots of 10 ± 1.0 mg of dry leaf powders were weighed in 1.1 mL Micronic tubes (Micronic, Lelystad, The Netherlands) and extracted at room temperature with a robotized Star/Starlet platform (Hamilton, Reno, NV, US) using ethanol/water (80:20, v/v) added with 0.1% formic acid and 1.37 mM methyl vanillate as solvent. Methyl vanillate was used as internal standard to verify the quality of injection for LC-MS.

Two successive extractions (1 min shaking followed by 15 min ultra-sonication) were performed with 300 μL of extraction solvent. The two supernatants were combined and filtered using 0.22 μm hydrophilic Durapore filtering microplates (Merck Millipore, Carrigtwohill, Ireland). Several blank extracts were prepared using the same procedure and without sample powder. A quality control (QC) sample was prepared by pooling 10 μL of each sample extract.

LC-MS profiling was performed using the ethanol supernatant extracts. The sample injection order was randomized, and QC samples were injected every 10 samples to correct for the signal intensity drift. The extracts were analyzed using an LTQ Orbitrap Elite mass spectrometer (Thermo Scientific) interfaced to an UltiMate 3000L UHPLC system (Thermo Scientific) using a C18 chromatographic column (C18-Gemini 2.0 \times 150 mm, 3 μm , 110 \AA , Phenomenex, Torrance, CA, USA). An 18-min acetonitrile gradient in acidified water (solvent A: ultrapure water + 0.1% formic acid, solvent B: LC-MS grade acetonitrile) was used with a 300 $\mu\text{L}/\text{min}$ flow rate and the following elution gradient: 0–0.5 min, 3% B; 0.5–1 min, 3–10% B; 1–9 min, 10–50% B; 9–13 min, 50–100% B; 13–14 min, 100% B; 14–14.5 min 100–3% B; 14.5–18 min, 3% B. The column temperature was set

at 30°C and the injection volume was 5 μL . The LC-MS instrument was equipped with an electrospray ionization (ESI) source operated in the positive ion mode. Source parameters were set as follows: source voltage, 3.2 kV; sheath gas, 45 arbitrary units (a.u.); auxiliary gas, 15 a.u.; sweep gas, 0 a.u.; capillary temperature, 350°C ; heater temperature, 350°C . Full scan MS spectra were acquired at 240 k resolution power at 200 *m/z* with a 50–1000 *m/z* range. All the chemicals used for LC-MS were purchased from Sigma Aldrich (Saint Louis, MO, USA) and Extrasynthese (Genay, France).

LC-MS data were processed using the 'XCMS' R package [76]. Variables detected in blank extracts, with *m/z* values varying by more than 0.005 Da or with RT varying by more than 40 s between different samples were filtered out. Variables with intensity coefficients of variation in QCs greater than 20% were also removed. This resulted in a matrix of 3507 metabolite features. Intensity drift was corrected using support vector regression [77], and intensities were normalized according to the sample powder mass used for extraction. After a final step of quality assessment, the LC-MS data corresponding to 450 hybrids and 64 control hybrids were retained.

Processing, annotation and exploratory analysis of metabolome data

Biases occurring because of the spatial variation in the field trial were adjusted based on the information obtained from randomly replicated control hybrids using a script based on the 'ASReml-R' R package v 3.0 [78]. Data from control hybrids were discarded and the spatially corrected matrix was used for the subsequent steps of analysis.

Isotopes and adducts were searched for among the initial 3507 LC-MS features using the 'Binner' software [79]. A total of 950 redundant variables were removed, thus bringing the final LC-MS dataset to 2557 features. The most intense ions were then annotated using RT and accurate *m/z* values and the information available from previous studies [16, 18, 27]. This resulted in the putative annotation of 21 compounds (Table S02), whose MSI levels were attributed according to [80]. Eventually, to gather information about the structure of metabolome data, a PCA was carried out after scaling and mean centering using the 'pca' function of the 'mixOmics' R package v 6.16.3 [81].

Genotyping

The genotyping of the hybrids was carried out within the frame of the sunflower genome sequencing project [74]. Briefly, the parent lines were genotyped by whole genome resequencing, and the genotype of each hybrid was then obtained from those of its parents [74].

Initially, 14,127,553 SNPs were detected using the XRQ v1.0 assembly of the sunflower genome. Then, all the sets of SNPs in complete linkage disequilibrium among them, called ‘co-inherited SNPs sets’, were identified, and only one SNP was kept for each set. Subsequently, SNPs presenting a minor allelic frequency (MAF) < 0.1 or only detected in the male or female panel were filtered out. A final number of 350,052 SNPs, referred to as ‘reference SNPs’, were used for GWAS and are available through the Heliagene XRQ v1.0 genome portal (<https://www.heliagene.org/HanXRQ-SUNRISE/>). The genotypes of hybrids were coded as ‘0’, ‘1’, or ‘2’ for homozygous XRQ, heterozygous and variant homozygous, respectively. Both the additive (A) and the dominant (D) centered genotyping matrices were produced [73].

Association analysis

Association analysis was performed in two steps (Fig. 2). First, a multi-locus with forward selection GWAS was carried out with the ‘mlmm.gwas’ R package v 1.0.6 [82] and using the 2557 filtered LC-MS features and the 350,052 reference SNPs. Both the additive (A) and the dominant (D) effects of SNP markers were considered and 20 maximum steps were imposed for the fitting of the linear mixed model, which had an equation of this form:

$$y_i = \mu + x_i^l \theta_a^l + w_i^l \theta_d^l + A_i + D_i + e_i$$

Where x_i^l is the centered genotype of the i^{th} hybrid at the l^{th} marker locus; w_i^l is defined later; θ_a^l is the additive effect of the l^{th} locus; θ_d^l is the dominance effect of the l^{th} locus; and e_i denotes error.

A_i is the random additive effect of the i^{th} hybrid with the vector $A \sim \mathcal{N}(0, \sigma_a^2 K_a)$, D_i is the random dominant effect of the i^{th} hybrid with the vector $D \sim \mathcal{N}(0, \sigma_d^2 K_d)$, e_i is the residual error of the i^{th} hybrid with the vector $e \sim \mathcal{N}(0, \sigma_e^2 Id)$ and Id the identity matrix. K_a is the additive and K_d is the dominance kinship matrix calculated using the alike in state (AIS) relatedness criterion as indicated by [83]; σ_a^2 , σ_d^2 and σ_e^2 are additive, dominance and residual variances, respectively; and w_i^l is calculated as already described [83].

The best GWAS model was chosen using the extended Bayesian information criterion (eBIC) as proposed by [84]. The value of genomic heritability (h_g^2) for each LC-MS feature was calculated using the general purpose solver function ‘mixed.solve’ of the ‘rrBLUP’ R package v 4.6.1 [85].

The second step of the analysis consisted in linking reference SNPs to their corresponding co-inherited SNPs sets, which included all the SNPs in complete linkage disequilibrium with them (Fig. 2). This step was similar to

the ‘block analysis’ conducted by Temme and coworkers [25], although in our case blocks were defined by requiring complete LD among the different SNPs.

Unsupervised identification and enrichment analysis of putative candidate genes

To identify the candidate genes putatively involved in the biosynthesis of metabolites in an unsupervised way, i.e. without any further biological information, SNPs from co-inherited sets were mapped to exons, introns, and intergenic regions using a gft file corresponding to the annotation of the XRQ v1.0 sunflower genome (<http://www.heliagene.org/HanXRQ-SUNRISE/>). A gene was then considered associated to an LC-MS feature if at least one SNP from a co-inherited set mapped to one of its exons. The corresponding XRQ v2.1 genes were identified using the synonymy table available at the ‘Download’ section of the XRQ v2.1 portal at (<https://www.heliagene.org/HanXRQr2.0-SUNRISE/>).

The putative candidate XRQ v2.1 genes were used to perform enrichment analyses using the software ClueGO 2.5.8 [86]. A two-tailed hypergeometric test was performed to identify enriched ontology terms. Significance was set at a Benjamini-Hochberg-adjusted p -value of 0.05, the ‘GO fusion’ option was used and the k -score was fixed at 0.4. Three custom sunflower ontologies were used for the analysis, corresponding to the two Gene Ontology (GO) subsets ‘biological process’ and ‘molecular function’ and to the KEGG pathways.

The GO sub-ontology files were built using the Blast2GO output files available on the Heliagene website (<https://www.heliagene.org/HanXRQr2.0-SUNRISE/>), while the KEGG ontology file was created by performing a double best hit search using the XRQ v2.1 sunflower protein sequences on the KAAS automatic annotation server (<https://www.genome.jp/kegg/kaas/>). The KO codes thus obtained were then manually inspected in order to remove ontologies spuriously related to bacteria, fungi and animals.

Hot spots of metabolite-associated SNPs and genes

To describe the patterns of localization of metabolite-associated SNPs and genes along the sunflower genome and detect the potential presence of hot spots, a sliding windows approach was chosen. The window length was set at 5 Mb, and the window was incrementally advanced along the chromosomes using a pass of 1 Mb. For each window, the following measures were calculated: (i) the absolute number of metabolite-associated SNPs and their frequency respect to the total number of SNPs, and (ii) the absolute number of metabolite-associated genes and their frequency respect to the total number of genes. The values of the frequencies thus obtained were then plotted

against the sunflower chromosomes and visualized with the ‘Circlize’ R package v 0.4.14 [87].

A sliding window was considered as being part of a hot spot if it presented a frequency of metabolite-associated genes higher than 0.075 and an absolute number of genes higher than 10, and adjacent windows were merged in order to obtain the final hot spots. Eventually, the identification of potential metabolic clusters as defined by [33] was performed by visually inspecting the identified regions.

Identification of the SNPs co-localizing with known abiotic stress-related QTLs and of the associated LC-MS features

To study the genetic control of specialized metabolites linked to abiotic stresses, we followed a strategy based on co-localization with QTLs of interest that had been discovered in prior works. First, SNPs from co-inherited sets obtained from our GWAS analysis were tested for co-localization with two groups of QTLs, i.e.: (i) QTLs related to drought, cold, and nutrient stress tolerance which had been detected on other field trials; (ii) QTLs related to productivity and development traits which had been detected either on the same field trial or on other trials [15, 34]. An SNP was considered as co-localizing with a QTL if it fell in an interval of 50 kb downstream or upstream respect to an SNP belonging to the QTL itself. Second, all the SNPs in complete LD with co-localizing SNPs were identified and subsequently used to identify the putative candidate genes for metabolite biosynthesis. Eventually, using the previously defined associations among co-inherited SNPs and reference SNPs, reference SNPs co-localizing with QTLs were found. These reference SNPs were then used to determine which LC-MS features could be considered as co-localizing with QTLs, and which therefore had to be annotated.

Annotation of LC-MS features co-localizing with known QTLs

None of the LC-MS features co-localizing with known QTLs possessed an annotation based on previous works (see previous paragraph). Therefore, the tentative annotation of these features was based on their raw chemical formulas and on the comparison with MS-related information available from KNApSACk (<http://www.cheminfo.org/Chemistry/Database/Knapsack/index.html>) and Dictionary of natural products (<https://dnp.chemnetbase.com>). This resulted in the putative annotation of 30 other compounds belonging to 11 compound families (Table 1).

Identification of the most likely candidate genes

The putative candidate genes previously identified through GWAS and co-localization analysis were further characterized in order to identify the most likely

candidates for the associations with putative metabolites. This was carried out in a functional perspective and following a two-step procedure. The first step focused on the annotated metabolites and consisted in a systematic bibliographic search through the NCBI PubMed database (<https://pubmed.ncbi.nlm.nih.gov>). The names of the metabolites and of the corresponding biochemical families were used as inputs.

The second step focused on the genes, and consisted in (i) retrieving the gene descriptions available for each gene using a gft file corresponding to the XRQ v2.1 of the sunflower genome; (ii) carrying out tBLASTn searches through the NCBI BLAST website (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>); (iii) performing a systematic bibliographic search through the NCBI PubMed database, using both the gene descriptions and the gene names obtained with the tBLASTn search as inputs.

Genotypic boxplots were produced for all the reference SNPs associated to the tentatively annotated LC-MS features using the ‘genotypes.boxplot’ function of the ‘mlmm.gwas’ R package v 1.0.6. Eventually, we characterized all the co-inherited SNPs found in exons by determining if they represented synonymous or missense mutations using the information available for the XRQ v1.0 sunflower genome (<http://www.heliogene.org/HanXRQ-SUNRISE/>).

Abbreviations

ROS	Reactive oxygen species
LC-MS	Liquid chromatography coupled to mass spectrometry
RT	Retention time
<i>m/z</i>	Mass over charge ratio
SNP	Single nucleotide polymorphism
GWAS	Genome-wide association study
PCA	Principal component analysis
QTL	Quantitative trait locus
LD	Linkage disequilibrium

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12864-024-10104-9>.

- Supplementary Material 1.
- Supplementary Material 2.
- Supplementary Material 3.
- Supplementary Material 4.
- Supplementary Material 5.
- Supplementary Material 6.
- Supplementary Material 7.
- Supplementary Material 8.
- Supplementary Material 9.
- Supplementary Material 10.
- Supplementary Material 11.
- Supplementary Material 12.

Acknowledgments

We would like to thank the RAGT group and Innolea for the help provided in setting up the experimental field and in collecting the plant material used for our study.

Authors' contributions

NBL, AM and YG conceived and designed the experiment. NB and NBL participated in the sampling. HD supervised and coordinated the production of data. SB, TB and OF performed the LC-MS analyses. MM performed the GWAS and all of the other statistical analyses. MM and NBL interpreted the results and wrote the original draft. AM and SB critically read and edited the manuscript. All the authors reviewed and accepted the manuscript.

Funding

This work was supported by the French National Research Agency (SUNRISE ANR-11-BTBR-0005, PHENOME ANR-11-INBS-0012 and MetaboHUB ANR-11-INBS-0010 projects). This research was part of the French Laboratory of Excellence project "TULIP" (ANR-10-LABX-41; ANR-11-IDEX-0002-02).

Availability of data and materials

The genotype and the kinship matrices needed to perform GWAS and the information concerning previously known QTLs are found in the corresponding cited references. All the genomic information concerning candidate genes are available on the Heliagene portal (<http://www.heliagene.org>). LC-MS data are reported in Tables S01-S02. The results of all the other analyses are reported in Tables S03-S10 and in Figs. S01-S02.

Declarations

Ethics approval and consent to participate

All the methods used in this study on field-grown sunflowers complied with national regulations.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Author details

¹UMR LIPME, INRAE, CNRS, Université de Toulouse, 31326 Castanet Tolosan, France. ²UMR BFP, INRAE, Université de Bordeaux, 33140 Villenave d'Ornon, France. ³Bordeaux Metabolome, MetaboHUB, PHENOME-EMPHASIS, 33140 Villenave d'Ornon, France. ⁴UMR BioForA, INRAE, ONF, Orléans 45075, France. ⁵UMR MYCSA, INRAE, 33140 Villenave d'Ornon, France. ⁶USC RIBP, INRAE, Université de Reims, 51100 Reims, France.

Received: 6 July 2023 Accepted: 9 February 2024

Published online: 20 February 2024

References

- Cramer GR, Urano K, Delrot S, Pezzotti M, Shinozaki K. Effects of abiotic stress on plants: a systems biology perspective. *BMC Plant Biol.* 2011;11:1–14.
- He M, He CQ, Ding NZ. Abiotic stresses: general defenses of land plants and chances for engineering multistress tolerance. *Front Plant Sci.* 2018;9:1771.
- Petrov V, Hille J, Mueller-Roeber B, Gechev TS. ROS-mediated abiotic stress-induced programmed cell death in plants. *Front Plant Sci.* 2015;6:69.
- Das K, Roychoudhury A. Reactive oxygen species (ROS) and response of antioxidants as ROS-scavengers during environmental stress in plants. *Front Environ Sci.* 2014;2:53.
- Miller G, Suzuki N, Ciftci-Yilmaz S, Mittler R. Reactive oxygen species homeostasis and signalling during drought and salinity stresses. *Plant Cell Environ.* 2010;33(4):453–67.
- Michalak A. Heavy metals toxicity phenolic compounds and their antioxidant activity in plants growing under heavy metal stress. *Pol J Environ Stud.* 2006;15:523–30.
- Graßmann J. Terpenoids as plant antioxidants. *Vitam Horm.* 2005;72:505–35.
- Munné-Bosch S, Alegre L. Changes in carotenoids, tocopherols and diterpenes during drought and recovery, and the biological significance of chlorophyll loss in *Rosmarinus officinalis* plants. *Planta.* 2000;210(6):925–31 Available from: <https://pubmed.ncbi.nlm.nih.gov/10872224/>.
- Munné-Bosch S, Mueller M, Schwarz K, Alegre L. Diterpenes and antioxidative protection in drought-stressed *Salvia officinalis* plants. *J Plant Physiol.* 2001. Available from: <http://www.urbanfischer.de/journals/jpp;158:1431–7>.
- Zhao M, Zhang N, Gao T, Jin J, Jing T, Wang J, Wu Y, Wan X, Schwab W, Song C. Sesquiterpene glucosylation mediated by glucosyltransferase UGT91Q2 is involved in the modulation of cold stress tolerance in tea plants. *New Phytol.* 2020;226(2):362–72.
- Nakabayashi R, Saito K. Integrated metabolomics for abiotic stress responses in plants. *Curr Opin Plant Biol.* 2015;24:10–6.
- Debaeke P, Casadebaig P, Flenet F, Langlade N. Sunflower crop and climate change: vulnerability, adaptation, and mitigation potential from case-studies in Europe. *OCL.* 2017;24(1):15.
- Debaeke P, Casadebaig P, Langlade NB. New challenges for sunflower ideotyping in changing environments and more ecological cropping systems. *OCL.* 2021;28:29.
- Allinne C, Ghoribi N, Maury P, Maougal R, Sarrafi A, Ykhlef N, et al. Crop production-physiology early sowing as a means of drought escape in sunflower: effects on vegetative and reproductive stages. 2008.
- Mangin B, Casadebaig P, Cadic E, Blanchet N, Boniface MC, Carrère S, Gouzy J, Legrand L, Mayjonade B, Pouilly N, André T. Genetic control of plasticity of oil yield for combined abiotic stresses using a joint approach of crop modelling and genome-wide association. *Plant Cell Environ.* 2017;40(10):2276–91.
- Berton T, Bernillon S, Fernandez O, Duruflé H, Flandin A, Cassan C, Langlade NB, Gibon Y, Moing A. Leaf metabolomic data of eight sunflower lines and their sixteen hybrids under water deficit. *OCL.* 2021;28(42):42. <https://doi.org/10.15454/1.5572412770331912E12>.
- Balliau T, Duruflé H, Blanchet N, Blein-Nicolas M, Langlade NB, Zivy M. Proteomic data from leaves of twenty-four sunflower genotypes under water deficit. *OCL.* 2021;28:2020–6.
- Fernandez O, Urrutia M, Berton T, Bernillon S, Deborde C, Jacob D, Moing A. Metabolomic characterization of sunflower leaf allows discriminating genotype groups or stress levels with a minimal set of metabolomic markers. *Metabolomics.* 2019;15(4):1–14.
- Moschen S, Di Rienzo JA, Higgins J, Tohge T, Watanabe M, González S, Rivarola M, García-García F, Dopazo J, Hopp HE, Hoefgen R. Integration of transcriptomic and metabolic data reveals hub transcription factors involved in drought stress response in sunflower (*Helianthus annuus* L.). *Plant Mol Biol.* 2017;94(4–5):549–64.
- Wu Y, Wang Y, Shi H, Hu H, Yi L, Hou J. Time-course transcriptome and WGCNA analysis revealed the drought response mechanism of two sunflower inbred lines. *PLoS ONE.* 2022;17:e0265447.
- Liang C, Wang W, Wang J, Ma J, Li C, Zhou F, Zhang S, Yu Y, Zhang L, Li W, Huang X. Identification of differentially expressed genes in sunflower (*Helianthus annuus*) leaves and roots under drought stress by RNA sequencing. *Bot Stud.* 2017;58(1):1–11.
- Wu Y, Shi H, Yu H, Ma Y, Hu H, Han Z, et al. Combined GWAS and transcriptome analyses provide new insights into the response mechanisms of sunflower against drought stress. *Front Plant Sci.* 2022;13:847435.
- Rengel D, Arribat S, Maury P, Martin-Magniette ML, Hourlier T, Laporte M, et al. A gene-phenotype network based on genetic variability for drought responses reveals key physiological processes in controlled and natural environments. *PLoS ONE.* 2012;7(10):e45249.
- Barnhart MH, Masalia RR, Mosley LJ, Burke JM. Phenotypic and transcriptomic responses of cultivated sunflower seedlings (*Helianthus annuus* L.) to four abiotic stresses. *PLoS ONE.* 2022;17:e0275462.
- Temme AA, Kerr KL, Masalia RR, Burke JM, Donovan LA. Key traits and genes associate with salinity tolerance independent from vigor in cultivated sunflower. *Plant Physiol.* 2020;184(2):865–80.

26. Chernova AI, Gubaev RF, Singh A, Sherbina K, Goryunova SV, Martynova EU, et al. Genotyping and lipid profiling of 601 cultivated sunflower lines reveals novel genetic determinants of oil fatty acid content. *BMC Genomics*. 2021;22(1):1–15.
27. Stelzner J, Roemhild R, Garibay-Hernández A, Harbaum-Piayda B, Mock HP, Bilger W. Hydroxycinnamic acids in sunflower leaves serve as UV-A screening pigments. *Photochem Photobiol Sci*. 2019;18(7):1649–59.
28. Hu C, Shi J, Quan S, Cui B, Kleessen S, Nikoloski Z, et al. Metabolic variation between japonica and indica rice cultivars as revealed by non-targeted metabolomics. *Sci Rep*. 2014;4:5067.
29. Chan EKF, Rowe HC, Hansen BG, Kliebenstein DJ. The complex genetic architecture of the metabolome. *PLoS Genet*. 2010;6(11):e1001198.
30. Chen W, Gao Y, Xie W, Gong L, Lu K, Wang W, et al. Genome-wide association analyses provide genetic and biochemical insights into natural variation in rice metabolism. *Nat Genet*. 2014;46(7):714–21.
31. Piasecka A, Sawikowska A, Kuczyńska A, Ogródowicz P, Mikołajczak K, Krystkowiak K, et al. Drought-related secondary metabolites of barley (*Hordeum vulgare* L.) leaves and their metabolomic quantitative trait loci. *Plant J*. 2017;89(5):898–913.
32. Wahyuni Y, Stahl-Hermes V, Ballester AR, de Vos RCH, Voorrips RE, Maharijaya A, et al. Genetic mapping of semi-polar metabolites in pepper fruits (*Capsicum* sp.): towards unravelling the molecular regulation of flavonoid quantitative trait loci. *Mol Breed*. 2014;33(3):503–18.
33. Nützmann HW, Huang A, Osbourn A. Plant metabolic clusters – from genetics to genomics. *New Phytol*. 2016;211(3):771–89.
34. Gosseau F, Blanchet N, Varès D, Burger P, Campergue D, Colombet C, et al. Heliaphen, an outdoor high-throughput phenotyping platform for genetic studies and crop modeling. *Front Plant Sci*. 2019;9:1908.
35. Agati G, Azzarello E, Pollastri S, Tattini M. Flavonoids as antioxidants in plants: location and functional significance. *Plant Science*. 2012;196:67–76.
36. Nakabayashi R, Yonekura-Sakakibara K, Urano K, Suzuki M, Yamada Y, Nishizawa T, et al. Enhancement of oxidative and drought tolerance in *Arabidopsis* by overaccumulation of antioxidant flavonoids. *Plant J*. 2014;77(3):367–79.
37. Schulz E, Tohge T, Zuther E, Fernie AR, Hincha DK. Flavonoids are determinants of freezing tolerance and cold acclimation in *Arabidopsis thaliana*. *Sci Rep*. 2016;23:6.
38. Lillo C, Lea US, Ruoff P. Nutrient depletion as a key factor for manipulating gene expression and product formation in different branches of the flavonoid pathway. *Plant Cell Environ*. 2008;31:587–601.
39. Yadav B, Jogawat A, Rahman MS, Narayan OP. Secondary metabolites in the drought stress tolerance of crop plants: a review. *Gene Rep*. 2021;23:101040.
40. Lee S, Oh DG, Singh D, Lee JS, Lee S, Lee CH. Exploring the metabolomic diversity of plant species across spatial (leaf and stem) components and phylogenetic groups. *BMC Plant Biol*. 2020;20(1):1–10.
41. Tamagnone L, Merida A, Stacey N, Plaskitt K, Parr A, Chang CF, et al. Inhibition of Phenolic Acid Metabolism Results in Precocious Cell Death and Altered Cell Morphology in Leaves of Transgenic Tobacco Plants. *The Plant Cell*. 1998. www.plantcell.org;10:1801–16.
42. Vanholme R, De Meester B, Ralph J, Boerjan W. Lignin biosynthesis and its integration into metabolism. *Curr Opin Biotechnol*. 2019;56:230–9.
43. Le Gall H, Philippe F, Domon JM, Gillet F, Pelloux J, Rayon C. Cell wall metabolism in response to abiotic stress. Vol. 4, *Plants*; 2015. p. 112–166.
44. Konovalov DA. Polyacetylene compounds of plants of the Asteraceae Family (review). Vol. 48, *Pharm Chem J*; 2014. p. 613–631.
45. Minto RE, Blacklock BJ. Biosynthesis and function of polyacetylenes and allied natural products. *Prog Lipid Res*. 2008;47:233–306.
46. Champagne D, Arnason J, Philogène B, Morand P, Lam J. Light-mediated allelochemical effects of naturally occurring polyacetylenes and thiophenes from asteraceae on herbivorous insects. *J Chem Ecol*. 1986;12(4):835–57.
47. Blokhina O, Virolainen E, Fagerstedt KV. Antioxidants, oxidative damage and oxygen deprivation stress: a review. *Ann Bot*. 2003;91:179–94.
48. Janeczko A, Skoczowski A, Janeczko A. Mammalian sex hormones in plants. *Folia Histochem Cytobiol*. 2005;43:71–9.
49. Tarkowská D. Plants are capable of synthesizing animal steroid hormones. *Molecules*. 2019;24:2585.
50. Ackah M, Shi Y, Wu M, Wang L, Guo P, Guo L, et al. Metabolomics response to drought stress in *Morus alba* l. variety yu-711. *Plants*. 2021;10(8):1636.
51. Ferraz ABF, Bordignon SAL, Staats C, Schripsema J, Lino von Poser G. Benzopyrans from *Hypericum polyanthemum*. *Phytochemistry*. 2001;57(8):1227–30.
52. Proksch P, Rodriguez E. Chromenes and benzofurans of the Asteraceae, their chemistry and biological significance. *Phytochemistry*. 1983;22(11):2335–48.
53. Fernandez X, Lizzani-Cuvelier L, Loiseau AM, Perichet C, Delbecq C, Arnaudo JF. Chemical composition of the essential oils from Turkish and Honduras styrax. *Flavour Fragr J*. 2005;20(1):70–3.
54. Yahya M, Rachmany D, Shaltiel-Harpaz L, Nawade B, Sadeh A, Ibdah M, et al. A *Pyrus communis* gene for p-hydroxystyrene biosynthesis, has a role in defense against the pear psylla *Cacopsylla biden*. *Phytochemistry*. 2019;161:107–16.
55. Tienda-Parrilla M, López-Hidalgo C, Guerrero-Sanchez VM, Infantes-González Á, Valderrama-Fernández R, Castillejo MÁ, et al. Untargeted MS-based metabolomics analysis of the responses to drought stress in *Quercus ilex* L. leaf seedlings and the identification of putative compounds related to tolerance. *Forests*. 2022;13(4):551.
56. Sun S, Fang J, Lin M, Hu C, Qi X, Chen J, et al. Comparative Metabolomic and transcriptomic studies reveal key metabolism pathways contributing to freezing tolerance under cold stress in kiwifruit. *Front Plant Sci*. 2021;12:628969.
57. Li D, Heiling S, Baldwin IT, Gaquerel E. Illuminating a plant's tissue-specific metabolic diversity using computational metabolomics and information theory. *Proc Natl Acad Sci USA*. 2016;113(47):E7610–8.
58. Das A, Begum K, Akhtar S, Ahmed R, Tamuli P, Kulkarni R, et al. Genome-wide investigation of cytochrome P450 superfamily of *Aquilaria agallocha*: association with terpenoids and phenylpropanoids biosynthesis. *Int J Biol Macromol*. 2023;234:123758.
59. Ayabe SI, Akashi T. Cytochrome P450s in flavonoid metabolism. *Phytochem Rev*. 2006;5:271–82.
60. Werck-Reichhart D, Bak S, Paquette S. Cytochromes P450. *Arabidopsis Book*. 2002;1:e0028.
61. Isin EM, Guengerich FP. Complex reactions catalyzed by cytochrome P450 enzymes. *Biochim Biophys Acta Gen Subj*. 2007;1770:314–29.
62. Wang H, Wang Q, Liu Y, Liao X, Chu H, Chang H, et al. PCPD: plant cytochrome P450 database and web-based tools for structural construction and ligand docking. *Synth Syst Biotechnol*. 2021;6(2):102–9.
63. Seitz C, Eder C, Deiml B, Kellner S, Martens S, Forkmann G. Cloning, functional identification and sequence analysis of flavonoid 3'-hydroxylase and flavonoid 3',5'-hydroxylase cDNAs reveals independent evolution of flavonoid 3',5'-hydroxylase in the Asteraceae family. *Plant Mol Biol*. 2006;61(3):365–81.
64. Yonekura-Sakakibara K, Hanada K. An evolutionary view of functional diversity in family 1 glycosyltransferases. *Plant J*. 2011;66(1):182–93.
65. Ross J, Li Y, Lim EK, Bowles DJ. Higher plant glycosyltransferases. *Genome Biol*. 2001;2:1–6.
66. Bowles D, Lim EK, Poppenberger B, Vaistij FE. Glycosyltransferases of lipophilic small molecules. *Annu Rev Plant Biol*. 2006;57:567–97.
67. Li P, Li Y, Zhang FJ, Zhang GZ, Jiang XY, Yu HM, et al. The *Arabidopsis* UDP-glycosyltransferases UGT79B2 and UGT79B3, contribute to cold, salt and drought stress tolerance via modulating anthocyanin accumulation. *Plant J*. 2017;89(1):85–103.
68. Zhao M, Jin J, Gao T, Zhang N, Jing T, Wang J, et al. Glycosyltransferase CsUGT78A14 regulates Flavonols accumulation and reactive oxygen species scavenging in response to cold stress in *Camellia sinensis*. *Front Plant Sci*. 2019;27:10.
69. Li X, Xu Y, Shen S, Yin X, Klee H, Zhang B, et al. Transcription factor Cit-ERF71 activates the terpene synthase gene CitTPS16 involved in the synthesis of e-geraniol in sweet orange fruit. *J Exp Bot*. 2017;68(17):4929–38.
70. Wang M, Gao M, Zhao Y, Chen Y, Wu L, Yin H, et al. LcERF19, an AP2/ERF transcription factor from *Litsea cubeba*, positively regulates geraniol and neral biosynthesis. *Hortic Res*. 2022;9:uhac093.
71. Wu S, Tohge T, Cuadros-Inostroza Á, Tong H, Tenenboim H, Kooke R, et al. Mapping the *Arabidopsis* metabolic landscape by untargeted metabolomics at different environmental conditions. *Mol Plant*. 2018;11(1):118–34.
72. Zhang F, Wu J, Sade N, Wu S, Egbaria A, Fernie AR, et al. Genomic basis underlying the metabolome-mediated drought adaptation of maize. *Genome Biol*. 2021;22(1):1–26.

73. Mangin B, Bonnafous F, Blanchet N, Boniface MC, Bret-Mestries E, Carrère S, et al. Genomic prediction of sunflower hybrids oil content. *Front Plant Sci.* 2017;8:1633.
74. Badouin H, Gouzy J, Grassa CJ, Murat F, Staton SE, Cottret L, et al. The sunflower genome provides insights into oil metabolism, flowering and Asterid evolution. *Nature.* 2017;546(7656):148–52.
75. Penouilh-Suzette C, Pomies L, Duruflé H, Blanchet N, Bonnafous F, Dinis R, et al. RNA expression dataset of 384 sunflower hybrids in field condition. *OCL.* 2020;27:36.
76. Smith CA, Want EJ, O'Maille G, Abagyan R, Siuzdak G. XCMS: processing mass spectrometry data for metabolite profiling using nonlinear peak alignment, matching, and identification. *Anal Chem.* 2006;78(3):779–87.
77. Shen X, Gong X, Cai Y, Guo Y, Tu J, Li H, et al. Normalization and integration of large-scale metabolomics data using support vector regression. *Metabolomics.* 2016;12(5):1–12.
78. Butler D, Cullis B, Gilmour AR, Gogel BJ. ASReml-R reference manual, release 3.0. Australia: Queensland Department of Primary Industries; 2009.
79. Kachman M, Habra H, Duren W, Wigginton J, Sajjakulnukit P, Michailidis G, et al. Deep annotation of untargeted LC-MS metabolomics data with Binner. *Bioinformatics.* 2020;36(6):1801–6.
80. Sumner LW, Amberg A, Barrett D, Beale MH, Beger R, Daykin CA, et al. Proposed minimum reporting standards for chemical analysis: chemical analysis working group (CAWG) metabolomics standards initiative (MSI). *Metabolomics.* 2007;3(3):211–21.
81. Rohart F, Gautier B, Singh A, Lê Cao KA. mixOmics: an R package for 'omics feature selection and multiple data integration. *PLoS Comput Biol.* 2017;13(11):e1005752. <https://doi.org/10.1371/journal.pcbi.1005752>.
82. Segura V, Vilhjálmsson BJ, Platt A, Korte A, Seren Ü, Long Q, et al. An efficient multi-locus mixed-model approach for genome-wide association studies in structured populations. *Nat Genet.* 2012;44(7):825–30.
83. Bonnafous F, Fievet G, Blanchet N, Boniface MC, Carrère S, Gouzy J, et al. Comparison of GWAS models to identify non-additive genetic control of flowering time in sunflower hybrids. *Theor Appl Genet.* 2018;131(2):319–32.
84. Chen J, Chen Z. Extended Bayesian information criteria for model selection with large model spaces. *Biometrika.* 2008;95(3):759–71.
85. Endelman JB. Ridge regression and other kernels for genomic selection with R package rrBLUP. *Plant Genome.* 2011;4(3):250–5.
86. Bindea G, Mlecnik B, Hackl H, Charoentong P, Tosolini M, Kirilovsky A, et al. ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology and pathway annotation networks. *Bioinformatics.* 2009;25(8):1091–3.
87. Gu Z, Gu L, Eils R, Schlesner M, Brors B. Circlize implements and enhances circular visualization in R. *Bioinformatics.* 2014;30(19)

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.