



**HAL**  
open science

## From data on gross activity to the characterization of animal behaviour: which metrics for which purposes

Ingrid van Dixhoorn, Lydiane Aubé, Coenraad van Zyl, Rudi de Mol, Joop van der Werf, Romain Lardy, Marie-Madeleine Mialon, Kees van Reenen, Isabelle Veissier

### ► To cite this version:

Ingrid van Dixhoorn, Lydiane Aubé, Coenraad van Zyl, Rudi de Mol, Joop van der Werf, et al.. From data on gross activity to the characterization of animal behaviour: which metrics for which purposes. 2024. hal-04469855

**HAL Id: hal-04469855**

**<https://hal.inrae.fr/hal-04469855>**

Preprint submitted on 21 Feb 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1                   **From data on gross activity to the characterization of animal behaviour: which**  
2                   **metrics for which purposes**

3                   December 21, 2023

4  
5 Ingrid van Dixhoorn\*<sup>1</sup>, Lydiane Aubé<sup>2</sup>, Coenraad van Zyl<sup>3</sup>, Rudi de Mol<sup>1</sup>, Joop van der Werf<sup>1</sup>, Romain  
6 Lardy<sup>2</sup>, Marie Madeleine Mialon<sup>2</sup>, C.G. van Reenen<sup>1</sup>, and Isabelle Veissier<sup>2</sup>

7  
8 <sup>1</sup>Wageningen UR, Livestock Research, Wageningen, P.O. Box 338, 6700 AH Wageningen, the Netherlands

9 <sup>2</sup>Université Clermont Auvergne, INRAE, VetAgro Sup, UMR Herbivores, 63122 Saint-Genes-Champanelle,  
10 France

11 <sup>3</sup>Wageningen Adaptation Physiology, Wageningen University & Research, Wageningen, the Netherlands

12  
13 \*Corresponding author: Ingrid D. E. van Dixhoorn, +31 (0) 320 293 514, ingrid.vandixhoorn@wur.nl

14  
15  
16 **Key words:** sensors, time budget, animal welfare, health, activity metrics, cow

17  
18  
19                   **Abstract**

20 The behaviour of an animal is closely linked to its internal state. Various metrics can be calculated from  
21 activity data. Complex patterns of activity within or between individuals, such as cyclic patterns and  
22 synchrony, can inform on the biological functioning, the health status, or the welfare of an animal. These  
23 patterns are now available thanks to sensors that continuously monitor the activity of individual animals  
24 over long periods. Data processing and calculations, however, should be clarified and harmonised across  
25 studies for the results to be comparable. We present metrics describing activity patterns, we discuss their  
26 significance and relevance for behavioural and welfare studies and their limits, and we detail how they  
27 can be calculated. Four groups of metrics are distinguished: metrics related to overall activity (e.g., time  
28 spent in each activity per unit of time), metrics related to fluctuations around mean activity, metrics  
29 related to the cyclicity of activity, and metrics related to the synchrony between animals. Metrics may  
30 take statistical approaches (e.g., average and variance) or modelling approaches (e.g., Fourier Transform).  
31 Examples are taken essentially from cattle for which individual activity sensors are easily available at  
32 present. The calculations, however, can be applied to other species and can be performed on data  
33 obtained from sensors as well as visual observations. The present methodological article will help  
34 researchers to obtain the most benefit from activity data and will support the decision of which metric  
35 can be used to address a given purpose.

36

38 The behaviour of an animal can inform about the internal state of that animal, in relation to biological  
39 functioning, health and welfare. Activity measurements have long been used to identify differences in  
40 walking behaviour to detect lameness (Pastell et al., 2009) or differences in the amount of activity to detect  
41 oestrus (Saint-Dizier & Chastant-Maillard, 2012) or periparturient disorders (Rutten et al., 2017; Rutten et  
42 al., 2013; Weary et al., 2009). Additionally, comparing the activity of a focus animal to its baseline or to pen-  
43 mates allows identification of deviations that potentially indicate a change in internal state, e.g., an animal  
44 becoming ill (Kok et al., 2023). The continuous measurements of activity allow for complex patterns to be  
45 highlighted, such as circadian components and regularity, and these patterns can be used to identify  
46 animals at risk of diseases (van Dixhoorn et al., 2023; Van Dixhoorn et al., 2018; Wagner et al., 2021).

47 Ethologists, and other researchers usually calculate several metrics from activity data to characterise an  
48 animal's behaviour, e.g., time spent on each activity, fragmentation of activity, 24-h patterns or proportion  
49 of animals engaged in the same activity. However, the way these metrics are calculated vary between  
50 studies. For instance, the synchrony between animals, that reflects the functioning of a group, can be  
51 calculated at individual level as the percentage of animals performing the same activity as the focus animal,  
52 then at group level as a mean of that percentage or using concordance indices such as Kappa coefficients  
53 or overdispersion index (Raussi et al., 2011; Tuomisto et al., 2019; Veissier et al., 1990). The pros and cons  
54 of each metric are rarely explained.


55 The activities can be documented by observers, from direct observation or from video recordings, or  
56 obtained using sensors. Direct observation allows precise activities to be recorded. For instance, lying can  
57 be divided into lying head down versus head up with corresponding arousal levels inferred (Veissier et al.,  
58 2001). Direct observation (both real time or from video recordings) requires a high workload and data at  
59 night are often missing due to difficulties in observing animals in the dark. The last ten years have seen a  
60 boom in the development of sensor technologies, which can provide data along time series more easily  
61 than direct observation. Activities (especially in large domestic animals like cattle) can now be recorded  
62 continuously on individual animals and for very long periods with little workload, using accelerometers,  
63 image analysis from videos, or Real Time Locating System (RTLS) (Buller et al., 2020). Most of the  
64 commercially available sensors that monitor cattle activity show excellent performance in validation studies  
65 (Lee & Seo, 2021). They usually provide information on gross activities such as lying, standing, moving,  
66 feeding, and ruminating or the position of animals in the barn. From the organisation of these activities,  
67 specific patterns can be detected, especially those indicative of animal malaise due to illness or stress  
68 (Wagner et al., 2021) or related to the social organization of animal groups (Rocha et al., 2020), allowing  
69 new insight into animal behaviour. However, to date the flood of data obtained from sensors seem under-  
70 utilised (Koltes et al., 2019).

71 Access to data on activity is not only facilitated for ethologists, but also for researchers from other  
72 discipline, e.g., animal health and animal production and for non-scientists like users of precision farming  
73 systems that are not necessarily used to process data on behaviour (Borchers et al., 2016). It is thus crucial  
74 to provide harmonised metrics to analyse behaviour and to discuss what they are supposed to measure and  
75 their limits. This would allow comparisons between studies, help the interpretation of results, extend the  
76 use of activity data beyond ethologists, and ease the re-use of datasets.

77 In the present paper, we consider the metrics that describe different aspects of behaviour from data  
78 collected by sensors or by direct observations (Figure 1). We consider metrics describing 1- overall activity,  
79 2- fluctuations around mean activity, 3- cyclicity of activity, and 4- synchrony between animals. For each of  
80 these four groups of metrics, we identify the main metrics in use, the calculation methods, the limitations  
81 or the difficulties that can be encountered to calculate them, and their biological meaning (i.e. what it  
82 implies for animal welfare or health or the functioning of social groups). Examples are taken essentially  
83 from cattle, for which individual activity sensors are easily available at present. The calculations, however,  
84 can apply to datasets from other species (including other animal-specific activities) and can be performed  
85 on data coming from sensors as well as from observation. This paper does not focus on the validation of  
86 the sensors or the observers (to assess inter or intra observer variation) but on the processing of the data  
87 acquired by the sensors or observers.

88 We believe such a methodological review is pivotal and timely given the boom of activity data obtained  
89 by sensors. The review will hopefully support researchers by improving the use of activity data to answer

90 their research questions. In addition, it should facilitate the consideration of animal behaviour by non-  
 91 ethologists especially in Precision Livestock Farming (PLF), with a view to help phenotyping animals for  
 92 selection, monitoring them for the detection of changes due to specific states such as oestrus, disease, or  
 93 stress, or evaluating housing conditions and management aspects.



94

95 **Figure 1** - Visualisation of the data acquisition and processing proces. The scope of this paper focusses  
 96 on the possible calculations of the data that describe the four metrics: overall activity, fluctuations  
 97 around the mean activity, cyclicality of activity and synchrony. The calculations can be done on data  
 98 that is acquired by observations and/or with sensors.

99

## 100 The data

101 In the present paper, we focus on gross activities such as feeding behaviour (including eating and  
 102 drinking), active behaviour (apart from eating or drinking) including walking, running or other movements,  
 103 or inactive behaviour including standing still and lying. More specific activities include grazing and  
 104 ruminating in ruminants (e.g., cattle, sheep, goats), rooting in pigs, and foraging behaviour in poultry. All  
 105 these activities are characterised by lasting for some time. They are sometime referred to as 'states', as  
 106 opposed to brief behaviours (e.g., interactions between animals) that are referred to as 'events'.

107 Observers can perform focal (and continuous) sampling i.e., they note changes in activity with reference  
 108 to the time of the change occurrence. Alternatively, observers can perform scan sampling i.e. they note the  
 109 animal's activity as detected at first glance at regular intervals (Bateson, 2021). Examples of the continuous  
 110 and scan sampling data are shown in the supplementary materials. A mix of the two is also possible: the  
 111 activity is recorded continuously and at the end of each interval (e.g. 5 min) the observer notes the  
 112 predominant activity; the format of data will then be similar to that of scan sampling. Sensors generally  
 113 produce a signal that is nearly continuous; the data on gross activity is usually delivered as time spent in  
 114 each activity per time intervals (e.g. minutes per hour or per 15 min) or as predominant behaviour per time  
 115 interval. The metrics that can be calculated, depend on the formats of data (see next sections).

116

117

## Metrics to address overall activity

### 118 Definitions

119 The overall activity refers to the time spent performing specific activities during a certain time period.  
 120 Each activity such as feeding, drinking, walking, standing idling or lying, can be characterised by the duration  
 121 it is performed, the number of bouts (where a bout is defined by the continuous expression of an activity),  
 122 and the average duration of bouts. The overall fractioning of activity refers to how many bouts of activity  
 123 are noticed, in other words how often the animal changes of activity. The level of activity reflects how much  
 124 an animal is active, that is walking, running, or feeding rather than lying or standing immobile. The time  
 125 period on which these metrics are calculated varies between studies, e.g. an hour or a day. The term 'time  
 126 budget' is specifically used to describe how an animal divides its day (or shorter period) into the various  
 127 activities.

### 128 Calculations

129 *Proportion of time spent in an activity.*

130 The proportion of time spent in a given activity  $a$  in a collection of activity bouts  $B$  (whatever the  
 131 activity) is calculated as follows:

$$132 \quad \text{Proportion Of Activity}_a^B = \left( \sum_{b \in B, A_b=a} D_b \right) / \sum_{b \in B} D_b \quad (1)$$

133 where  $D_b$  is the duration of the bout  $b$  and  $A_b$  is the activity in bout  $b$ .

134 In case of scan sampling, the time spent in an activity is estimated from the number of scans per  
 135 activity, multiplied by the interval between scans - although what the animal has done between scans  
 136 remains unknown -. The proportion of time spent in an activity  $a$  in a period  $P$  can be calculated as follows:

$$137 \quad \text{Proportion Of Activity}_a^P = \left( \sum_{p \in P, A_p=a} L_p \right) / \sum_{p \in P} L_p \quad (1)$$

138 where  $p$  a subperiod of  $P$ ,  $L_p$  is the length of the subperiod  $p$ , and  $A_p$  is the activity in the subperiod  
 139  $p$ .

140 Duration of activities (or proportion of time spent in activities) can be calculated for each activity  
 141 separately and whatever the time period. If the experimenter decides to group two activities (e.g., lying  
 142 ruminating and standing ruminating), the duration/proportion of the new activity (here ruminating) is  
 143 obtained by summing up those of the individual activities.  
 144

### 145 *Number and duration of activity bouts.*

146 Continuous and scan sampling also allow detecting when the activity changes, so that the number  
 147 of activity bouts can be obtained, and the mean duration of bouts can be calculated. The average bout  
 148 duration of activity  $a$  in bouts collection  $B$  is calculated as:

$$149 \quad \text{AvgBoutDuration}_a^B = \left( \sum_{b \in B, A_b=a} D_b \right) / N_B(a) \quad (2)$$

150 where  $D_b$  is the duration of the bout  $b$  and  $N_B(a)$  the number of bouts of the collection  $B$  where the  
 151 activity equals  $a$ .

152 Sensor-based systems, however, sometimes provide the time spent in each activity per time period  
 153 and not the exact timing of a change if any; in this case, the number and the duration of bouts of activity  
 154 remain unknown.  
 155

156 Calculation of the number of bouts of an activity and of the mean duration of the bouts requires  
 157 that the recording is done on long periods to avoid edge effects. Indeed, when the monitoring starts, the

158 animal is observed in a given activity, but one does not know for how long the animal has been performing  
159 the activity. It is common practice to remove the first and the last activity bout observed during the time  
160 period studied. The time period must thus be long enough so that several entire bouts of activity can be  
161 recorded. In practice, the number of bouts and their mean duration are often calculated per day (see for  
162 instance Veissier et al., (2004)).

163 Assessing the total number of activity bouts in a day, in other words the number of times an animal  
164 changes of activity, requires that the ethogram consists of activities described with the same level of details,  
165 so that the number of bouts does not depend on what activity an animal performs most during the day.  
166 The gross activities detected by sensors are usually adequate. This is not necessarily the case with direct  
167 observations. For instance, to answer a specific question, an experimenter may want to sort activities into  
168 lying, standing inactive, feeding (all of these lasting for minutes or hours) and other activities, that can  
169 consist of walking, running, exploring the environment, interacting with other animals or self-directed  
170 activities (all of them lasting for few seconds or minutes). In that case, the number of times an animal  
171 changes of activity will largely depend on whether the animal performs the short-lasting activities  
172 frequently. The grouping of activities into gross activities should be done before the total number of activity  
173 bouts is calculated. Indeed, in the example given above on ruminating while lying or standing, the number  
174 of ruminating bouts during a day cannot be calculated by adding the number of lying-ruminating bouts and  
175 that of standing-ruminating bouts, because the two activities can be performed in the same bout.

176 Duration and number of bouts also largely depend on how bouts are defined. The most common  
177 practice is to consider each change of activity as the beginning of a new bout. For example, with continuous  
178 observations the number of lying bouts of a cow can be assessed by the number of times the animal lays  
179 down; with scan sampling, an eating bout can be defined when eating is observed on at least one scan  
180 (Tucker et al., 2009). When scan sampling is used, the interval between scans should be smaller than the  
181 duration of activities to not miss bouts. An animal switching from an activity to another can still be  
182 considered in the same bout if it returns quickly to the initial activity. One needs to define how long the  
183 animal must stop an activity between two instances of that activity so that separate bouts are identified  
184 (Yeates et al., 2001).

185 For instance, eating bouts are usually combined into meals if the interval between successive eating  
186 bouts is less than the meal criterion. The minimum interval between bouts can be determined by different  
187 methods (Tolkamp et al., 1998, Yeates et al., 2001, Dado & Allen, 1993), for example, using log-survivorship  
188 and log-frequency analysis (see Tolkamp et al., (1998), for a description of these methods).  
189

#### 190 *Activity level.*

191 The overall activity of an animal can be summarised into an activity level by assigning a weight to  
192 each activity, the weight expressing the contribution of the activity to the arousal of the animal (Veissier et  
193 al., 2001). The level of activity of the period  $P$  is calculated by the sum of the time spent in each activity  
194 multiplied by the weight of the activity:

$$195 \quad \text{ActivityLevel}_a^P = \left( \sum_{\substack{p \in P \\ a \in A}} T_{ap} \cdot W_a \right) \quad (3)$$

196 where  $T_{ap}$  is the time spend on activity  $a$  in period  $p$ ,  $W_a$  is the weight of activity  $a$ , the summations  
197 are over subperiods  $p$  in period  $P$  (and over all activities in  $A$ ).

198 The weights can be assigned *a priori* by the experimenter or elicited from observations. Veissier et al. (2001)  
199 performed a Factorial Analysis of Correspondence (**FAC**) on the number of instances (scans x calves) of each  
200 of five activities per hour; the first axis - that summarises most of the variations between the 24 hours of  
201 the day - brought decreasing weights to feeding (1.438), walking (0.763), standing immobile (- 0.085), lying  
202 head up (- 0.261), then lying head down (- 0.541), ordering the activities as one would intuitively do to  
203 express the decreasing arousal. The FAC is based on associations between activities and therefore, the  
204 outcome of the FAC strongly depends on the level of detail of the activities that are included. Experience  
205 told us that the ethogram should not be split in too many (short lasting) activities to elicit meaningful  
206 weights.

207 **Biological meaning**

208 The time budget of animals depends on their living conditions and the farm management. Cows  
209 spend about half of the day lying but this can vary from 8 h to 13 h (reviewed by Tucker et al., 2021). Lying  
210 time is reduced in case of uncomfortable lying areas (poorly designed or too hard, wet, small, hot) or not  
211 enough resting places for the size of the herd (Tucker et al., 2021). Cow may nevertheless spend more time  
212 lying in case of short cubicles preventing them to stand properly in a cubicle and thus forcing them to lie  
213 down as soon as they enter a cubicle (Veissier et al., 2004). The time spent feeding and walking also largely  
214 depends on housing and management conditions: cows grazing at pasture spend much more time eating  
215 and walking than cows kept indoors and fed herbage harvested from the same pasture (527 min/d eating  
216 and 311 min/d walking at pasture vs. 398 min/d and 133 min/d indoor (Dohme-Meier et al., 2014)); bulls  
217 and sheep spend less time eating when the diets contain a large proportion of fibre than when the diets  
218 contain a large proportion of starch (- 67% in bull fed a 45% starch diet and - 18% in sheep fed a 38% starch  
219 diet, compared to animals fed diets with less than 20% starch (Commun et al., 2012; Mialon et al., 2008)).  
220 The effects described in the above paragraph are typically observed on all animals from a herd.

221 Modifications are also observed between individuals in a herd. The overall activity of an animal  
222 varies over time due to its physiological state. Young cows are often more active and change more often of  
223 activity than adult ones (Solano et al., 2016). At the time of oestrus, cows are agitated, spending less time  
224 eating but more time active in other ways (more walking, less lying) (Reith & Hoy, 2018). Changes are also  
225 observed due to gestation and parturition: the time spent lying by cows decreases from one month before  
226 to one month after calving and slowly increases thereafter up to end of lactation, with 2 h of amplitude of  
227 variations in multiparous cows (Hut et al., 2022). The changes in activity are generally well marked and short  
228 lasting around oestrus but less marked and gradual around calving, making calving detection from gross  
229 activity more difficult than that of oestrus (Benaissa et al., 2020).

230 The overall activity of an animal can also change due to a pathological state (e.g. due to  
231 inflammation (Dittrich et al., 2019). These modifications are called sickness behaviour, characterised –  
232 among others – by a low activity of the animal (Weary et al., 2009). Metabolic disorders (e.g., hypocalcemia,  
233 ketosis, acidosis) are generally accompanied by an increase in the time spent lying and a corresponding  
234 decrease of the time spent active and feeding (Weary et al., 2009; Belaid et al. 2021). These modifications  
235 are more marked in hypocalcemia than in other metabolic diseases, hence the name ‘downer cow’  
236 syndrome for hypocalcemia (Wadhwa & Prasad, 2002). Acidosis can be accompanied by a higher fractioning  
237 of activity: sheep suffering from acidosis often change their posture from lying to standing (Commun et al.,  
238 2012), as if they do not feel comfortable in either of these postures. Infectious diseases are also associated  
239 with an increase in time spent lying down and a decrease in time spent feeding (Weary et al., 2009). Mastitis  
240 however may result in a decreased time spent lying down, compensated by an increased time spent  
241 standing (Fogsgaard et al., 2015; Medrano-Galarza et al., 2012), presumably due to the pain on the udder  
242 which is increased by the pressure on it when the animal is lying. Lameness is also accompanied by sickness  
243 behaviour and a specific pattern of lying behaviour with less lying bouts but of longer duration (Solano et  
244 al., 2016, de Mol et al., 2013). Sick cows, whatever the origin of the disorders, usually spend less time  
245 ruminating (Calamari et al., 2014).

246 Stress is another factor affecting the activity of animals. Stressed animals usually spend less time  
247 lying down and change more often of activity. For instance, suckling calves separated from their dam and  
248 moved from pasture to a barn respond to these changes by spending less time lying down, more time  
249 standing still or walking, and by fractionating their activities to a larger extent (Veissier et al., 1989). These  
250 modifications fade within days or weeks, indicating habituation to the new environment. Similar changes  
251 are observed when primiparous cows join the lactating herd. Around calving, primiparous cows decrease  
252 their time spent lying to a larger extent than multiparous cows (Hut et al., 2022) and we suspect that this is  
253 due to the many changes undergone by them around calving: they are milked for the first time and they  
254 are introduced to the lactating herd and so exposed to social partners and a pen, both novel to them.

255 Monitoring the overall activity can thus help to check if the animals are managed adequately and  
256 to detect physiological states (especially oestrus), pathological states, or stress. Feeding, ruminating and  
257 lying seem especially sensitive to variations in the animal or its environment. Lying is generally reduced in  
258 case of stress or an uncomfortable lying area, so it is often considered that a prolonged time spent lying  
259 indicates good welfare (Piñeiro et al., 2019a, 2019b). However, lying time is often increased in case of  
260 disease too. In any case, the value obtained for duration and frequency of activities should be interpreted

261 considering the context in which they are obtained, for instance the type and management of feeding, the  
262 housing or grazing conditions, and the timing in relation to calving.

## 263 Metrics to address fluctuations around the mean activity

### 264 Definition

265 The activity of an animal varies within and between days (Hut et al., 2022). The variations are  
266 described by metrics calculating how far values, obtained on a given time period, are spread around the  
267 mean value across several time periods. The calculations are generally applied to the duration of activities  
268 or the level of activity, less often to the number of bouts or their duration, with all values obtained at animal  
269 level. The time period is often the hour within the day (Mialon et al 2008 (eating duration in bulls), Lardy et  
270 al., 2023 (level of activity in cows)) or the day within a period of at least several days (Hut et al., 2022  
271 (duration of each activity); Solano et al., 2016 (number of bouts)).

### 272 Calculations

273 The metrics used to describe variations of an animal activity across time periods (e.g., hours within  
274 a day) are similar to those traditionally used in descriptive statistics except that they are applied at  
275 individual level (to characterise the variability of the activity of a given animal and not the variability  
276 between animals):

277 *Minimum (Min)* refers to the minimum value observed/recorded

278 *Maximum (Max)* refers to the maximum value observed/recorded

279 *Range* corresponds to the difference between Min and Max

280

### 281 Quantiles

282 Quantiles are cut points dividing the dataset into continuous intervals with equal probabilities. The  
283 most commonly used quantiles are quartiles, which divide the number of data points into four parts, where  
284 the first quartile (Q1, 25<sup>th</sup> percentile) is the maximum value on the 25% of the dataset that obtained the  
285 lowest values, the second quartile (Q2) corresponds to the median, and the third quartile (Q3, 75<sup>th</sup>  
286 percentile) is the minimum value on the 25% of the dataset that obtained the highest values.

287

### 288 Variance

289 The sum of the squares of the differences between each value and the mean (see formula below);  
290 and *Standard deviation (SD)*: the square root of the variance.

291 
$$\text{Variance}(x) = \frac{1}{N-1} \sum_{i=1}^N (x_i - \mu)^2 \quad (4)$$

292 where  $x$  is the sequence of activities (expressed for example as the level of activity) composed of  $N$   
293 observations,  $x_i$  is the  $i^{\text{th}}$  observation, and  $\mu$  the mean of the activities.

294

### 295 Root Mean Square of the Successive Differences (RMSSD):

296 RMSSD measures the variations from one interval to the next one. RMSSD is calculated as follows:

297

298 
$$\text{RMSSD}(x) = \sqrt{\frac{1}{N-1} \sum_{i=1}^{N-1} (x_i - x_{i+1})^2} \quad (5)$$

299 where  $x$  is the sequence of activities (expressed for example as the level of activity) composed of  
300  $N$  observations,  $x_i$  is the  $i^{\text{th}}$  observation.



301 *Irregularity.*  
302 An index of irregularity of behaviour can be calculated as proposed for feed intake (Salgado et al.,  
303 2021a, 2021b). First a regression line of cumulative data (e.g., cumulative time spent in a given activity) is  
304 drawn, then the differences between observed values and the regression line are calculated. The sum of  
305 these differences brings the irregularity index.

### 306 **Biological meaning**

307 Maximum activity during the day can be relevant to identify cow states: in a study where many  
308 metrics were used for a random forest classification, Lardy et al. (2023) found that the maximum activity  
309 during the day and Quantile 90 (two metrics closely linked) were the most important features to  
310 discriminate pathological and physiological states of cows.

311 Within-day variations of activity are well marked in healthy and non-stressed animals. For instance,  
312 lying is predominant at some times during the day whereas eating and other activities are predominant at  
313 other times. Therefore, the activity level is not constant from hour to hour during the day. Within-day  
314 variations (measured by SD and RMSSD) are less marked in sick animals than in healthy ones; this is the  
315 case for cows affected by mastitis and to a lesser extent by lameness (Veissier et al., 2017). Within-day  
316 variations are also affected by oestrus: variations rise above baseline at the beginning of oestrus then  
317 decrease below baseline for at least two days (Veissier et al., 2017). RMSSD slightly differs from SD. With  
318 the same amount of variations during the day (same SD), an activity that varies smoothly between  
319 successive hours results in a low RMSSD while an activity that fluctuates between successive hours results  
320 in a high RMSSD. In Veissier et al. (2017) the decrease in within day variations under mastitis, lameness and  
321 oestrus was more marked when assessed by RMSSD than by SD, suggesting that cows change of activity  
322 more often when diseased or in oestrus. Stress may also be associated with a reduction in within-day  
323 variation in activity, as observed in cattle moved from pasture to indoor housing (Veissier et al., 1989; using  
324 the difference between night and early morning). Spreading activities throughout the day can also be a way  
325 to adapt to specific conditions. Bulls fed fibrous diets eat in few meals during the day whereas bulls fed high  
326 starch diets spread their eating activity over the entire day, which results in a low SD of eating duration  
327 (Mialon et al., 2008). Dispersing small meals over the day seems a strategy to avoid ruminal acidosis due to  
328 high amounts of starch in a diet.

329 Between-day variations of activity can increase when animals are disturbed. For instance, cattle  
330 and sheep affected by acidosis or ketosis have more variable activities across days than healthy animals  
331 (Commun et al., 2012; González et al., 2008). At least in case of ketosis, an effective treatment eliminates  
332 the effect (Goldhawk et al., 2009). Lame cows can also display high between-days variations in the duration  
333 of lying bouts (Ito et al., 2010; Solano et al., 2016). A quick return to normal or baseline values after small  
334 disturbances (micro-recoveries) results in low variance, and is considered as a sign of good resilience  
335 (Scheffer et al., 2018). Animals that spontaneously (i.e., apart from diseases or other challenges) have a  
336 variable activity are less prone to further diseases (Van Dixhoorn et al., 2018).

337

## 338 **Metrics to address the cyclicity of activity**

### 339 **Definition**

340 Cyclicity indicates fluctuations at regular intervals around activity trends. The most common cycle  
341 is the circadian cycle, which is the 24-h rhythm due to the alternance of day and night.

### 342 **Calculations**

343 Several metrics can be used for addressing the cyclicity of activity (Kok et al., 2023). Autocorrelation  
344 and Fourier transform are basic calculations. Other indices are derived from them such as non-periodicity  
345 index (Van Dixhoorn et al., 2023; Van Dixhoorn et al., 2018) and degree of functional coupling (Berger et  
346 al., 2003; Scheibe et al., 1999).

### 347 *Autocorrelation.*

348 The autocorrelation measures the correlation between successive values of a signal. The  
349 autocorrelation depends on the chosen delay, e.g., for hourly data the lag-1 corresponds to the correlation

350 between successive hours and lag-24 corresponds to the correlation between an hour of the day and the  
 351 same hour of the next day, therefore depending on the circadian cycle. For a sequence  $x$  of size  $N$ , the  
 352 autocorrelation with a lag  $l$  is calculated as following:  
 353

$$354 \quad ACF(x, l) = \frac{1}{(N - l)\sigma^2} \sum_{i=1}^{N-l} (x_i - \mu)(x_{i+l} - \mu) \quad (6)$$

355 Where  $\sigma$  is the variance of the sequence  $x$ ,  $\mu$  is the average value of the sequence  $x$  and  $x_i$  the  $i^{th}$   
 356 element of the sequence  $x$ .  
 357

### 358 *Non-periodicity.*

359 Non-periodicity is calculated by plotting the correlogram of the raw data (which is a graphical  
 360 display of a correlation matrix of the data) over a sinusoid with an amplitude of 0.25 and a 24-hour cycle  
 361 and assessing the difference between the correlogram and the sinusoid by calculating the mean squared  
 362 error (Figure 2) (Van Dixhoorn et al., 2023; Van Dixhoorn et al., 2018). The value of an amplitude of 0.25 is  
 363 chosen as it gave the best fit and might be adjusted in other situations where the autocorrelation shows a  
 364 circadian rhythm.

$$365 \quad Nonperiodicity(x) = \frac{(\sum_{l=1...100} (ACF(x, l) - 0.25 \cdot \cos(2\pi \cdot l/24))^2)}{100} \quad (9)$$

366 where  $x$  is a variable measured at hourly (or other chosen time interval) time intervals,  $ACF(x, l)$  is  
 367 the autocorrelation function for variable  $x$  at lag  $l$  (ranging from 1 to 100),  $0.25 \cdot \cos(2\pi \cdot l/24)$  is the cosine  
 368 function with a 24-h cycle and an amplitude of 0.25 that is used as a fit function.  
 369

### 370 *Fourier transform.*

371 Fourier transform represents the sinusoids that compose the original variations. Each sinusoid is  
 372 defined by a frequency and an amplitude. The contribution of each sinusoid to explain the original variations  
 373 is expressed in absolute or relative power, usually referring to the frequency of the sinusoid or to frequency  
 374 bands. For instance, the contribution of the circadian cycle and of ultradian cycles can be calculated. When  
 375 variations within 24 h time series are analysed by Fourier transform, the fundamental (h0) refers to the  
 376 average activity during 24 h, harmonic 1 (h1) refers to variations following a 24 h cycle; h2, to a 12 h cycle;  
 377 h3, to an 8 h cycle; h4 to a 6 h cycle, etc. The main cycle is the circadian one; the activity of an animal can  
 378 therefore be modelled by its overall activity (mean during 24 h) and the variations around overall activity  
 379 following a 24 h cycle, in other words into h0 and h1. The Fourier-based approximation with thresholding  
 380 (FBAT) method was developed to compare such models obtained on successive time series (Wagner et al.,  
 381 2021). An alternative to the Fourier transform is the Cosinor method. Cosinor and Fourier are analogous in  
 382 formulation, but differ in operation (see (Chkeir et al., 2019), for a comparison of the two modelling  
 383 approaches). An example of Fourier transform is visualised in Figure 2.

### 384 *Degree of functional coupling (DFC).*

385 The degree of functional coupling is obtained by calculating autocorrelations, then applying Fourier  
 386 transform to the correlogram, extracting the significant harmonics and calculating the power of each  
 387 significant harmonics out of the power of all significant harmonics (DFC). More specifically, the relative  
 388 power of the harmonic corresponding to a 24 h cycle expresses how much the variations are due to the  
 389 circadian cycle: when DFC equals 100%, the variation in activity follows strictly a circadian cycle vs. when  
 390 DFC equals 0%, the activity not at all depends on the 24 h cycle (Berger et al., 2003).

391 In theory, data on 24 h only can be used to identify a circadian cycle. In practice, activity data usually  
 392 contains noise, i.e., erratic fluctuations so that more than one day is necessary to identify correctly cyclic  
 393 components. The number of days required depends on the amount of noise vs. cyclic components.


394 In general, the activity of an animal peaks several times during the day, e.g., depending on the  
 395 timing of feeding or milking, whereas the activity stays low during whole night. Autocorrelations and Fourier  
 396 transform hardly take the dissymmetry between day and night activity into account. Wavelet methods  
 397 could be used to overcome this problem (discussed in Wagner et al., 2021).

398 **Biological meaning**

399 Daily periodicities are influenced by internal clocks (endogenous driven biological cycles) and  
400 external factors such as temperature, light, humidity, feeding time (exogenously driven biological cycles)  
401 (Saper et al., 2005). Internal clocks generate a rhythm of about 24 h period. In mammals, the central internal  
402 clock situated in the brain (suprachiasmatic nuclei) coordinates peripheral clocks in the body (Honma,  
403 2018). Among external cues (or “Zeitgebers), light is known to be the most powerful one, impacting both  
404 on behaviour and physiological functions (Honma, 2018). For instance, cows are typically diurnal animals:  
405 they eat essentially between dawn and dusk and they predominantly rest at night (DeVries et al., 2003;  
406 Hafez et al., 1969). Variations can nevertheless be observed between cows in the cyclicity of their  
407 behaviour. Competition for resources (e.g. feed, lying area), due to overstocking or ambiguity in the ranking  
408 order because of frequent change in group composition, can cause a misalignment with the circadian  
409 rhythm (McCabe et al., 2021; Van Dixhoorn et al., 2023; Van Dixhoorn et al., 2018; Van Erp et al., 2020).

410 Cows that have more marked circadian patterns of activity seem more resistant to health disorders.  
411 For instance, cows with marked cyclicity of eating, walking or lying before calving are less affected by post-  
412 partum health disorders (including inflammatory and metabolic problems (Van Dixhoorn et al., 2023; Van  
413 Dixhoorn et al., 2018)). Indeed, dairy cows need to have their physiological mechanisms fine-tuned to be  
414 able to produce large quantities of milk while avoiding nutritional and metabolic deficiencies (negative  
415 energy balance or mineral deficiencies such as hypocalcemia). We hypothesise that good cyclicity, aligned  
416 with circadian rhythm can help to avoid such dysfunctions, especially in the high-risk postpartum period.

417 Any change of the internal state of an animal - due to stress, disease or some specific reproductive  
418 status (parturition, oestrus) – can in turn affect the cyclicity of activities. For instance, the difference  
419 between activity during the day and at night is less marked in heifers experiencing a large change in their  
420 environment – weaning and turning from pasture to indoors – and in cows affected by mastitis or lameness  
421 (Veissier et al., 1989; Veissier et al., 2017). Variations during the day can be modelled, e.g., thanks to Fourier  
422 transform in the FBAT method. The distance between models obtained on successive 24 h series increases  
423 when cows are stressed, diseased, in oestrus or about to calve, expressing a change in daily patterns  
424 (Wagner et al., 2021). These effects may be due to the release of glucocorticoids during stress, disease or  
425 even calving. Indeed, glucocorticoids, the secretion of which follows a circadian pattern, help to coordinate  
426 peripheral clocks with the brain pacemaker (Dumbell et al., 2016).



427

428  
 429  
 430  
 431  
 432  
 433  
 434  
 435  
 436

**Figure 2** - Examples of non-periodicity of eating behaviour visualised through correlograms (top), and Fourier transform pattern (bottom) (frequency of activity performed per day). In the correlograms, dotted lines represent the expected cyclicality of the specific behaviour, while the solid lines indicate the observed behaviours. Non-periodicity is assessed by calculating the RMSE of the correlogram as compared to the sinusoid. In the Fourier transform pattern the amplitude is given per frequency, expressing the strength of the cycles in activity for that frequency. The peak at frequency 1 shows that this cow has a strong circadian pattern. The sum of the amplitudes at frequency 1, 2, 3 and 4 represents the strength of the cycles in activities with a 24, 12, 8, and 6 h cycles and is used as a measure of the cyclicity of the cow.

437

## Metrics to address the synchrony between animals

### 438 Definition

439 The synchrony measures the extent to which animals of a given group perform the same activity at  
440 the same time. The synchrony can be assessed between two animals, between an animal and the group it  
441 belongs to, or at the group level.

### 442 Calculations

443 The synchrony is to be assessed based on what animals do at certain points in time spaced at  
444 regular intervals. Collecting data using scan sampling therefore is appropriate for calculations of synchrony  
445 metrics. In case of continuous observations, which is changes in behaviour are noted for each animal exactly  
446 when they occur, a pre-processing of data is necessary: at each time interval the instantaneous activity of  
447 each animal of the group need to be extracted (resampling procedure). When data are expressed as main  
448 activity of each animal during a certain interval, as often done with sensors, synchrony can only be  
449 approximated: if the intervals at which the activity is noted are short enough (e.g. 5 min) then it may be  
450 considered that the animal kept the same activity during the whole interval, and the data can then be  
451 processed as scan sampling.

452 Several metrics can be found in the literature to calculate synchrony. We cite here the most  
453 common ones.

454 (a) The synchrony between two animals is calculated as the proportion of the scans during  
455 which they are engaged in the same activity (see example in Veissier et al., 1989):

456 
$$\text{synchrony} = \frac{1}{n} \sum_{i=1}^n S_i \quad (7)$$

457 where  $n$  is the number of scans,  $S_i$  equals 1 if the two animals have the same activity at scan  $i$  and  
458 0 if they do not.

460 The synchrony of a focus animal with the rest of the group or a certain sub-group of animals in the  
461 group can be calculated as:

462 (b) the average of the proportion of animals from the group or the sub-group performing the  
463 same activity as the focus animal across scans. This also corresponds to the average of synchrony indices  
464 (a) obtained for a focus animal and any other animal from the herd or the sub-group (Veissier et al., 1989):

465 
$$\text{synchrony} = \frac{1}{n(m-1)} \sum_{j=1}^{m-1} \sum_{i=1}^n S_{ij} \quad (8)$$

466 where  $n$  is the number of scans,  $m$  is the number of animals in the group,  $S_{ij}$  equals 1 if the focus  
467 animal and the other animal  $j$  have the same activity at scan  $i$  and 0 if they do not.

468 (c) the proportion of scans during which the focal individual performs the same activity as  
469 most individuals of the rest of the group (Ruckstuhl, 1999). The calculation is similar to that for the  
470 synchrony between two animals given in (a) above but with  $S_i$  equals 1 when the focus animal has the same  
471 activity as most of the group and 0 when it does not.

472 At group level, the synchrony can be calculated with several indices:

473 (d) The proportion of scans where all animals of the group perform the same activity; Again,  
474 similar calculations as for (a) are used with  $S_i$  equals 1 if all animals perform the same activity and 0 if not.

475 (e) The average of metrics (b) (Veissier et al., 1989) or (c) (Asher & Collins, 2012).

476 The metrics presented above depend largely on the number of activity categories and the number  
477 of animals in the group (especially metrics (d)): when the group is large and the number of activity  
478 categories is high, there is little chance that animals perform the same activity at the same time. Asher and

479 Collins (2012) thus recommend comparing the distribution of activity observed with the one obtained at  
480 random. This can be done with:

481 (f) Kappa coefficient of agreement. For instance, the proportion of pairs of animals observed  
482 with the same activity is calculated (Rook & Penning, 1991):

$$483 \quad P(O) = \frac{1}{np} \sum_{i=1}^n S_i \quad (9)$$

484 where  $n$  is the number of scans,  $p$  is the total number of pairs of animals in the groups, and  $S_i$  the  
485 number of pairs of animals performing the same activity at scan  $i$ .

486 The expected proportion of pairs that would perform the same activity by chance is then  
487 calculated:

$$488 \quad P(E) = \frac{1}{(nm)^2} \sum_{k=1}^l C_k^2 \quad (10)$$

489 where  $n$  is the number of scans,  $m$  is the group size,  $l$  is the number of activity categories, and  $C_k$   
490 the frequency of observation of activity category  $k$  (i.e., total number of scan x animals occurrence of the  
491 activity).

492 Then

$$493 \quad \text{Kappa coefficient} = \frac{P(O) - P(E)}{1 - P(E)} \quad (11)$$

494 The Kappa coefficient equals 1 if all animals always perform the same activity at the same time (full  
495 synchronization). It equals 0 when animals are not synchronised more than at random.

496 (g) Other methods can be found in the literature to compare the synchrony observed to that  
497 expected on a randomised dataset: calculating a dispersion index (Raussi et al., 2011) or applying Monte-  
498 Carlo methods (Whitehead, 1999). These two options are rarely used in the literature on behaviour (e.g.,  
499 we did not find studies using the dispersion index apart from that of Raussi et al. (2011)). In most case, they  
500 actually may not have added value compared to the Kappa coefficient of agreement.

501 The activity categories should be carefully chosen. If there are too many categories (e.g., detailing  
502 precisely what the animals do when standing active: walking, scratching, interacting with each other), the  
503 animals will seem little synchronised whereas if there are too few categories (e.g., active vs. inactive) they  
504 will seem very synchronised. Asher and Collins (2012) recommend using 5 activity categories in laying hens.  
505 It must be considered whether we expect that there can be or not social facilitation of an activity (that is an  
506 animal engaging in an activity encourages the other to do so). In ruminants, ruminating appears as a reflex  
507 activity, ruminating thus should not be used as a separate category but rather included in lying and standing  
508 idling (i.e., postures when ruminating can occur) because we do not expect social facilitation of ruminating.

509 To be interpreted in terms of positive relationship, the synchrony needs to be estimated between  
510 animals that have about the same time budget (i.e., same amount of time spent in each activity per day).  
511 For instance, although a cow has a strong bond to its new-born calf, the apparent synchrony between  
512 them may be low because the cow spends lot of the time foraging whereas the calf spends more time lying  
513 (Veissier et al., 1990).

#### 514 **Biological meaning**

515 Animals may be synchronised because the activity of an animal is influenced by that of other  
516 animals. Social facilitation has been described in many species and contexts (Clayton, 1978). Animals may  
517 be synchronised also because they adopt a similar rhythm of activity: activities follow a circadian rhythm  
518 triggered at least in part by external cues such as light or timing of food distribution (or milking in dairy  
519 cows) so if animals are subjected to the same cues, their activity will tend to be similar (Flury & Gygas,  
520 2016).

521           The synchrony between two animals (fighting excluded) gives us an estimate of how closely (and  
522 positively) they are related to each other. Two animals bound by a positive social relationship have more  
523 chances than unrelated animals to express the same activity at the same time. For instance, when calves  
524 stay with their dam after weaning they keep preferential relations that are shown by proximity, exchanges  
525 of positive interactions, and also synchrony (Veissier et al., 1990).

526           Synchrony at the group level reflects social cohesion, i.e. the strength of the bonds between the  
527 animals in the group (Clayton, 1978). Groups newly formed are usually less cohesive, animals exchanging  
528 aggressive interactions and being less synchronised (Mounier et al., 2005). The synchrony can also decrease  
529 if there is competition for the access to a resource, e.g., food and lying places. For instance, in cows and  
530 sheep, synchrony of lying decreases when lying space is limited (Bøe et al., 2006; Winckler et al., 2015). The  
531 synchrony between an animal and the rest of its group reflects its familiarity with the group: synchrony may  
532 be low in case of newly introduced animals until the organization of the group is stabilised (Arey, 1999;  
533 Boyle et al., 2013). A variation in synchrony can also be caused by a health disorder: the activity of an animal  
534 is modified in case of disease (Dantzer et al., 2008), so that it can depart from the rest of the group (e.g., a  
535 cow isolates and stay standing idling or lying for longer when ill (Proudfoot & Habing, 2015; Proudfoot et  
536 al., 2012).

537  
538

539  
540

**Table 1** - Summary of metrics to describe the activity of animals with their condition of use

Category	Metrics	Raw data	Purpose	Limitations	Reference
<b>Overall activity</b>	Proportion of time spent in an activity	Collection of activity bouts with duration	Detection of physiological states (e.g., oestrus), pathological states (lameness), or stress	Depends on their living conditions and the farm management Modifications are also observed between individuals	Veissier 2004
	Number and duration of activity bouts	Collection of activity bouts with duration		There is a need to : - record on long periods to avoid edge effects. - describe activity with similar details - have a clear determination of when a new activity starts Weights of each activity to be elicited.	Tucker et al., 2009. Ledgerwood et al., 2010. Yeates et al., 2001
	Activity level	Time spent on activities			
<b>Fluctuations around the mean</b>	Min/max/range/quantiles/variance/RMSSD	Timeseries	To identify pathological or physiological states of cows		Hut et al., 2022; Lardy et al., 2023; Mialon et al. 2008.
<b>Cyclicity</b>	Irregularity Autocorrelation Non-periodicity Cosinor method Fourier transform Degree of functional coupling (DFC)	Continuous sampling Continuous sampling Continuous sampling Continuous sampling Continuous sampling	Detection of physiological states (e.g., oestrus), pathological states (lameness), or stress. To predict resilience.	Recording on long periods (longer than the cycle to be detected)	Salgado et al. 2021a&b Dixhoorn et al., 2023, 2018; Chkeir et al., 2019
<b>Synchrony between individuals</b>	Proportion of animals from the group or the sub-group performing the same activity as a focal individual.	Scan sampling	Bonds between animals and / or availability of resources	Depends on the number of animals and number of activity categories	Berger et al., 2003; Scheibe et al., 1999 Veissier et al., 1989
<b>Synchrony at herd level</b>	Proportion of scans during which a focal individual performs the same activity as most individuals of the rest of the group	Scan sampling	Inclusion of an animal in a group. Detection of health disorder		Ruckstuhl, 1999 Asher & Collins 2012
	The average of metrics taken at individual level.	Scan sampling and continuous sampling	Social cohesion Competition for resources		Veissier et al., 1989; Stoye et al. 2012 Arsher and Collins, 2012 Arsher and Collins, 2012 Stoye et al., 2012
	The proportion of scans where all animals of the group perform the same activity Kappa coefficient of agreement				Rook and Penning, 1991

541  
542



544 In this review we present metrics that can be calculated from data on gross activity and identify  
545 their conditions for use (summarised in Table 1). We divided the metrics into four groups that all represent  
546 a different aspects of animal behaviour: the overall activity, the fluctuations around the mean activity, the  
547 cyclicity and the synchrony between animals.

548 The overall activity is the most often studied aspect. It represents the total duration and  
549 organisation of the activity in bouts. The fluctuations of activity during the day or across days and the  
550 organisation of these fluctuations according to (circadian) cycles are less often addressed but are gaining  
551 attention in research. Synchrony between animals is generally used to study the social organisation of a  
552 group of animals, but is also more and more used as specific indicators (e.g. diseases or evaluation of the  
553 management) and constitutes also a promising indicator of positive welfare (Keeling et al., 2021; Napolitano  
554 et al., 2009).

555 These four groups of metrics can be used to analyse the impact of housing and management  
556 procedures. In that case, the values at herd level are evaluated, all animals within the herd are taken into  
557 account. In most cases, the average herd level values per metric are expected to not vary much in time,  
558 unless management or housing change.

559 These metrics can also be used to compare animals within a herd individually (with same  
560 management and housing). Animals are usually consistent with time, so that the data can be used to  
561 phenotype them (Bacher et al., 2022; Poppe et al., 2022). In turn the behavioural phenotype can inform  
562 about other traits. For instance, a cow that shows low regularity in activity is likely to be more susceptible  
563 to post-partum diseases, when she has to cope with metabolic constraints (van Dixhoorn et al., 2023).

564 Transient changes in activity can be observed under certain circumstances at animal level,  
565 especially when an animal is sick, in a specific physiological state (oestrus, calving), or stressed. When such  
566 transient changes are observed on most animals of the herd and at the same time, it is likely that the herd  
567 has been disturbed by external events (e.g., handling to apply a treatment, hoof trimming, period of heat  
568 stress). When the transient change is observed in only one or few animals, it is more likely an individual  
569 case of disease or a reproductive event.

570 Combining several behavioural metrics is usually necessary to have a comprehensive overview of  
571 the internal state of an animal. For instance, to infer the internal state of a cow (diseased, in oestrus, about  
572 to calve, or stressed), metrics on overall activity, fluctuations and cyclicity are necessary to be able to classify  
573 the cows into the corresponding state (Lardy et al., 2023).

574 In this paper, only metrics describing overall activity, fluctuations around mean activity, cyclicity  
575 and synchrony between animals are described and discussed. The data can nevertheless be further  
576 processed to extract more information. For instance, a network analysis could be performed with links  
577 between individuals of a group estimated from their synchrony. Such an approach could probably be  
578 applied to study group effects such as social facilitation, leadership, or cooperation among individuals.  
579 Markov chains can be used to analyse sequences of activities to better understand of the organisation of  
580 activities (Rugg & Buech, 1990; Schafer et al., 2020). Machine learning applied to metrics describing activity  
581 can also help to classify animals according to their phenotype or to detect changes in activity on specific  
582 animals and days (Lardy et al., 2023; Wagner et al., 2020, Debauche et al., 2021).

583 In conclusion, activity data provide the raw material for the calculation of several metrics that  
584 describe animal behaviour. The choice of which metrics to use, depends on the research question or  
585 potential application. A clear research question helps to select the metrics that best characterise an animal  
586 behaviour in relation to the research or applied question. We believe that clarification of the metrics and  
587 on how they should be calculated will help to standardise these metrics, making them easier to use and  
588 allowing comparisons between studies.

589  
590

591

## **Funding**

592

593

594

The study received financial support via the SmartCow project funded by the European Commission (Horizon 2020 research and innovation program, grant agreement No. 730924) and is part of IRC-SAE (CAP 20–25).

595

## **Data, scripts, code, and supplementary information availability**

596

597

598

599

600

Supplementary material : Examples of calculations of metrics using R software are available online: Supplementary materials belonging to From data on gross activity to the characterization of animal behaviour: which metrics for which purposes: <https://doi.org/10.6084/m9.figshare.24891252>.

601

## **Conflict of interest disclosure**

602

603

604

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article. Isabelle Veissier is recommender of PCI Animal Science.

- 605 Arey, D. S. (1999). Time course for the formation and disruption of social organisation in group-housed sows  
 606 [Article]. *Applied Animal Behaviour Science*, 62(2-3), 199-207. <https://doi.org/10.1016/S0168->  
 607 [1591\(98\)00224-X](https://doi.org/10.1016/S0168-1591(98)00224-X)  
 608
- 609 Asher, L., & Collins, L. M. (2012). Assessing synchrony in groups: Are you measuring what you think you are  
 610 measuring? [Article]. *Applied Animal Behaviour Science*, 138(3-4), 162-169.  
 611 <https://doi.org/10.1016/j.applanim.2012.02.004>  
 612
- 613 Bacher, L. M., Prieur, V., Veissier, I., & Boivin, X. (2022). Association between breeding bulls' reactivity to  
 614 humans or handling and their daily behaviour and growth. *Animal*, 16(7), 100568.  
 615 <https://doi.org/10.1016/j.animal.2022.100568>  
 616
- 617 Bateson, M. M., Paul. (2021). *Measuring Behaviour: An Introductory Guide*. Cambridge University Press.  
 618
- 619 Benaissa, S., Tuytens, F. A. M., Plets, D., Trogh, J., Martens, L., Vandaele, L., Joseph, W., & Sonck, B. (2020).  
 620 Calving and estrus detection in dairy cattle using a combination of indoor localization and  
 621 accelerometer sensors. *Computers and Electronics in Agriculture*, 168, 105153.  
 622 <https://doi.org/10.1016/j.compag.2019.105153>  
 623
- 624 Berger, A., Scheibe, K. M., Michaelis, S., & Streich, W. J. (2003). Evaluation of living conditions of free-ranging  
 625 animals by automated chronobiological analysis of behavior. *Behavior Research Methods,*  
 626 *Instruments, & Computers*, 35(3), 458-466. <https://doi.org/10.3758/BF03195524>  
 627
- 628 Bøe, K. E., Berg, S., & Andersen, I. L. (2006). Resting behaviour and displacements in ewes—effects of  
 629 reduced lying space and pen shape. *Applied Animal Behaviour Science*, 98(3), 249-259.  
 630 <https://doi.org/10.1016/j.applanim.2005.10.001>  
 631
- 632 Borchers, M. R., Chang, Y. M., Tsai, I. C., Wadsworth, B. A., & Bewley, J. M. (2016). A validation of  
 633 technologies monitoring dairy cow feeding, ruminating, and lying behaviors [Article]. *Journal of*  
 634 *Dairy Science*, 99(9), 7458-7466. <https://doi.org/10.3168/jds.2015-10843>  
 635
- 636 Boyle, A. R., Ferris, C. P., & O'Connell, N. E. (2013). Does housing nulliparous dairy cows with multiparous  
 637 animals prior to calving influence welfare- and production-related parameters after calving?  
 638 [Article]. *Applied Animal Behaviour Science*, 143(1), 1-8.  
 639 <https://doi.org/10.1016/j.applanim.2012.11.004>  
 640
- 641 Buller, H., Blokhuis, H., Lokhorst, K., Silberberg, M., & Veissier, I. (2020). Animal welfare management in a  
 642 digital world [Review]. *Animals*, 10(10), 1-12, Article 1779. <https://doi.org/10.3390/ani10101779>  
 643
- 644 Calamari, L., Soriani, N., Panella, G., Petrera, F., Minuti, A., & Trevisi, E. (2014). Rumination time around  
 645 calving: An early signal to detect cows at greater risk of disease [Article]. *Journal of Dairy Science*,  
 646 97(6), 3635-3647. <https://doi.org/10.3168/jds.2013-7709>  
 647
- 648 Chkeir, A., Abdallah, M., Soubra, R., & Nassereddine, M. (2019, 9-11 April 2019). A mathematical approach  
 649 using a thoracic temperature sensor for detecting human circadian rhythms. 2019 IEEE Jordan  
 650 International Joint Conference on Electrical Engineering and Information Technology (JEEIT),  
 651 Clayton, D. A. (1978). Socially Facilitated Behavior. *The quarterly review of biology*, 53(4), 373-392.  
 652 <http://www.jstor.org.ezproxy.library.wur.nl/stable/2826580>  
 653
- 654 Commun, L., Silberberg, M., Mialon, M. M., Martin, C., & Veissier, I. (2012). Behavioural adaptations of  
 655 sheep to repeated acidosis challenges and effect of yeast supplementation [Article]. *Animal*, 6(12),  
 656 2011-2022. <https://doi.org/10.1017/S1751731112001309>  
 657
- 658 Dado, R. G., & Allen, M. S. (1993). Continuous Computer Acquisition of Feed and Water Intakes, Chewing,  
 659 Reticular Motility, and Ruminant pH of Cattle. *Journal of Dairy Science*, 76(6), 1589-1600.  
 660 [https://doi.org/10.3168/jds.S0022-0302\(93\)77492-5](https://doi.org/10.3168/jds.S0022-0302(93)77492-5)  
 661
- 662 Dantzer, R., O'Connor, J. C., Freund, G. G., Johnson, R. W., & Kelley, K. W. (2008). From inflammation to  
 663 sickness and depression: when the immune system subjugates the brain. *Nat Rev Neurosci*, 9(1),  
 664 46-56. <https://doi.org/10.1038/nrn2297>  
 665
- 666 De Mol, R. M., André, G., Bleumer, E. J. B., van der Werf, J. T. N., de Haas, Y., & van Reenen, C. G. (2013).  
 667 Applicability of day-to-day variation in behavior for the automated detection of lameness in dairy  
 668 cows [Article]. *Journal of Dairy Science*, 96(6), 3703-3712. <https://doi.org/10.3168/jds.2012-6305>  
 669
- 670 Debauche, O., Elmoulat, M., Mahmoudi, S., Bindelle, J., & Lebeau, F. (2021). Farm animals' behaviors and  
 671 welfare analysis with ia algorithms: A review [Review]. *Revue d'Intelligence Artificielle*, 35(3), 243-  
 672 253. <https://doi.org/10.18280/ria.350308>  
 673

658 DeVries, T. J., Von Keyserlingk, M. A. G., & Beauchemin, K. A. (2003). Short communication: Diurnal feeding  
659 pattern of lactating dairy cows [Article]. *Journal of Dairy Science*, 86(12), 4079-4082.  
660 [https://doi.org/10.3168/jds.S0022-0302\(03\)74020-X](https://doi.org/10.3168/jds.S0022-0302(03)74020-X)

661 Dittrich, I., Gertz, M., & Krieter, J. (2019). Alterations in sick dairy cows' daily behavioural patterns [Review].  
662 *Heliyon*, 5(11), Article e02902. <https://doi.org/10.1016/j.heliyon.2019.e02902>

663 Dohme-Meier, F., Kaufmann, L. D., Görs, S., Junghans, P., Metges, C. C., Van Dorland, H. A., Bruckmaier, R.  
664 M., & Münger, A. (2014). Comparison of energy expenditure, eating pattern and physical activity  
665 of grazing and zero-grazing dairy cows at different time points during lactation [Article]. *Livestock*  
666 *Science*, 162(1), 86-96. <https://doi.org/10.1016/j.livsci.2014.01.006>

667 Dumbell, R., Matveeva, O., & Oster, H. (2016). Circadian Clocks, Stress, and Immunity. *Front Endocrinol*  
668 *(Lausanne)*, 7, 37. <https://doi.org/10.3389/fendo.2016.00037>

669 Flury, R., & Gygax, L. (2016). Daily patterns of synchrony in lying and feeding of cows: Quasi-natural state  
670 and (anti-) synchrony factors. *Behavioural Processes*, 133, 56-61.  
671 <https://doi.org/10.1016/j.beproc.2016.11.004>

672 Fogsgaard, K. K., Bennedsgaard, T. W., & Herskin, M. S. (2015). Behavioral changes in freestall-housed dairy  
673 cows with naturally occurring clinical mastitis [Article]. *Journal of Dairy Science*, 98(3), 1730-1738.  
674 <https://doi.org/10.3168/jds.2014-8347>

675 Goldhawk, C., Chapinal, N., Veira, D. M., Weary, D. M., & von Keyserlingk, M. A. G. (2009). Prepartum  
676 feeding behavior is an early indicator of subclinical ketosis. *Journal of Dairy Science*, 92(10), 4971-  
677 4977. <https://doi.org/10.3168/jds.2009-2242>

678 González, L. A., Tolkamp, B. J., Coffey, M. P., Ferret, A., & Kyriazakis, I. (2008). Changes in Feeding Behavior  
679 as Possible Indicators for the Automatic Monitoring of Health Disorders in Dairy Cows. *Journal of*  
680 *Dairy Science*, 91(3), 1017-1028. <https://doi.org/10.3168/jds.2007-0530>

681 Hafez, E., Schein, M., & Ewbank, R. (1969). *The behaviour of Domestic Animals*. Baillere, Tindall & Casell.  
682 <https://wellcomecollection.org/works/zvgxb37r>

683 Hut, P. R., Kuiper, S. E. M., Nielen, M., Hulsen, J. H. J. L., Stassen, E. N., & Hostens, M. M. (2022). Sensor  
684 based time budgets in commercial Dutch dairy herds vary over lactation cycles and within 24 hours.  
685 *PLoS One*, 17(2). <https://doi.org/10.1371/journal.pone.0264392>

686 Ito, K., von Keyserlingk, M. A. G., LeBlanc, S. J., & Weary, D. M. (2010). Lying behavior as an indicator of  
687 lameness in dairy cows [Article]. *Journal of Dairy Science*, 93(8), 3553-3560.  
688 <https://doi.org/10.3168/jds.2009-2951>

689 Jud, C., Schmutz, I., Hampp, G., Oster, H., & Albrecht, U. (2005). A guideline for analyzing circadian wheel-  
690 running behavior in rodents under different lighting conditions [Article]. *Biological Procedures*  
691 *Online*, 7(1), 101-116. <https://doi.org/10.1251/bpo109>

692 Koltjes, J. E., Cole, J. B., Clemmens, R., Dilger, R. N., Kramer, L. M., Lunney, J. K., McCue, M. E., McKay, S. D.,  
693 Mateescu, R. G., Murdoch, B. M., Reuter, R., Rexroad, C. E., Rosa, G. J. M., Serão, N. V. L., White, S.  
694 N., Woodward-Greene, M. J., Worku, M., Zhang, H., & Reecy, J. M. (2019). A Vision for Development  
695 and Utilization of High-Throughput Phenotyping and Big Data Analytics in Livestock [Review].  
696 *Frontiers in Genetics*, 10(1197). <https://doi.org/10.3389/fgene.2019.01197>

697 Lardy, R., Ruin, Q., & Veissier, I. (2023). Discriminating pathological, reproductive or stress conditions in  
698 cows using machine learning on sensor-based activity data [Article]. *Computers and Electronics in*  
699 *Agriculture*, 204, Article 107556. <https://doi.org/10.1016/j.compag.2022.107556>

700 Ledgerwood, D. N., Winckler, C., & Tucker, C. B. (2010). Evaluation of data loggers, sampling intervals, and  
701 editing techniques for measuring the lying behavior of dairy cattle [Article]. *Journal of Dairy*  
702 *Science*, 93(11), 5129-5139. <https://doi.org/10.3168/jds.2009-2945>

703 Lee, M., & Seo, S. (2021). Wearable Wireless Biosensor Technology for Monitoring Cattle: A Review.  
704 *Animals*, 11(10), 2779. <https://doi.org/10.3390/ani11102779>

705 McCabe, C. J., Aryal, U. K., Casey, T., & Boerman, J. (2021). Impact of exposure to chronic light–dark phase  
706 shifting circadian rhythm disruption on muscle proteome in periparturient dairy cows. *Proteomes*,  
707 9(3), Article 35. <https://doi.org/10.3390/proteomes9030035>

708 Medrano-Galarza, C., Gibbons, J., Wagner, S., de Passillé, A. M., & Rushen, J. (2012). Behavioral changes in  
709 dairy cows with mastitis [Article]. *Journal of Dairy Science*, 95(12), 6994-7002.  
710 <https://doi.org/10.3168/jds.2011-5247>

- 711 Mialon, M. M., Martin, C., Garcia, F., Menassol, J. B., Dubroeuq, H., Veissier, I., & Micol, D. (2008). Effects  
712 of the forage-to-concentrate ratio of the diet on feeding behaviour in young Blond d'Aquitaine  
713 bulls. *Animal*, 2(11), 1682-1691. <https://doi.org/10.1017/S1751731108002905>
- 714 Mounier, L., Veissier, I., & Boissy, A. (2005). Behavior, physiology, and performance of bulls mixed at the  
715 onset of finishing to form uniform body weight groups. *Journal of Animal Science*, 83(7), 1696-1704.  
716 <https://doi.org/10.2527/2005.8371696x>
- 717 Pastell, M., Tiusanen, J., Hakojärvi, M., & Hänninen, L. (2009). A wireless accelerometer system with wavelet  
718 analysis for assessing lameness in cattle [Article]. *Biosystems Engineering*, 104(4), 545-551.  
719 <https://doi.org/10.1016/j.biosystemseng.2009.09.007>
- 720 Poppe, M., Mulder, H. A., van Pelt, M. L., Mullaart, E., Hogeveen, H., & Veerkamp, R. F. (2022). Development  
721 of resilience indicator traits based on daily step count data for dairy cattle breeding [Article].  
722 *Genetics Selection Evolution*, 54(1), Article 21. <https://doi.org/10.1186/s12711-022-00713-x>
- 723 Proudfoot, K., & Habing, G. (2015). Social stress as a cause of diseases in farm animals: Current knowledge  
724 and future directions. *Veterinary Journal*, 206(1), 15-21. <https://doi.org/10.1016/j.tvjl.2015.05.024>
- 725 Proudfoot, K. L., Weary, D. M., & von Keyserlingk, M. A. G. (2012). Linking the social environment to illness  
726 in farm animals. *Applied Animal Behaviour Science*, 138(3), 203-215.  
727 <https://doi.org/10.1016/j.applanim.2012.02.008>
- 728 Raussi, S., Jauhainen, L., Saastamoinen, S., Siivonen, J., Hepola, H., & Veissier, I. (2011). A note on  
729 overdispersion as an index of behavioural synchrony: a pilot study in dairy cows. *Animal*, 5(3), 428-  
730 432. <https://doi.org/10.1017/S1751731110001928>
- 731 Reith, S., & Hoy, S. (2018). Review: Behavioral signs of estrus and the potential of fully automated systems  
732 for detection of estrus in dairy cattle. *Animal*, 12(2), 398-407.  
733 <https://doi.org/10.1017/S1751731117001975>
- 734 Rocha, L. E. C., Terenius, O., Veissier, I., Meunier, B., & Nielsen, P. P. (2020). Persistence of sociality in group  
735 dynamics of dairy cattle [Article]. *Applied Animal Behaviour Science*, 223, Article 104921.  
736 <https://doi.org/10.1016/j.applanim.2019.104921>
- 737 Rook, A. J., & Penning, P. D. (1991). Synchronisation of eating, ruminating and idling activity by grazing  
738 sheep. *Applied Animal Behaviour Science*, 32(2), 157-166. [https://doi.org/10.1016/S0168-  
739 1591\(05\)80039-5](https://doi.org/10.1016/S0168-1591(05)80039-5)
- 740 Ruckstuhl, K. E. (1999). To synchronise or not to synchronise: A dilemma for young bighorn males? [Article].  
741 *Behaviour*, 136(6), 805-818. <https://doi.org/10.1163/156853999501577>
- 742 Rugg, D. J., & Buech, R. R. (1990). Analyzing Time Budgets with Markov Chains. *Biometrics*, 46(4), 1123-  
743 1131. <https://doi.org/10.2307/2532453>
- 744 Rutten, C. J., Kamphuis, C., Hogeveen, H., Huijps, K., Nielen, M., & Steeneveld, W. (2017). Sensor data on  
745 cow activity, rumination, and ear temperature improve prediction of the start of calving in dairy  
746 cows [Article]. *Computers and Electronics in Agriculture*, 132, 108-118.  
747 <https://doi.org/10.1016/j.compag.2016.11.009>
- 748 Rutten, C. J., Velthuis, A. G. J., Steeneveld, W., & Hogeveen, H. (2013). Invited review: Sensors to support  
749 health management on dairy farms. *Journal of Dairy Science*, 96(4), 1928-1952.  
750 <https://doi.org/10.3168/jds.2012-6107>
- 751 Saint-Dizier, M., & Chastant-Maillard, S. (2012). Towards an Automated Detection of Oestrus in Dairy Cattle  
752 [Review]. *Reproduction in Domestic Animals*, 47(6), 1056-1061. [https://doi.org/10.1111/j.1439-  
753 0531.2011.01971.x](https://doi.org/10.1111/j.1439-0531.2011.01971.x)
- 754 Salgado, H. H., Méthot, S., Remus, A., Létourneau-Montminy, M. P., & Pomar, C. (2021a). Elucidating the  
755 impact of feeding behaviour on body composition in finishing pigs fed ad libitum using an  
756 integrative feeding behaviour index [Article]. *Livestock Science*, 251, Article 104650.  
757 <https://doi.org/10.1016/j.livsci.2021.104650>
- 758 Salgado, H. H., Méthot, S., Remus, A., Létourneau-Montminy, M. P., & Pomar, C. (2021b). A novel feeding  
759 behavior index integrating several components of the feeding behavior of finishing pigs [Article].  
760 *Animal*, 15(7), Article 100251. <https://doi.org/10.1016/j.animal.2021.100251>
- 761 Saper, C. B., Scammell, T. E., & Lu, J. (2005). Hypothalamic regulation of sleep and circadian rhythms. *Nature*,  
762 437(7063), 1257-1263. <https://doi.org/10.1038/nature04284>
- 763 Schafer, T. L. J., Wikle, C. K., VonBank, J. A., Ballard, B. M., & Weegman, M. D. (2020). A Bayesian Markov  
764 Model with Pólya-Gamma Sampling for Estimating Individual Behavior Transition Probabilities from



765 Accelerometer Classifications [Article]. *Journal of Agricultural, Biological, and Environmental*  
766 *Statistics*, 25(3), 365-382. <https://doi.org/10.1007/s13253-020-00399-y>

767 Scheffer, M., Bolhuis, J. E., Borsboom, D., Buchman, T. G., Gijzel, S. M. W., Goulson, D., Kammenga, J. E.,  
768 Kemp, B., van de Leemput, I. A., Levin, S., Martin, C. M., Melis, R. J. F., van Nes, E. H., Romero, L.  
769 M., & Olde Rikkert, M. G. M. (2018). Quantifying resilience of humans and other animals.  
770 *Proceedings of the National Academy of Sciences*. <https://doi.org/10.1073/pnas.1810630115>

771 Scheibe, K. M., Berger, A., Langbein, J., Streich, W. J., & Eichhorn, K. (1999). Comparative Analysis of  
772 Ultradian and Circadian Behavioural Rhythms for Diagnosis of Biorhythmic State of Animals.  
773 *Biological Rhythm Research*, 30(2), 216-233. <https://doi.org/10.1076/brhm.30.2.216.1420>

774 Solano, L., Barkema, H. W., Pajor, E. A., Mason, S., LeBlanc, S. J., Nash, C. G. R., Haley, D. B., Pellerin, D.,  
775 Rushen, J., de Passillé, A. M., Vasseur, E., & Orsel, K. (2016). Associations between lying behavior  
776 and lameness in Canadian Holstein-Friesian cows housed in freestall barns. *Journal of Dairy Science*,  
777 99(3), 2086-2101. <https://doi.org/10.3168/jds.2015-10336>

778 Stoye, S., Porter, M. A., & Stamp Dawkins, M. (2012). Synchronized lying in cattle in relation to time of day  
779 [Article]. *Livestock Science*, 149(1-2), 70-73. <https://doi.org/10.1016/j.livsci.2012.06.028>

780 Tolkamp, B. J., Allcroft, D. J., Austin, E. J., Nielsen, B. L., & Kyriazakis, I. (1998). Satiety Splits Feeding  
781 Behaviour into Bouts. *Journal of Theoretical Biology*, 194(2), 235-250.  
782 <https://doi.org/10.1006/jtbi.1998.0759>

783 Tucker, C. B., Cox, N. R., Weary, D. M., & Špinka, M. (2009). Laterality of lying behaviour in dairy cattle.  
784 *Applied Animal Behaviour Science*, 120(3), 125-131.  
785 <https://doi.org/10.1016/j.applanim.2009.05.010>

786 Tucker, C. B., Jensen, M. B., de Passillé, A. M., Hänninen, L., & Rushen, J. (2021). Lying time and the welfare  
787 of dairy cows. *Journal of Dairy Science*, 104(1), 20-46. <https://doi.org/10.3168/jds.2019-18074>

788 Tuomisto, L., Huuskonen, A., Jauhiainen, L., & Mononen, J. (2019). Finishing bulls have more synchronised  
789 behaviour in pastures than in pens [Article]. *Applied Animal Behaviour Science*, 213, 26-32.  
790 <https://doi.org/10.1016/j.applanim.2019.02.007>

791 Van Dixhoorn, I. D. E., de Mol, R. M., Schnabel, S. K., van der Werf, J. T. N., van Mourik, S., Bolhuis, J. E.,  
792 Rebel, J. M. J., & van Reenen, C. G. (2023). Behavioral patterns as indicators of resilience after  
793 parturition in dairy cows. *Journal of Dairy Science*, 106(9), 6444-6463.  
794 <https://doi.org/10.3168/jds.2022-22891>

795 Van Dixhoorn, I. D. E., de Mol, R. M., van der Werf, J. T. N., van Mourik, S., & van Reenen, C. G. (2018).  
796 Indicators of resilience during the transition period in dairy cows: A case study. *J Dairy Sci*, 101(11),  
797 10271-10282. <https://doi.org/10.3168/jds.2018-14779>

798 Van Erp, R. J. J., De Vries, S., Van Kempen, T. A. T. G., Den Hartog, L. A., & Gerrits, W. J. J. (2020). Circadian  
799 misalignment imposed by nocturnal feeding tends to increase fat deposition in pigs [Article]. *British*  
800 *Journal of Nutrition*, 123(5), 529-536. <https://doi.org/10.1017/S0007114519003052>

801 Veissier, I., Boissy, A., dePassillé, A. M., Rushen, J., van Reenen, C. G., Roussel, S., Andanson, S., & Pradel, P.  
802 (2001). Calves' responses to repeated social regrouping and relocation. *J Anim Sci*, 79(10), 2580-  
803 2593. <https://doi.org/10.2527/2001.79102580x>

804 Veissier, I., Capdeville, J., & Delval, E. (2004). Cubicle housing systems for cattle: Comfort of dairy cows  
805 depends on cubicle adjustment [Review]. *Journal of Animal Science*, 82(11), 3321-3337.  
806 <https://doi.org/10.2527/2004.82113321x>

807 Veissier, I., Lamy, D., & Le Neindre, P. (1990). Social behaviour in domestic beef cattle when yearling calves  
808 are left with the cows for the next calving [Article]. *Applied Animal Behaviour Science*, 27(3), 193-  
809 200. [https://doi.org/10.1016/0168-1591\(90\)90056-J](https://doi.org/10.1016/0168-1591(90)90056-J)

810 Veissier, I., & Le Neindre, P. (1989). Weaning in calves: Its effects on social organization [Article]. *Applied*  
811 *Animal Behaviour Science*, 24(1), 43-54. [https://doi.org/10.1016/0168-1591\(89\)90124-X](https://doi.org/10.1016/0168-1591(89)90124-X)

812 Veissier, I., Le Neindre, P., & Trillat, G. (1989). The use of circadian behaviour to measure adaptation of  
813 calves to changes in their environment. *Applied Animal Behaviour Science*, 22(1), 1-12.  
814 [https://doi.org/10.1016/0168-1591\(89\)90075-0](https://doi.org/10.1016/0168-1591(89)90075-0)

815 Veissier, I., Mialon, M. M., & Sloth, K. H. (2017). Short communication: Early modification of the circadian  
816 organization of cow activity in relation to disease or estrus. *J Dairy Sci*, 100(5), 3969-3974.  
817 <https://doi.org/10.3168/jds.2016-11853>

- 818 Wagner, N., Antoine, V., Koko, J., Mialon, M. M., Lardy, R., & Veissier, I. (2020). Comparison of Machine  
819 Learning Methods to Detect Anomalies in the Activity of Dairy Cows. In *Lecture Notes in Computer*  
820 *Science (including subseries Lecture Notes in Artificial Intelligence and Lecture Notes in*  
821 *Bioinformatics)* (Vol. 12117 LNAI, pp. 342-351).
- 822 Wagner, N., Mialon, M. M., Sloth, K. H., Lardy, R., Ledoux, D., Silberberg, M., de Boyer des Roches, A., &  
823 Veissier, I. (2021). Detection of changes in the circadian rhythm of cattle in relation to disease,  
824 stress, and reproductive events [Article]. *Methods*, 186, 14-21.  
825 <https://doi.org/10.1016/j.ymeth.2020.09.003>
- 826 Weary, D. M., Huzzey, J. M., & Von Keyserlingk, M. A. G. (2009). Board-invited Review: Using behavior to  
827 predict and identify ill health in animals. *Journal of Animal Science*, 87(2), 770-777.  
828 <https://doi.org/10.2527/jas.2008-1297>
- 829 Whitehead, H. A. L. (1999). Testing association patterns of social animals. *Animal Behaviour*, 57(6), F26-F29.  
830 <https://doi.org/10.1006/anbe.1999.1099>
- 831 Winckler, C., Tucker, C. B., & Weary, D. M. (2015). Effects of under- and overstocking freestalls on dairy  
832 cattle behaviour. *Applied Animal Behaviour Science*, 170, 14-19.  
833 <https://doi.org/10.1016/j.applanim.2015.06.003>
- 834 Yeates, M. P., Tolkamp, B. J., Allcroft, D. J., & Kyriazakis, I. (2001). The use of mixed distribution models to  
835 determine bout criteria for analysis of animal behaviour [Article]. *Journal of Theoretical Biology*,  
836 213(3), 413-425. <https://doi.org/10.1006/jtbi.2001.2425>

837