



HAL
open science

Promising parameters to foressee intake and feed efficiency at pasture - a meta-analysis approach

Maryline Boval, Daniel Sauvant

► To cite this version:

Maryline Boval, Daniel Sauvant. Promising parameters to foressee intake and feed efficiency at pasture - a meta-analysis approach. 70. Annual Meeting of the European Federation of NAimal Science, Aug 2019, Ghent, Belgium. pp.Session 45/Theatre 7. hal-04483728

HAL Id: hal-04483728

<https://hal.inrae.fr/hal-04483728>

Submitted on 29 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.


Book of Abstracts of the 70th Annual Meeting of the European Federation of Animal Science

Ghent, Belgium, 26th-30th August, 2019



EAAAP Scientific Committee:

E. Strandberg
G. Savoini
H.A.M. Spolder
H. Sauerwein
M. Lee
J.F. Hocquette
J. Conington
E.F. Knol
A.S. Santos
T. Veldkamp
I. Halachmi
G. Pollott



OASES

Online Academic Submission and Evaluation System

Promising parameters to foresee intake and feed efficiency at pasture – a meta-analysis approach

M. Boval and D. Sauvant

INRA, PHASE, INRA, AgroParisTech, Université Paris-Saclay, 75005, Paris, France, France;
maryline.boval@agroparistech.fr

In a context of major advances in electronic/computer technologies, identifying key and routinely measurable criterial to detect animals that are likely to be more efficient in various feeding environments, is essential. For that purpose, a meta-analysis carried out on 110 publications (n) and gathering 905 experiments (nexp) involving feeding behaviour of grazing cattle and small ruminants, stressed the importance of the bite mass (BM, mg DM/bite/kg BW, $0.11 < 1.79 \pm 1.27 < 7.41$, n=582), as an essential unitary criterion for dry matter intake (DMI, $0.25 < 2.79 \pm 1.06 < 8.01\%$ BW, n=238). The intra-experiment equation is $DMI = 2.16 + 0.56 BM$ (RMSE=0.42, n=180, nexp=67). This meta-analysis also highlighted the importance of the incisive arcade (IA, $2.22 < 5.23 \pm 2.11 < 8.60$ cm, n=112), which is measurable and related both to BW ($IA = 0.91 BW^{0.346}$, RMSE=0.27, n=20) and to BM ($BM = 0.015 IA^{1.88}$, RMSE=0.10, n=45, nexp=21). In addition, we have tested several predictive regressions of DMI and notably one based on BM combined with ruminating time (RU, $0.52 < 6.06 \pm 2.15 < 9.57$ h/day, n=265). The equation is $DMI = 1.395 + 0.652 BM + 0.097 RU$, (RMSE=0.19, n=87, nexp=33). It must be emphasized that grazing time (GT, $3.33 < 8.79 \pm 2.43 < 18.00$ h/day, n=142) has no influence in the latter equation; moreover, the association of BM and GT in order to predict DMI is less accurate than the association of BM and RU (RMSE of 0.28 vs 0.19, respectively). Otherwise, biting rate (BR, $11.18 < 46.89 \pm 14.65 < 106.71$ bites/min, n=559), which is also an easily measurable parameter, presents no intra experiment relationship with DMI. As the BM can be estimated from the IA, measurable with a simple Vernier caliper, and that RU is easily measurable with various technologies such as acoustic recordings or accelerometers, these findings are very promising for assessing feed efficiency and individually.

Session 45

Theatre 8

Validation of sensor on rumination and feeding behaviour of dairy heifers in two feedlot systemsR.D. Kliemann¹, E.M. Nascimento¹, S.R. Fernandes¹, M.M. Campos², T.R. Tomich², L.G.R. Pereira² and A.F. Garcez Neto¹¹Federal University of Paraná, Palotina-Paraná, 85.950-000, Brazil, ²EMBRAPA, Dairy Cattle, Juiz de Fora-Minas Gerais, 36.038-330, Brazil; americo.garcez@ufpr.br

The aim of this study was to validate the sensor HEATIME® HR SYSTEM for rumination activity and evaluate the behaviour traits of dairy heifers in tie-stall and loose-housing systems. Eleven Gir heifers with 10 months of age and 179 ± 26 kg of body weight were housed in both systems following a crossover design with two periods of five consecutive days for data recording. A total mixed ration was offered *ad libitum* in both systems. Before each period, the animals were adapted to the collars with the sensors and systems during seven days. Each day of evaluation regarded 8 hours of observation (08:00-12:00 and 14:00-18:00). The time spent in feeding, water intake, rumination in standing and lying, and idleness in standing and lying were assessed by visual evaluation, whereas the sensor monitored only the rumination activity. In the visual evaluation the activities were recorded every three minutes, and the data collected by the sensor were recorded in 2-h interval. For the sensor validation, the data from visual evaluation were adjusted to 2-h interval. The regression of rumination recorded by the sensor to the visual evaluation was significant only for loose-housing system ($P=0.0002$), but the Pearson correlation between both measures was negative and low ($r=-0.25$; $P=0.0002$). The sensor overestimated the rumination by 27.3% in loose-housing (28 vs 22 min/2 h) and 38.5% in tie-stall (36 vs 26 min/2 h) and, therefore, the sensor was not validated. Regarding the systems comparison by visual evaluation, the time spent drinking, idleness in standing, rumination in standing and lying, and the total rumination time not differed between systems. However, the heifers in tie-stall spent more ($P<0.05$) time in feeding (41 vs 29 min/2 h), idleness in lying (19 vs 15 min/2 h) and total idleness (22 vs 17 min/2 h), whereas those in loose-housing spent more ($P<0.05$) time in other activities (50 vs 30 min/2 h). The sensor is not effective to record the rumination activity in Gir heifers. Loose-housing leads animals to greater activity not related to feeding compared to tie-stall.