



HAL
open science

The bii4africa dataset of faunal and floral population intactness estimates across Africa's major land uses

Hayley S. Clements, Emmanuel Do Linh San, Gareth Hempson, Birthe Linden, Bryan Maritz, Ara Monadjem, Chevonne Reynolds, Frances Siebert, Nicola Stevens, Reinette Biggs, et al.

► To cite this version:

Hayley S. Clements, Emmanuel Do Linh San, Gareth Hempson, Birthe Linden, Bryan Maritz, et al.. The bii4africa dataset of faunal and floral population intactness estimates across Africa's major land uses. *Scientific Data*, 2024, 11, pp.191. <10.1038/s41597-023-02832-6>. <hal-04484669>

HAL Id: hal-04484669

<https://hal.inrae.fr/hal-04484669v1>

Submitted on 29 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC 4.0 - Attribution - Non-commercial use - International License



OPEN

DATA DESCRIPTOR

The bii4africa dataset of faunal and floral population intactness estimates across Africa's major land uses

Hayley S. Clements *et al.*[#]

Sub-Saharan Africa is under-represented in global biodiversity datasets, particularly regarding the impact of land use on species' population abundances. Drawing on recent advances in expert elicitation to ensure data consistency, 200 experts were convened using a modified-Delphi process to estimate 'intactness scores': the remaining proportion of an 'intact' reference population of a species group in a particular land use, on a scale from 0 (no remaining individuals) to 1 (same abundance as the reference) and, in rare cases, to 2 (populations that thrive in human-modified landscapes). The resulting bii4africa dataset contains intactness scores representing terrestrial vertebrates (tetrapods: $\pm 5,400$ amphibians, reptiles, birds, mammals) and vascular plants ($\pm 45,000$ forbs, graminoids, trees, shrubs) in sub-Saharan Africa across the region's major land uses (urban, cropland, rangeland, plantation, protected, etc.) and intensities (e.g., large-scale vs smallholder cropland). This dataset was co-produced as part of the Biodiversity Intactness Index for Africa Project. Additional uses include assessing ecosystem condition; rectifying geographic/taxonomic biases in global biodiversity indicators and maps; and informing the Red List of Ecosystems.

Background & Summary

Accelerating socio-economic development over the past century has caused a dramatic transformation of ecosystems, through human activities such as cultivation, urbanisation, resource extraction and infrastructure development^{1,2}. Human land use activities are major drivers of biodiversity loss^{3,4}. As awareness grows about the scale and pace of biodiversity loss, so does our understanding of the importance of biodiversity to human well-being⁵. Despite this increased awareness, development agendas and policy interventions persistently overlook the critical support-system role of biodiversity in sustainable development and as a source of resilience in times of change⁶⁻¹⁰. This under-representation is perpetuated by the scarcity of suitable biodiversity data and the difficulty of consistently quantifying biodiversity at the scales relevant for policy, in metrics that indicate its support-system role and the impacts of human activities on that support system¹⁰⁻¹².

Existing biodiversity datasets that could be used to assess human impacts on biodiversity and the support system it provides have significant limitations that hamper the mainstreaming of biodiversity into policy and planning. Firstly, these datasets are biased across taxa (towards larger, more conspicuous species, especially large mammals and birds¹³⁻¹⁵), regions (towards North America and Europe, with Africa being particularly under-represented¹³⁻¹⁵), and land uses (towards more intact land uses, notably protected areas^{14,16}). Secondly, to consistently assess and compare anthropogenic impacts on biodiversity, the current state of biodiversity ideally needs to be compared to a reference state¹⁷. This comparison requires biodiversity data to be collected using comparable methods either over time (e.g., McRae *et al.*¹⁶) or across contemporary human-modified and unmodified 'intact' landscapes (e.g., Schipper *et al.*¹⁸), further curtailing data availability. Finally, the connection between common biodiversity metrics (e.g., global or regional threat status of species; representation of species distributions within protected areas^{14,19}) and the functional dimensions of biodiversity relevant to its

[#]A full list of authors and their affiliations appears at the end of the paper.

support-system roles is often unclear^{6,10}. Such roles typically manifest at local scales and depend on the population abundance of species within different functional groups, and how they are impacted by human land uses in these areas^{11,20,21}. Exemplifying these limitations, the largest global dataset of species population abundances across different land uses (the ‘PREDICTS’ database²²) includes just 35 Afrotropical studies with population counts in an anthropogenically modified site compared with an ‘intact’ reference site. African decision-makers have noted data limitations as a major constraint to mainstreaming biodiversity into national sustainable development efforts²³, and such limitations can bias international decision-making towards Global North solutions²⁴.

We address this data gap through a structured expert elicitation process (Fig. 1) involving 200 experts in Afrotropical biodiversity on mainland sub-Saharan Africa. Expert elicitation is used widely in conservation and natural resource management when data are insufficient or absent²⁵. Notable examples of where expert elicitation has been used include the International Union for Conservation of Nature (IUCN) Red List of Threatened Species²⁶, and assessments of the Intergovernmental Panels on Climate Change²⁷ and on Biodiversity and Ecosystem Services⁵. We employed the latest advances in expert elicitation to ensure data rigour and consistency²⁵. Expanding substantially on an earlier approach that involved 16 experts in Southern Africa²⁸, experts estimated ‘intactness scores’: the remaining proportion of an ‘intact’ reference population of a given species group (Table 1) in a particular land use (Table 2), on a scale from 0 (no remaining individuals) to 1 (same abundance as a reference population) and, in rare cases, 2 (populations that thrive in human-modified landscapes). The ‘intact’ reference is the population abundance that would likely have occurred in the area before alteration by modern industrial society. Because information on species populations from this era is virtually non-existent, standard protocol is to consider a remote wilderness area with a natural disturbance regime²⁸ (a Hybrid-Historical approach¹⁷) where necessary. The resulting bii4africa dataset contains standardised intactness scores representing terrestrial vertebrates (tetrapods: $\pm 5,400$ amphibians, reptiles, birds and mammals) and vascular plants ($\pm 45,000$ forbs, graminoids, trees and shrubs) in sub-Saharan Africa across nine major land uses of varying intensity.

The dataset was developed to enable the quantification of the Biodiversity Intactness Index²⁸ for sub-Saharan Africa (<https://bii4africa.org/>). It provides an African-led alternative to a previous attempt to map the Biodiversity Intactness Index globally based on a model²⁹, which produced inaccurate results for Africa³⁰ (likely in part because of the under-representation of African data in the model). The bii4africa dataset that we present here has a broad range of additional uses, including assessing and mapping ecosystem condition to inform national planning and reporting on Goal A in the post-2020 Global Biodiversity Framework; alleviating geographic and taxonomic biases in global biodiversity indicators and maps; parameterising models of biodiversity in a changing world; informing the IUCN Red List of Ecosystems; supporting the United Nations Decade on Ecosystem Restoration in identifying priority ecosystems and monitoring the impact of investments into restoration; identifying the properties of novel ecosystems; and informing future research and training in African biodiversity (Table 3). Importantly, the dataset—co-produced by 200 experts—embodies context-specific knowledge on African biodiversity that contributes to inclusivity in ecology³¹. It is also a positive response to the recent call by the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES) for African-led research that closes knowledge gaps by mobilising local data²⁴.

Methods

Elicitation planning and expert recruitment. We implemented a published, modified-Delphi protocol (multiple rounds of individual, independent expert estimation interspersed with group discussion and review; Fig. 1), which has been shown to improve the rigour of elicitation outcomes²⁵. The elicitation was limited to terrestrial vertebrates (tetrapods) and vascular plants – groups that comprise species with a high diversity of attributes and functions. Experts were sought with knowledge of the degree to which human land uses impact populations of these species groups in sub-Saharan Africa (or a region therein). Such expert knowledge is typically limited to one taxonomic class (e.g., birds or reptiles), or in the case of mammals, often just one or several orders (e.g., primates or bats). As such, multiple expert elicitation processes were run between November 2020 and January 2022, to cover the various broad taxonomic groups of species (see Table 1). An expert in each broad taxonomic group was invited to lead the elicitation for that taxonomic group. Each ‘lead expert’ was identified based on their relevant expertise and existing network across the continent or willingness to develop such a network. For example, several lead experts serve in the IUCN Species Survival Commission working groups and other regional networks (e.g., <https://ascaris.org/>; <https://www.birdlife.org/our-partners-africa/>). The lead expert was responsible for identifying experts and inviting them to participate, as well as proposing a draft list of species response groups and providing input into the development of the land use categories. These tasks are detailed in the subsequent sections.

Identifying and recruiting experts. A broad definition of expertise was used to identify experts, centred on experience of how sub-Saharan species are impacted by human land uses^{25,32}. Diverse types of people can have such experience (e.g., researchers, field or tour guides, park rangers, conservation practitioners, museum curators, and consultants), and inclusion was thus not limited to specific qualifications or institutional affiliations. The aim was to include about 20 experts for each broad taxonomic group, according to guidelines based on practicality and evidence of limited improvements in group performance above 6 to 12 experts²⁵.

The lead expert identified individuals known to have relevant expertise. If this activity did not achieve the target of 20 individuals, additional experts were identified through relevant publications (using appropriate search terms on Google Scholar) and websites (e.g., specialist nature guides or tours, conservation organisations). In some cases, participating experts were asked to recommend other experts (snowball sampling). An invitation was emailed by the lead expert to each identified expert, explaining the project and what would




Fig. 1 Overview of the structured expert elicitation process, based on the IDEA protocol (Investigate, Discuss, Estimate, Aggregate). This process was run for each broad taxonomic group (Table 1), between November 2020 and January 2022, to elicit from 200 experts the estimated impact of nine major African land uses (Table 2) on the relative population abundances of terrestrial vertebrates and vascular plants.

be required of them. Experts confirmed that they had the relevant experience and were willing to participate by returning a signed consent form, including demographic details (notably taxa, years and region of experience). Ethical clearance for the project was provided by Stellenbosch University (project number 15182).

Of the 248 experts who agreed to participate, a total of 200 (81%) participated in the full elicitation process (<https://bii4africa.org/category/experts/>), with an average of 24 experts per broad taxonomic group (Fig. 2). These experts were from 39 countries, including 23 African countries (59% of experts were African nationals). Most experts (72%) were resident in Africa at the time of elicitation, across 26 countries. Their experience spanned sub-Saharan Africa, with Southern Africa being best represented, followed by East Africa, and West Africa having the lowest representation. Experts worked in a range of sectors, most commonly at universities, followed by research institutes and conservation organisations. Collectively, experts had over 3,300 years of relevant accumulated experience, with a mean (\pm SD) of 18 ± 10 years per expert.

Identifying and describing species response groups. It was not possible for experts to estimate the impact of different human land uses on the abundance of every terrestrial vertebrate and vascular plant species in sub-Saharan Africa ($\pm 50,000$ species), both because of the large number of estimates that would be required, and because there are many species for which there is limited ecological knowledge. Therefore, species that were expected to respond in a similar way to human land uses were assigned to a ‘species response group’²⁸. These groups can be considered broadly synonymous to functional groups³³, though our focus is specifically on common responses of population abundance to human land use.

Within each broad taxonomic group (Table 1), the lead expert proposed a draft set of species response groups, based on their knowledge of the key organismal attributes likely to determine the impact of different land uses on populations. These draft species response groups were presented to the participating experts for that taxonomic group during an introductory planning meeting (see *Structured expert elicitation* section below) and revised based on experts’ feedback. Key organismal attributes that informed species response group categorisation varied between the taxonomic groups (Table 1; Supplementary Table 1). For terrestrial vertebrates, attributes commonly included habitat requirements (e.g., forest, grassland, generalist), body size, diet and stratum (e.g., arboreal, fossorial, rupicolous). Terrestrial vascular plants were divided into three broad groups: trees and shrubs, graminoids, and forbs, and species response groups were described within each broad group based on a framework of life-history strategies shaped by disturbance³⁴. Species response groups were thus ecologically defined, not necessarily taxonomically defined. Large mammals were an exception to the response-group approach. The elicitation was done at species level for the seven large carnivores and 95 large herbivores. While the large herbivores were initially categorised into response groups, experts decided during the introductory meeting to estimate per species rather than per species response group. Thus, for large herbivores, we present species response groups, as well as species-level estimates. Details of each response group are included in the ‘Sp_Groups’ spreadsheet in the bii4africa dataset³⁵.

For each broad taxonomic group of terrestrial vertebrates, the lead expert allocated each IUCN-listed species (up to date at the time of the elicitation) in sub-Saharan Africa into the appropriate species response group, with input from participating experts where necessary (‘Sp_Vert’ spreadsheet in the bii4africa dataset³⁵). Determining an average intactness score across species for a given region (country, ecoregion, biome, etc.) can thus be done including only the species that occur in that region, based on species range maps available through the IUCN Red List²⁶. By contrast, allocating each plant species into its respective response group was not possible given the

Taxonomic group	Response group allocation
Birds (Aves) - 1970 species - 17 response groups	Large, taxonomically unique species were allocated into response groups first: vultures and raptors (large and small), large terrestrial species (e.g., cranes) and waterbirds. Two additional wide-ranging groups were added: aerial feeders (swifts and swallows) and opportunistic species that thrive in human-dominated landscapes. The remaining groups were classified based on their habitat nesting location (cavity, ground, other) and body size.
Amphibians (Amphibia) - 799 species - 7 response groups	Experts emphasised the influence of breeding habitat in how amphibians respond to human-modified landscapes. Six groups were thus defined by the predominant habitat in which each species breeds (direct developers, water: permanent or ephemeral and flowing or still, seep, tree hollow).
Reptiles (Reptilia) - 1481 species - 13 response groups	Habitat utilisation, taxonomy, and body size (in a hierarchical manner) influence likely responses of reptiles. First, species occupying strictly aquatic, fossorial, rupicolous, or arboreal habitats were grouped. Because of their derived life history traits, the remaining chelonians were allocated to their own group. Remaining species were grouped on the basis of taxonomy (snake vs lizard), body size, and degree of habitat specialisation.
Mammals (Mammalia)	
Bats - 214 species - 10 response groups	Bats were grouped by their foraging strategy (clutter, edge or open environment insect foragers; fruit-eaters), their roosting location (cave, crevice, foliage) and degree of roost flexibility.
Insectivores - 205 species - 9 response groups	Insectivores were grouped by a combination of taxonomy, body size, habitat (forest, montane, savanna) and stratum (aquatic, arboreal, fossorial, terrestrial).
Rodents, rabbits, hyraxes - 426 species - 16 response groups	Rodents were grouped based on a combination of their taxonomy, habitat preference (arid, forest, montane, savanna/grassland, wetland), body size, diet (herbivorous, granivorous, generalist), stratum (aquatic, arboreal, fossorial, rupicolous, terrestrial) and habits (diurnal, nocturnal).
Small carnivores, aardvark, pangolins - 70 species - 16 response groups	Small carnivores were grouped based on range size (esp. when restricted vs large); stratum (arboreal, terrestrial, aquatic); habitat(s) (notably to differentiate forest, savanna, rocky); diet (esp. for specialised vs omnivorous species); foraging socio-ecology (solitary vs social); body size; and taxonomic affiliation. Five non-carnivorous mammalian species were also included because of similar ecological characteristics, relatively small body size, and similar phylogenetic relationships.
Large carnivores - 7 species	The elicitation was done at species level; no response groups were provided.
Large herbivores - 95 species	The elicitation was done at species level; with each large mammal herbivore species also allocated into one of 12 response groups based on diet (grazer, browser, mixed, frugivore) and body size, with sociality, habitat (e.g., arid, montane, water-dependent), movement (e.g., migratory) and taxonomic groupings also considered.
Primates - 106 species - 6 response groups	Primates were grouped using a combination of habitat (primary/secondary rainforest, woodland/savanna, grassland, generalist found in moist and more arid forests/woodlands); stratum (strictly arboreal, semi-terrestrial, terrestrial); and diet (omnivorous, specialist).
Vascular plants: graminoids, forbs, trees and shrubs ~45,000 species - 33 response groups	Plants were first differentiated by growth form: graminoids (grasses, sedges, rushes); forbs (herbaceous annuals and perennials, geophytes, geoxyles, tubers, herbaceous climbers, dwarf shrubs, succulents, suffrutices); and trees and shrubs (including woody lianas and epiphytes). Within each of these broad groups, plants were grouped based on an assembly of traits that facilitate survival by avoiding, promoting, resisting or tolerating dominant abiotic or biotic limitations to growth (e.g., water, light, herbivory, fire). The potential for humans to modify these abiotic and biotic drivers contributed to the decision to assess plant groups within biomes. E.g., deforestation for agriculture removes light competition constraints, potentially allowing an influx of savanna grasses. By contrast, clearing a savanna environment for agriculture would have much less effect on light availability, and different shifts in plant groups would be anticipated.

Table 1. Broad taxonomic groups encompassing all sub-Saharan African terrestrial vertebrates and vascular plants, with rationale for how they were allocated into species response groups. An expert elicitation process was run for each broad taxonomic group, with participants estimating intactness scores for each response group. Supplementary Table 1 includes details of each response group.

large number of species (~45,000). Because of this limitation, a biogeographical delineation was included in the elicitation process. Experts were asked to provide estimates for the plant species response groups in eight biogeographical units (forest, Caesalpinoid-miombo humid savanna, mixed-acacia savanna, grassland, shrubland, thicket, desert and fynbos). These represent the major sub-Saharan African biomes, with savanna—the most extensive biome—treated as two biomes differentiated by distinct vegetation types. This delineation ensured that only plants present in each biome were considered in the estimation process and in any subsequent data aggregations. Because biome-specific environmental and evolutionary assembly processes act as a strong filter to the set of ‘available traits’, not all plant response groups occur in all biomes (e.g., there are no fire-tolerant forbs in the forest biome). Plant experts were also asked to estimate, within the broad groups of trees and shrubs, forbs and graminoids, the proportion of species within each species response group in each biome (‘Sp_Plant’ spreadsheet in the bii4africa dataset³⁵). These estimates provide a proxy for the distribution of species richness across the plant groups in each biome (e.g., the proportional richness of the graminoid response groups in the desert biome sums to 1). To enable data users to consistently spatialise these biomes, each ecoregion (based on Ecoregions2017© Resolve map³⁶) was allocated to one (or two, if considered a mosaic) of these eight biomes informed by the literature^{37–39} and expert opinion (‘Biome’ spreadsheet in the bii4africa dataset³⁵).

Identifying and describing land use categories. Intactness scores were estimated for nine distinct land uses (Table 2). These land uses with varying intensities of human modification were selected to capture the major land covers, uses and associated activities relevant to sub-Saharan Africa, while being broadly comparable with other land-use maps (e.g., Ellis *et al.*⁴⁰; Goldewijk *et al.*⁴¹; Hurtt *et al.*⁴²). Experts made estimates at a ‘landscape’ scale (i.e., several square kilometres). A finer ‘patch’ scale (i.e., several square metres) would be inappropriate

Land use	Description
Dense urban	Densely built-up environments with high human population densities and limited green space – city centres, dense townships, industrial areas, transformed mining areas (e.g., open cast mines, quarries, dumps). Most ecological processes are highly modified. There are few remaining near-natural patches in the landscape, except for e.g., roadside trees, small parks.
Mixed settlements	Suburban areas, smaller towns and rural settlements with large but fragmented human populations interspersed with gardens, parks and near-natural patches of open space, potentially with low densities of cattle, goats, sheep or chickens, or small-scale croplands.
Non-intensive smallholder croplands	Lands used mainly for smallholder agriculture in small fields (<2 ha), consisting of a diversity of short-duration and long-duration crops (e.g., maize, millet, cassava, beans, squashes, as well as scattered fruit, shade or timber trees). Agricultural inputs of fertilisers and pesticides are very low if any, cultivation is usually manual, there is little or no ploughing or irrigation, and harvest is staggered in time. Fields and homesteads are interspersed with patches of near-natural vegetation. These lands often also support low densities of livestock or smallstock, which are partly free-roaming, and may have semi-natural grazing areas in addition to eating crop residues and cut forage.
Intensive large-scale croplands	Lands used mainly for short duration, monocultural crops in large fields (e.g., staple cereal crops, soybeans, sugar cane). Land use activities usually include several of the following: annual ploughing, inorganic fertiliser application, pesticide application, irrigation, mechanisation. When the crop is harvested, the entire biomass is removed and the next crop is planted, perhaps after a fallow period. There are few remaining near-natural patches in the landscape, except for instance on drainage lines, field boundaries and contour strips, or some woodlots or windbreaks of trees.
Tree crop (fruit) plantations	Lands used mainly for tree crops including fruit-bearing tree or shrub plantations (e.g., bananas, coffee, oil palm, cacao, oranges, vineyards, nuts). Non-transformational harvest, usually only the fruit is taken, and trees may be replaced at some stage. Includes limited remnant forest, riparian or grassland patches between plantation compartments.
Timber plantations	Lands used for growing trees, typically exotic species, for saw timber, poles or pulp. Harvested by clear-cut every 10 to 30 years, and replanted or regrown from coppice. Includes limited remnant forest, riparian or grassland patches between plantation compartments.
Intensive rangelands	Lands used mainly for livestock grazing either with input of fertiliser or pesticide, or with high stock density relative to what the land can sustain (high enough to cause some disturbance or to stop regeneration of vegetation, or to have done so in the recent past). Domesticated stock such as cattle, sheep, goats are typical, but could also include intensive use of indigenous species such as ostrich.
Near-natural lands	Lands (which could be forests, savannas, arid lands, mountainous lands, grasslands) remote from infrastructure, having only minor transformational land use such as crops, planted trees, livestock and human settlements. The human population is relatively low, and livestock or crop-based agriculture or harvest of resources is not at levels that substantially alter natural ecological processes or habitats.
Strictly protected areas	Strictly protected areas that generally do not allow for permanent settlements or resource use, though sometimes allow tourism including limited accommodation and road infrastructure (World Database on Protected Area categories I-III or equivalent). Minimal recent human impact on structure, composition and function of the ecosystem.

Table 2. Descriptions of the nine sub-Saharan African land uses presented to experts to estimate the remaining proportion of an ‘intact’ reference population, for diverse groups of species. Supplementary Table 2 includes representative images of each land use.

for an expert-elicited approach because experts need to consider multiple activities in an area that impact species. For example, a patch of vegetation in a city is likely to have a different impact on a species than a patch of vegetation in a smallholder cropland or a protected area (particularly for large-bodied or wide-ranging species). Instead, the landscape of land covers, uses and activities characteristic of each land use category were described to experts (Table 2), who were asked to consider the collective impact on the abundance of a given group of species within that land use²⁸.

Experts were provided with photographic examples of these land uses to help visualise them and promote consistency in scoring between experts (Supplementary Table 2). For each land use, experts were asked to visualise a specific landscape with which they are familiar that matched the description provided, or several such landscapes across which they could consider an average. Experts were instructed to consider the integrated impact of all characteristics of that landscape (e.g., habitat loss and fragmentation, land cover change, disturbance by people, their domestic animals, infrastructure, pollution, harvesting, persecution, introduced species and diseases, elimination of mutualistic species, altered fire and herbivory regimes) on each species response group. Integrated information at a landscape scale is also likely to be more useful for certain decision-maker needs than fine-scale, species-level data²³.

Structured expert elicitation. The IDEA (‘Investigate’, ‘Discuss’, ‘Estimate’, and ‘Aggregate’) structured expert elicitation protocol²⁵ was used (Fig. 1). It is a modified-Delphi procedure that treats each step as a process of formal data acquisition, incorporating research from mathematics, psychology, and decision theory to help reduce the influence of biases and enhance the transparency, accuracy, and repeatability of the resulting estimates²⁵. The protocol was implemented as follows for each of the broad taxonomic groups:

1. Investigate: A one-hour online meeting was held to introduce the project and explain what was expected of participating experts. Experts had an opportunity to ask questions. The lead expert also presented the draft species response groups and received feedback from the participating experts that, in many cases, led to revisions of these groupings. Afterwards, experts were emailed a recording of the meeting, written instructions, and a survey spreadsheet in which to provide estimates. Experts could provide estimated intactness scores for all species response groups (ranging from six groups for primate experts to 33 for plant experts; Table 1) in all nine land uses (Table 2) or for a subset of species response groups and land uses, depending on the extent of their knowledge. Experts were encouraged to provide any comments relevant

Use	Details
Quantifying ecosystem integrity/condition across space and through time	The first goal of the post-2020 Global Biodiversity Framework is to increase the 'area, connectivity and integrity of natural ecosystems'. The dataset can be used towards assessing ecosystem condition, e.g., mapping the Biodiversity Intactness Index ²⁸ .
Assessing the severity of functional decline for the IUCN Red List of Ecosystems	Aggregated indices of ecosystem health or condition are proposed as one option for quantifying functional decline for the IUCN Red List of Ecosystems ⁵⁴ . The dataset could enable such a quantification.
Quantifying relative population abundance and biodiversity composition indicators	This dataset could be used towards quantifying several composite biodiversity indicators, e.g., Essential Biodiversity Variables ⁵⁵ ; Multidimensional Biodiversity Index ¹² ; Ecosystem Integrity Index ^{26,57} ; Biodiversity Intactness Index ^{28,29} ; and Mean Species Abundance metric (GLOBIO) ^{18,43} . Several of these are also proposed indicators in the Global Biodiversity Framework.
Setting conservation and restoration goals and/or monitoring progress towards these goals	The dataset could be used to assess progress towards restoring 'intactness' in a region. The data could also be used in prioritisation exercises to identify ecosystems for restoration action to maximise improvements in biodiversity intactness.
Assessing the impact of regional development plans	Large-scale infrastructure and agriculture projects are planned across sub-Saharan Africa (e.g., Laurance <i>et al.</i> ⁵⁸). This dataset could be used to predict the impacts of such development plans on biodiversity intactness.
Considering biodiversity sensitivity to development	The data could identify the types of taxa that are particularly sensitive to development, to inform Environmental Impact Assessments and other development plans.
Identifying indicator species groups	Species groups with lower intactness scores are more vulnerable to environmental or developmental change, and monitoring their populations could give early warnings of system degradation.
Assessing trends in how diverse species respond to land use activities	The data could be analysed to test hypotheses and explore trends across species groups and/or land uses.
Species ecological (as opposed to taxonomic) classifications	The species response groups presented in this dataset (Supplementary Table 1) may be useful for a range of applications that require species to be organised into 'functional' (as opposed to purely taxonomic) categories.
Zoonotic disease risk and mitigation assessments	The dataset could be used in identifying and monitoring species groups (and areas, if spatialised) to prevent zoonotic and epizootic disease outbreaks.
Characterising novel ecosystems	Intactness scores >1 depict species groups that respond positively to human land use activities, thus contributing to understanding novel ecosystems ⁵⁹ .
Parameterising, calibrating and validating models of biodiversity in a changing world	Biodiversity models are used to predict biodiversity patterns across space and through time (e.g., Di Marco <i>et al.</i> ⁶⁰ ; Harfoot <i>et al.</i> ⁶¹ ; Schipper <i>et al.</i> ¹⁸) under changing land use conditions. This dataset could be used to parameterise, calibrate, or validate such models.
Climate change research	The approach taken in this paper offers opportunities for natural and/or experimental designs to test interactions of biodiversity, land use and climate change across variable spatial and temporal scales.
Informing future research and training in biodiversity	The species groups and land uses for which there were either few scores or large expert score variability highlight knowledge gaps that require further study. These knowledge gaps could also be used to guide future scientist training efforts.
Comparison with other regions, taxonomic groups or time periods	A similar expert-elicited approach could be used to estimate intactness scores for other regions/taxonomic groups (e.g., invertebrates), allowing for comparison with this dataset. The approach could be repeated in the future to assess how knowledge on land use impacts on biodiversity abundance has changed.
List of biodiversity experts to contact for data, collaboration, etc.	The 200 participating experts (see author list and contributions, and https://bii4africa.org/category/experts/) can serve as points of contact for global initiatives looking to aggregate data or build collaborations.

Table 3. A non-exhaustive list of potential uses of the bii4africa dataset. Supplementary Table 3 includes further details on these uses.

- to each estimate (e.g., land use characteristics that could influence their score, assumptions that they made, uncertainties, the likely score range across species in a group, and any other explanatory information). Experts could also add general reflections to a comments box at the bottom of the spreadsheet. This qualitative information was useful when aggregating the data (step 2), to gain insight into the reasoning behind experts' scoring, and detect potential inconsistencies between experts. Experts were asked not to talk to other participants about their estimates to ensure that they did not influence each other's initial scoring. Experts were encouraged to use other means available to them to inform their scoring, such as talking to colleagues, consulting literature and relevant species reference lists, drawing on experience, and acquiring and interpreting data. They were given two weeks to email their 'Round 1' spreadsheet to the project lead.
2. Discuss: A one to 1.5-hour online meeting was convened, where aggregated (anonymised) results from Round 1 were presented to participating experts. These aggregated results included boxplots showing the range of estimates that experts provided for each response group and land use (and biome, for plants), and 95% confidence interval plots showing trends in mean intactness scores across experts for each response group–land use combination (see examples in the right-hand panels of Figs. 3, 4). Project and expert leads reflected on key trends and sources of variability, and any insights or discrepancies (e.g., when it was apparent from experts' spreadsheet comments that they were interpreting a given land use in different ways). The project lead then facilitated a discussion among the experts, where they were encouraged to share their experiences from Round 1 (e.g., with what did they struggle; what helped them) and to reflect on the aggregated results (e.g., their insights for the species response groups they know well, or any results that surprised them). They were encouraged to discuss outlying (anonymous) expert estimates and why they may have occurred. The project lead emphasised that the purpose of the discussion was not to reach




Fig. 2 Attributes of the 200 participating experts. All values in white font (and black font on the cord plot) represent the number of experts. Numbers do not add up to 200 when categories are not mutually exclusive (region, taxonomic group, employment sector) or when experts did not report a certain attribute (unk = unknown; Mamm = Mammals; org. = organisation).

consensus. Rather, it was to interrogate sources of variability, improve the consistency with which experts were interpreting the species response groups and land uses, and cross-examine reasoning, assumptions and evidence, thereby sharing insights between experts to promote learning²⁵. (We found this discussion meeting to be a particularly useful step in the process, with many experts providing feedback saying that they found it an enjoyable opportunity to learn from each other.)

3. Estimate: An email was sent to participating experts with instructions for Round 2 of the elicitation. This email included a recording of the discussion meeting, written summary of key points, and the Round 1 aggregated results plots. Based on the meeting discussion and summary, experts were asked to revisit their initial scores and independently revise any of these scores if they deemed it necessary (again without discussing their individual scores with other participants). It was emphasised that the objective was not to revise their scores to be closer to the ‘mean’, but rather to revise scores if the discussion gave the expert additional insight that caused them to reconsider their initial estimates (which may or may not result in their revised scores being closer to the mean). Experts had one week to either send revised estimates to the project lead, or to confirm that they did not need to make any revisions.
4. Aggregate: Final ‘Round 2’ estimates were aggregated by calculating the mean expert score and confidence interval across expert scores, for each species response group and landscape (and biome for plants). These




Fig. 3 Intactness scores depicting the remaining proportions of ‘intact’ reference populations of terrestrial vertebrates (tetrapods) in different land uses, where 0 indicates that no individuals remain and 1 indicates the same number of individuals as in an ‘intact’ reference population. Average scores across experts (\pm 95% confidence intervals; CI) are shown. The left panel (a) depicts an aggregated score for each taxonomic group and land use – an average across species response groups, weighted by species richness (i.e., response groups representing a higher number of species in a taxonomic group count more towards its aggregated score). The right panels show examples of the scores for species response groups in two taxonomic groups in different land uses: (b) reptiles in non-intensive, smallholder croplands and (c) bats in intensive rangelands.

final confidence interval plots were shared with experts via email. The average change from Round 1 to 2 in variation between expert estimates was assessed (see *Technical Validation*). The resulting dataset of intactness scores³⁵ includes both individual experts’ Round 2 estimates and associated comments (‘Scores_Raw’ spreadsheet) and aggregated scores across experts (‘Scores_Agg’ spreadsheet).

Data Records

The bii4africa dataset is presented in a multi-spreadsheet .xlsx (Microsoft Excel Spreadsheet) file, which is freely accessible in Figshare³⁵. The raw data spreadsheet (‘Scores_Raw’) includes 31,313 individual expert estimates of the impact of a sub-Saharan African land use (Table 2) on a species response group of terrestrial vertebrates or vascular plants (Table 1). Estimates are reported as intactness scores – the remaining proportion of an ‘intact’ reference (pre-industrial or contemporary wilderness area) population of a species response group in a land use,




Fig. 4 Intactness scores depicting the remaining proportions of ‘intact’ reference populations of terrestrial vascular plants in different land uses, where 0 indicates that no individuals remain and 1 indicates the same number of individuals as in an ‘intact’ reference population. Average scores across experts ($\pm 95\%$ confidence intervals; CI) are shown. The left panel (a) depicts an aggregated score for plants in each land use in each biome – an average across species response groups, weighted by species richness (i.e., response groups representing a higher number of species in a biome count more towards its aggregated score). The right panel (b) shows an example: the scores for plant species response groups in intensive rangelands in the mixed-acacia savanna biome. (Thicket and fynbos biomes are not shown because of low sample sizes: < 3 expert scores across all land uses; USO = underground storage organ).

on a scale from 0 (no individuals remain) through 0.5 (half the individuals remain), to 1 (same as the reference population) and, in limited cases, 2 (two or more times the reference population). For species that thrive in human-modified landscapes, scores could be greater than 1 but not exceeding 2 to avoid extremely large scores biasing aggregation exercises. Such truncation is common in standardised biodiversity metrics^{18,28,43}. Expert comments are included alongside respective estimates.

The raw dataset links, via unique species response group codes, to a spreadsheet (‘Sp_Groups’) describing the species response groups for which experts provided intactness scores (see summary in Table 1). For terrestrial vertebrates, the response group codes also link to a spreadsheet (‘Sp_Vert’) containing all IUCN-listed species in sub-Saharan Africa. Each species has been assigned to one species response group. As large herbivores were initially allocated to response groups but ultimately scored at the species level (see *Identifying and describing species response groups* section), each species has been assigned the appropriate response group code,




Fig. 5 Change in the variation (standard error) of estimated intactness scores for each species response group and land use, between the first and second round of the expert elicitation process. Boxplots show median (horizontal line in the box), interquartile range (box), and max/min values within 1.5 times the interquartile range (vertical lines). Values less than 0 (below the red-dashed horizontal line) show a decrease in score variability between experts. For terrestrial vertebrates, results are shown per taxonomic group; for terrestrial vascular plants they are shown per biome. (C-m = Caesalpinoid-miombo; Mamm = Mammals).

with a unique letter differentiating species (e.g., African elephant and black rhinoceros, both in the ‘megaherbivore’ response group LH1 in Supplementary Table 1, are assigned codes LH1A and LH1B, respectively). For terrestrial vascular plants, the response group codes link to a spreadsheet (‘Sp_Plant’) containing aggregated expert-elicited estimates of the proportion of species in each biome that constitute each response group, within the broad groups of trees and shrubs, forbs and graminoids. Biome names also link to a spreadsheet (‘Biomes’) that includes a list of the 89 mainland Afrotropical ecoregions³⁶, each allocated to one biome (or two biomes if considered a mosaic).

The raw dataset links via unique, anonymous expert codes to a spreadsheet (‘Experts’) listing the region(s) of expertise of each expert (Central Africa, East Africa, Horn of Africa, Southern Africa, West Africa). A summary of the attributes of the 200 participating experts (including their regions of expertise) is provided in Fig. 2. Expert codes are also linked to a spreadsheet (‘Comments’) containing general comments made by experts. The number of intactness scores provided by each expert (‘Scores_Raw’) varied based on the elicitation that they participated in (e.g., primate experts were asked to score six species response groups, while bird experts were asked to score 17 such groups, see Table 1) as well as their extent of expertise (i.e., some experts provided scores for only a subset of species response groups and/or land uses). On average, each expert provided 155 intactness scores.

The number of experts estimating an intactness score for a given species response group in a given land use (and biome for plants) varied from one to 28, with an average (\pm SD) of ten (\pm 7) experts providing independent scores for a particular combination. Arithmetic means are the most widely used form of data aggregation in applications of the IDEA expert elicitation protocol²⁵. For each species response group in each land use (per biome for plants), we report the mean intactness score across experts, the number of experts providing a score (sample size), as well as the variability in scores between experts (standard deviation, standard error, 95% confidence interval; ‘Scores_Agg’ spreadsheet). Figs. 3, 4 visualise the data at two levels of aggregation. The right-hand panels display examples of the aggregated (‘Scores_Agg’) data for reptiles and bats (Fig. 3) and mixed acacia-savanna plants (Fig. 4). These plots depict the variation in how different species response groups within a given taxonomic group are expected to be impacted by a given land use (differences between response group mean scores). They further depict variation between estimates of individual experts (95% confidence intervals around each response group mean score).

The left-hand panels in Figs. 3, 4 display a further level of data aggregation, in which intactness scores for each taxonomic group in each land use are presented. Each score is an average across the associated species response group means and confidence intervals (‘Scores_Agg’), weighted relative to the proportion of species in that response group (‘Sp_Vert’ and ‘Sp_Plant’). In other words, each species in sub-Saharan Africa counts equally in the aggregated intactness score. Such an aggregation could similarly be performed for all species in

a given ecoregion, country, or other spatial unit. The dataset can be downloaded together with an R code script for performing such aggregations.

Notably, the ‘strictly protected areas’ land use does not always have an intactness score of 1 (i.e., equivalent to a pre-industrial ‘intact’ reference population; Figs. 3, 4). These lower scores are because (a) some species benefit from human-related disturbances that would have been present in a pre-industrial landscape but are no longer present in a strictly managed protected area and (b) for particularly large, wide-ranging species, even the best-managed protected area is unlikely to contain a ‘pre-industrial’ reference population (e.g., Prins *et al.*⁴⁴; Balme *et al.*⁴⁵). It is also worth noting that not all protected areas in the region are strictly protected in practice – these protected areas were not considered in this land use category, and intactness scores for such protected areas would likely be closer to the ‘near-natural lands’ land use or even some of the other land use categories (e.g., intensive rangelands or non-intensive smallholder croplands; Table 2), depending on the activities occurring in these areas.

Technical Validation

The reliability of expert judgement will always be sensitive to which experts participate and how questions are asked. Each step of the IDEA structured expert elicitation protocol makes use of procedures that have been demonstrated to improve elicitation rigour²⁵. These include the identification and recruitment of experts, the framing of questions, the two rounds of independent estimation and the aggregation, review and critical appraisal of expert judgements during a facilitated group discussion (see *Methods*).

The purpose of the facilitated group discussion is not to reach consensus²⁵. In our case, experts could disagree on how species are likely to respond to land uses, particularly for lesser-known species groups. Rather, the discussion in a modified-Delphi process aims to resolve linguistic ambiguity, promote critical thinking, share evidence, and improve the consistency with which experts interpret the questions²⁵. Thus, we would expect some but not all of the variability in expert scores to reduce in the second round of the elicitation process, subsequent to the group discussion. The standard error around expert scores for each species response group in each land use (and in each biome for terrestrial vascular plants) was lower in the second, compared with the first, round of the elicitation for 85% and 88% of terrestrial vertebrate and plant intactness scores respectively (Fig. 5). On average, there was a reduction in standard error of 0.01 for vertebrates and 0.03 for vascular plants (Fig. 5). Thus, the variability in estimates between experts was generally lower following the group discussions, indicating that the elicitation process resulted in improved scoring consistency between experts.

We would also expect a reasonable degree of consensus between experts, and therefore for the variability between their scores to be significantly lower than that of chance. To test this expectation, we generated a random estimate within the allowed range (0 to 2) for each Round 2 expert estimate. We then determined the standard error around these random estimates for each species response group and land use (and biome for terrestrial vascular plants). This process ensured comparable sample sizes between expert estimates and random estimates. The standard error in expert estimates was significantly lower than that expected by chance (paired *t* test: $t = -54.58$, $d.f. = 2796$, $p < 0.001$). On average, the standard error in expert estimates in Round 2 was 63% lower than expected by chance (0.075 compared with 0.200). These results suggest that (1) the IDEA protocol served to promote consistency in scoring between experts and (2) experts were significantly more consistent in their scoring than expected by chance.

Validating this dataset using available field-collected data is limited by a spatial and temporal scale mismatch. Most field data are collected at the ‘patch’ scale, while the landscape scale was appropriate for producing this dataset, as explained in the methods. The experts considered diverse land covers and activities characteristic of each land use type (Table 2) to estimate their collective impact on a population in that land use. For example, in an agricultural landscape there is likely to be a higher abundance of fossorial reptiles in the habitat remnants than in the surrounding croplands. This landscape composition was considered by experts when estimating the overall remaining proportion of a reference population of these reptiles in a landscape characterised by non-intensive, smallholder croplands (or intensive, large-scale croplands, with fewer remnant habitat patches). In contrast, field studies tend to report land use at the scale of the habitat or cropland patch. For example, in the PREDICTS database²²—the largest global dataset of species abundances in different land uses—the only relevant cropland data⁴⁶ for Afrotropical birds contains bird counts in ‘private farmhouse gardens surrounded by agricultural matrix’. This ‘patch’ scale, focused on a small subset of land covers and activities in an agricultural landscape, is incompatible with the cropland landscapes that experts considered (Table 2). The spatial and temporal scale of the reference site can also be mismatched. For example, the only relevant plantation data⁴⁷ for Afrotropical birds in the PREDICTS database has a reference site of ‘forest fragments in timber plantations’, which is incompatible with the landscape-scale pre-industrial/large wilderness area reference state that experts considered in this study. While this scale mismatch limits validation using existing field data across multiple regions and taxa, future research aimed at validation could design landscape-scale data collection protocols that are more comparable (i.e., developing landscape-scale data collection protocols with multiple multi-taxa sampling sites evenly distributed across different land uses across the region). However, the absence of a ‘pre-industrial’ reference in field-collected data can still impede comparison¹⁷.

Usage Notes

A non-exhaustive list of potential uses of the bii4africa dataset is provided in Table 3. The data are best suited to broad-scale, multi-species applications, rather than finer-scale applications for which site-specific, field-collected data would be more appropriate. The standardised nature of the intactness scores (0–2 scale) means that the data (or a subset thereof) can be aggregated in several ways to meet a user’s needs, e.g., by taxonomic group, functional type, land use and/or spatial (e.g., biogeographical or political) unit (see R code provided for aggregating

the data³⁵). To assist with data aggregation, scores for response groups can be linked back to individual species for terrestrial vertebrates (noting that large mammal scores are already at species level). Spatial distributions of these species are available from the IUCN Red List²⁶. A species list can thus be obtained for the area of interest, and those species scores then extracted from the bii4africa dataset. While the scores for terrestrial vascular plants are not connected to individual species, scores are provided per biome, meaning the data can be extracted at biome-scale, informed for example by the WWF ecoregion maps³⁶. The proportions of terrestrial vascular plant species in each response group per biome are also provided, enabling the weighting of score aggregations relative to the proportion of species that those scores represent. Data can also be aggregated according to the regional expertise of the contributing experts (e.g., using scores from only West African experts for a West African data application, or for testing differences between regions). For some species groups such as large mammals and birds, regional considerations such as whether bushmeat harvesting is prevalent can have an influence on experts' intactness scores.

While the data presented here are non-spatial, they can be made spatially explicit by linking the scores for different land uses to a map of those land uses (encompassing sub-Saharan Africa or a region therein). As the land uses were selected to reflect those most common in sub-Saharan Africa, the map used to spatialise the data should include those classes (urban, crop, plantation, rangeland, near-natural, protected). See Scholes and Biggs²⁸, Newbold *et al.*²⁹ and Schipper *et al.*¹⁸ for examples of mapping Biodiversity Intactness based on intactness scores for different land uses. Different land use intensities (e.g., dense urban vs mixed settlements; smallholder vs large-scale cropland; rangeland vs near-natural land) could also be mapped using proxies such as percentage urban cover⁴⁸ and population density⁴⁹ in human settlements; percentage crop cover⁵⁰, nitrogen input⁵¹ and field size⁵² in croplands; livestock density⁵³ in rangelands; etc. Importantly, as land use changes across the region, estimates and maps of intactness can be updated using the bii4africa dataset.

Following the IDEA protocol recommendations²⁵, outliers were not removed from the data when determining mean intactness scores across experts ('Scores_Agg' spreadsheet in the dataset). Rather, anonymised outliers were flagged in the discussion meeting, after which experts who provided such scores could reconsider if they were appropriate and revise them if not. Equally weighted data aggregations (i.e., arithmetic means) can be sensitive to outliers in small groups, and we thus recommend careful consideration regarding the use of mean scores that are based on a low number of experts. This consideration is most relevant for large mammals and terrestrial vascular plants – the only groups with mean scores for some species (large mammals) or species response groups (plants) in some land uses based on fewer than the recommended six experts²⁵. With an average of 10 contributing experts per mean score, our dataset has a considerably larger 'sample size' than other similar processes (e.g., $n = 3$)²⁸. We report sample size and standard deviation in our aggregated dataset, as well as provide the raw data, enabling users to assess whether the degree of variability in scores is acceptable for their purposes. We also think the scores with higher variability (i.e., less consensus between experts) could identify important knowledge gaps regarding how species respond to land uses, or important regional differences, thus informing future empirical research.

Code availability

R code for calculating aggregated intactness scores for a focal region (e.g., ecoregion or country) and/or taxonomic group can be downloaded with the bii4africa dataset on Figshare³⁵; see Data Records section.

Received: 29 June 2023; Accepted: 7 December 2023;

Published online: 12 February 2024

References

1. Steffen, W., Broadgate, W., Deutsch, L., Gaffney, O. & Ludwig, C. The trajectory of the anthropocene: The great acceleration. *Anthr. Rev.* **2**, 81–98 (2015).
2. Watson, J. E. M. & Venter, O. Mapping the continuum of humanity's footprint on land. *One Earth* **1**, 175–180 (2019).
3. Maxwell, S. L., Fuller, R. A., Brooks, T. M. & Watson, J. E. M. Biodiversity: The ravages of guns, nets and bulldozers. *Nature* **536**, 143–145 (2016).
4. Dullinger, I. *et al.* Biodiversity models need to represent land-use intensity more comprehensively. *Glob. Ecol. Biogeogr.* **30**, 924–932 (2021).
5. IPBES. *Global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services*. (Secretariat of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem services, Bonn, Germany, 2019).
6. Mace, G. M. *et al.* Approaches to defining a planetary boundary for biodiversity. *Glob. Environ. Chang.* **28**, 289–297 (2014).
7. Steffen, W. *et al.* Planetary boundaries: guiding human development on a changing planet. *Science* **347**, 1259855 (2015).
8. Manning, P. *et al.* Redefining ecosystem multifunctionality. *Nat. Ecol. Evol.* **2**, 427–436 (2018).
9. Otu, A. *et al.* Africa needs to prioritize One Health approaches that focus on the environment, animal health and human health. *Nat. Med.* **27**, 943–946 (2021).
10. Reyers, B. & Selig, E. R. Global targets that reveal the social–ecological interdependencies of sustainable development. *Nat. Ecol. Evol.* **4**, 1011–1019 (2020).
11. Mace, G. M. *et al.* Aiming higher to bend the curve of biodiversity loss. *Nat. Sustain.* **1**, 448–451 (2018).
12. Soto-Navarro, C. A. *et al.* Towards a multidimensional biodiversity index for national application. *Nat. Sustain.* **4**, 933–942 (2021).
13. Hortal, J. *et al.* Seven Shortfalls that Beset Large-Scale Knowledge of Biodiversity. *Annu. Rev. Ecol. Syst.* **46**, 523–549 (2015).
14. Stephenson, P. J. & Stengel, C. An inventory of biodiversity data sources for conservation monitoring. *PLoS One* **15**, e0242923 (2020).
15. Tydecks, L., Jeschke, J. M., Wolf, M., Singer, G. & Tockner, K. Spatial and topical imbalances in biodiversity research. *PLoS One* **13**, e0199327 (2018).
16. McRae, L., Deinet, S. & Freeman, R. The diversity-weighted living planet index: Controlling for taxonomic bias in a global biodiversity indicator. *PLoS One* **12**, e0169156 (2017).
17. McNellie, M. J. *et al.* Reference state and benchmark concepts for better biodiversity conservation in contemporary ecosystems. *Glob. Chang. Biol.* **26**, 6702–6714 (2020).
18. Schipper, A. *et al.* Projecting terrestrial biodiversity intactness with GLOBIO 4. *Glob. Chang. Biol.* **26**, 760–771 (2020).

19. Tittensor, D. P. *et al.* A mid-term analysis of progress toward international biodiversity targets. *Science* **346**, 241–248 (2014).
20. Newbold, T. *et al.* Global effects of land use on local terrestrial biodiversity. *Nature* **520**, 45–50 (2015).
21. Oliver, T. H. *et al.* Declining resilience of ecosystem functions under biodiversity loss. *Nat. Commun.* **6**, 1–8 (2015).
22. Hudson, L. N. *et al.* The database of the PREDICTS (Projecting Responses of Ecological Diversity In Changing Terrestrial Systems) project. *Ecol. Evol.* **4**, 4701–4735 (2016).
23. Stephenson, P. J. *et al.* Unblocking the flow of biodiversity data for decision-making in Africa. *Biol. Conserv.* **213**, 335–340 (2017).
24. IPBES. *The IPBES regional assessment report on biodiversity and ecosystem services for Africa*. (Secretariat of the Intergovernmental Science–Policy Platform on Biodiversity and Ecosystem services, Bonn, Germany, 2018).
25. Hemming, V., Burgman, M. A., Hanea, A. M., McBride, M. F. & Wintle, B. C. A practical guide to structured expert elicitation using the IDEA protocol. *Methods Ecol. Evol.* **9**, 169–180 (2018).
26. IUCN. *The IUCN Red List of Threatened Species. Version 2022-1*. (2022).
27. Pörtner, H.-O. *et al.* *Climate change 2022: Impacts, adaptation, and vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. (2022).
28. Scholes, R. J. & Biggs, R. A biodiversity intactness index. *Nature* **434**, 45–49 (2005).
29. Newbold, T. *et al.* Has land use pushed terrestrial biodiversity beyond the planetary boundary? A global assessment. *Science* **351**, 600–604 (2016).
30. Martin, P. A., Green, R. E. & Balmford, A. The biodiversity intactness index may underestimate losses. *Nat. Ecol. Evol.* **3**, 862–863 (2019).
31. Trisos, C. H., Auerbach, J. & Katti, M. Decoloniality and anti-oppressive practices for a more ethical ecology. *Nat. Ecol. Evol.* **5**, 1205–1212 (2021).
32. Krueger, T., Page, T., Hubacek, K., Smith, L. & Hiscock, K. The role of expert opinion in environmental modelling. *Environ. Model. Softw.* **36**, 4–18 (2012).
33. de Bello, F. *et al.* Towards an assessment of multiple ecosystem processes and services via functional traits. *Biodivers. Conserv.* **19**, 2873–2893 (2010).
34. Archibald, S., Hempson, G. P. & Lehmann, C. A unified framework for plant life-history strategies shaped by fire and herbivory. *New Phytol.* **224**, 1490–1503 (2019).
35. Clements, H. S. *et al.* Faunal and floral population intactness estimates across sub-Saharan African land uses: The bii4africa dataset, <https://doi.org/10.6084/m9.figshare.c.6710463.v1> (2023).
36. Olson, D. M. *et al.* Terrestrial Ecoregions of the World: A New Map of Life on Earth. *Bioscience* **51**, 933 (2001).
37. Armani, M. *et al.* *Enhancing climate change adaptation and mitigation actions on land in Africa*. (Future Ecosystems for Africa, University of the Witwatersrand, Johannesburg, South Africa, 2022).
38. Sankaran, M. *et al.* Determinants of woody cover in African savannas. *Nature* **438**, 846–849 (2005).
39. White, F. The vegetation of Africa, a descriptive memoir to accompany the UNESCO/AETFAT/UNSO vegetation map of Africa. UNESCO. *Nat. Resour. Res.* **20**, 1–356 (1983).
40. Ellis, E. C., Beusen, A. H. W. & Goldewijk, K. K. Anthropogenic biomes: 10,000 BCE to 2015 CE. *Land* **9**, 8–10 (2020).
41. Goldewijk, K. K., Beusen, A., Doelman, J. & Stehfest, E. Anthropogenic land use estimates for the Holocene - HYDE 3.2. *Earth Syst. Sci. Data* **9**, 927–953 (2017).
42. Hurtt, G. *et al.* Harmonization of Global Land-Use Change and Management for the Period 850–2100 (LUH2) for CMIP6. *Geosci. Model Dev.* **13**, 5425–5464 (2020).
43. Alkemade, R. *et al.* GLOBIO3: A framework to investigate options for reducing global terrestrial biodiversity loss. *Ecosystems* **12**, 374–390 (2009).
44. Prins, H. H. & de Jong, J. F. The Ecohistory of Tanzania's Northern Rift Valley – Can One Establish an Objective Baseline as an Endpoint for Ecosystem Restoration? in *Tarangire: Human-Wildlife Coexistence in a Fragmented Ecosystem* 129–161 (Springer International Publishing, 2022).
45. Balme, G. A., Slotow, R. & Hunter, L. T. B. Edge effects and the impact of non-protected areas in carnivore conservation: leopards in the Phinda-Mkhuzi Complex, South Africa. *Anim. Conserv.* **13**, 315–323 (2010).
46. Neuschulz, E. L., Botzat, A. & Farwig, N. Effects of forest modification on bird community composition and seed removal in a heterogeneous landscape in South Africa. *Oikos* **120**, 1371–1379 (2011).
47. Farwig, N., Sajita, N. & Böhning-Gaese, K. Conservation value of forest plantations for bird communities in western Kenya. *For. Ecol. Manage.* **255**, 3885–3892 (2008).
48. Pesaresi, M. & Politis, P. GHS-BUILT-S R2023A - GHS built-up surface grid, derived from Sentinel2 composite and Landsat, multitemporal (1975–2030). *Joint Research Centre Data Catalogue* <https://doi.org/10.2905/9F06F36F-4B11-47EC-ABB0-4F8B7B1D72EA> (2023).
49. Schiavina, M., Freire, S., Carioli, A. & MacManus, K. GHS-POP R2023A - GHS population grid multitemporal (1975–2030). *Joint Research Centre Data Catalogue* <https://doi.org/10.2905/2FF68A52-5B5B-4A22-8F40-C41DA8332CFE> (2023).
50. Potapov, P. *et al.* Global maps of cropland extent and change show accelerated cropland expansion in the twenty-first century. *Nat. Food* **3**, 19–28 (2022).
51. Lu, C. & Tian, H. Global nitrogen and phosphorus fertilizer use for agriculture production in the past half century: Shifted hot spots and nutrient imbalance. *Earth Syst. Sci. Data* **9**, 181–192 (2017).
52. Lesiv, M. *et al.* Estimating the global distribution of field size using crowdsourcing. *Glob. Chang. Biol.* **25**, 174–186 (2019).
53. Gilbert, M. *et al.* Global distribution data for cattle, buffaloes, horses, sheep, goats, pigs, chickens and ducks in 2010. *Sci. Data* **5**, 1–11 (2018).
54. Keith, D. A. *et al.* The IUCN red list of ecosystems: Motivations, challenges, and applications. *Conserv. Lett.* **8**, 214–226 (2015).
55. Jetz, W. *et al.* Essential biodiversity variables for mapping and monitoring species populations. *Nat. Ecol. Evol.* **3**, 539–551 (2019).
56. Hill, S. L. L. *et al.* The Ecosystem Integrity Index: a novel measure of terrestrial ecosystem integrity with global coverage. *bioRxiv* 2022.08 (2022).
57. Hansen, A. J. *et al.* Towards monitoring forest ecosystem integrity within the Post-2020 Global Biodiversity Framework. *Conserv. Lett.* **14**, e12822 (2021).
58. Laurance, W. F., Sloan, S., Weng, L. & Sayer, J. A. Estimating the Environmental Costs of Africa's Massive 'Development Corridors'. *Curr. Biol.* **25**, 3202–3208 (2015).
59. Hobbs, R. J., Higgs, E. & Harris, J. A. Novel ecosystems: implications for conservation and restoration. *Trends Ecol. Evol.* **24**, 599–605 (2009).
60. Di Marco, M. *et al.* Projecting impacts of global climate and land-use scenarios on plant biodiversity using compositional-turnover modelling. *Glob. Chang. Biol.* **25**, 2763–2778 (2019).
61. Harfoot, M. B. J. *et al.* Emergent Global Patterns of Ecosystem Structure and Function from a Mechanistic General Ecosystem Model. *PLoS Biol.* **12**, e1001841 (2014).

Acknowledgements

This work was funded by a Jennifer Ward Oppenheimer Research Grant.

Author contributions

H.S.C. led project conceptualisation and the expert elicitation processes, aggregated the data, ran the workshops, and drafted the manuscript. E.D.L.S. (small carnivores), G.H. (large mammals and graminoids), B.L. (primates), B.Ma. (reptiles and amphibians), A.Mon. (bats, rodents, insectivores), F.S. (forbs), C.R. (birds), and N.S. (trees and shrubs) contributed as lead experts: conceptualising and leading an elicitation process for their respective taxonomic groups, contributing their own independent expert estimates, as well as giving input during a feedback workshop and on the draft paper. R.Bi. and A.D.V. contributed to initial project conceptualisation, provided input during an inception and two feedback workshops, and provided input on the draft paper. R.Bl., M.Ch., K.J.E., M.Ham., B.R. and O.S. provided input during an inception and feedback workshop, and input on the draft manuscript. T.T. provided input during an inception and feedback workshop and assisted with data compilation. T.L. provided feedback on data structure, wrote the data aggregation code, and provided input during a feedback workshop. All other authors contributed as participating experts (see *Structured expert elicitation* section) and read and approved the draft manuscripts, with M.F.B., E.B., F.C., M.Co., P.M.K.C., C.De., F.D., M.D., L.G., P.Ga., J.S.G., M.Hau., C.K., M.T.K., A.I.R., A.T.K.L., A.M.L., K.Z.M., L.M., E.E.N., K.N., A.J.P., F.P., R.J.P., F.G.T.R., T.R., J.T.S., J.A.S., M.J.S., I.T., A.A.T., J.T.d.T., K.A.T., V.V.C., H.vd.M., J.A.V., N.W. and P.A.W. also providing input on the draft manuscript.

Competing interests

All authors declare that they have no competing interests as defined by Nature Research, or other interests that might be perceived to influence the results and/or discussion reported in this paper.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41597-023-02832-6>.

Correspondence and requests for materials should be addressed to H.S.C.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2024

Hayley S. Clements^{1,2}✉, Emmanuel Do Linh San³, Gareth Hempson^{4,5}, Birthe Linden⁶, Bryan Maritz⁷, Ara Monadjem^{8,9}, Chevonne Reynolds¹⁰, Frances Siebert¹¹, Nicola Stevens¹², Reinette Biggs^{1,13}, Alta De Vos^{1,14}, Ryan Blanchard^{1,15}, Matthew Child¹⁶, Karen J. Esler¹⁷, Maike Hamann^{1,18}, Ty Loft¹⁹, Belinda Reyers²⁰, Odirilwe Selomane^{1,21}, Andrew L. Skowno^{16,22}, Tshogofatso Tshoke^{1,17}, Diarrassouba Abdoulaye²³, Thierry Aebischer²⁴, Jesús Aguirre-Gutiérrez²⁵, Graham J. Alexander¹⁰, Abdullahi H. Ali²⁶, David G. Allan²⁷, Esther E. Amoako²⁸, Samuel Angedakin²⁹, Edward Aruna³⁰, Nico L. Avenant^{31,32}, Gabriel Badjedjea³³, Adama Bakayoko³⁴, Abraham Bamba-kaya³⁵, Michael F. Bates^{36,37}, Paul J. J. Bates³⁸, Steven R. Belmain³⁹, Emily Bennitt⁴⁰, James Bradley⁴¹, Chris A. Brewster⁴², Michael B. Brown⁴³, Michelle Brown⁴⁴, Josef Bryja⁴⁵, Thomas M. Butynski⁴⁶, Filipe Carvalho^{3,47}, Alan Channing¹¹, Colin A. Chapman⁴⁸, Callan Cohen⁴⁹, Marina Cords⁵⁰, Jennifer D. Cramer⁵¹, Nadine Cronk¹⁰, Pamela M. K. Cunneyworth⁵², Fredrik Dalerum^{9,53,54}, Emmanuel Danquah⁵⁵, Harriet T. Davies-Mostert^{9,56}, Andrew D. de Blocq⁵⁷, Yvonne A. De Jong⁴⁶, Terrence C. Demos⁵⁸, Christiane Denys⁵⁹, Chabi A. M. S. Djagoun⁶⁰, Thomas M. Doherty-Bone⁶¹, Marine Drouilly^{62,63,64}, Johan T. du Toit^{65,66}, David A. Ehlers Smith⁶⁷, Yvette C. Ehlers Smith^{67,68}, Seth J. Eiseb⁶⁹, Peter J. Fashing⁷⁰, Adam W. Ferguson⁷¹, José M. Fernández-García⁷², Manfred Finckh⁷³, Claude Fischer⁷⁴, Edson Gandiwa⁷⁵, Philippe Gaubert⁷⁶, Jerome Y. Gaugris⁷⁷, Dalton J. Gibbs⁷⁸, Jason S. Gilchrist⁷⁹, Jose M. Gil-Sánchez⁸⁰, Anthony N. Githitho⁸¹, Peter S. Goodman⁸², Laurent Granjon⁸³, J. Paul Grobler⁸⁴, Bonginkosi C. Gumbi⁸⁵, Vaclav Gvozdk^{45,86}, James Harvey⁸⁷, Morgan Hauptfleisch⁸⁸, Firas Hayder³, Emmanuel M. Hema⁸⁹, Marna Herbst⁹⁰, Mariano Hougbedji⁹¹, Brian J. Huntley⁹², Rainer Hutterer⁹³, Samuel T. Ivande⁹⁴, Kate Jackson⁹⁵, Gregory F. M. Jongsma⁹⁶, Javier Juste⁹⁷, Blaise Kadjo⁹⁸, Prince K. Kaleme⁹⁹, Edwin Kamugisha¹⁰⁰, Beth A. Kaplin¹⁰¹,

Humphrey N. Kato¹⁰², Christian Kiffner^{103,104}, Duncan M. Kimuyu¹⁰⁵, Robert M. Kityo¹⁰⁶, N'goran G. Kouame¹⁰⁷, Marcel Kouete¹⁰⁸, Aliza le Roux¹⁰⁹, Alan T. K. Lee¹¹⁰, Mervyn C. Lötter¹⁰, Anne Mette Lykke¹¹¹, Duncan N. MacFadyen¹¹², Gacheru P. Macharia¹¹³, Zimkitha J. K. Madikiza¹⁰, Them'b'alilahlwa A. M. Mahlaba⁸, David Mallon¹¹⁴, Mnqobi L. Mamba⁸, Claude Mande¹¹⁵, Rob A. Marchant¹¹⁶, Robin A. Maritz^{7,117}, Wanda Markotter¹¹⁸, Trevor McIntyre¹¹⁹, John Measey^{120,121,122}, Addisu Mekonnen¹²³, Paulina Meller⁷³, Haemish I. Melville¹²⁴, Kevin Z. Mganga¹²⁵, Michael G. L. Mills¹²⁶, Liaan Minnie^{126,127}, Alain Didier Missouf¹²⁸, Abubakr Mohammad¹²⁹, Nancy N. Moinde¹³⁰, Bakwo Fils E. Moise¹³¹, Pedro Monterroso^{132,133,134}, Jennifer F. Moore¹³⁵, Simon Musila¹³⁶, Sedjro Gilles A. Nago¹³⁷, Maganizo W. Namoto¹³⁸, Fatimata Niang¹³⁹, Violaine Nicolas¹⁴⁰, Jerry B. Nkenku¹⁴¹, Evans E. Nkrumah⁵⁵, Gonwouo L. Nono¹⁴², Mulawwa M. Norbert¹⁴³, Katarzyna Nowak¹⁴⁴, Benneth C. Obitte^{145,146}, Arnold D. Okoni-Williams¹⁴⁷, Jonathan Onongo¹⁴⁸, M. Justin O'Riain¹⁴⁹, Samuel T. Osinubi¹⁴⁴, Daniel M. Parker¹²⁶, Francesca Parrini¹⁰, Mike J. S. Peel^{10,150,151}, Johannes Penner^{152,153}, Darren W. Pietersen⁹, Andrew J. Plumptre¹⁵⁴, Damian W. Ponsoby¹⁰, Stefan Porembski¹⁵⁵, R. John Power¹⁵⁶, Frans G. T. Radloff¹⁵⁷, Ramugondo V. Rambau¹⁵⁸, Tharmalingam Ramesh¹⁵⁹, Leigh R. Richards¹⁶⁰, Mark-Oliver Rödel¹⁶¹, Dominic P. Rollinson⁴⁹, Francesco Rovero¹⁶², Mostafa A. Saleh¹⁶³, Ute Schmiedel, M. Corrie Schoeman¹⁶⁴, Paul Scholte¹⁶⁵, Thomas L. Serfass¹⁶⁶, Julie Teresa Shapiro¹⁶⁷, Sidney Shema¹⁶⁸, Stefan J. Siebert¹¹, Jasper A. Slingsby^{15,169}, Alexander Sliwa¹⁷⁰, Hanneline A. Smit-Robinson^{171,172}, Etotepe A. Sogbohossou¹⁷³, Michael J. Somers¹⁷⁴, Stephen Spawls¹⁷⁵, Jarryd P. Streicher⁶⁷, Lourens Swanepoel¹⁷⁶, Irero Tanshi^{145,177}, Peter J. Taylor¹⁰⁹, William A. Taylor¹⁷⁸, Mariska te Beest^{127,179}, Paul T. Telfer¹⁸⁰, Dave I. Thompson^{10,181}, Elie Tobi¹⁸², Krystal A. Tolley¹⁶, Andrew A. Turner^{183,184}, Wayne Twine¹⁰, Victor Van Cakenberghe^{185,186}, Frederik Van de Perre¹⁸⁷, Helga van der Merwe^{22,188}, Chris J. G. van Niekerk¹⁸⁹, Pieter C. V. van Wyk¹⁹⁰, Jan A. Venter¹⁹¹, Luke Verburgt⁶⁶, Geraldine Veron¹⁹², Susanne Vetter¹⁹³, Maria S. Vorontsova¹⁹⁴, Thomas C. Wagner¹⁹⁵, Paul W. Webala¹⁹⁶, Natalie Weber^{197,198}, Sina M. Weier¹⁹⁹, Paula A. White²⁰⁰, Melissa A. Whitecross^{10,201}, Benjamin J. Wigley^{202,203,204}, Frank J. Willems²⁰⁵, Christiaan W. Winterbach²⁰⁶ & Galena M. Woodhouse²⁰⁷

¹Centre for Sustainability Transitions, Stellenbosch University, Stellenbosch, South Africa. ²Helsinki Lab of Interdisciplinary Conservation Science, University of Helsinki, Helsinki, Finland. ³Department of Zoology and Entomology, University of Fort Hare, Alice, South Africa. ⁴Centre for African Ecology, School of Animal, Plant and Environmental Sciences, University of the Witwatersrand, Johannesburg, South Africa. ⁵Institute of Biodiversity, One Health and Veterinary Medicine, University of Glasgow, Glasgow, Scotland, United Kingdom. ⁶Chair in Biodiversity Value & Change, Faculty of Science, Engineering & Agriculture, University of Venda, Thohoyandou, South Africa. ⁷Department of Biodiversity and Conservation Biology, University of the Western Cape, Bellville, South Africa. ⁸Biological Sciences, University of Eswatini, Kwaluseni, Eswatini. ⁹Mammal Research Institute, Department of Zoology and Entomology, University of Pretoria, Pretoria, South Africa. ¹⁰School of Animal, Plant and Environmental Sciences, University of the Witwatersrand, Johannesburg, South Africa. ¹¹Unit for Environmental Sciences and Management, North-West University, Potchefstroom, South Africa. ¹²Environmental Change Institute, University of Oxford, Oxford, United Kingdom. ¹³Stockholm Resilience Centre, Stockholm University, Stockholm, Sweden. ¹⁴Department of Environmental Sciences, Rhodes University, Makhanda, South Africa. ¹⁵Fynbos Node of the South African Environmental Observation Network, Cape Town, South Africa. ¹⁶South African National Biodiversity Institute, Cape Town, South Africa. ¹⁷Department of Conservation Ecology & Entomology, Stellenbosch University, Stellenbosch, South Africa. ¹⁸Centre for Geography and Environmental Science, University of Exeter, Penryn, Cornwall, United Kingdom. ¹⁹School of Geography and the Environment, Environmental Change Institute, University of Oxford, Oxford, United Kingdom. ²⁰Centre for Environmental Studies, University of Pretoria, Pretoria, South Africa. ²¹Department of Agricultural Economics, Extension and Rural Development, University of Pretoria, Pretoria, South Africa. ²²Department of Biological Sciences, University of Cape Town, Cape Town, South Africa. ²³Office Ivoirien des Parcs et Réserves, Soubré, Côte d'Ivoire. ²⁴African Parks, Bangui, Central African Republic. ²⁵Environmental Change Institute, School of Geography and the Environment, University of Oxford, Oxford, United Kingdom. ²⁶Hirola Conservation Programme, Garissa, Kenya. ²⁷Bird Department, Durban Natural Science Museum, Durban, South Africa. ²⁸Department of Environmental and Sustainability Sciences, University for Development Studies, Tamale, Ghana. ²⁹Department of Environmental Management, Makerere University, Kampala, Uganda. ³⁰Biodiversity Conservation, Reptile and Amphibian Program - Sierra Leone, Freetown, Sierra Leone. ³¹Department of Mammalogy, National Museum, Bloemfontein, South Africa. ³²Centre for Environmental Management, University of the Free State, Bloemfontein, South Africa. ³³Aquatic Ecology, University of Kisangani/Biodiversity Monitoring Center, Kisangani, Democratic Republic of the Congo. ³⁴UFR Sciences de la Nature, Université Nanguil Abrogoua, Abidjan, Côte d'Ivoire. ³⁵Institut de Recherches Agronomiques et Forestières (IRAF), Centre National de la Recherche Scientifique et Technologique (CENAREST), Libreville, Gabon. ³⁶Department of Animal and Plant Systematics, National Museum, Bloemfontein, South Africa. ³⁷Department of Zoology & Entomology, University of the Free State, Bloemfontein, South Africa. ³⁸Harrison Institute, Sevenoaks, Kent, United Kingdom. ³⁹Agriculture, Health and

Environment, Natural Resources Institute, University of Greenwich, Chatham, Maritime, United Kingdom. ⁴⁰Okavango Research Institute, University of Botswana, Maun, Botswana. ⁴¹Kalahari Research and Conservation, Botswana, Botswana. ⁴²Independent, Gaborone, Botswana. ⁴³Giraffe Conservation Foundation, Windhoek, Namibia. ⁴⁴Department of Anthropology, University of Minnesota - Twin Cities, Minneapolis, MN, USA. ⁴⁵Institute of Vertebrate Biology of the Czech Academy of Sciences, Brno, Czech Republic. ⁴⁶Eastern Africa Primate Diversity and Conservation Program, Nanyuki, Kenya. ⁴⁷BIOPOLIS-CIBIO/InBIO, University of Porto, Porto, Portugal. ⁴⁸Biology, Vancouver Island University, Nanaimo, Canada. ⁴⁹FitzPatrick Institute of African Ornithology, University of Cape Town, Rondebosch, South Africa. ⁵⁰Department of Ecology, Evolution & Environmental Biology, Columbia University, New York, NY, USA. ⁵¹ROC USA, Concord, USA. ⁵²Colobus Conservation, Diani, Kenya. ⁵³Biodiversity Research Institute (CSIC-UO-PA), Mieres, Spain. ⁵⁴Department of Zoology, Stockholm University, Stockholm, Sweden. ⁵⁵Department of Wildlife and Range Management, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana. ⁵⁶Conserve Global, London, United Kingdom. ⁵⁷BirdLife South Africa, Johannesburg, South Africa. ⁵⁸Negaunee Integrative Research Center, The Field Museum, Chicago, United States of America. ⁵⁹Institut de Systématique, Evolution, Biodiversité (ISYEB), Muséum national d'Histoire naturelle, CNRS, Sorbonne Université, EPHE, Université des Antilles, Paris, France. ⁶⁰Faculty of Agronomic Sciences, Laboratory of Applied Ecology, University of Abomey Calavi, Cotonou, Benin. ⁶¹Conservation Programs, Royal Zoological Society of Scotland, Edinburgh, United Kingdom. ⁶²Institute for Communities and Wildlife in Africa (iCWild), University of Cape Town, Cape Town, South Africa. ⁶³Centre for Social Science Research (CSSR), University of Cape Town, Cape Town, South Africa. ⁶⁴Panthera, New York, USA. ⁶⁵Institute of Zoology, Zoological Society of London, London, United Kingdom. ⁶⁶Department of Zoology and Entomology, University of Pretoria, Pretoria, South Africa. ⁶⁷Centre for Functional Biodiversity, School of Life Sciences, University of KwaZulu-Natal, Pietermaritzburg, South Africa. ⁶⁸Ezemvelo KZN Wildlife, Pietermaritzburg, South Africa. ⁶⁹Department of Environmental Science, School of Science, University of Namibia, Windhoek, Namibia. ⁷⁰Anthropology Department & Environmental Studies Program, California State University Fullerton, Fullerton, United States of America. ⁷¹Gantz Family Collection Center, Field Museum of Natural History, Chicago, USA. ⁷²Hazi Foundation, Arcaute, Spain. ⁷³Institute of Plant Science and Microbiology, Universität Hamburg, Hamburg, Germany. ⁷⁴Nature Management, University of Applied Sciences of Western Switzerland, Geneva, Jussy, Switzerland. ⁷⁵Scientific Services, Zimbabwe Parks and Wildlife Management Authority, Harare, Zimbabwe. ⁷⁶Laboratoire Evolution et Diversité Biologique, IRD/CNRS/UPS, Université Toulouse III Paul Sabatier, Toulouse, cedex, 9, France. ⁷⁷Flora Fauna & Man, Ecological Services Limited, Tortola, British Virgin Islands. ⁷⁸Independent, Cape Town, South Africa. ⁷⁹School of Applied Sciences, Edinburgh Napier University, Edinburgh, Scotland, UK. ⁸⁰Departamento de Zoología, Universidad de Granada, Granada, Spain. ⁸¹Centre for Biodiversity, National Museums of Kenya, Nairobi, Kenya. ⁸²Wildlife Conservation Solutions, Mkuze, South Africa. ⁸³CBGP, IRD, CIRAD, INRAE, Institut Agro, University of Montpellier, Montpellier, France. ⁸⁴Genetics, University of the Free State, Bloemfontein, South Africa. ⁸⁵Evolution, Ecology, and Organismal Biology, University of California, Riverside, Riverside, CA, USA. ⁸⁶Department of Zoology, National Museum of the Czech Republic, Prague, Czech Republic. ⁸⁷Harvey Ecological, Howick, South Africa. ⁸⁸Biodiversity Research Centre, Namibia University of Science and Technology, Windhoek, Namibia. ⁸⁹Unité de Formation et de Recherche en Sciences Appliquées et Technologies (UFR-SAT), Université de Dédougou, Dédougou, Burkina Faso. ⁹⁰Conservation Services, South African National Parks, Pretoria, South Africa. ⁹¹Organisation pour le Développement Durable et la Biodiversité, Cotonou, Benin. ⁹²CIBIO-Centro de Investigação em Biodiversidade e Recursos Genéticos, University of Porto, Vairao, Portugal. ⁹³LIB Museum Koenig, Bonn, Germany. ⁹⁴A.P. Leventis Ornithological Research Institute (APLORI), University of Jos, Jos, Nigeria. ⁹⁵Biology Department, Whitman College, Walla Walla, WA, USA. ⁹⁶Department of Natural History, New Brunswick Museum, Saint John, Canada. ⁹⁷Evolutionary Biology, Estación Biológica de Doñana (CSIC), Seville, Spain; CIBER, CIBERESP, Madrid, Spain. ⁹⁸Natural habitats and biodiversity management, University Félix Houphouët-Boigny, Abidjan, Côte d'Ivoire. ⁹⁹Department of Biology, CRSN/ LWIRO, DS Bukavu, DR Congo, Bukavu, Democratic Republic of the Congo. ¹⁰⁰Nature Tanzania, Arusha, Tanzania. ¹⁰¹Center of Excellence in Biodiversity and Natural Resource Management, University of Rwanda, Huye, Rwanda. ¹⁰²Biology, Mbarara University of Science and Technology, Mbarara, Uganda. ¹⁰³Department of Human Behavior, Ecology and Culture, Max Planck Institute for Evolutionary Anthropology, Leipzig, Germany. ¹⁰⁴Department of Anthropology, University of California, Los Angeles, USA. ¹⁰⁵Department of Natural Resources, Karatina University, Karatina, Kenya. ¹⁰⁶Zoology, Entomology and Fisheries Sciences, Makerere University, Kampala, Uganda. ¹⁰⁷UFR Environnement, Laboratoire de Biodiversité et Ecologie Tropicale, Université Jean Lorougnon Guédé, Daloa, Côte d'Ivoire. ¹⁰⁸Department of Natural History, Florida Museum of Natural History, University of Florida, Gainesville, USA. ¹⁰⁹Zoology and Entomology, University of the Free State, Qwaqwa campus, Phuthaditjhaba, South Africa. ¹¹⁰School of Life Sciences, University of KwaZulu-Natal, Scottsville, South Africa. ¹¹¹Department of Ecoscience, Aarhus University, Aarhus C, Denmark. ¹¹²Research and Conservation, Oppenheimer Generations, Parktown, Johannesburg, South Africa. ¹¹³Species and Sites Program, Nature Kenya, Nairobi, Kenya. ¹¹⁴Department of Natural Sciences, Manchester Metropolitan University, Manchester, United Kingdom. ¹¹⁵Department of Ecology and Wildlife Management, University of Kisangani, Kisangani, Democratic Republic of the Congo. ¹¹⁶York institute for Tropical Ecosystems, University of York, York, United Kingdom. ¹¹⁷Conservation Alpha, Cape Town, South Africa. ¹¹⁸Department of Microbiology and Plant Pathology, University of Pretoria, Pretoria, South Africa. ¹¹⁹Department of Life and Consumer Sciences, University of South Africa, Roodepoort, South Africa. ¹²⁰Centre for Invasion Biology, Department of Botany and Zoology, Stellenbosch University, Stellenbosch, South Africa. ¹²¹Centre for Invasion Biology, Institute of Biodiversity, Yunnan University, Kunming, UMR7179, China. ¹²²MECADEV CNRS/MNHN, Département Adaptations du Vivant, Muséum National d'Histoire Naturelle, Bâtiment d'Anatomie Comparée, Paris, France. ¹²³Department of Anthropology and Archaeology, University of Calgary, Calgary, Canada. ¹²⁴Department of Environmental Sciences, University of South Africa, Florida, South Africa. ¹²⁵Copernicus Institute of Sustainable Development, Utrecht University, Utrecht, The Netherlands. ¹²⁶School of Biology and Environmental Science, University of Mpumalanga, Mbombela, South Africa. ¹²⁷Centre for African Conservation Ecology, Nelson Mandela University, Gqeberha, South Africa. ¹²⁸Faculty of Science, Laboratory of Biology and Physiology of Animal

Organisms, Zoology Unit, University of Douala, Douala, Cameroon. ¹²⁹Researcher, Conflict and Environmental Observatory, Manchester, United Kingdom. ¹³⁰Conservation Biology, Institute of Primate Research-National Museums of Kenya, Nairobi, Kenya. ¹³¹Environmental Sciences, University of Ebolowa, Ebolowa, Cameroon. ¹³²Wildlife Conservation Ecology Research Group, CIBIO/InBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, Universidade do Porto, Vairã, Portugal. ¹³³BIOPOLIS Program in Genomics, Biodiversity and Land Planning, CIBIO, Campus de Vairão, Vairão, Portugal. ¹³⁴African Parks, Johannesburg, South Africa. ¹³⁵Moore Ecological Analysis and Management, LLC, Gainesville, USA. ¹³⁶Mammalogy Section-Department of Zoology, National Museums of Kenya, Nairobi, Kenya. ¹³⁷Laboratoire d'Ecologie, de Botanique et de Biologie végétale, University of Parakou, Parakou, Benin. ¹³⁸Indigenous Woodland Strategy Area, Forestry Research Institute of Malawi, Zomba, Malawi. ¹³⁹Institute of Environmental Sciences, Faculty of Technology and Sciences, University Cheikh Anta Diop de Dakar, Dakar, Sénégal. ¹⁴⁰Institut de Systématique, Evolution, Biodiversité (ISYEB), Muséum national d'Histoire naturelle, CNRS, Sorbonne Université, EPHE, Université des Antilles, Paris, France. ¹⁴¹Departement of Biology, Faculty of Science, University of Kinshasa, Kinshasa, Democratic Republic of the Congo. ¹⁴²Department of Animal Biologie and Physiologie, University of Yaounde I, Yaounde, Cameroon. ¹⁴³Primates, Center for Research in Ecology and Forestry (CREF), Bikoro, Democratic Republic of the Congo. ¹⁴⁴Białowieża Geobotanical Station, Faculty of Biology, University of Warsaw, Białowieża, Poland. ¹⁴⁵Small Mammal Conservation Organization, Benin City, Nigeria. ¹⁴⁶Biological Sciences, Texas Tech University, Lubbock, United States of America. ¹⁴⁷Department of Biological Sciences, University of Sierra Leone, Freetown, Sierra Leone. ¹⁴⁸Nature Uganda, Kampala, Uganda. ¹⁴⁹Institute for Communities and Wildlife in Africa, University of Cape Town, Cape Town, South Africa. ¹⁵⁰Animal Production Institute, Rangeland Ecology, Agricultural Research Council, Pretoria, South Africa. ¹⁵¹College of Agriculture and Environmental Sciences: Department of Environmental Sciences (ABEERU), University of South Africa, Pretoria, South Africa. ¹⁵²Frogs & Friends, Berlin, Germany. ¹⁵³Chair of Wildlife Ecology & Management, University of Freiburg, Freiburg, Germany. ¹⁵⁴KBA Secretariat, c/o BirdLife International, Cambridge, United Kingdom. ¹⁵⁵Institute of Biosciences, Department of Botany, University of Rostock, Rostock, Germany. ¹⁵⁶Department of Economic Development, Environment, Conservation & Tourism, North West Provincial Government, Mahikeng, South Africa. ¹⁵⁷Department of Conservation and Marine Sciences, Cape Peninsula University of Technology, Cape Town, South Africa. ¹⁵⁸Department of Botany and Zoology, Stellenbosch University, Stellenbosch, South Africa. ¹⁵⁹Division of Conservation Ecology, Sâlim Ali Centre for Ornithology and Natural History, Coimbatore, India. ¹⁶⁰Mammalogy Department, Durban Natural Science Museum, Durban, South Africa. ¹⁶¹Herpetology, Museum für Naturkunde - Leibniz Institute for Evolution and Biodiversity Science, Berlin, Germany. ¹⁶²Department of Biology, University of Florence, Sesto Fiorentino, Italy. ¹⁶³Department of Zoology, Al-Azhar University, Cairo, Egypt. ¹⁶⁴School of Life Sciences, University of KwaZulu Natal, Durban, South Africa. ¹⁶⁵Gesellschaft fuer Internationale Zusammenarbeit (GIZ), Addis Ababa, Ethiopia. ¹⁶⁶Department of Biology and Natural Resources, Frostburg State University, Frostburg, USA. ¹⁶⁷CIRI, Centre International de Recherche en Infectiologie, Université de Lyon, Lyon, France. ¹⁶⁸Ornithology Section, Zoology Department, National Museums of Kenya, Nairobi, Kenya. ¹⁶⁹Biological Sciences and Centre for Statistics in Ecology, Environment and Conservation, University of Cape Town, Cape Town, South Africa. ¹⁷⁰Cologne Zoo, Cologne, Germany. ¹⁷¹Conservation Division, BirdLife South Africa, Johannesburg, South Africa. ¹⁷²Applied Behavioural Ecological & Ecosystem Research Unit (ABEERU), University of South Africa, Florida, South Africa. ¹⁷³Laboratory of Applied Ecology, University of Abomey-Calavi, Abomey-Calavi, Benin. ¹⁷⁴Mammal Research Institute, Centre for Invasion Biology, Department of Zoology and Entomology, University of Pretoria, Pretoria, South Africa. ¹⁷⁵Independent researcher, London, United Kingdom. ¹⁷⁶Department of Biology, University of Venda, Thohoyandou, South Africa. ¹⁷⁷Biology, University of Washington, Seattle, USA. ¹⁷⁸Endangered Wildlife Trust, Midrand, South Africa. ¹⁷⁹Grasslands-Forests-Wetlands Node of the South African Environmental Observation Network, Pietermaritzburg, South Africa. ¹⁸⁰Wildlife Conservation Society, Bronx, USA. ¹⁸¹Ndlovu Node of the South African Environmental Observation Network, Phalaborwa, South Africa. ¹⁸²Gabon Biodiversity Program, Smithsonian National Zoo and Conservation Biology Institute, Center for Conservation and Sustainability, Gamba, Gabon. ¹⁸³Biodiversity Capabilities Directorate, CapeNature, Cape Town, South Africa. ¹⁸⁴Department of Biodiversity and Conservation Biology, University of the Western Cape, Cape Town, South Africa. ¹⁸⁵FunMorph Lab, Department of Biology, University of Antwerp, Antwerp, Belgium. ¹⁸⁶AfricanBats NPC, Centurion, South Africa. ¹⁸⁷Evolutionary Ecology Group, University of Antwerp, Wilrijk, Belgium. ¹⁸⁸Arid Lands Node of the South African Environmental Observation Network, Kimberley, South Africa. ¹⁸⁹NWU Botanical Garden, School of Biological Sciences, North-West University, Potchefstroom, South Africa. ¹⁹⁰Richtersveld Desert Botanical Gardens, Richtersveld National Park, SANParks, Sendelingsdrift, South Africa. ¹⁹¹Department of Conservation Management, Nelson Mandela University, George, South Africa. ¹⁹²Institut de Systématique, Evolution, Biodiversité, Muséum National d'Histoire Naturelle, Paris, France. ¹⁹³Department of Botany, Rhodes University, Makhanda, South Africa. ¹⁹⁴Accelerated Taxonomy, Royal Botanic Gardens, Kew, Richmond, United Kingdom. ¹⁹⁵Restoration Ecology, Technische Universität München, Freising, Germany. ¹⁹⁶Department of Forestry and Wildlife Management, Maasai Mara University, Narok, Kenya. ¹⁹⁷Department of Migration, Max Planck Institute of Animal Behavior, Radolfzell, Germany. ¹⁹⁸Ecological Consultant, Fürth, Germany. ¹⁹⁹SARChI (NRF-DST) Research Chair on Biodiversity Value and Change, University of Venda, Thohoyandou, South Africa. ²⁰⁰Center for Tropical Research, Institute of the Environment and Sustainability, University of California Los Angeles, Los Angeles, USA. ²⁰¹Landscape Conservation Programme, BirdLife South Africa, Johannesburg, South Africa. ²⁰²Plant Ecology, University of Bayreuth, Bayreuth, Germany. ²⁰³School of Natural Resource Management, Nelson Mandela University, George, South Africa. ²⁰⁴Scientific Services, South African National Parks, Skukuza, South Africa. ²⁰⁵Kigelia Solutions, Lusaka, Zambia. ²⁰⁶Tau Consultants (Pty) Ltd, Maun, Botswana. ²⁰⁷University of Edinburgh, Edinburgh, Scotland, UK. ✉e-mail: clementshayley@gmail.com