



**HAL**  
open science

## Recent advances in the long-distance transport of sugars by the phloem and consequences on source-sink relations

Françoise Vilaine, Fabien Chardon, Noémie Vignolles, Rozenn Le Hir,  
Catherine C. Bellini, Sylvie Dinant

### ► To cite this version:

Françoise Vilaine, Fabien Chardon, Noémie Vignolles, Rozenn Le Hir, Catherine C. Bellini, et al.. Recent advances in the long-distance transport of sugars by the phloem and consequences on source-sink relations. *Plant Biology Europe*, Christophe Robaglia, Jul 2023, Marseilles, France. hal-04489028

**HAL Id: hal-04489028**

**<https://hal.inrae.fr/hal-04489028>**

Submitted on 4 Mar 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Recent advances in the long-distance transport of sugars by the phloem and consequences on source-sink relations

Françoise Vilaine<sup>1</sup>, Fabien Chardon<sup>1</sup>, Noémie Vignolles<sup>1</sup>, Rozenn Le Hir<sup>1</sup>, Catherine Bellini<sup>1,2</sup>, Sylvie Dinant<sup>1</sup>

<sup>1</sup> Université Paris-Saclay, INRAE, AgroParisTech, Institut Jean-Pierre Bourgin (IJPB), 78000 Versailles, France

<sup>2</sup> Umeå Plant Science Centre, Department of Plant Physiology, Umeå University, 90183 Umeå, Sweden

The distribution of photoassimilates in the plant relies on the phloem functions from source to sink organs. Phloem sugar transport is regulated by SUC/SUT and SWEET sugar transporters. We observed that deregulation of SUT1 or SUT2 in Tomato lines had various consequences on plant growth, sugar phloem transport rate and phloem sap composition. Similar effects were observed in response to the deregulation of SUC2 in Arabidopsis mutants and impairs source-sink relations, with consequences on root development and leaf evapotranspiration. Meanwhile, we observed natural variations in the metabolite profiles of phloem exudates of Arabidopsis accessions grown in low or non-limiting N supply. Ectopic deregulation of SUC2 in Arabidopsis impairs source-sink relationships, with consequences on root development and evapotranspiration from the leaf. Nutrient phloem transport thus also depends on downstream regulation of the central metabolism and water fluxes.

## Références :

Chardon F, De Marco F, Marmagne A, Le Hir R, Vilaine F, Bellini C, Dinant S. 2022.

Natural variation in the long-distance transport of nutrients and photoassimilates in response to N availability. *Journal of Plant Physiology* 273: 153707.

De Marco F, Batailler B, Thorpe MR, Razan F, Le Hir R, Vilaine F, Bouchereau A, Martin-Magniette ML, Eveillard S, Dinant S. 2021. Involvement of SUT1 and SUT2 sugar transporters in the impairment of sugar transport and changes in phloem exudate contents in phytoplasma-infected plants. *International Journal of Molecular Sciences* 22: 1–23.