



**HAL**  
open science

## Starch digestion in *H. illucens* conversion: exploring the role of amylases from larvae and substrate

Jérémy Guillaume, S. Mezdour, Frédéric Marion-Poll, Cécile Terrol, Chloé M C Brouzes, Philippe Schmidely

### ► To cite this version:

Jérémy Guillaume, S. Mezdour, Frédéric Marion-Poll, Cécile Terrol, Chloé M C Brouzes, et al.. Starch digestion in *H. illucens* conversion: exploring the role of amylases from larvae and substrate. 74. Annual Meeting of the European Federation of Animal Science, Aug 2023, Lyon, France. hal-04495576

**HAL Id: hal-04495576**

**<https://hal.inrae.fr/hal-04495576>**

Submitted on 8 Mar 2024


**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License

# Book of Abstracts of the 74<sup>th</sup> Annual Meeting of the European Federation of Animal Science



**Book of abstracts No. 29 (2023)**  
**Lyon, France**  
**26 August – 1 September, 2023**

**Starch digestion in *H. illucens* conversion: exploring the role of amylases from larvae and substrate**J.B. Guillaume<sup>1,2,3</sup>, S. Mezdour<sup>4</sup>, F. Marion-Poll<sup>2,5</sup>, C. Terrol<sup>1</sup>, C. Brouzes<sup>1</sup> and P. Schmidely<sup>3</sup>

<sup>1</sup>Agronutris, R&D Department, 35 blv du Libre Echange, 31650 Saint-Orens de Gameville, France, <sup>2</sup>CNRS, IRD, Université Paris-Saclay, Laboratoire EGCE, IDEEV, 12 route 128, 91190 Gif-sur-Yvette, France, <sup>3</sup>Université Paris-Saclay, INRAE, AgroParisTech, UMR Modélisation Systémique Appliquée aux Ruminants, 22 place de l'Agronomie, 91120 Palaiseau, France, <sup>4</sup>Université Paris-Saclay, INRAE, AgroParisTech, Sayfood, Campus Palaiseau, 22 place de l'Agronomie, 91120 Palaiseau, France, <sup>5</sup>Université Paris-Saclay, AgroParisTech, 22 place de l'Agronomie, 91120 Palaiseau, France; jeremy.guillaume@agroparistech.fr

The capacity of black soldier fly larvae (BSFL; *Hermetia illucens*) to transform organic substrates into body proteins and lipids suitable for animal nutrition is receiving growing attention. Carbohydrate content of BSFL diet is positively related to larval growth, fat body content and adult egg production, but underlying mechanisms of feed conversion remain to be explained. Previous studies reported amylase activity in larval midgut, and it has recently been shown that BSFL were highly efficient at digesting starch. This study focused on the effect of starch content and type on BSFL amylase activity and starch Estimated Digestibility (ED). BSFL were fed on five plant-based diets with different starch contents and types, and larvae and substrates were sampled after 4, 7 and 11 days of feeding, along with initial BSFL and diets. Each sample was ground in phosphate-buffered saline with protease inhibitor and centrifuged to collect water phase. Amylase activity was assessed using the Bernfeld technique and reported to total soluble protein measured according to Bradford. For substrate samples that could contain enzymes from the plant material, microbes or larvae, BSFL amylases were detected by Western-Blot using antibodies specific to insect amylases. This approach offers insight into larval amylase regulation mechanisms and the role of extra-oral digestion of starch in BSFL conversion systems.

**Investigating the nutritional requirements of black soldier fly larvae using artificial substrates**

L. Broeckx, L. Frootinckx, A. Wuyts, S. Berrens, M. Van Peer and S. Van Miert

Thomas More University of Applied Sciences, RADIUS, Kleinhofstraat, 4 2440 Geel, Belgium; laurens.broeckx@thomasmore.be

Due to the rising need for alternative protein, the black soldier fly gained interest of researchers worldwide. One reason for this is the insect's remarkable efficiency at converting feed into larval biomass, rich in high-quality protein. Moreover, they are able to grow on a high variety of organic side- and waste streams, allowing local production of high-quality protein, while also valorising organic waste. Even though their remarkable plasticity towards growing on a wide range of organic substrates has been proven in many studies, optimization is still necessary to increase the sector's viability. In order to optimize the sector, one important approach is to tailor the nutritional requirements of black soldier fly larvae. By doing so, models can be developed that predict larval growth based on substrate nutrient compositions, which can help combine different organic side-streams to create a more nutritionally complete substrate for black soldier fly rearing. Therefore in this study, multiple designs were developed and tested to study the effects of macronutrient contents on the growth of black soldier fly larvae through the use of artificial substrates. Substrates were composed of casein, sunflower oil, potato starch and cellulose, which allowed precise formulation of substrates, and could reduce the background noise such as the presence of complex sugars and unbalances in amino acid profiles. In this study, a model was built, showing significant main, interaction and quadratic effects of substrate protein and fat contents. A maximal larval growth was acquired at a substrate protein content of 30.12% and a fat content of 8.75%. The effect of carbohydrates on larval growth was not significant in this study, however this could be due to poor digestibility of raw potato starch by the larvae. While larvae tend to grow best on high-protein substrates, research shows that the protein conversion efficiency is inversely proportional to substrate protein content, meaning we may need to reconsider our approach to substrate optimization if we want to balance between maximizing larval growth and improving protein conversion efficiency.