



Influence of larval density on estimated digestibility of a standard diet in black soldier fly (*Hermetia illucens*)

Jérémy Guillaume, S. Mezdour, Frédéric Marion-Poll, Cécile Terrol, Philippe Schmidely

► To cite this version:

Jérémy Guillaume, S. Mezdour, Frédéric Marion-Poll, Cécile Terrol, Philippe Schmidely. Influence of larval density on estimated digestibility of a standard diet in black soldier fly (*Hermetia illucens*). Insects to Feed the World 2022, Jun 2022, Québec (CA), Canada. hal-04498139

HAL Id: hal-04498139

<https://hal.inrae.fr/hal-04498139v1>

Submitted on 11 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.




Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License



Journal of
Insects
as Food and Feed

ISSN 2352-4588

2022 – VOLUME 8 – SUPPLEMENT 1





Insects to Feed the World 2022

June 12 – 16, 2022

Can we ensure a humane death for black soldier fly larvae used in food and feed?

E.C. Rowe, E. De Castro and M. Pipan

Better Origin, Kings Hedges Road, CB4 2HY, United Kingdom; lizzie.rowe@betterorigin.co.uk

The International Platform of Insects for Food and Feed (IPIFF) states that 'We are committed to promoting good welfare practices in husbandry, transport and at the point of death, caring for insects' well-being.' Good welfare at the point of death means a death that is free from pain or suffering, which can be achieved by killing the animal instantly. Blanching is often cited as a method to kill black soldier fly larvae (BSFL) instantly, and yet it appears that there is no empirical data to support this. This study piloted methodology to answer the question: what is the lowest temperature of water needed to cause instantaneous death in black soldier fly larvae. Instantaneous was defined as within one second, and death was defined as lack of response to mechanical stimulation and loss of turgidity. Results indicated that the proxy measures used to determine death need further refinement, in part through a clearer understanding as to whether post-mortem reflex activity can occur in BSFL. Through presenting this pilot work we wish to stimulate discussion and collaboration on the important topic of humane killing of insects used in food and feed, so that insect farming companies can meet the growing societal expectation that insects, as living creatures, are treated with care and respect.

Influence of larval density on estimated digestibility of a standard diet in black soldier fly (*Hermetia illucens*)

J.B. Guillaume^{1,2,3}, S. Mezdour⁴, F. Marion-Poll^{2,5}, C. Terrol³ and P. Schmidely¹

¹Univ Paris-Saclay, INRAE, AgroParisTech, UMR Modélisation Systémique Appliquée aux Ruminants, 16 rue Claude Bernard, 75005 Paris, France, ²Univ Paris-Saclay, CNRS, IRD, UMR Evolut, Genomes, Comportement & Ecol, 1 Av de la Terrasse, 91198 Gif Sur Yvette, France, ³Agronutris, R&D Department, 35 Blv du Libre-Echange, 31650 Saint-Orens de Gameville, France, ⁴Univ Paris Saclay, SayFood, AgroParisTech, INRAE, 1 Rue des Olympiades, 91300 Massy, France, ⁵Univ Paris-Saclay, AgroParisTech, 16 rue Claude Bernard, 75005 Paris, France; jeremy.guillaume@agronutris.com

Black soldier fly larvae (BSFL) efficiency at converting organic by-products into insect biomass suitable for livestock nutrition depends on the nature and macronutrient profile of their feed. Digestibility is a key parameter to the development of a diet formulation approach, but methods to assess digestibility in most insects are unsuitable for BSFL as ingesta is not measurable and pure faeces can not be obtained and analysed. This work defined estimated digestibility (ED) as the difference between feed and frass nutrient weight, divided by the weight of nutrient in feed, and explored its variation with larval density. ED of dry matter (DM), starch, nitrogen, ether extract (EE), neutral detergent fibre (NDF), acid detergent fibre (ADF), acid detergent lignin (ADL), ash and energy were evaluated using the same amount of chicken feed, same feeding time and densities from 0 to 29 larvae/cm². ED of DM, starch, nitrogen, EE and energy increased with larval density following an asymptotic trend. ED of ash also increased between 0 and 11.4 larvae/cm², but decreased for higher densities. For NDF, ADF and ADL, no clear results were obtained presumably because the fibre dosing procedure used was not specific enough and also accounted for chitin in the frass. ED of all nutrients but fibre fractions were modelled using a logistic equation. This method was able to propose asymptotic ED values for some major nutrients. This new indicator corresponds to the maximal fraction of each nutrient that BSFL can digest and shows potential for the development of a BSFL diet formulation approach based on digestible fractions rather than crude nutrient content. Future studies should compare asymptotic ED of various substrates to provide insight on BSFL ability to digest different ingredients.