



## Investigating the role of amylases from *Hermetia illucens* larvae and their substrate in starch digestion

Jérémy Guillaume, S. Mezdour, Frédéric Marion-Poll, Cécile Terrol, Chloé M C Brouzes, Philippe Schmidely

### ► To cite this version:

Jérémy Guillaume, S. Mezdour, Frédéric Marion-Poll, Cécile Terrol, Chloé M C Brouzes, et al.. Investigating the role of amylases from *Hermetia illucens* larvae and their substrate in starch digestion. 12. European Congress of Entomology, Oct 2023, Heraklion - Crete, Greece. hal-04498179


**HAL Id: hal-04498179**

**<https://hal.inrae.fr/hal-04498179>**

Submitted on 11 Mar 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



**BOOK OF ABSTRACTS**



# **ECE 2023** **CRETE**

## European Congress of Entomology

### **XII European Congress of Entomology**

16-20.10.2023

Cultural Conference Center of Heraklion  
Crete, Greece

[www.ece2023.com](http://www.ece2023.com)

Organised by



Under the Auspices



HELLENIC REPUBLIC  
Ministry of Rural Development  
and Food



HELLENIC REPUBLIC  
MINISTRY OF TOURISM



ΠΕΡΙΦΕΡΕΙΑ ΚΡΗΤΗΣ  
REGION OF CRETE



MUNICIPALITY OF  
HERAKLION



biotransformation efficiency of the Organic Fraction of Municipal Solid Waste (OFMSW) by BSF larvae with the final aim to obtain biobased materials, such as bioplastics and biodiesel from insect proteins and lipids, respectively. In particular, in this study we evaluated the growth performance of BSF larvae and their efficiency in reducing two substrates with different nutritional content that mimic the composition of OFMSW. Although the growth rate of larvae reared on both substrates was comparable, differences in the reduction of waste and the conversion efficiency into insect body mass were observed. As the midgut plays a fundamental role in the insect-mediated bioconversion processes, we evaluated the digestion capability of BSF larvae reared on the two substrates. Our results showed that the larvae can compensate variations in nutrient composition of the substrate by post-ingestion responses, through the regulation of gene expression and activity of digestive enzymes. Indeed, despite differences in diet composition, we observed similar growth performance and chemical composition of the insect.

**Keywords:** *Hermetia illucens* larvae, waste bioconversion indexes, midgut physiology, digestive enzymes

---

### OC063. Investigating the role of amylases from *Hermetia illucens* larvae and their substrate in starch digestion

J.B. Guillaume<sup>\*1,2,3</sup>, S. Mezdour<sup>4</sup>, F. Marion-Poll<sup>2,5</sup>, C. Terrol<sup>3</sup>, C. Brouzes<sup>3</sup>, P. Schmidely<sup>1</sup>

<sup>1</sup>Université Paris-Saclay, INRAE, AgroParisTech, UMR Modélisation Systémique Appliquée aux Ruminants, Palaiseau, France

<sup>2</sup>Laboratoire Évolution, Génomes, Comportement et Écologie, CNRS, IRD, Université Paris-Saclay, Institut Diversité, Ecologie et Evolution du Vivant (IDEEV), Gif-sur-Yvette, France

<sup>3</sup>Agronutris, R&D Department, Saint-Orens de Gameville, France

<sup>4</sup>Université Paris-Saclay, Sayfood, AgroParisTech, INRAE, Palaiseau, France

<sup>5</sup>Université Paris-Saclay, AgroParisTech, Palaiseau, France

\*Corresponding author: [jeremy.guillaume@agroparistech.fr](mailto:jeremy.guillaume@agroparistech.fr)

Black soldier fly larvae (BSFL; *Hermetia illucens*) are increasingly studied for their ability to transform organic substrates into body proteins and lipids that can be used for animal or human nutrition. Carbohydrate content of their diet has been associated with larval growth and fat body content, and it has recently been reported that BSFL had high starch digestion efficiency and that their midgut presented high amylase activity. This study investigated the effect of starch content and type on BSFL amylase activity and starch Estimated Digestibility (ED). BSFL were fed various plant-based diets with different starch content. Both larvae and substrates were sampled after feeding periods of 4, 7 and 11 days, along with initial BSFL and diets. Each sample was ground in phosphate-buffered saline with protease inhibitor and centrifuged to collect water phase. Amylase activity was assessed in each solution via spectrophotometry using soluble starch at pH 6.5 and dinitrosalicylic acid. Values were reported as enzymatic activity units per gram of total soluble protein measured according to Bradford. Western-blot was used to detect BSFL amylases in substrate samples that could contain enzymes from plant material, microbes, or larvae. This approach provides insight into larval amylase regulation mechanisms and the role of extra-oral digestion of starch in BSFL conversion systems.

**Keywords:** insect nutrition, extra-oral digestion, alternative protein, insects as feed, Diptera, digestive enzymes

---