

## Editorial: Metallic micronutrient homeostasis in plants, volume II

Hannetz Roschzttardtz, Diego Gomez-Casati, Christian Dubos, Julia Quintana

### ▶ To cite this version:

Hannetz Roschzttardtz, Diego Gomez-Casati, Christian Dubos, Julia Quintana. Editorial: Metallic micronutrient homeostasis in plants, volume II. Frontiers in Plant Science, 2023, 14, pp.1329190. 10.3389/fpls.2023.1329190. hal-04523580

### HAL Id: hal-04523580 https://hal.inrae.fr/hal-04523580v1

Submitted on 27 Mar 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.







### **OPEN ACCESS**

EDITED AND REVIEWED BY
M. J. I. Shohag,
University of Florida, United States

\*CORRESPONDENCE

Hannetz Roschzttardtz

Diego Gomez-Casati

Christian Dubos

□ christian.dubos@inrae.fr

Julia Quintana

RECEIVED 28 October 2023 ACCEPTED 24 November 2023 PUBLISHED 01 December 2023

#### CITATION

Roschzttardtz H, Gomez-Casati D, Dubos C and Quintana J (2023) Editorial: Metallic micronutrient homeostasis in plants, volume II. *Front. Plant Sci.* 14:1329190. doi: 10.3389/fpls.2023.1329190

### COPYRIGHT

© 2023 Roschzttardtz, Gomez-Casati, Dubos and Quintana. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

# Editorial: Metallic micronutrient homeostasis in plants, volume II

Hannetz Roschzttardtz<sup>1\*</sup>, Diego Gomez-Casati<sup>2\*</sup>, Christian Dubos<sup>3\*</sup> and Julia Quintana<sup>4,5\*</sup>

<sup>1</sup>Facultad de Ciencias Biológicas, Pontificia Universidad Católica de Chile, Santiago, Chile, <sup>2</sup>Centro de Estudios Fotosintéticos y Bioquímicos (CEFOBI-CONICET), Universidad Nacional de Rosario, Rosario, Argentina, <sup>3</sup>Institut des Sciences des Plantes de Montpellier (IPSiM), University Montpellier, CNRS, INRAE, Montpellier, France, <sup>4</sup>Center for Plant Biotechnology and Genomics, Universidad Politécnica de Madrid, Madrid, Spain, <sup>5</sup>Department of Biology and Geology, Physics and Inorganic Chemistry, Universidad Rey Juan Carlos, Madrid, Spain

### KEYWORDS

plant nutrition, metal accumulation, transcriptomics, seed nutrient composition, metal homeostasis, iron deficiency

### Editorial on the Research Topic

Metallic micronutrient homeostasis in plants, volume II

"Metallic Micronutrient Homeostasis in Plants, Volume II" represents an initiative that responds to the need for a research area enriched with its unique historical context and challenges, all of which are critically relevant in the context of the ever-evolving global environmental landscape. The development and use of amenable techniques for metal localization in vivo remains a major experimental challenge in the study of metal nutrient homeostasis. Moreover, analysis of the interactions between micronutrient and macronutrient homeostasis is gaining more and more attention within the scientific community. Lastly, there is a pressing need to expand the knowledge in iron deficiency responses in non-model crops. This Research Topic agglutinates knowledge in these directions. In one of the contributions,

Grant-Grant et al. have elegantly demonstrated that alterations in the spatial distribution of iron within plant embryos do not exert a significant influence on the overall iron content within the seeds. This suggests that the regulation of iron levels within plants is not determined by how iron is distributed within the embryo. From a biotechnological perspective this discovery is interesting as it demonstrates the difficulty in designing strategies to increase micronutrients in seeds.

Xu et al., have used different maize inbred lines in order to identify potential important genes in iron deficiency tolerance in maize, which affect a significant proportion of cultivable land on the planet. This study provides valuable insights into unravelling the molecular mechanisms that underlie maize's resilience to iron-deficiency stress.

In other work, Yuan et al. identified potential genes for maintaining iron homeostasis under iron-deficient stress in non-heading Chinese cabbage. This could provide a robust theorical foundation for the research on molecular mechanisms governing the adaptation

Roschzttardtz et al. 10.3389/fpls.2023.1329190

to iron stress and could also serve to design new strategies for the development of iron-tolerant crop varieties.

How plants respond to iron deficiency stress is an important area of research in plant biology. Tran et al. demonstrated that the alteration of porphyrin biosynthesis via the introduction of the *Arabidopsis thaliana* Mg-chelatase H subunit gene (AtCHLH) into transgenic rice effectively mitigates stress induced by iron deficiency. It has been proposed the existence of an ABA-independent pathway as a potential mechanism for alleviating this stress through protoporphyrin synthesis.

In summary, the work presented in this Research Topic showed the importance of different genes in the distribution and the maintenance of iron homeostasis, specially under iron stress conditions. Interestingly, is has been proposed that regulatory mechanism for porphyrin metabolism would be part of the complex protective systems against Fe deficiency stress. Moreover, the landscape of iron deficiency transcriptomes has been enriched with an interesting analysis of gene expression changes in Chinese cabbage. All in all, this research aims to ultimately aid the development of new solutions to mitigate fertilizer overuse and while increasing crop nutritional value.

### **Author contributions**

HR: Writing – original draft. DG-C: Writing – original draft. CD: Writing – original draft. JQ: Writing – original draft.

### Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

### Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.