



HAL
open science

BRAHMA, a SWI/SNF Complex Regulating Seed Longevity in Arabidopsis

Julia Zinsmeister, Magdalena Wrona, Hubert Schaller, Pierre Mercier, Claire Villette, Dimitri Heintz, Loïc Rajjou, Ruslan Yatusевич, Szymon Swiezewski

► **To cite this version:**

Julia Zinsmeister, Magdalena Wrona, Hubert Schaller, Pierre Mercier, Claire Villette, et al.. BRAHMA, a SWI/SNF Complex Regulating Seed Longevity in Arabidopsis. 14. ISSS biennial conference, Sorbonne Université, Jul 2023, Paris, France. hal-04527250

HAL Id: hal-04527250

<https://hal.inrae.fr/hal-04527250v1>

Submitted on 29 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

BRAHMA, A SWI/SNF COMPLEX REGULATING SEED LONGEVITY IN ARABIDOPSIS

Julia Zinsmeister¹, Magdalena Wrona², Hubert Schaller³, Pierre Mercier³, Claire Villette³, Dimitri Heintz³, Loïc Rajjou¹, Ruslan Yatusевич², Szymon Swiezewski²

¹Université Paris-Saclay, INRAE, AgroParisTech, Institut Jean-Pierre Bourgin (IJPB), 78000, Versailles, France.

²Institute of Biochemistry and Biophysics PAS, Pawinskiego 5A,02-106 Warsaw, Poland

³Institut de Biologie Moléculaire des Plantes du CNRS, Université de Strasbourg, 67084 Strasbourg, France

Corresponding author: Julia Zinsmeister (julia.zinsmeister@inrae.fr)

Seed longevity, the time a seedlot can be kept at the dry state without loss of viability, is key in agriculture as it directly ensures field establishment and thus yields, in addition to its main role in biodiversity preservation. Acquisition of seed longevity is tightly regulated at molecular level during seed development. A role of DELAY OF GERMINATION1 (DOG1) in positively regulating seed longevity in *Arabidopsis thaliana* has been previously characterized[1]. In addition, further research has revealed that BRAHMA (BRM), an SNF2 (Sucrose NonFermenting) – type ATPase from SWI/SNF (SWItch/Sucrose Non-Fermentable) chromatin remodelling complex – shows a 3' localized pattern of binding at 1759 genes, including *DOG1*[2]. Here, we investigated the role of BRM in relation with *DOG1* in seed longevity. Using physiological assessments, RNAseq analysis, and untargeted metabolomics approach, we conducted a deep analysis of the *brm-3*, *dog1-4* and double *dog1-4xbrm-3* mutants. Our results indicate that BRM negatively regulates seed longevity. In order to identify putative pathways involved in the regulation of seed longevity by BRM, we analyzed both RNAseq and metabolomics data of the mutants. Our findings indicate a role of BRM in balancing redox status by negatively regulating antioxidants pathways at gene expression level as well as by inhibiting glutathione accumulation in seeds, an important antioxidant in seed longevity[3]. Finally, our results show also a role of BRM in other agronomical traits of interest such as yield and nutritional quality by regulating seed size, weight and lipid accumulation.

[1] Dekkers, Bas JW, et al. *The Plant Journal*. (2016): 451-465.

[2] Archacki R, et al. *Nucleic Acids Res* (2016) 45: 3116-3129.

[3] Kranner, Ilse, et al. *Free Radical Biology and Medicine* (2006): 2155-2165.

Acknowledgements & Funding

This project has received funding from the European Union's Horizon H2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 101038075 and from EMBO scientific exchange grant No 9107. The IJPB benefits from the support of Saclay Plant Sciences-SPS (ANR-17-EUR-0007).

Topic: From genes to seed biology, novel issues in molecular mechanisms and beyond
Oral application