

Fetal sex differences in placental LCPUFA ether and plasmalogen phosphatidylethanolamine and phosphatidylcholine contents in pregnancies complicated by obesity

Theresa L Powell, Charis Uhlson, Lana Madi, Karin Zemski Berry, Stephanie S Chassen, Thomas Jansson, Veronique Ferchaud-Roucher

▶ To cite this version:

Theresa L Powell, Charis Uhlson, Lana Madi, Karin Zemski Berry, Stephanie S Chassen, et al.. Fetal sex differences in placental LCPUFA ether and plasmalogen phosphatidylethanolamine and phosphatidylcholine contents in pregnancies complicated by obesity. Biology of Sex Differences, 2023, 14 (1), pp.66. 10.1186/s13293-023-00548-1. hal-04530595

HAL Id: hal-04530595 https://hal.inrae.fr/hal-04530595

Submitted on 3 Apr 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Fetal sex differences in placental LCPUFA ether and plasmalogen phosphatidylethanolamine and phosphatidylcholine contents in pregnancies complicated by obesity



Theresa L. Powell^{1,2}, Charis Uhlson², Lana Madi¹, Karin Zemski Berry³, Stephanie S. Chassen², Thomas Jansson¹ and Veronique Ferchaud-Roucher^{4,5*}

Abstract

Background We have previously reported that maternal obesity reduces placental transport capacity for lysophosphatidylcholine-docosahexaenoic acid (LPC-DHA), a preferred form for transfer of DHA (omega 3) to the fetal brain, but only in male fetuses. Phosphatidylethanolamine (PE) and phosphatidylcholine (PC), have either sn-1 ester, ether or vinyl ether (plasmalogen) linkages to primarily unsaturated and monounsaturated fatty acids and DHA or arachidonic acid (ARA, omega 6) in the sn-2 position. Whether ether and plasmalogen PC and PE metabolism in placenta impacts transfer to the fetus is unexplored. We hypothesized that ether and plasmalogen PC and PE containing DHA and ARA are reduced in maternal–fetal unit in pregnancies complicated by obesity and these differences are dependent on fetal sex.

Methods In maternal, umbilical cord plasma and placentas from obese women (11 female/5 male infants) and normal weight women (9 female/7 male infants), all PC and PE species containing DHA and ARA were analyzed by LC– MS/MS. Placental protein expression of enzymes involved in phospholipid synthesis, were determined by immunoblotting. All variables were compared between control vs obese groups and separated by fetal sex, in each sample using the Benjamini–Hochberg false discovery rate adjustment to account for multiple testing.

Results Levels of ester PC containing DHA and ARA were profoundly reduced by 60–92% in male placentas of obese mothers, while levels of ether and plasmalogen PE containing DHA and ARA were decreased by 51–84% in female placentas. PLA2G4C abundance was lower in male placentas and LPCAT4 abundance was lower solely in females in obesity. In umbilical cord, levels of ester, ether and plasmalogen PC and PE with DHA were reduced by 43–61% in male, but not female, fetuses of obese mothers.

*Correspondence: Veronique Ferchaud-Roucher veronique.ferchaud-roucher@univ-nantes.fr Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.gr/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.gr/licenses/by/4.0/.

Conclusions We found a fetal sex effect in placental PE and PC ester, ether and plasmalogen PE and PC containing DHA in response to maternal obesity which appears to reflect an ability of female placentas to adapt to maintain optimal fetal DHA transfer in maternal obesity.

Highlights

- Ether and plasmalogen phosphatidylcholine (PC) and phosphatidylethanolamine (PE) are considered DHA reservoirs and whether their metabolism in placenta and DHA transfer to the fetus are altered in obesity is unexplored, especially regarding the fetal sex.
- We found that in obesity, (1) pregnancies with male fetuses decrease ester PC with DHA and ARA in placentas but not plasmalogen PC species and the levels of ester, ether and plasmalogen PC and PE with DHA is decreased in fetal umbilical circulation, and (2) pregnancies with female fetuses decrease ether and plasmalogen PE with DHA and ARA in placentas and maintain the levels of ester, ether and plasmalogen PC and PE with DHA in fetal circulation.
- Both males and females have compensatory mechanisms that attempt to maintain an adequate supply of DHA species, particularly for brain development, in obesity, however, the male placental adaptation failed to maintain fetal levels to those found in normal weight pregnancies.

Keywords Sexual dimorphism, De novo placental phospholipid synthesis, Remodeling phospholipid pathway, Maternal obesity

Plain language summary

Docosahexaenoic acid (DHA) is a critical omega 3 long chain polyunsaturated fatty acid (LCPUFA) for fetal brain development. We have recently reported that maternal obesity reduces placental transport capacity for Lysophos-PhatidylCholine-DHA (LPC-DHA), a preferred form for transfer of DHA to the fetal brain, but only in male fetuses. Other important lipids, the plasmalogen phosphatidylcholine (PC) and phosphatidylethanolamine (PE) are considered DHA reservoirs, but its roles in the maternal–fetal unit are largely unexplored. We examined these lipid species in maternal and fetal circulation and in placental tissue to uncover potential novel roles for ether and plasmalogen lipids in the regulation of placenta delivery of these vital nutrients in pregnancies complicated by obesity depending of fetal sex. We demonstrated for the first time, that female fetuses of obese mothers decrease placental ether and plasmalogen PE containing DHA and arachidonic acid (ARA, omega 6), and show a high fetal–placental adaptability and placental reserve capacity that can maintain the PC-LCPUFA synthesis and the transfer of these crucial species to the fetus to preserve brain development. Our study also demonstrated that male fetuses, in response to maternal obesity, reduce the placental ester PC species containing DHA and ARA and reduce the ether and plasmalogen PE reservoir of DHA and ARA in fetal circulation. Our findings support a fetal sex effect in placental ester, ether and plasmalogen PE and PC containing DHA in response to maternal obesity which appears to reflect an ability of female placentas to adapt to maintain optimal fetal DHA transfer in maternal obesity.

Introduction

The worldwide prevalence of maternal obesity continues to increase and affects not only maternal health, but also negatively impacts fetal growth and development by modulating placental function [1]. Placental dysfunction has been recently shown to be associated with changes in fat mass at birth and with risks of metabolic disorders later in life [2]. Pre-gravid maternal obesity is associated with an elevated risk of obesity in boys from age one [3]. In addition, pregnancies complicated by overweight and obesity are associated with long-term cognitive issues in the children, such as attention deficit and hyperactivity disorder (ADHD) [4] and autism spectrum disorder (ASD) [5] with male children being more frequently affected than female children [6]. For example, boys aged 7–11 years have reduced hippocampal volume when their mothers were obese during pregnancy but this was not observed in girls [7]. Similar sex differences in brain development have been described in children of mothers with gestational diabetes mellitus (GDM) [8]. Numerous studies found correlations between maternal obesity and child neurodevelopment disorders [9, 10] suggesting in utero programming of brain development. However, the mechanisms linking maternal obesity in pregnancy and impaired neurodevelopment in children remain unclear.

Long chain polyunsaturated fatty acids (LCPUFA), such as docosahexaenoic acid (DHA, C22:6n-3), are considered essential nutrients for fetal development, particularly for the brain. Supplied by the mother's dietary intake, DHA is preferentially taken up by the placenta [11, 12], metabolized and transferred to the fetus through the trophoblast basal plasma membrane and the fetal endothelium mediated by specific lipid transporters [13, 14]. Maternal n-3 LCPUFA deficiency has been reported in pre-gestational obese women [15] and is associated with decreased visual acuity, hypertension, diabetes, ADHD and ASD in the offspring [16]. Whether the DHA deficit in maternal circulation of obese mothers is related to her own metabolism and thus results in decreased transplacental supply of LCPUFAs is not well known. If lower maternal supply was the cause of decreased LCP-UFA transfer through the placenta, the fetal sex differences on neurodevelopment are more difficult to explain. However, placental function is dependent on fetal sex [17] and may contribute to differences in delivery of DHA to the developing fetus.

DHA and arachidonic acid (ARA, C20:4n-6) are highly abundant in circulating phospholipids, specifically phosphatidylcholine (PC) and phosphatidylethanolamine (PE), and increase in the maternal circulation over gestation. Whereas PE displays the most significant increase during pregnancy compared to the other phospholipid classes [18], the level of DHA and ARA is higher in PC than in PE. PC is synthesized de novo by two pathways in the endoplasmic reticulum: (1) the Kennedy pathway, a part of the cytidine diphosphocholine (CDP-choline) pathway, which produces the majority of cellular PC from dietary choline and non-esterified fatty acids (NEFAs) and at a lesser extend the conversion of PE to PC through the phosphatidylethanolamine methyltransferase (PEMT) pathway; (2) the acyl-chain composition of PC can be modified through Land's cycle (remodeling pathway), by replacement of fatty acids at the sn-2 position by the combined action of the calcium-independent Group VIA phospholipase A2 (iPLA2) and reacylation by lysophosphatidylcholine acyltransferase (LPCAT) [19].

We have previously demonstrated that levels of nonesterified DHA and lysophosphatidylcholine containing DHA (LPC-DHA) are lower in placentas of obese women carrying a male fetus and reduced LPC-DHA levels were also found in cord blood, indicating reduced delivery to the fetus. This was in spite of an up-regulation of the protein expression of MFSD2a, LPC-DHA transporter, in the basal syncytiotrophoblast plasma membrane of male placentas, suggesting that the adaptation in transporter expression was insufficient to maintain normal DHA transfer to the fetus [20]. We speculate that reduced DHA transfer to the fetal circulation may negatively impact brain development in male fetuses of obese mothers.

The physicochemical characteristics of phospholipids affect the structure and function of cell membranes. PC and PE can have either ester (common diacyl phospholipids), ether (O=alkyl glycerophospholipid) or vinyl ether (P=plasmalogen glycerophospholipid) linkages to primarily unsaturated and monounsaturated fatty acids at the sn-1 position of the glycerol backbone. The phospholipid fraction in most cell membranes contains 15-20% plasmalogens with vinyl ether linked to sn-1 position [21]. Despite the high abundance of plasmalogens in cell membranes in many organs such as brain, heart and kidney [22], the role of ether and plasmalogen linkages in phospholipids remains unclear and literature on lipidomic analysis often do not specifically differentiate plasmalogen phospholipids from ether linked phospholipids. Because plasmalogens are preferentially enriched in ARA and DHA at the sn-2 position [23], they may function as reservoirs for these important LCPUFAs. Indeed, due to the presence of the vinyl ether at the sn-1 position, the plasmalogens cannot be hydrolyzed by phospholipase A1. However, LCPUFAs at the sn-2 position can be released from plasmalogens by PLA₂ hydrolysis during the remodeling process [24]. Release of ARA is vital as a precursor for important lipid signaling molecules (prostaglandin, leukotriene, thromboxane) and DHA is mainly esterified into ester forms of phospholipids such as PC, which are crucial components of cell membranes in the brain and neural tissues, or transported as LPC-DHA by MFSD2a [25].

Ether phospholipids and particularly plasmalogens are thought to also act as endogenous antioxidants that protect critical cell membrane components such as LCP-UFA from oxidative stress [3, 26]. A protective role for LCPUFA containing plasmalogens in atherosclerosis was demonstrated in LDLR-/- mice [27] and APOE-deficient mice [28]. Consistent with these data, lower levels of plasmalogen PC of HDL are associated with coronary disease [29] and decreased serum ether lipid levels have been implicated in hypertension and obesity [30, 31]. In addition, lower plasmalogen proportion in phospholipids is associated with neurodegenerative diseases such as Alzheimer, with specific decreases in plasmalogen PC and PE containing DHA [32]. Conversely, serum plasmalogen levels increase in men with aerobic training and in patients after a healthy dietary intervention [33, 34]. Although altered circulating levels of these compounds are associated with higher risks to develop cardiovascular and degenerative diseases, few clinical studies have investigated their role over healthy pregnancy and in pregnancy complicated by obesity and their impact on fetal development. One study has observed increased plasma ether PC containing DHA (PC O-16:0 22:6) in healthy women at 26-28 weeks of pregnancy compared to postnatal 4–5 years after delivery [18]. We have recently identified some ether linked phospholipid species in human placenta across gestation and found PE containing LCPUFA, such as PE O-16:1 22:6 and PE O-16:1_20:4, are enhanced in the third trimester compared to the first and second trimesters [35]. These data suggest a potential implication of those phospholipids in the placental LCPUFA metabolism and transfer to the fetus. A specific role of placental ether and plasmalogen PC and PE containing DHA and ARA particularly in the context of maternal obesity as how they may relate to transfer to the fetus depending of fetal sex is currently unknown. We therefore hypothesized that ester, ether and plasmalogen linked PC and PE containing DHA and ARA are reduced in maternal-fetal unit in pregnancies complicated by obesity and these differences are dependent on fetal sex.

Materials and methods

Subjects and sample collections

Pregnant women were enrolled before delivery to donate their blood, placenta and umbilical cord blood to a data/ bio-repository for use in multiple research studies under a protocol approved by the Institutional Review Board at University of Colorado, Denver (COMIRB 14-1073). All participants gave their informed written consent for the use of their biological samples and protected health information. Inclusion criteria for the repository included information on pre-pregnancy/early pregnancy BMI, ultrasound confirmation of gestational age at 14-18 weeks, singleton pregnancy, and maternal age 18-45 years. Exclusion criteria included concurrent inflammatory, vascular, or metabolic disease, current use of tobacco, street drugs, or medications, fetal malformations, history of pregnancy loss or pre-term delivery, any pregnancy pathology (gestational diabetes, hypertension, pre-eclampsia) other than elevated BMI. Anonymized biological samples such as maternal blood, placenta and umbilical cord vein and artery were transferred to the lab for preparation and storage at - 80 °C for batch analysis. Pregnant women involved in the current study were either normal weight (BMI, range 18.5–24.9 kg/m²), i.e., control group (n=16; n=9 females, n=7 males) or obese (BMI, range 30–45 kg/m²) (n=16; n=11 females, n = 5 males). Maternal and infant characteristics for study subjects are shown in Additional file 1: Table S1A and have been previously reported [20]. For placental protein expression analysis, placentas were collected immediately after delivery and homogenized in buffer D (250 mM sucrose, 10 mM hepes, pH 7.4), including a 1:100 dilution of both phosphatase and protease inhibitors (Sigma-Aldrich, St. Louis, MO) and stored at - 80 °C until batch processing. The number of studied placentas was increased to 38 (control group n=18; n=10 females and n=8 males, obese group n=20; n=10 females and n=10 males) to increase power to allow studies of the influence of fetal sex (Additional file 1: Table S1B).

Quantification of protein expression by western blot and simple western

Total protein content of placental homogenates was determined using BCA assay (Pierce BCA Protein Assay Kit, Thermo Scientific, MA, USA). Twenty µg total protein was loaded and separated on Bis-Tris gels (4-20%) using a pre-cast gel systems (Bio-Rad) as previously described [36]. After electrophoresis and transfer onto a PVDF membrane (Bio-Rad Laboratories Inc.), we stained for total protein using Amido Black Stain (Sigma-Aldrich). Incubation of primary antibody LysoPhosphatidylCholine Acyl Transferase 4 (LPCAT4), PhosphoLipase A2 Group IVC (PLA2G4C or iPLA2) and 1-acylglycerol-3-phosphate O-acyltransferase 4 (AGPAT4) was carried out overnight at 4 °C (details on primary antibodies are provided in Additional file 1: Table S2), and secondary antibody (peroxidase labeled anti-Rabbit IgG, diluted 1:3000, Cell Signaling Technology (Danvers, MA)) for 1 h at room temperature as previously depicted [35]. Immunolabeling was made visible with SuperSignal West Pico Plus detection solution (Thermo Scientific) in a G:Box ChemiXL1.4 (SynGene, Cambridge, UK). Densitometry analysis of target protein bands was performed with GeneTools (SynGene) and target protein expression was adjusted for Amido Black staining to account for any variation in protein loading and transfer.

We also used the JESS (capillary-based immunoblotting, Simple Western with total protein normalization— ProteinSimple) [37] to measure the abundance of target proteins: fatty acyl-CoA reductase 1 (FAR1), 1-acylglycerol-3-phosphate O-acyltransferase 2 (AGPAT2), and glycerol-3-phosphate O-acyltransferase 3 (GPAT3) (Additional file 1: Table S2). We ran the plates based on the recommended manufacturer's settings (separation voltage of 375 for 25 min) with the placental homogenates at a 1 μ g/ μ l protein concentration per well. An equalizer sample was used between each plate to correct for variations, and positive controls were run with each antibody to confirm the presence of the target.

Analysis of phosphatidylcholine and phosphatidylethanolamine species containing LCPUFA

in plasma and placental tissue by LC-MS/MS Phosphatidylcholine (PC) and phosphatidylethanolamine (PE) were extracted and analyzed in placental homogenates and plasma samples [maternal vein (MV), umbilical vein (UV) and artery (UA)], by LC-MS/MS in the Mass Spectrometry Lipidomics Core facility at the University of Colorado using a protocol previously described [20, 36]. Briefly, 1 µl of a commercial internal standard mixture containing the major phospholipid classes (SPLASH Lipidomix, from Avanti Polar Lipids, Inc.) was added to 50 µl of each plasma and placental homogenate sample and lipids were extracted according to the method of Bligh and Dyer [38]. LC–MS/MS system equipped with a HPLC system (Shimadzu, Kyoto, Japan) with a normal phase chromatography (silica HPLC column, Ascentis, 150×2.1 mm, 5 µm, Supelco, Bellefonte, PA) and a 5500 QTRAP mass spectrometer (SCIEX, Framingham, MA) was used. PC and PE containing esterified ARA or DHA chains were analyzed by scanning precursor ions of the m/z 303 or of the m/z 327 product anion, respectively. Results were analyzed using MultiQuant software (SCIEX) and all detected ARA and DHA containing PC and PE results are reported as the ratio between the integrated area of each analyte and the integrated area of the corresponding internal standard (IS) for PC class (PC 15:0 [²H₇]18:1 and LPC [²H₇]18:1) and PE class (PE 15:0 $[{}^{2}H_{7}]$ 18:1 and LPE $[{}^{2}H_{7}]$ 18:1).

In order to distinguish vinyl ether (P=plasmalogen) from 1-O-alkyl (O=ether) linkages at the sn-1 position of PC and PE species, acid labile plasmalogens were hydrolyzed by exposure to 12N HCl fumes [39]. This was achieved by drying down the placenta total lipid extract under a stream of nitrogen followed by exposure to hydrochloric acid fumes for 1 h. Upon acid hydrolyzed but the 1-O-alkyl or 1-acyl phospholipid species remained intact. The lipids were reconstituted in initial HPLC conditions for LC–MS/MS analysis and plasmalogen identity was determined by disappearance of the molecular species by comparing pre- vs post-HCl exposed samples.

Statistical analyses

All statistical analyses were performed using GraphPad Prism version 9.01. Results are expressed as $mean \pm SD$ for each sample type; maternal vein (MV), placenta, umbilical vein (UV) and artery (UA). Chi-square test was used to compare qualitative data in obese and control groups. Data distribution was assessed by the Shapiro–Wilk normality test. All lipid variables were compared between two independent groups, control vs

obese groups and separated by fetal sex, in each sample. We applied the Benjamini–Hochberg false discovery rate adjustment to account for multiple testing. All tests were two-sided. A significance threshold set at $q \le 0.05$ was used for the targeted lipidomic analysis [40]. Correlation analysis with spearman test was performed to assess correlations between maternal obesity and ether and plasmalogen PC and PE species in placentas and in umbilical cord. *p* values < 0.05 were considered as statistically significant.

In addition, for each phospholipid variable showing a statistical difference between groups, effect size was calculated as following: ((Mean PC ob – Mean PC ctr)/ mean PC ctr) \times 100).

Results

Clinical parameters of mothers and their infants are provided in Additional file 1: Table S1A and separated by fetal sex in Additional file 1: Table S1B. By design, maternal BMI in the obese group were greater than in the normal weight group. Maternal age at delivery, gestational age, placenta and birth weights were not significantly different between the obese vs control groups for either sex.

Plasma from obese mothers have lower levels of sn-1 ether and plasmalogen PC and PE containing DHA but not ARA

We analyzed all the species of sn-1 ester (common form), ether and plasmalogen PC and PE containing DHA and ARA in maternal venous plasma, in placenta and in umbilical cord venous and arterial plasma by LC–MS/ MS.

Metabolic profile of obese mothers was characterized by lower ester, ether and plasmalogen PC and PE species containing DHA compared to normal weight mothers. Differences achieving statistical significance are presented in Fig. 1A. Those PC and PE species with ARA were not affected by maternal obesity (Fig. 1B).

Then, we studied the influence of the fetal sex on maternal circulating phospholipids, by comparing the levels of all PC and PE species containing DHA and ARA detected in plasma of obese women to those in plasma of normal weight women carrying either a male or a female fetus. Interestingly, only LPE 22:6 was significantly decreased in obese mothers compared to normal weight mothers carrying a male (p=0.025, reduced by 50%) (Additional file 1: Table S3).

Fetal sex differentially influences the levels of PC and PE containing DHA and ARA in placenta of obese mothers

The evaluation of fetal sex differences in the placenta revealed distinct changes in pregnancies complicated by maternal obesity compared to healthy pregnancies. We observed decreases in the ester linked PC containing



Fig. 1 Differences in PC and PE containing DHA and ARA variables in maternal plasma between the two groups (control/obese) of mothers including both male and female fetuses. **A** Significant differences in PC and PE containing DHA in maternal plasma between control and obese groups. **B** PC and PE species containing ARA were not changed between control and obese mothers. Relative levels are expressed in mean \pm SD and significant differences pass at false-discovery rate-adjusted p value (Benjamini–Hochberg false discovery rate adjustment to account for multiple testing). **p* < 0.05

DHA and ARA in male placentas whereas the levels of almost all PE containing DHA and ARA including those with sn-1 ether and plasmalogen linkages were reduced in female placentas (Table 1). These findings suggest differing mechanisms for handling of LCPUFA in phospholipids in the placenta of male and female fetuses exposed to the obesogenic environment. These sex differences are all the more relevant because the levels of those phospholipid species were similar between the female and male placentas of mothers with normal weight (control) (Additional file 1: Figure S1). However, the specific lipids species impacted was highly influenced by fetal sex with the male placentas primarily and profoundly reduced in ester PC forms by 60–92% while the female placentas demonstrated significant decreases of 51–84% in PE forms.

Levels of PC and PE containing DHA and ARA were reduced in the circulation of male fetus exposed to maternal obesity but not in females

Interestingly, only male fetuses exposed to obesity demonstrated a lower level of phospholipids containing DHA and ARA in umbilical cord venous plasma (Table 2). In particular, we found reduced ester linked PC containing DHA (PC 18:0_22:6, PC 18:1_22:6) by 43 and 51%, ether and plasmalogen linked PE containing DHA (PE O-18:0_22:6, PE P-18:0_22:6) by 55 and 61%, and ARA (PE P-16:0_20:4, PE P-18:0_20:4) by 56 and 58%, respectively. We previously reported reduced PC 16:0_22:6 and LPC 22:6 (LPC-DHA) levels in male fetus of mothers with obesity [20]. The level of ether and plasmalogen PE species containing DHA and ARA remained unchanged in umbilical circulation of female fetuses (Table 2) despite a significant decrease in PE species containing DHA and ARA in female placentas of obese mothers. Similar to the phospholipid analysis in placenta of normal weight subjects, we observed no change in any DHA and ARA PC and PE species in umbilical cord venous plasma when comparing females and males of mothers with normal weight (controls) (Additional file 1: Figure S2).

Conversely to umbilical cord venous plasma, we found decreased sn-1 ester, ether and plasmalogen PC with ARA in umbilical cord arterial plasma of female fetuses exposed to obesity (Additional file 1: Table S4), and no change in those same species in males. Interestingly, LPC 16:0 and LPC 18:2 levels were increased in umbilical artery in the female not in male fetuses in pregnancy complicated by obesity.

Table 1 Profile of phosphatidylcholine (PC) and phosphatidylethanolamine (PE) containing DHA and ARA in placenta of obese (n = 16) compared to normal BMI women (n = 16) regarding the fetal sex

	Placenta						
Relative level/µg prot	Female			Male			
	Ctr (<i>n</i> =9)	Ob (<i>n</i> = 11)	p value (FDR)	Ctr (<i>n</i> =7)	Ob (<i>n</i> = 5)	p value (FDR)	
PC-DHA							
PC 16:0_22:6	0.03 ± 0.02	0.01 ± 0.008	0.126	0.030 ± 0.010	0.012 ± 0.005	0.045	
PC O-16:0_22:6	0.01 ± 0.01	0.006 ± 0.004	0.179	0.012 ± 0.007	0.004 ± 0.002	0.129	
PC P-16:0_22:6	0.009 ± 0.007	0.004 ± 0.003	0.126	0.008 ± 0.005	0.003 ± 0.001	0.202	
PC 18:0_22:6	0.02 ± 0.02	0.009 ± 0.005	0.101	0.025 ± 0.015	0.008 ± 0.003	0.039	
PC 18:1_22:6	0.01 ± 0.007	0.004 ± 0.004	0.098	0.010 ± 0.006	0.005 ± 0.002	0.157	
PC O-18:0_22:6	0.01 ± 0.01	0.003 ± 0.002	0.126	0.010 ± 0.008	0.003 ± 0.001	0.129	
PC O-18:1_22:6	0.02 ± 0.02	0.004 ± 0.002	0.101	0.013±0.010	0.003 ± 0.002	0.101	
PC 20:4_22:6	0.005 ± 0.004	0.003 ± 0.002	0.284	0.003 ± 0.002	0.002 ± 0.002	0.157	
LPC 22:6	0.0007 ± 0.0003	0.0004 ± 0.00008	0.098	0.001 ± 0.0001	0.0002 ± 0.00005	0.039	
PC-ARA							
PC 16:0_20:4	0.76 ± 0.58	0.24±0.18	0.098	0.74 ± 0.49	0.06 ± 0.01	0.029	
PC 16:1_20:4	0.02 ± 0.01	0.001 ± 0.001	0.101	0.02 ± 0.02	nd		
PC O-16:0_20:4	0.14 ± 0.11	0.05 ± 0.02	0.101	0.13 ± 0.09	0.03 ± 0.01	0.045	
PC P-16:0_20:4	0.13 ± 0.09	0.03 ± 0.02	0.098	0.12 ± 0.09	0.03 ± 0.02	0.101	
PC 18:0_20:4	0.56 ± 0.46	0.17±0.12	0.098	0.56 ± 0.41	0.15 ± 0.10	0.045	
PC 18:1_20:4	0.21 ± 0.14	0.05 ± 0.04	0.098	0.21±0.15	0.03 ± 0.03	0.039	
PC 18:2_20:4	0.08 ± 0.06	0.02 ± 0.01	0.098	0.09 ± 0.05	0.005 ± 0.003	0.029	
PC O-18:0_20:4	0.12 ± 0.12	0.05 ± 0.02	0.152	0.11 ± 0.08	0.06 ± 0.04	0.222	
PC O-18:1_20:4	0.28 ± 0.23	0.09 ± 0.04	0.165	0.25 ± 0.18	0.11±0.07	0.222	
PC 20:1_20:4	0.02 ± 0.01	0.006 ± 0.003	0.101	0.01 ± 0.01	0.006±0.003	0.222	
PC 20:2_20:4	0.01 ± 0.009	0.005 ± 0.003	0.152	0.01 ± 0.007	0.005 ± 0.005	0.129	
PC 20:3_20:4	0.03 ± 0.01	0.009 ± 0.006	0.098	0.02 ± 0.01	0.009 ± 0.005	0.280	
PC 20:4_20:4	0.02 ± 0.01	0.006 ± 0.003	0.098	0.02 ± 0.01	0.003 ± 0.002	0.039	
LPC 20:4	0.09 ± 0.07	0.06 ± 0.05	0.359	0.07 ± 0.06	0.04 ± 0.01	0.471	
LPC							
LPC 16:0	0.13 ± 0.05	0.171±0.12	0.534	0.06 ± 0.03	0.14±0.15	0.434	
LPC 16:1	0.002 ± 0.001	0.005 ± 0.003	0.165	0.002 ± 0.0002	0.002 ± 0.001	0.825	
LPC 18:0	0.09 ± 0.07	0.06 ± 0.05	0.567	0.07 ± 0.06	0.04 ± 0.01	0.471	
LPC 18:1	0.02 ± 0.01	0.02 ± 0.009	0.746	0.01 ± 0.008	0.02 ± 0.02	0.646	
LPC 18:2	0.01 ± 0.02	0.01 ± 0.008	0.699	0.01 ± 0.009	0.01 ± 0.01	0.471	
LPC 20:3	0.01 ± 0.07	0.005 ± 0.003	0.301	0.009 ± 0.007	0.003 ± 0.002	0.210	
PE-DHA							
PE 16:1_22:6	0.04 ± 0.02	0.01 ± 0.006	0.034	0.03 ± 0.03	0.02 ± 0.01	0.650	
PE 16:0_22:6	0.07 ± 0.03	0.03 ± 0.01	0.030	0.06 ± 0.03	0.05 ± 0.03	0.704	
PE O-16:0_22:6	0.40 ± 0.28	0.14 ± 0.07	0.037	0.34 ± 0.24	0.22±0.15	0.649	
PE P-16:0_22:6	0.45 ± 0.29	0.14 ± 0.08	0.030	0.41±0.27	0.22 ± 0.16	0.297	
PE 18:0_22:6	0.19 ± 0.09	0.08 ± 0.04	0.030	0.18±0.10	0.11 ± 0.08	0.274	
PE O-18:0_22:6	0.10 ± 0.08	0.02 ± 0.01	0.037	0.09 ± 0.09	0.03 ± 0.02	0.560	
PE P-18:0_22:6	0.31±0.26	0.05 ± 0.02	0.030	0.29±0.28	0.07 ± 0.05	0.234	
PE 18:1_22:6	$0.11 \pm 0.06 (n = 4)$	0.06±0.03	0.133	0.10±0.03 (n=4)	0.09 ± 0.07	0.767	
PE 18:2_22:6	0.03 ± 0.01	0.01 ± 0.007	0.030	0.03 ± 0.02	0.02 ± 0.01	0.369	
LPE 22:6	0.005 ± 0.002	0.003 ± 0.002	0.119	0.005 ± 0.002	0.003 ± 0.002	0.283	
PE-ARA							
PE 16:1_20:4	0.26 ± 0.15	0.09 ± 0.04	0.037	0.21 ± 0.12	0.09 ± 0.02	0.234	
PE 16:0_20:4	1.65 ± 0.95	0.98 ± 0.37	0.120	1.43 ± 0.87	0.80 ± 0.44	0.297	

Relative level/µg prot	Placenta						
	Female			Male			
	Ctr (<i>n</i> =9)	Ob (<i>n</i> = 11)	p value (FDR)	Ctr (<i>n</i> =7)	Ob (<i>n</i> = 5)	p value (FDR)	
PE O-16:0_20:4	0.50±0.27	0.23±0.09	0.037	0.41±0.20	0.21±0.07	0.202	
PE P-16:0_20:4	2.42 ± 1.43	1.11 ± 0.45	0.037	2.06 ± 1.02	0.97 ± 0.28	0.202	
PE 18:0_20:4	2.26 ± 1.19	1.10 ± 0.36	0.037	2.03 ± 1.03	0.99 ± 0.34	0.202	
PE O-18:0_20:4	0.66 ± 0.45	0.22 ± 0.09	0.037	0.59 ± 0.44	0.17 ± 0.04	0.202	
PE P-18:0_20:4	2.95 ± 1.93	1.01 ± 0.43	0.037	2.67 ± 1.93	0.77 ± 0.17	0.195	
PE 18:1_20:4	0.75 ± 0.37	0.36 ± 0.13	0.037	0.63 ± 0.27	0.32 ± 0.13	0.202	
PE 18:2_20:4	0.18 ± 0.09	0.08 ± 0.03	0.037	0.16 ± 0.05	0.07 ± 0.04	0.195	
PE 20:4_20:4	0.02 ± 0.009	0.02 ± 0.007	0.114	0.02 ± 0.004	0.01 ± 0.005	0.202	
LPE 20:4	0.06 ± 0.08	0.01 ± 0.01	0.093	0.06 ± 0.03	0.002 ± 0.002	0.234	

Table 1 (continued)

20:4 = ARA, 22:6 = DHA, PC = phosphatidylcholine, PE = phosphatidylethanolamine, LPC = lysophosphatidylcholine, LPE = lysophosphatidylethanolamine. In grey all "O" = ether, "P" = plasmalogen species, nd = not detected. Values are the mean ± SD. In bold, *p* value < 0.05 after FDR calculation

Placental remodeling pathway of phospholipids is differently affected by the fetal sex in pregnancy complicated by obesity

Enzymes involved in the de novo phospholipid synthesis (Kennedy pathway and Lands cycle) were evaluated in placenta homogenates (Additional file 1: Table S2). Additional file 1: Table S5 reports three placental enzymes that were not altered at protein level in association to maternal obesity in male and in female placentas. Relative protein expression of placental Fatty Acyl-CoA Reductase 1 (FAR1), the rate-limiting step for ether lipid biosynthesis in peroxisome is presented in Fig. 2A and was also unchanged in both fetal sexes exposed to maternal obesity.

We also examined the phospholipid remodeling pathway by evaluating expression of PhosphoLipase A_2 Group IVC (PLA₂G4C or iPLA₂) which hydrolyzes acyl groups from PC to form the lysophospholipid (LPC) form and found that PLA₂G4C abundance was lower in male placentas of obese compared to control mothers (Fig. 2B). We determined the protein expression of LysoPhosphatidylCholine Acyl Transferase 4 (LPCAT4) which catalyzes the re-esterification of acyl chains to LPC to regenerate PC with a unique acyl chain combination and observed a reduction of LPCAT4 abundance in female placentas in pregnancy complicated by obesity (Fig. 2C).

Relationship between maternal obesity and placental and fetal levels of ether and plasmalogen PE phospholipids

In order to examine whether maternal obesity has an incremental impact on placental or cord ether and plasmalogen phospholipids containing DHA or ARA, we tested the relationship between maternal BMI and lipid levels in placenta and umbilical cord. We found an inverse correlation between maternal BMI and the placental tissue levels of the eight ether and plasmalogen PE species containing DHA and ARA in female placentas (Fig. 3). We also found significant inverse correlations between maternal BMI and the levels of four circulating ether and plasmalogen PE species with DHA and ARA in umbilical cord venous plasma of male fetuses (Fig. 4).

We provide a summary of the aforementioned results, demonstrating the influence of fetal sex on maternal–placental–fetal adaptation of the DHA and ARA for ester, ether and plasmalogen PC and PE in maternal obesity in Fig. 5.

Discussion

Our study demonstrates a strong influence of pre-pregnancy maternal BMI on phosphatidylcholine and phosphatidylethanolamine species containing LCPUFA (DHA and ARA) in three compartments: maternal circulation, placenta and fetal umbilical circulation. These changes may have consequences for placental function, birth weight, and fetal brain development as well as lipid signaling regarding the fetal sex. We found significant differences in levels of sn-1 ester, and particularly sn-1 ether and plasmalogen PC and PE containing DHA and ARA in all three compartments in pregnancies complicated by obesity and changes in placenta and cord plasma were highly sex specific. This is the first report to suggest that sn-1 ether and plasmalogen phospholipids containing crucial LCPUFA may play a key role in maternal to fetal transfer by the placenta, and possibly act as a reservoir of vital DHA and ARA for the fetus. We also speculate that the female placenta is able to compensate in the obese environment to more effectively deliver vital lipids to the fetal compartment.

Table 2 Profile of phosphatidylcholine (PC) and phosphatidylethanolamine (PE) containing DHA and ARA in umbilical cord vein from obese (n = 16) compared to normal BMI women (n = 16) regarding the fetal sex

	Umbilical cord vein							
Relative level	Female			Male				
	Ctr (<i>n</i> =9)	Ob (<i>n</i> = 11)	q value (FDR)	Ctr (<i>n</i> = 7)	Ob (<i>n</i> = 5)	q value (FDR)		
PC-DHA								
PC 14:0_22:6	0.02 ± 0.01	0.01 ± 0.008	> 0.999	0.01 ± 0.007	0.007 ± 0.005	0.258		
PC 16:0_22:6	2.79 ± 1.10	2.14 ± 0.76	0.724	3.25 ± 0.67	1.86 ± 0.54	0.046		
PC O-16:0_22:6	0.19 ± 0.08	0.12 ± 0.06	0.675	0.16 ± 0.04	0.17±0.21	0.242		
PC P-16:0_22:6	0.08 ± 0.03	0.04 ± 0.02	0.673	0.09 ± 0.03	0.04 ± 0.03	0.057		
PC 18:0_22:6	1.87 ± 0.88	1.35 ± 0.50	0.675	1.97±0.23	0.97±0.32	0.041		
PC 18:1_22:6	0.14 ± 0.05	0.11 ± 0.03	0.675	0.14±0.02	0.10 ± 0.02	0.057		
PC O-18:0_22:6	0.17±0.11	0.13 ± 0.05	0.813	0.17±0.06	0.12 ± 0.09	0.363		
PC O-18:1 22:6	0.14 ± 0.09	0.09 ± 0.04	0.675	0.12 ± 0.07	0.09 ± 0.09	0.475		
LPC 22:6	0.35 ± 0.10	0.30 ± 0.10	0.813	0.39 ± 0.04	0.19 ± 0.05	0.041		
PC-ARA								
PC 14:0 20:4	0.009 ± 0.002	0.01 ± 0.003	0.813	0.01 ± 0.005	0.01 ± 0.005	0.914		
PC 16:0 20:4	9.47 ± 6.90	10.7 ± 2.50	0.813	12.02±3.97	7.03 ± 5.10	0.242		
PC 16:1 20:4	0.15 ± 0.17	0.04 ± 0.01	0.831	0.15 ± 0.15	0.06 ± 0.08	0.305		
PC O-16:0 20:4	0.89 ± 0.59	1.25 ± 0.39	0.675	1.51±0.73	0.77 ± 0.45	0.216		
PC P-16:0 20:4	0.58 ± 0.51	0.67 ± 0.29	0.813	0.95 ± 0.49	0.39 ± 0.24	0.082		
PC 18:0 20:4	9.20+6.14	10.3 + 1.97	0.813	12.4+2.64	6.19+4.29	0.057		
PC 18:1 20:4	0.70 ± 0.55	0.82 ± 0.23	0.813	0.96 ± 0.30	0.67 ± 0.22	0.482		
PC 18:2 20:4	0.07 ± 0.03	0.07 ± 0.02	0.813	0.10 ± 0.03	0.05 ± 0.02	0.113		
PC Q-18:0 20:4	1.41 + 1.19	1.29+0.57	0.813	1.47+0.92	1.25 + 1.30	0.743		
PC Q-18:1_20:4	1 32 + 0 74	143+061	0.813	166+0.89	0.88+0.31	0.299		
PC 20:1 20:4	0.07 ± 0.03	0.08 ± 0.02	0.675	0.08+0.03	0.07 ± 0.01	0.951		
PC 20:2 20:4	0.04 ± 0.02	0.05 ± 0.02	0.813	0.00 ± 0.00	0.05 ± 0.01	0.840		
PC 20:3 20:4	0.05 ± 0.01	0.05 ± 0.01	>0.999	0.04 ± 0.01	0.04 ± 0.02	0.943		
PC 20:4 20:4	0.03 ± 0.07	0.04 ± 0.01	0.813	0.04 ± 0.01	0.02 ± 0.002	0.055		
I PC 20:4	936+351	836+236	0.813	8 26 + 1 37	5 97 + 1 36	0.082		
I PC	2100 - 210 1	0.00 _ 2.00	0.010	0.20 2 1.57	5157 _ 1150	0.002		
LPC 16:0	538+256	669+177	0.675	814+477	516+92	0.639		
LPC 16:1	287+150	351+093	0.724	249+10	277+045	0.920		
LPC 18:0	930 + 594	641+153	0.675	665+155	5 25 + 1 78	0.242		
LPC 18:1	151+512	165+398	0.813	137+424	108+219	0.242		
LPC 18:2	1999+89	2314+739	0.813	258+132	135+422	0.113		
LPC 18:3	033+020	031+012	>0.999	0.25+0.03	0.20 ± 0.05	0.159		
LPC 20:3	4.16+1.02	4.62 + 1.55	0.813	4.37+0.96	2.68+0.81	0.055		
LPC 20:5	0.30 ± 0.14	0.29 ± 0.14	0.934	033+010	0.13 ± 0.05	0.055		
LPC 22:0	0.20 ± 0.13	0.11+0.03	0.813	0.17+0.06	0.09 ± 0.03	0.057		
PF-DHA	0.20 ± 0.15	0.11 ± 0.05	0.015	0.17 ± 0.00	0.09 ± 0.05	0.007		
PF 16.1 22.6	0.06+0.03	0.05+0.02	>0.999	0.06+0.01	0.04 ± 0.01	0.074		
PE 16:0 22:6	0.85 ± 0.34	0.05 ± 0.02	0.804	1 26 + 0 37	0.74 ± 0.33	0.074		
PE Q-16:0_22:0	0.03 ± 0.01	0.30 + 0.07	0.456	0.51+0.37	0.30 ± 0.08	0.545		
PE P-16:0_22:6	0.81+0.42	0.50 ± 0.07 0.52 ± 0.23	0.329	0.98 + 0.59	0.30 ± 0.00	0.074		
PE 18:0 22:6	0.57 ± 0.12 0.52 ± 0.35	0.52 ± 0.23 0.45 ± 0.21	>0.999	0.66 ± 0.29	0.17 ± 0.13 0.35 ± 0.12	0.082		
PF ()-18-0 22-0	0.07+0.03	0.07+0.02	> 0.999	0.09+0.02	0.04 + 0.008	0.024		
PF P-18:0 22:6	0.24+0.14	0.07 ± 0.02 0.15 ± 0.06	0.329	0.34 + 0.24	0.13+0.03	0.024		
PF 18·1 22:0	0.07+0.01	0.08+0.04	>0.999	0.11+0.02	0.06+0.02	0.048		
L PE 22:6	0.12+0.05	0.00 ± 0.04	> 0.999	0.14+0.07	0.11 + 0.02	0.906		
LPC 16:0 LPC 16:1 LPC 18:0 LPC 18:1 LPC 18:2 LPC 18:3 LPC 20:3 LPC 20:5 LPC 22:4 PE-DHA PE 16:1_22:6 PE 0-16:0_22:6 PE 0-16:0_22:6 PE P-16:0_22:6 PE 0-18:0_22:6 PE 0-18:0_22:6 PE 0-18:0_22:6 PE 0-18:0_22:6 PE 18:1_22:6 LPC 22:6	53.8 ± 25.6 2.87 ± 1.50 9.30 ± 5.94 15.1 ± 5.12 19.99 ± 8.9 0.33 ± 0.20 4.16 ± 1.02 0.30 ± 0.14 0.20 ± 0.13 0.06 ± 0.03 0.85 ± 0.34 0.43 ± 0.21 0.81 ± 0.42 0.52 ± 0.35 0.07 ± 0.03 0.24 ± 0.14 0.07 ± 0.01 0.12 ± 0.05	$b 0.9 \pm 17.7$ 3.51 ± 0.93 6.41 ± 1.53 16.5 ± 3.98 23.14 ± 7.39 0.31 ± 0.12 4.62 ± 1.55 0.29 ± 0.14 0.11 ± 0.03 0.05 ± 0.02 0.95 ± 0.29 0.30 ± 0.07 0.52 ± 0.23 0.45 ± 0.21 0.07 ± 0.02 0.15 ± 0.06 0.08 ± 0.04 0.12 ± 0.05	0.875 0.724 0.675 0.813 0.999 0.813 0.934 0.813 > 0.999 0.804 0.456 0.329 > 0.999 > 0.999 0.329 > 0.999 > 0.999 > 0.999	81.4 ± 47.7 2.49 ± 1.0 6.65 ± 1.55 13.7 ± 4.24 25.8 ± 13.2 0.25 ± 0.03 4.37 ± 0.96 0.33 ± 0.10 0.17 ± 0.06 0.06 ± 0.01 1.26 ± 0.37 0.51 ± 0.37 0.98 ± 0.59 0.66 ± 0.29 0.09 ± 0.02 0.34 ± 0.24 0.11 ± 0.03 0.14 ± 0.07	51.6 ± 9.2 2.77 ± 0.45 5.25 ± 1.78 10.8 ± 2.19 13.5 ± 4.22 0.20 ± 0.05 2.68 ± 0.81 0.13 ± 0.05 0.09 ± 0.03 0.04 ± 0.01 0.74 ± 0.33 0.30 ± 0.08 0.47 ± 0.15 0.35 ± 0.12 0.04 ± 0.008 0.13 ± 0.03 0.06 ± 0.02 0.11 ± 0.03	0.639 0.920 0.242 0.113 0.159 0.055 0.055 0.055 0.057 0.074 0.074 0.074 0.074 0.074 0.074 0.024 0.024 0.048 0.048 0.048		

Relative level	Umbilical cord vein						
	Female			Male			
	Ctr (<i>n</i> =9)	Ob (<i>n</i> = 11)	q value (FDR)	Ctr (<i>n</i> =7)	Ob (<i>n</i> = 5)	q value (FDR)	
PE-ARA							
PE 16:1_20:4	0.14 ± 0.10	0.08 ± 0.03	0.329	0.12 ± 0.06	0.06 ± 0.02	0.232	
PE 16:0_20:4	5.29 ± 2.03	6.97 ± 1.84	0.329	6.02 ± 1.45	6.22 ± 2.02	0.805	
PE O-16:0_20:4	0.68 ± 0.28	0.68 ± 0.28	> 0.999	0.76 ± 0.23	0.48 ± 0.16	0.082	
PE P-16:0_20:4	2.40 ± 1.18	2.12 ± 0.86	> 0.999	2.75 ± 1.08	1.21 ± 0.46	0.049	
PE 18:0_20:4	4.19 ± 1.46	4.76 ± 1.06	0.771	5.17 ± 1.67	3.82 ± 1.80	0.294	
PE O-18:0_20:4	0.56 ± 0.33	0.45 ± 0.18	> 0.999	0.64 ± 0.16	0.35 ± 0.17	0.082	
PE P-18:0_20:4	1.83 ± 1.24	1.37±0.51	> 0.999	2.19 ± 0.75	0.90 ± 0.50	0.048	
PE 18:1_20:4	0.63 ± 0.22	0.74 ± 0.25	> 0.999	0.64 ± 0.19	0.64 ± 0.19	0.957	
PE 18:2_20:4	0.04 ± 0.01	nd		0.08 ± 0.07	0.06 ± 0.007	0.945	
LPE 20:4	1.24±0.19	1.86±0.57	0.329	1.26±0.29	1.39±0.91	0.892	

Table 2 (continued)

20:4 = ARA, 22:6 = DHA, PC = phosphatidylcholine, PE = phosphatidylethanolamine, LPC = lysophosphatidylcholine, LPE = lysophosphatidylethanolamine. In grey all "O" = ether, "P" = plasmalogen species, nd = not detected. Values are the mean ± SD. In bold, *p* value < 0.05 after FDR calculation

In the maternal circulation of obese women, we found lower levels in several phospholipid species with DHA, including sn-1 ester linked PE 16:0_22:6 and PE 18:0_22:6, sn-1 ether linked PE O-18:0_22:6 and sn-1 plasmalogen linked PC P-16:0_22:6 and PE P-18:0_22:6 but we found no differences in ARA linked phospholipids (Fig. 1). Interestingly, the lower maternal levels of PC and PE species containing DHA were not influenced by fetal sex (Additional file 1: Table S3) and are more likely to be a direct dietary origin. Also, the reduced DHA content in the maternal circulation is related to the obesogenic phenotype and not to the sex of the fetus. These results are in line with our previous study showing reduced DHA in the total circulating phospholipid fraction in plasma of obese women [20]. Lower circulating DHA levels of obese women reflect changes in phospholipid metabolism related to her diet which is often poor in n-3 LCPUFA and high n-6 LCPUFA including ARA. Lower DHA levels and higher n-6 LCPUFA/n-3 LCPUFA ratio have been previously reported in obese pregnant women [41]. In addition, estrogen enhanced the conversion of linolenic acid (ALA, n-3 LCPUFA) into DHA, by increasing the expression of FADS2, the rate-limiting enzyme in the conversion of ALA into DHA, and other transcription factors like PPAR- α , synergically with the low linoleic acid (LA, n-6 LCPUFA) diet [42]. In pregnant obese women, lower estrogen level as well as higher LA diet [41] are described and could explain in part the lower DHA level in circulating phospholipids in the mother.

Among the five PC and PE species containing DHA and decreased in maternal plasma from obese mothers, three are ether and plasmalogen species. Plasmalogens are a subset of ether glycerophospholipids that are characterized by a *cis* double bond adjacent to the ether linkage, forming a vinyl ether linkage. They are considered the most abundant and biologically active class of ether lipids [21, 43]. The distribution of these lipids is variable among tissues with a high content in the brain and paradoxically low levels in liver. The low ether lipid content, including plasmalogens, in the liver may be due to rapid release and incorporation into lipoproteins for transport to other tissues, such as the brain [44]. Several studies have reported that lower ether lipid content in serum is associated with hypertension and obesity [30, 31]. In the context of pregnancy, lower sn-1 ether PC species with DHA suggests reduced DHA enrichment of lipoproteins (HDL, LDL, VLDL) which are essential for lipid transfer across the placenta.

We also observed that maternal obesity affects the placental level of sn-1 ester, ether and plasmalogen PC and PE containing both DHA and ARA but with differential effects of fetal sex (Table 1). In the same cohort, we have previously shown that the protein expression of major fatty acid transporters (FATP1-4, FATP6, FABPM, FAT/CD36) was unchanged in microvillous membrane of the placenta (MVM, maternal facing plasma membrane) in both fetal sexes [20]. Gene (mRNA) expression of placental FATP1 and FATP4 has been correlated with the percent of phospholipids containing DHA in maternal plasma in normal pregnancy indicating a key role of those transporters for uptake of DHA from the mother [45]. However, these fatty acid transporters can also transport other fatty acids such as the saturated and monounsaturated fatty acids (SFA and MUFA). With lower DHA in circulation of obese mothers, the action



Fig. 2 Representative image of capillary electrophoresis bands and relative protein abundance **A** of fatty acyl-CoA reductase 1 (FAR1) in placental homogenates in obese vs control women in male and female groups obtained by simple West, **B** of PhosphoLipase A2 Group IVC (PLA2G4C or iPLA2) and **C** of LysophosPhatidylCholine Acyl Transferase 4 (LPCAT4). **B** and **C** were analyzed by western blot. Ctr-M=control-Male, Ob-M=Obese-Male, Ctr-F=control-Female, Ob-F=Obese-Female. Data are means \pm SD. *p < 0.05

of these transporters in the placenta is likely to take up additional SFA or MUFA or n-6 LCPUFA all of which are highly abundant in maternal plasma. In our study, a

decrease of DHA esterification into PC (PC 16:0_22:6, PC 18:0_22:6 for males) and PE (PE 16:0_22:6, PE 16:1_22:6, PE 18:0_22:6, PE 18:2_22:6 and similar PE species with



Fig. 3 Inverse correlations between maternal BMI and levels of ether and plasmalogen linked PE with DHA and ARA in placentas of female fetus. Spearman test was performed to assess correlations, r and p values are in the figure, *p < 0.05, **p < 0.01, **p < 0.005



Fig. 4 Inverse correlations between maternal BMI and levels of ether and plasmalogen linked PE with DHA and ARA in umbilical cord venous plasma of male fetus. Spearman test was performed to assess correlations, r and p values are in the figure, *p < 0.05, **p < 0.01, ***p < 0.005

ether and plasmalogen linkages for females) (Table 1) in the placental phospholipid fraction could, in part, be explained by lower DHA substrate availability rather than a deficient transport capacity in MVM for uptake of DHA by both male and female placentas [46]. However, obese mothers with both male and female fetuses had lower DHA content in their plasma but placental phospholipid species levels were distinct in the sexes suggesting that metabolism and transfer of DHA differs by fetal sex and that maternal supply is not the only parameter of interest. This suggestion is also valid for ARA which is not altered in maternal plasma by obesity but is differentially reduced in placental PC and PE when analyzing the data based on fetal sex (Table 1). Decreased placental PC and PE with DHA and ARA in pregnancies complicated by maternal obesity may be due to increased phospholipid synthesis with SFA and MUFA at the sn-2 position to maintain phospholipid content in the expanding exchange surface area of the placenta across gestation. That process may take precedence over delivery of LCPUFA to the fetus when DHA is not readily available. In this study, we did not quantify the PC and PE species with specific SFA and/or MUFA in both sn-1 and sn-2 positions, therefore we provide this as a speculation for how placentas might respond to low DHA availability. Placental sex differences in PC and PE containing DHA and ARA were not found in pregnancies with normal weight mothers (Additional file 1: Figure S1) suggesting the findings in this report are a specific response to maternal obesity that differs by fetal sex.

Another possible explanation for the placental decreases in preferred forms of PC and PE containing LCPUFA could be that other lipid complexes such are triacylglycerols (TG), cellular storage form, are synthesized at the expense of incorporation of DHA and ARA into phospholipids. In our previous study, we have shown that DHA and ARA levels in total TG remained stable in both male and female placentas of obese women while the non-esterified form of both FAs were decreased along with a trend to be decreased in total phospholipid pool only in male placenta. This indicates a possible preferential mobilization of DHA and ARA toward the triglyceride form [20]. Since obesity during pregnancy is associated with placental inflammation and hypoxic stress [47], increased storage of LCPUFA in TG may be similar to what has been previously reported for intrauterine growth restriction (IUGR) placentas [48, 49].

Interestingly, we show a specific decrease of particular sn-1 ether and plasmalogen PE species with DHA and ARA in placentas from obese mothers carrying female



*PC 16:0_22:6, LPC-22:6 published in Powell et al BBA lipids 2021

Fig. 5 Summary of the influence of fetal sex in ester, ether and plasmalogen PC and PE containing DHA and ARA in the maternal–fetal unit in response to maternal obesity. The phenotype of obese mothers includes reduced ester, ether and plasmalogen PC and PE containing DHA but not those phospholipids containing ARA regardless of fetal sex. In obese mother carrying a male baby: in placenta, all the ester linked PC containing DHA and ARA are decreased by obesity and only one ether PC with DHA species while all ester linked PE were unchanged suggesting downregulated PEMT enzyme activity to convert PE into PC. In addition, decreased LPC-DHA is likely due to decreased iPLA2 protein expression. In fetal circulation, several ester PC DHA and ARA species and ester, ether and plasmalogen PE with DHA and ARA were reduced suggesting a defect of LCPUFA transfer capacity. In obese mother carrying a female baby: in placenta, all PE species with DHA and ARA were decreased but not PC species and LPCAT4 expression was lower. These results suggest an upregulated PEMT activity to maintain PC production. No change was observed in female umbilical cord. By decreasing all ether and plasmalogen PE, species considered as reservoir for DHA and ARA in the placenta, female fetuses of obese mothers show a high fetal–placental adaptability and placental reserve capacity that can maintain the PC-LCPUFA synthesis and the transfer of these crucial species to the fetus to preserve growth and brain development

fetuses (Table 1). Since ether and plasmalogen lipids are specifically synthesized in the peroxisome [43], lower levels in the female placentas of obese women in our study may suggest a defect of peroxisomal ether linked phospholipid production. Peroxisomes are membrane bound organelles that perform multiple functions, including ether lipid synthesis and fatty acid oxidation [50]. In the same cohort, we have previously shown a down-regulation of beta-oxidation in female placentas of obese women suggesting a mitochondrial dysfunction and perhaps a peroxisome deficiency [20]. Mitochondrial dysfunction has been observed in trophoblast cells of female placentas in response to maternal obesity resulting from an activation of signaling from inflammation via NFkB1 and miR-210 [51]. Peroxisomal dysfunction is characterized by reduced levels of plasmalogens contributing to various metabolic pathologies such as dyslipidemia, obesity, NAFLD and T2D possibly through multiple pathways, including the disruption of cellular membranes, increased oxidative stress, endoplasmic reticulum (ER) stress and inflammation [50].

The production of fatty alcohols from fatty acyl-CoA is needed for generation of the ether bond and is mediated by FAR1. This has been proposed as rate-limiting step for ether lipid biosynthesis in the peroxisome [52]. Substrate availability and activity of FAR1 are essential regulators of the entire ether phospholipid synthesis pathway [53]. FAR1 accounting for C16:0, C18:0, and C18:1 fatty alcohol synthesis is post-translationally regulated by the accumulation of plasmalogen PE in the inner leaflet of the plasma membrane [54]. In our study, placental FAR1 protein expression was not modified by maternal obesity in either fetal sex (Fig. 2A) while the placental plasma membrane levels of plasmalogen PE containing DHA and ARA were decreased. Since DHA and ARA are transferred to PE during the remodeling pathway (land's cycle), FAR1 activity is likely upstream regulated by PE containing other FAs in sn-2 position than DHA or ARA.

Two other peroxisomal enzymes play an important role in the initiation of the ether lipid synthesis; the glyceronephosphate O-acyltransferase (GNPAT), that acylates dihydroxyacetone phosphate (DHAP) with a fatty acyl-CoA at the sn-1 position and the alkylglycerone phosphate synthase (AGPS) that catalyzes the exchange of the fatty acyl group of 1-acyl-DHAP for a fatty alcohol [43]. However, we were not able to quantify the expression of those enzymes at the protein level in the placenta due to lack of specific antibodies.

Despite the unchanged expression in FAR1 enzyme, lower DHA supplied in the plasma of obese mothers may alter the ether and plasmalogen PE and PC synthesis outside the peroxisome. Indeed, phospholipid synthesis is completed at the ER mediated by specific enzymes involved in the de novo Kennedy pathway and remodeling pathways (Land's cycle) [43]. We found lower levels of sn-1 ether and plasmalogen linked PE with DHA and ARA, and no change in sn-1 ether and plasmalogen PC with DHA and ARA in female placentas of obese women suggesting an enhanced phosphatidylethanolamine methyl transferase (PEMT) enzyme activity to maintain de novo synthesis of ether and plasmalogen PC in the ER. This may act as a reservoir of DHA and ARA to be transferred to the fetus through generation of PC species which can be transporters as LPC through the MFSD2a transporter in the basal plasma membrane. This placental response to obesity may be a compensatory mechanism in response to deficient phospholipid remodeling pathway as indicated by the decreased expression of LPCAT4 in female placentas (Fig. 2C). Furthermore, the unchanged levels of PC and PE containing DHA and ARA in the umbilical cord venous plasma in pregnancies carrying female fetuses (Table 2) as well as our previous study showing that LPC-DHA was not changed in female placentas and in female fetal umbilical circulation exposed to obesity [20] is consistent with the concept that the fetus and placenta have the ability to adapt to preserve the transfer capacity of LCPUFA to the developing female fetus. We observed an inverse correlation between the placental level of sn-1 ether and plasmalogen PE containing DHA and ARA species and maternal BMI (Fig. 3) combined with altered expression of LPCAT4 enzyme, which potentially reduces the re-esterification of those PE species through incorporation of a LCPUFA into the sn-2 position of LPC. This complexity suggests a possible interplay between peroxisome derived ether phospholipids and the specific enzymes involved in the phospholipid remodeling pathway in placentas of obese mothers carrying a female fetus. We have attempted to summarize these findings in Fig. 5 for clarity.

In obese pregnant mothers carrying males, we observed levels of sn-1 ester linked PC with DHA and

ARA profoundly reduced in placenta, and no change in PE forms (Table 1). The lower PC DHA and ARA level in males is likely dependent on the PEMT pathway that converts PE into PC. While unable to quantify PEMT protein expression in the placenta, we speculate that the PEMT enzyme activity is downregulated in male placentas exposed to maternal obesity. Moreover, lower levels of major PC-DHA species (PC 16:0 22:6, PC 18:0 22:6) likely lead to decreased LPC-DHA in male placentas of obese mothers as we have recently reported [20]. This may be indirectly mediated by decreased PLA2G4C (iPLA2) protein expression in male but not in female placentas (Fig. 2B). Indeed, iPLA₂ cleaves at the sn-2 position of PC and preferentially released LCPUFA. Less DHA released in the remodeling phospholipid pathway may be reflected in decreased non-esterified DHA [20]. In addition, lower levels of PC containing DHA species in placenta may cause a defect of choline and DHA released and transferred to the male fetus with potential consequences for the brain development, since choline promotes the DHA transport as LPC through the blood brain barrier mediated by the mammalian family super domain 2a, MSFD2a transporter [25].

Lower circulating choline may be related to the observation of a reduced levels of ether and plasmalogen PC and PE containing DHA and ARA in the fetal circulation of males of obese mothers. Indeed, choline is a substrate of all forms of PC synthesis, and in case of ester PC linkage, can be converted into sphingomyelin (SM) metabolites including SM containing very long chain fatty acid, such as nervonic acid (24:1). This fatty acid is exclusively catabolized in the peroxisome and its breakdown is considered a source for alkyl groups in plasmalogen synthesis [55]. Inversely, a supplementation of choline to pregnant mice fed with a high-fat diet increases the relative abundance of hepatic plasmalogen and sphingomyelin d42:2 (SM d18:1_24:1, containing nervonic acid) in fetus at 17.5 day of gestation and after 6 weeks postnatal in male offspring, indicating an antioxidative response to protect the mouse liver from damages due to high-fat feeding [56]. Another study has shown that choline supplementation increased the incorporation of DHA into cholinecontaining phospholipids in a mouse model of maternal obesity [57].

Lower levels of ether and plasmalogen phospholipids at birth in male infants (Table 2) may be a signature for higher risk to develop metabolic diseases later in life. In children diagnosed with type 1 diabetes, reduced ether lipids (such as PE O-18:1_20:4) were detected in the serum prior to the detection of autoantibodies [58]. Male fetuses have higher nutritional requirements as evidenced by studies observing that women carrying male babies consume more nutrients, especially lipids [59]. Thus, male fetal growth is particularly sensitive to maternal dietary content and a diet deficient in n-3 LCPUFA in obese mothers appears to more profoundly affect male rather than female fetuses with respect to circulating DHA in the umbilical circulation.

Although the effect of decreased DHA supply on brain growth and development remains to be fully established, we speculate that altered placental PC-DHA metabolism and less DHA availability in circulation of male fetuses exposed to obesity may negatively influence the development of the brain, programming early life neurodevelopmental disorders. Indeed, dietary n-3 fatty acid deficiency reduces DHA-regulated PLA₂ and calcium-independent iPLA₂ in the frontal cortex in rodents and subsequently downregulates the brain phospholipid remodeling pathway [60].

The mechanisms that drive the sex-specific adaptations that we show in the placental phospholipid metabolism including the ether linked PC and PE containing DHA and ARA in the setting of obesity (Fig. 5) remain largely unexplored. Literature evidences suggest that the fetoplacental hormonal environment could play a major contributor. An important role of androgens was described in regulation of fetal growth and development [61] and changes in the placental–fetal hormonal environment due to maternal obesity may contribute to sex-specific differences in placental lipid metabolism. Further studies need to be developed to determine the hormonal effect on ether lipid in placenta and in fetal circulation in both sexes in normal pregnancy and in pregnancy complicated by obesity.

Limitations of this study include the small number of enrolled women and smaller sample size in each fetal sex group and lack of dietary information from the pregnant women. In addition, we were not able to find acceptable antibodies for many peroxisome and lipid synthesis enzymes involved in ether and plasmalogen synthesis. One of the major strengths of the study is that we have specifically identified the plasmalogen and ether linked phospholipids and analyzed those species in maternal and umbilical cord plasma as well as placentas of healthy mothers and those complicated by obesity and that those maternal–placental–fetal samples provide from the same pregnancy.

Perspectives and significance

We show that maternal obesity reduces the content of placental PC and PE containing LCPUFA differentially in women carrying male compared to female infants. This difference has potential consequences on supply of this preferred lipid form for fetal brain development with short- and long-term consequences. This study is the first to explore the two important phospholipids; PC and PE containing DHA and ARA, and examined sn-1 ester, ether and plasmalogen linkages, in maternal and fetal circulation and in placental tissue to uncover potential novel roles for ether and plasmalogen lipids in the regulation of placenta delivery of these vital nutrients in pregnancies complicated by obesity. By decreasing all ether and plasmalogen PE, species which are considered a reservoir for DHA and ARA in the placenta, female fetuses of obese mothers show a high fetal-placental adaptability and placental reserve capacity that can maintain the PC-LCPUFA (including LPC-DHA) synthesis and the transfer of these crucial species to the fetus to preserve growth and brain development. Male fetuses, in response to maternal obesity, reduce the placental ester PC species containing DHA and ARA as well as reduce the ether and plasmalogen PE reservoir of DHA and ARA in fetal circulation. Both males and females appear to have compensatory mechanisms that attempt to maintain an adequate supply of LCPUFA species for fetal organ growth, in particular the brain. The adaptation in females maintains normal levels of DHA and ARA in the umbilical circulation while lower plasmalogen PE LCPUFA contents in the male umbilical circulation at birth appears to be a failure of the adaptive responses. This could explain, in part, the male fetus being more vulnerable to the longterm detrimental consequences of gestation in the environment of maternal obesity. Further molecular studies will help to better understand the mechanism of ether and plasmalogen phospholipids in maternal-fetal unit and the role of fetal sex.

Supplementary Information

The online version contains supplementary material available at https://doi.org/10.1186/s13293-023-00548-1.

Additional file 1. Supplemental data.

Acknowledgements

The authors thank Dr. Robert Murphy who was the chief of the Mass Spectrometry Lipidomics Core Facility at the University of Colorado Anschutz Medical Campus, Aurora, Colorado where all the targeted lipidomic analyses were performed in plasma and placental tissue. We also thank Anita Kramer and Kathryn Erickson who helped for collection and processing blood and placenta samples. We are grateful to staff at the Clinical and Translational Research Center (CTRC) at the University of Colorado Hospital for collection of placental tissue.

Author contributions

TLP: funding acquisition, writing—review and editing. CU: formal analysis. LM: formal analysis. KZB: lipid data validation—review and editing. SSC: review and editing TJ: review and editing. VF-R: conceptualization, investigation, supervision, writing—review and editing. All authors have read and approved the final manuscript.

Funding

This work was supported by the 1R01HD104644 Grant. Contents are the authors' sole responsibility and do not necessarily represent official NIH views.

Availability of data and materials

All the data that support the findings are presented in the manuscript and in the supplementary data.

Declarations

Ethics approval and consent to participate

The study was conducted in accordance with the Declaration of Helsinki, and approved by the Institutional Review Board (or Ethics Committee) of the University of Colorado (COMIRB 14-1073). Informed consent was obtained from all subjects involved in the study.

Consent for publication

Not applicable.

Competing interests

The authors have no disclosures.

Author details

¹Department of Obstetrics and Gynecology, Division of Reproductive Sciences, University of Colorado Anschutz Medical Campus, Aurora, CO, USA. ²Department of Pediatrics, Section of Neonatology, University of Colorado Anschutz Medical Campus, Aurora, CO, USA. ³Department of Medicine, Division of Endocrinology, Metabolism, and Diabetes, University of Colorado Anschutz Medical Campus, Aurora, CO, USA. ⁴Nantes University of Colorado INRAE UMR 1280 PhAN, CRNH Ouest, 44000 Nantes, France. ⁵Nantes Université, INRAE, UMR 1280 PhAN, CHU Hôtel Dieu, HNB1, 1 place Alexis Ricordeau, 44093 Nantes, France.

Received: 15 May 2023 Accepted: 13 September 2023 Published online: 28 September 2023

References

- 1. Jansson T, Powell TL. Role of the placenta in fetal programming: underlying mechanisms and potential interventional approaches. Clin Sci (Lond). 2007;113:1–13.
- Kelly AC, Powell TL, Jansson T. Placental function in maternal obesity. Clin Sci (Lond). 2020;134:961–84.
- Bridgman SL, Azad MB, Persaud RR, Chari RS, Becker AB, Sears MR, et al. Impact of maternal pre-pregnancy overweight on infant overweight at 1 year of age: associations and sex-specific differences. Pediatr Obes. 2018;13:579–89.
- Fuemmeler BF, Zucker N, Sheng Y, Sanchez CE, Maguire R, Murphy SK, et al. Pre-pregnancy weight and symptoms of attention deficit hyperactivity disorder and executive functioning behaviors in preschool children. Int J Environ Res Public Health. 2019;16:667.
- Moss BG, Chugani DC. Increased risk of very low birth weight, rapid postnatal growth, and autism in underweight and obese mothers. Am J Health Promot. 2014;28:181–8.
- Baron-Cohen S, Lombardo MV, Auyeung B, Ashwin E, Chakrabarti B, Knickmeyer R. Why are autism spectrum conditions more prevalent in males? PLoS Biol. 2011;9: e1001081.
- 7. Alves JM, Luo S, Chow T, Herting M, Xiang AH, Page KA. Sex differences in the association between prenatal exposure to maternal obesity and hippocampal volume in children. Brain Behav. 2020;10: e01522.
- Lynch KM, Alves JM, Chow T, Clark KA, Luo S, Toga AW, et al. Selective morphological and volumetric alterations in the hippocampus of children exposed in utero to gestational diabetes mellitus. Hum Brain Mapp. 2021;42:2583–92.
- 9. Edlow AG. Maternal obesity and neurodevelopmental and psychiatric disorders in offspring. Prenat Diagn. 2017;37:95–110.
- Contu L, Hawkes CA. A review of the impact of maternal obesity on the cognitive function and mental health of the offspring. Int J Mol Sci. 2017;18:1093.
- Haggarty P, Ashton J, Joynson M, Abramovich DR, Page K. Effect of maternal polyunsaturated fatty acid concentration on transport by the human placenta. Biol Neonate. 1999;75:350–9.

- 12. Gil-Sanchez A, Larque E, Demmelmair H, Acien MI, Faber FL, Parrilla JJ, et al. Maternal-fetal in vivo transfer of [13C]docosahexaenoic and other fatty acids across the human placenta 12 h after maternal oral intake. Am J Clin Nutr. 2010;92:115–22.
- Perazzolo S, Hirschmugl B, Wadsack C, Desoye G, Lewis RM, Sengers BG. The influence of placental metabolism on fatty acid transfer to the fetus. J Lipid Res. 2017;58:443–54.
- Gazquez A, Prieto-Sanchez MT, Blanco-Carnero JE, van Harskamp D, Perazzolo S, Oosterink JE, et al. In vivo kinetic study of materno-fetal fatty acid transfer in obese and normal weight pregnant women. J Physiol. 2019;597:4959–73.
- Tomedi LE, Chang CC, Newby PK, Evans RW, Luther JF, Wisner KL, et al. Pre-pregnancy obesity and maternal nutritional biomarker status during pregnancy: a factor analysis. Public Health Nutr. 2013;16:1414–8.
- McNamara RK, Carlson SE. Role of omega-3 fatty acids in brain development and function: potential implications for the pathogenesis and prevention of psychopathology. Prostaglandins Leukot Essent Fatty Acids. 2006;75:329–49.
- 17. Mao J, Zhang X, Sieli PT, Falduto MT, Torres KE, Rosenfeld CS. Contrasting effects of different maternal diets on sexually dimorphic gene expression in the murine placenta. Proc Natl Acad Sci USA. 2010;107:5557–62.
- Mir SA, Chen L, Burugupalli S, Burla B, Ji S, Smith AAT, et al. Populationbased plasma lipidomics reveals developmental changes in metabolism and signatures of obesity risk: a mother-offspring cohort study. BMC Med. 2022;20:242.
- 19. Lands WE. Stories about acyl chains. Biochem Biophys Acta. 2000;1483:1–14.
- 20. Powell TL, Barner K, Madi L, Armstrong M, Manke J, Uhlson C, et al. Sexspecific responses in placental fatty acid oxidation, esterification and transfer capacity to maternal obesity. Biochim Biophys Acta Mol Cell Biol Lipids. 2021;1866: 158861.
- 21. Braverman NE, Moser AB. Functions of plasmalogen lipids in health and disease. Biochem Biophys Acta. 2012;1822:1442–52.
- 22. Lohner K. Is the high propensity of ethanolamine plasmalogens to form non-lamellar lipid structures manifested in the properties of biomembranes? Chem Phys Lipids. 1996;81:167–84.
- Dorninger F, Brodde A, Braverman NE, Moser AB, Just WW, Forss-Petter S, et al. Homeostasis of phospholipids - The level of phosphatidylethanolamine tightly adapts to changes in ethanolamine plasmalogens. Biochem Biophys Acta. 2015;1851:117–28.
- 24. Stables MJ, Gilroy DW. Old and new generation lipid mediators in acute inflammation and resolution. Prog Lipid Res. 2011;50:35–51.
- Nguyen LN, Ma D, Shui G, Wong P, Cazenave-Gassiot A, Zhang X, et al. Mfsd2a is a transporter for the essential omega-3 fatty acid docosahexaenoic acid. Nature. 2014;509:503–6.
- Murphy RC. Free-radical-induced oxidation of arachidonoyl plasmalogen phospholipids: antioxidant mechanism and precursor pathway for bioactive eicosanoids. Chem Res Toxicol. 2001;14:463–72.
- Ding L, Zhang LY, Shi HH, Wang CC, Jiang XM, Xue CH, et al. Eicosapentaenoic acid-enriched phosphoethanolamine plasmalogens alleviated atherosclerosis by remodeling gut microbiota to regulate bile acid metabolism in LDLR(-/-) mice. J Agric Food Chem. 2020;68:5339–48.
- Rasmiena AA, Barlow CK, Stefanovic N, Huynh K, Tan R, Sharma A, et al. Plasmalogen modulation attenuates atherosclerosis in ApoE- and ApoE/ GPx1-deficient mice. Atherosclerosis. 2015;243:598–608.
- Sutter I, Velagapudi S, Othman A, Riwanto M, Manz J, Rohrer L, et al. Plasmalogens of high-density lipoproteins (HDL) are associated with coronary artery disease and anti-apoptotic activity of HDL. Atherosclerosis. 2015;241:539–46.
- Pietilainen KH, Sysi-Aho M, Rissanen A, Seppanen-Laakso T, Yki-Jarvinen H, Kaprio J, et al. Acquired obesity is associated with changes in the serum lipidomic profile independent of genetic effects–a monozygotic twin study. PLoS ONE. 2007;2: e218.
- Graessler J, Schwudke D, Schwarz PE, Herzog R, Shevchenko A, Bornstein SR. Top-down lipidomics reveals ether lipid deficiency in blood plasma of hypertensive patients. PLoS ONE. 2009;4: e6261.
- 32. Dean JM, Lodhi IJ. Structural and functional roles of ether lipids. Protein Cell. 2018;9:196–206.
- 33. Lankinen M, Schwab U, Kolehmainen M, Paananen J, Nygren H, Seppanen-Laakso T, et al. A healthy nordic diet alters the plasma lipidomic

profile in adults with features of metabolic syndrome in a multicenter randomized dietary intervention. J Nutr. 2015;146:662–72.

- Felder TK, Ring-Dimitriou S, Auer S, Soyal SM, Kedenko L, Rinnerthaler M, et al. Specific circulating phospholipids, acylcarnitines, amino acids and biogenic amines are aerobic exercise markers. J Sci Med Sport. 2017;20:700–5.
- Bidne KL, Uhlson C, Palmer C, Zemski-Berry K, Powell TL. Human placental lipid content and lipid metabolic enzyme abundance in obesity and across gestation. Clin Sci (Lond). 2022;136:1389–404.
- Ferchaud-Roucher V, Kramer A, Silva E, Pantham P, Weintraub ST, Jansson T, et al. A potential role for lysophosphatidylcholine in the delivery of long chain polyunsaturated fatty acids to the fetal circulation. Biochim Biophys Acta Mol Cell Biol Lipids. 2019;1864:394–402.
- Castillo-Castrejon M, Yang IV, Davidson EJ, Borengasser SJ, Jambal P, Westcott J, et al. Preconceptional lipid-based nutrient supplementation in 2 low-resource countries results in distinctly different IGF-1/mTOR placental responses. J Nutr. 2021;151:556–69.
- Bligh EG, Dyer WJ. A rapid method of total lipid extraction and purification. Can J Biochem Physiol. 1959;37:911–7.
- Murphy EJ, Anderson DK, Horrocks LA. Phospholipid and phospholipid fatty acid composition of mixed murine spinal cord neuronal cultures. J Neurosci Res. 1993;34:472–7.
- 40. Storey JD. A direct approach to false discovery rates. J R Stat Soc Ser B Stat Methodol. 2002;64:479–98.
- Yu HT, Xu WH, Chen YR, Ji Y, Tang YW, Li YT, et al. Association of prepregnancy obesity and remodeled maternal-fetal plasma fatty acid profiles. Front Nutr. 2022;9: 897059.
- Kim D, Choi JE, Park Y. Low-linoleic acid diet and oestrogen enhance the conversion of alpha-linolenic acid into DHA through modification of conversion enzymes and transcription factors. Br J Nutr. 2019;121:137–45.
- Nagan N, Zoeller RA. Plasmalogens: biosynthesis and functions. Prog Lipid Res. 2001;40:199–229.
- Vance JE. Lipoproteins secreted by cultured rat hepatocytes contain the antioxidant 1-alk-1-enyl-2-acylglycerophosphoethanolamine. Biochem Biophys Acta. 1990;1045:128–34.
- Larque E, Krauss-Etschmann S, Campoy C, Hartl D, Linde J, Klingler M, et al. Docosahexaenoic acid supply in pregnancy affects placental expression of fatty acid transport proteins. Am J Clin Nutr. 2006;84:853–61.
- Fattuoni C, Mando C, Palmas F, Anelli GM, Novielli C, Parejo Laudicina E, et al. Preliminary metabolomics analysis of placenta in maternal obesity. Placenta. 2018;61:89–95.
- Wallace JG, Bellissimo CJ, Yeo E, Fei Xia Y, Petrik JJ, Surette MG, et al. Obesity during pregnancy results in maternal intestinal inflammation, placental hypoxia, and alters fetal glucose metabolism at mid-gestation. Sci Rep. 2019;9:17621.
- Chassen SS, Ferchaud-Roucher V, Palmer C, Li C, Jansson T, Nathanielsz PW, et al. Placental fatty acid transport across late gestation in a baboon model of intrauterine growth restriction. J Physiol. 2020;598:2469–89.
- Bildirici I, Schaiff WT, Chen B, Morizane M, Oh SY, O'Brien M, et al. PLIN2 is essential for trophoblastic lipid droplet accumulation and cell survival during hypoxia. Endocrinology. 2018;159:3937–49.
- Cipolla CM, Lodhi IJ. Peroxisomal dysfunction in age-related diseases. Trends Endocrinol Metab. 2017;28:297–308.
- Muralimanoharan S, Guo C, Myatt L, Maloyan A. Sexual dimorphism in miR-210 expression and mitochondrial dysfunction in the placenta with maternal obesity. Int J Obes. 2015;39:1274–81.
- 52. Honsho M, Asaoku S, Fukumoto K, Fujiki Y. Topogenesis and homeostasis of fatty acyl-CoA reductase 1. J Biol Chem. 2013;288:34588–98.
- Dorninger F, Forss-Petter S, Wimmer I, Berger J. Plasmalogens, plateletactivating factor and beyond—ether lipids in signaling and neurodegeneration. Neurobiol Dis. 2020;145: 105061.
- Honsho M, Yagita Y, Kinoshita N, Fujiki Y. Isolation and characterization of mutant animal cell line defective in alkyl-dihydroxyacetonephosphate synthase: localization and transport of plasmalogens to post-Golgi compartments. Biochem Biophys Acta. 2008;1783:1857–65.
- Engelmann B. Plasmalogens: targets for oxidants and major lipophilic antioxidants. Biochem Soc Trans. 2004;32:147–50.
- Korsmo HW, Kadam I, Reaz A, Bretter R, Saxena A, Johnson CH, et al. Prenatal choline supplement in a maternal obesity model modulates offspring hepatic lipidomes. Nutrients. 2023;15:965.

- 57. Nam J, Greenwald E, Jack-Roberts C, Ajeeb TT, Malysheva OV, Caudill MA, et al. Choline prevents fetal overgrowth and normalizes placental fatty acid and glucose metabolism in a mouse model of maternal obesity. J Nutr Biochem. 2017;49:80–8.
- Oresic M, Simell S, Sysi-Aho M, Nanto-Salonen K, Seppanen-Laakso T, Parikka V, et al. Dysregulation of lipid and amino acid metabolism precedes islet autoimmunity in children who later progress to type 1 diabetes. J Exp Med. 2008;205:2975–84.
- Tamimi RM, Lagiou P, Mucci LA, Hsieh CC, Adami HO, Trichopoulos D. Average energy intake among pregnant women carrying a boy compared with a girl. BMJ. 2003;326:1245–6.
- Rao JS, Ertley RN, DeMar JC Jr, Rapoport SI, Bazinet RP, Lee HJ. Dietary n-3 PUFA deprivation alters expression of enzymes of the arachidonic and docosahexaenoic acid cascades in rat frontal cortex. Mol Psychiatry. 2007;12:151–7.
- Meakin AS, Cuffe JSM, Darby JRT, Morrison JL, Clifton VL. Let's talk about placental sex, baby: understanding mechanisms that drive female- and male-specific fetal growth and developmental outcomes. Int J Mol Sci. 2021;22.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

