



HAL
open science

Some plasma biomarkers of residual feed intake in beef cattle remain consistent regardless of intake level

Gonzalo Cantalapiedra-Hijar, K. Nedelkov, P. Crosson, M. Mcgee

► To cite this version:

Gonzalo Cantalapiedra-Hijar, K. Nedelkov, P. Crosson, M. Mcgee. Some plasma biomarkers of residual feed intake in beef cattle remain consistent regardless of intake level. *Scientific Reports*, 2024, 14 (1), pp.8540. 10.1038/s41598-024-59253-7 . hal-04544483

HAL Id: hal-04544483

<https://hal.inrae.fr/hal-04544483>

Submitted on 16 Apr 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



OPEN

Some plasma biomarkers of residual feed intake in beef cattle remain consistent regardless of intake level

G. Cantalapiedra-Hijar^{1✉}, K. Nedelkov², P. Crosson³ & M. McGee³

This study investigated whether plasma biomarkers of residual feed intake (RFI), identified under ad libitum feeding conditions in beef cattle, remained consistent during feed restriction. Sixty Charolais crossbred young bulls were divided into two groups for a crossover study. Group A was initially fed ad libitum (first test) and then restricted (second test) on the same diet, while Group B experienced the opposite sequence. Blood samples were collected from the 12 most divergent RFI animals in each group at the end of the first test and again after the second test. 12 plasma variables consistently increased, while three consistently decreased during feed restriction (FDR < 0.05). Only two metabolites, α -amino adipic acid for Group A and 5-aminovaleric acid for Group B, were associated with RFI independent of feed intake level (FDR < 0.05), demonstrating moderate-to-high repeatability across feeding levels (intra-class correlation coefficient ≥ 0.59). Notably, both metabolites belong to the same metabolic pathway: lysine degradation. These metabolites consistently correlated with RFI, irrespective of fluctuations in feed intake, indicating a connection to individual metabolic processes influencing feed efficiency. These findings suggest that a portion of RFI phenotypic variance is inherent to an individual's metabolic efficiency beyond variations in feed intake.

Improving animal feed efficiency (FE) can significantly contribute to enhancing the sustainability of the beef cattle sector¹. In addition to established nutritional strategies aimed at improving the average FE of co-reared animal groups^{2,3}, there is, within the same breed, significant variability in FE among individual animals reared under identical conditions⁴. This variability in FE among animals, coupled with its moderate heritability, provides an opportunity for the genetic selection of superior animals for this trait⁵, a practice currently employed in cattle breeding programs⁶. In addition to genetic selection, understanding and forecasting the interindividual variability in FE could form the basis for innovative individual animal-centered feeding strategies or precision nutrition approaches^{4,7}. Precision feeding can further improve resource use efficiency, reduce animal excretions to the environment, and enhance the financial viability of livestock farming⁸. Indeed, as highlighted by Tedeschi et al.⁷, the identification of animals with superior FE is among the most influential factors towards achieving long-term livestock sustainability.

Among the metrics used for assessing beef cattle FE, residual feed intake (RFI) stands out as one of the most widely employed indices within animal breeding⁹. RFI enables the identification of individuals with lower feed intake while maintaining similar performance levels (i.e., superior FE)¹⁰. Nevertheless, due to the substantial cost and time associated with determining RFI in beef cattle, several blood biomarkers that correlate with the biological mechanisms underlying RFI have been proposed as alternative cost-effective tools for identifying efficient (i.e., low-RFI) cattle^{11,12}. In this regard, literature reviews^{4,13} highlighted that the primary biological mechanisms driving RFI appear to be intrinsically connected to animal metabolism. However, whether these metabolic regulations play a causal role or are merely associated with RFI due to covariance with feed intake remains uncertain. For instance, multiple studies have suggested that the plasma concentration of branched-chain amino acids, having signalling roles in key metabolic pathways such as protein and lipid synthesis, could serve as potential biomarkers of RFI in beef cattle^{11,14–17}. Likewise, a meta-analysis of genome-wide association studies conducted on beef cattle concluded that the only metabolic pathway genetically linked to RFI was the

¹Université Clermont Auvergne, INRAE, VetAgro Sup, UMR Herbivores, 63122 St-Genès-Champanelle, France. ²Faculty of Veterinary Medicine, Trakia University, Stara Zagora 6000, Bulgaria. ³Teagasc, Animal & Grassland Research and Innovation Centre, Grange, Dunsany, Co. Meath, Ireland. ✉email: gonzalo.cantalapiedra@inrae.fr

degradation pathway of branched-chain amino acids¹⁸. However, it is important to note that the plasma concentration of branched-chain amino acids in ruminants, much like other nutrients, is significantly influenced by feed protein intake and nutrient flow to the duodenum¹⁹. This confounding effect makes it challenging to determine whether these biomarkers capture processes that render individuals more metabolically efficient, or conversely, mirror reduced amino acid absorption resulting from the lower feed protein intake associated with efficient RFI phenotypes. Gaining insight into the manner in which blood metabolites are linked to phenotypes becomes pivotal before proposing them as robust biomarkers for assisting genetic selection and precision feeding. In other words, for the successful application of RFI biomarkers, it is crucial to comprehend the extent to which these blood metabolites are correlated with animal feed intake (feeding level) and host-related physiological differences independent of nutrient intake²⁰, such that metabolic feed efficiency is enhanced across contrasting conditions encompassing both feed abundance and scarcity.

We propose the hypothesis that numerous previously identified candidate biomarkers of RFI are associated with this phenotype primarily due to their blood or plasma concentrations covarying with nutrient intake. In contrast, certain other metabolites, such as branched-chain amino acids, may experience changes in their concentration when nutrient intake is altered, yet still retain a component linked to the animal's host physiology that these biomarkers can capture regardless of fluctuations in feeding level. Consequently, the aim of this study was to ascertain whether plasma biomarkers of RFI, identified under controlled ad libitum feeding conditions, remain consistent when the same animals are offered the same diet but at a restricted and identical feeding level.

Results

Mean metabolic body weight (BW) emerged as the sole consistent and significant predictor across all four RFI models (Supplementary Table S2), while average daily gain (ADG) was only significant in the models involving animals with ad libitum feeding. Nearly all the observed variability in dry matter (DM) intake within feed restriction conditions was explained through predictors in the RFI model ($r^2 = 0.98$ and 0.99), leading to minor fluctuations in RFI values, as indicated by their coefficients of variation (CV) of 0.59 and 0.64, respectively. Conversely, the RFI models accounted for only 81% and 51% of the explained variance within ad libitum conditions for the first and second test periods, respectively, resulting in RFI value CVs of 3.85% and 5.08%, respectively. As explained further, the RFI phenotype measured in ad libitum conditions was maintained as identical during feed restriction to assess whether the identified RFI biomarkers were still linked to the RFI phenotype when animals were fed at the same feeding level.

Animal performance per test period (first vs. second RFI test) and feeding level (ad-libitum vs. restriction) are presented in Table 1. There were statistically significant feeding level \times test period interactions for DM intake (kg/day and kg/100 kg BW), start and end BW, mean metabolic BW ($P < 0.001$), ADG ($P < 0.01$) and feed conversion efficiency (FCE) ($P < 0.05$). The expected increase in DM intake in ad libitum compared to feed restriction exhibited a more pronounced effect during the first (0–70 days) compared to the second (78–148 days) RFI test (42% vs. 24%, respectively). By design, the start BW did not differ between the feeding level groups during Test 1. However, due to greater growth performance, the end BW in Test 1 with ad libitum feeding (Group A) was

	Test 1 (0–70 days)		Test 2 (78–148 days)		SEM	P value		
	Ad-libitum (Group A)	Restriction (Group B)	Ad-libitum (Group B)	Restriction (Group A)		Feeding level	Feeding period	FL \times P
Number of animals	29	30	30	28				
Age, days	394	390	467	470	8.35	0.77	<0.001	0.69
DM intake, kg/days	9.72 ^b	6.84 ^d	10.4 ^a	8.44 ^c	0.114	<0.001	<0.001	<0.001
DM intake, kg/100 kg BW	1.91 ^a	1.46 ^c	1.78 ^b	1.41 ^c	0.015	<0.001	<0.001	<0.001
Body weight, kg								
Start	451 ^c	450 ^c	527 ^b	563 ^a	4.38	<0.001	<0.001	<0.001
End	562 ^b	502 ^c	632 ^a	627 ^a	5.36	<0.001	<0.001	<0.001
Mean metabolic BW, kg	107 ^b	102 ^c	118 ^a	120 ^a	0.76	0.09	<0.001	<0.001
Average daily gain ¹ , kg	1.59 ^a	0.74 ^c	1.49 ^a	0.92 ^b	0.0416	<0.001	0.32	0.001
Fat depth change ² , mm								
Rump	2.14	0.86	2.77	0.75	0.256	<0.001	0.29	0.15
Lumbar	0.631	0.391	0.620	0.156	0.0925	<0.001	0.19	0.23
Rib	1.51	0.410	1.12	0.212	0.161	<0.001	0.07	0.54
Muscle depth change ² , mm	11.1	6.67	6.06	0.59	0.878	<0.001	<0.001	0.54
Feed conversion efficiency, kg BW gain/kg DM intake	0.163 ^a	0.107 ^c	0.144 ^b	0.109 ^c	0.0039	<0.001	0.017	0.010

Table 1. Performance of growing-fattening bulls fed ad libitum and then restricted (Group A) or vice versa (Group B) during two consecutive 70-d feed efficiency tests. ^{a-d}Mean values with different superscripts are significantly different ($P < 0.05$). ¹Average daily live weight gain calculated as the difference between final and initial body weight divided by the duration (70 days). ²Calculated as the difference between depth values measured at the end minus those obtained at the beginning.

significantly greater ($P < 0.05$) compared to restricted feeding (Group B). Consequently, in Test 2, the initial BW of Group A animals (subjected to feed restriction) was also greater than those from Group B (subjected to ad libitum feeding). End BW and mean metabolic BW were greater for the ad libitum group than for the restricted group in Test 1 but did not differ between the feeding level groups in Test 2. Daily live weight gain for the ad libitum group did not differ between the two tests, whereas for the restricted group, it was lower in Test 1 than in Test 2. Feed conversion efficiency for the ad-libitum groups was greater during Test 1 compared to Test 2, whereas there was no difference between tests for the restricted groups. No significant interactions ($P > 0.05$) between feeding level and test period were observed for changes in ultrasound fat and muscle depth measurements; however, these parameters were all influenced by feeding level ($P < 0.001$), with greater changes observed with ad libitum feeding compared to feed restriction. The test period impacted the change in muscle depth ($P < 0.001$) and indicated a trend in rib fat depth change ($P = 0.07$), with increases in depth more substantial during Test 1 compared to Test 2.

Overall animal performances from day 0–148 per group (A vs. B) are presented in Table 2. Bulls assigned to Group A (starting the first RFI test with ad libitum feeding and then transitioning to restricted feeding) had a greater overall DM intake ($P = 0.001$) and similar overall ADG ($P = 0.36$), resulting in poorer FCE ($P = 0.002$) and RFI ($P < 0.001$) values compared to those assigned to Group B. Additionally, animals from Group A tended to have a lower overall change in rump fat depth ($P = 0.08$) and a higher kill-out proportion ($P = 0.09$). Slaughter weight, carcass weight and carcass conformation and fat scores did not differ ($P \geq 0.32$) between animals in Groups A and B.

Table 3 illustrates performance per test period for the 24 most divergent bulls for RFI (6 low-RFI and 6 high-RFI per feeding level), determined during the initial (0–70 days) RFI test. In Test 1, the low-RFI bulls in both Group A (ad-libitum) and Group B (restricted) did not differ ($P > 0.05$) in terms of DM intake (kg/day), mean metabolic BW, ADG, and FCE when compared to the high-RFI animals. In contrast, DM intake relative to BW was lower ($P < 0.05$) for the low- compared to the high-RFI animals in Group A, whereas there was no difference between the RFI groups in Group B. As expected, RFI values differed between the low-RFI and high-RFI groups ($P < 0.001$), with a mean difference of 0.99 kg DM/day for Group A (ad-libitum) and only 0.115 kg DM/day for Group B (restricted). In Test 2, the low- and high-RFI bulls in Group A (now restricted) did not differ in DM intake (kg/d or kg/100 kg BW), mean metabolic BW, ADG, FCE and RFI, whereas in Group B (now ad-libitum), the low-RFI bulls had a higher DM intake in kg/day or kg/100 kg BW and ADG ($P < 0.05$) and tended to have a greater FCE ($P = 0.06$) than the high-RFI animals. It is noteworthy to highlight that, although RFI values were less extreme in restricted feeding conditions compared to ad-libitum feeding for Group A, on average, they still exhibited negative values for low-RFI animals and positive ones for high-RFI values ($P = 0.13$). In contrast, feed-efficient animals (low-RFI) starting with feed restriction (Group B) showed, on average, positive values during ad-libitum feeding, and the reverse was observed for inefficient animals ($P = 0.24$).

Among the 74 plasma variables analysed in the same 24 RFI-divergent bulls across both test periods, 21 variables exhibited an increase ($P < 0.05$) in their concentration or values during feed restriction, while 12 variables

	Group A (Ad-libitum → Restriction)	Group B (Restriction → Ad-libitum)	SEM	P value
Number of animals	28	30		
Body weight at slaughter ¹ , kg	651	651	6.52	0.99
Carcass weight, kg	375	369	4.51	0.32
Kill-out proportion ² , %	57.7	56.7	0.42	0.09
Carcass conformation score (1–15) ³	10.0	10.0	0.24	1.00
Carcass fat score (1–15) ⁴	6.11	6.23	0.23	0.70
Performance from 0 to 148 days				
Total DM intake, kg	9.06	8.59	0.100	0.001
Total average daily gain, kg	1.20	1.24	0.029	0.36
Feed conversion efficiency, kg BW gain/kg DM intake	0.132	0.144	0.0026	0.002
Residual feed intake ⁵ , kg/days	0.160	-0.150	0.057	<0.001
Fat depth change ⁶ , mm				
Rump	2.92	3.63	0.286	0.08
Lumbar	0.779	1.01	0.105	0.12
Rib	1.73	1.52	0.194	0.46
Muscle depth change, mm	11.7	12.7	0.935	0.45

Table 2. Overall performance (0–148 days) of growing-fattening bulls fed ad libitum (0–70 days) and then restricted (78–148 days) (Group A) or vice versa (Group B). ¹Animals were slaughtered approximately 2 weeks after the end of the second feed efficiency test. ²Kill-out proportion: (Cold carcass weight/Body weight at slaughter) × 100. ³EU beef carcass classification scheme scale, where 1 represents the poorest and 15 the best conformation. ⁴EU beef carcass classification scheme scale, where 1 represents the leanest and 15 the fattest carcass. ⁵Residual feed intake calculated through the entire experimental period (0–148 days) with a model having significant DM intake predictors the block, pen and metabolic body weight ($r^2 = 0.64$). ⁶Calculated as the difference between depth values measured at the end minus those obtained at the beginning.

	First RFI test (0–70 days)							
	Group A—Ad-libitum				Group B—Restricted			
	Low-RFI	High-RFI	SEM	P value	Low-RFI	High-RFI	SEM	P value
Number of animals	6	6			6	6		
DM intake, kg/days	9.27	10.1	0.401	0.19	6.73	6.87	0.135	0.49
DM intake, kg/100 kg BW	1.80	1.98	0.055	0.04	1.46	1.46	0.004	0.42
Metabolic body weight, kg	105	107	1.5	0.37	101	101	1.5	0.90
Average daily gain, kg	1.61	1.54	0.136	0.72	0.64	0.69	0.073	0.63
FCE ¹ , kg BW gain/kg DM intake	0.173	0.152	0.010	0.17	0.095	0.100	0.010	0.72
Residual feed intake, kg DM/days	-0.56	0.44	0.077	<0.001	-0.059	0.057	0.013	<0.001
	Second RFI test (78–148 days)*							
	Group A—Restricted				Group B—Ad-libitum			
	Low-RFI	High-RFI	SEM	P value	Low-RFI	High-RFI	SEM	P value
Number of animals	6	6			6	6		
DM intake, kg/days	8.47	8.37	0.191	0.74	10.8	9.78	0.267	0.02
DM intake, kg/100 kg BW	1.41	1.42	0.005	0.16	1.86	1.72	0.041	0.04
Metabolic body weight, kg	119	121	2.0	0.50	118	116	2.0	0.41
Average daily gain, kg	0.93	0.91	0.076	0.91	1.66	1.31	0.086	0.02
FCE ¹ , kg BW gain/kg DM intake	0.110	0.109	0.0085	0.96	0.153	0.133	0.0064	0.06
Residual feed intake, kg DM/days	-0.027	0.015	0.018	0.13	0.140	-0.201	0.194	0.24

Table 3. Performances of the top 24 growing-fattening bulls with divergent residual feed intake (RFI) values (12 in group A and 12 in group B) identified during the initial 0–70 days feed efficiency test and selected for plasma metabolomics analysis conducted in the first and second feed efficiency tests. *Animals in the second RFI test kept the same RFI ranking (lowest vs highest) as in the first test. However, the RFI values shown in the table are the actual RFI values for those animals in the second RFI test. ¹FCE, feed conversion efficiency.

decreased ($P < 0.05$; Table 4). Following FDR correction ($FDR < 0.05$), among the plasma variables that exhibited a significant increase in concentration or values, there were seven amino acids or related metabolites (1-methylhistidine, 3-methylhistidine, 5-aminovaleric acid, β -aminobutyric acid, betaine, sarcosine, and glycine), four fatty acids (docosahexaenoic, eicosapentaenoic, octadecenoic, and octadecadienoic acids), as well as non-esterified fatty acids and the natural abundance of ¹⁵N. Conversely, following FDR corrections ($FDR < 0.05$), only β -alanine, hippuric acid, and β -hydroxybutyrate demonstrated a significant decrease during feed restriction. All the metabolites that demonstrated higher or lower concentrations/values during feed restriction exhibited consistent behaviour across test periods (i.e., no feeding level \times period interaction; $P > 0.05$). Of these metabolites, the feeding period exclusively affected sarcosine, docosahexaenoic acid, eicosapentaenoic acid, and the natural abundance of ¹⁵N (all displaying higher concentrations or values in the first test period) following FDR corrections ($FDR < 0.05$). The repeatability of plasma metabolites or variables analysed in the same 24 RFI-divergent bulls across two feeding levels is detailed in Table 5. Six plasma variables exhibited ‘high’ repeatability ($r > 0.70$), while 24 exhibited ‘moderate’ repeatability ($0.40 \geq r \geq 0.70$) across feeding periods. Notably, 3-methylhistidine and alkaline phosphatase exhibited repeatability values ≥ 0.90 across feeding levels, indicating that individuals are consistently ranked for these two variables regardless of the feeding level and test period.

The effect of RFI and feeding level on plasma parameters analysed from the 12 bulls identified as extreme RFI animals in group A and B are presented in Supplementary Tables 3 and 4. In group A animals, the plasma concentrations of three amino acid-related metabolites (alpha-amino adipic acid, asymmetric dimethylarginine and carnosine) along with three amino acids (methionine, alanine and isoleucine) and one hormone (IGF-1) were lower ($P < 0.05$) in efficient compared to inefficient RFI counterparts (Table 6). This pattern remained consistent whether they were fed ad libitum (Test 1) or under restriction (Test 2), as evidenced by the absence of an RFI \times feeding level interaction ($P \geq 0.39$). These metabolites were not impacted by feeding level ($P \geq 0.07$). Only the plasma concentration of alpha-amino adipic acid reached significant differences after FDR corrections ($FDR < 0.05$). In group B animals, only the concentrations of three plasma metabolites (5-amino valeric acid, serine and betaine) were significantly and positively correlated with ad-libitum RFI values in both test periods (Table 7). These three metabolites were also impacted by feeding level ($P < 0.001$), but they were correlated with ad-libitum RFI values in the same way across both feeding levels (RFI \times feeding level; $P \geq 0.11$). Among them, only the plasma concentration of 5-aminovaleric acid was significantly correlated after FDR corrections ($FDR < 0.05$).

Discussion

To distinguish the impact of RFI from feeding level on systemic plasma parameters, we conducted a comparative analysis involving feeding the same animals under ad-libitum and restricted feeding conditions. Instead of using feed restriction as a percentage of a previously recorded ad-libitum intake²¹, we applied it as a percentage of the animals’ body weight (i.e., a proxy of maintenance requirements). This approach ensured a uniform nutrient intake per unit of body weight across all evaluated animals during the restriction phase, effectively eliminating differences in feed intake between RFI groups during this phase. In this regard, during ad libitum feeding, we

	Test 1 (0–70 days)		Test 2 (78–148 days)		Fold change	SEM	P value		
	Ad-libitum (Group A)	Restricted (Group B)	Ad-libitum (Group B)	Restricted (Group A)			Feeding level	Feeding period	FL × P
Number of animals	12	12	12	12					
Increased during restriction									
Amino acid related									
1-Methyl-His	6.53	7.83	6.51	8.38	1.24	0.460	<0.0001*	0.256	0.642
3-Methyl-His	5.40	6.00	5.32	6.17	1.14	0.461	0.001*	0.739	0.843
5-Aminovaleric acid	1.19	1.78	1.02	1.41	1.44	0.153	0.0001*	0.012	0.595
β-aminobutyric acid	0.114	0.134	0.092	0.115	1.21	0.0091	0.0001*	0.002	0.898
Betaine	117	138	96.4	118	1.20	7.72	<0.0001*	<0.0001	0.972
Creatinine	139	152	131	143	1.09	7.71	0.009	0.062	0.974
Sarcosine	1.99	2.68	1.54	1.90	1.30	0.191	0.003*	0.0007*	0.467
Amino acids									
Glycine	409	528	356	482	1.32	23.7	<0.0001*	0.032	0.894
Histidine	53.1	66.2	53.4	61.3	1.20	4.12	0.006	0.519	0.587
Proline	66.0	75.4	66.9	68.3	1.08	2.85	0.045	0.230	0.218
Serine	91.9	138	71.2	98.4	1.45	7.73	<0.0001*	0.0002*	0.277
Bile acids									
Cholic acid	7.30	9.01	5.90	12.7	1.64	1.66	0.013	0.467	0.159
Chenodeoxycholic acid	0.386	0.449	0.259	0.558	1.56	0.086	0.043	0.915	0.194
Deoxycholic acid	0.868	0.983	0.634	2.089	2.05	0.358	0.038	0.233	0.078
Fatty acids									
Nonesterified fatty acids	88.9	143	105	266	2.14	26.0	0.0004*	0.016	0.059
Arachidonic acid	1.123	1.386	0.686	1.056	1.35	0.167	0.027	0.009	0.786
Docosahexaenoic acid	0.847	1.173	0.530	0.802	1.43	0.0871	0.0002*	0.0009*	0.811
Eicosapentaenoic acid	0.251	0.316	0.144	0.262	1.46	0.0241	0.0003*	0.001*	0.336
Octadecenoic acid	46.5	61.1	40.5	92.3	1.76	8.58	0.0005*	0.108	0.047
Octadecadienoic acid	4.84	6.59	5.13	9.84	1.65	0.886	0.0011*	0.050	0.113
Natural isotope abundance, ‰									
δ ¹⁵ N	5.64	5.88	5.29	5.61	1.02	0.095	0.0006*	0.0002*	0.768
Decreased during restriction									
Amino acid related									
Carnosine	20.7	18.0	19.7	19.4	0.93	1.29	0.018	0.678	0.495
Homoarginine	3.73	3.54	3.44	2.77	0.88	0.298	0.016	0.004	0.543
Amino acids									
Glutamine	406	382	423	389	0.93	14.3	0.032	0.353	0.741
Glutamate	88.3	78.7	79.0	70.8	0.89	6.23	0.032	0.039	0.929
Isoleucine	103	93.3	108	98.4	0.91	3.89	0.007	0.138	0.972
Tryptophane	43.4	37.3	43.2	42.0	0.92	2.52	0.036	0.177	0.439
Tyrosine	52.0	39.4	53.4	48.8	0.84	2.98	0.001	0.025	0.280
Biogenic amines									
β-alanine	3.95	2.80	3.49	3.00	0.78	0.291	0.008*	0.660	0.283
Carboxylic acids									
Hippuric acid	75.4	46.5	84.8	59.3	0.66	5.80	0.0001*	0.069	0.774
Indoles									
Indolepropionic acid	4.26	3.15	2.74	2.03	0.74	0.571	0.027	0.0024*	0.780
Ketone bodies									
β-hydroxybutyrate	389	273	445	308	1.44	22.8	<0.0001*	0.060	0.656
Hormones, ng/mL									
IGF-1	528	402	502	458	1.20	46.5	0.02	0.65	0.49

Table 4. Plasma metabolites (μM) and variables significantly ($P < 0.05$) affected by feeding level (FL) in most extreme RFI bulls fed ad libitum (0–70 days) and then restricted (78–148 days) (Group A) or vice versa (Group B). *P values with a star are also FDR < 0.05.

	Repeatability [#]
High repeatability ($0.7 \leq r \leq 0.9$) ¹	
3-Methyl-histidine	0.91
Alkaline phosphatase (ALP)	0.90
Betaine	0.78
Carnosine	0.78
1-Methyl-histidine	0.76
Indoleacetic acid*	0.76
Moderate repeatability ($0.4 \leq r < 0.7$) ¹	
Homoarginine	0.69
$\delta^{13}\text{C}$	0.68
Creatinine	0.66
Alpha-aminoadipic acid	0.65
Aspartate transaminase (AST)	0.64
Lactic acid	0.63
Glutamate	0.61
Glucose	0.60
5-Aminovaleric acid	0.59
Tryptophane	0.58
β -Aminobutyric acid*	0.57
Alanine transaminase (ALT)	0.56
Symmetric dimethylarginine	0.55
Indolepropionic acid*	0.55
Anserine	0.49
IGF-1	0.48
$\delta^{15}\text{N}$	0.48
Arginine	0.47
Trans-4-hydroxyproline	0.47
Kynurenine	0.47
Valine	0.44
Tyrosine	0.43
Alanine	0.42
Asymmetric dimethylarginine	0.41

Table 5. Repeatable plasma metabolites and parameters in growing-fattening bulls fed ad libitum (0–70 days) and then restricted (77–147 days) (Group A) or vice versa (Group B). [#]Repeatability defined as the tendency of individuals to maintain their ranking over time based on the plasma concentration of metabolites and calculated as the intraclass correlation coefficient (see “Methods”). ¹According to Martin and Bateson, 1986. *Microbial origin.

	Group A—Ad-libitum (0–70 days)		Group A—Restricted ¹ (78–148 days)		Fold change	SEM	P value		
	Low-RFI	High-RFI	Low-RFI	High-RFI			RFI	FL	RFI × FL
Number of animals	6	6	6	6					
Amino acid related, μM									
Alpha-Aminoadipic acid	1.13	1.83	0.983	1.48	1.57	0.142	0.003*	0.071	0.437
Asymmetric Dimethylarginine	0.933	1.10	0.933	1.03	1.14	0.050	0.023	0.520	0.520
Carnosine	18.0	23.7	16.8	22.2	1.32	1.85	0.042	0.249	0.881
Amino acids, μM									
Methionine	20.2	24.7	20.8	27.7	1.28	2.08	0.022	0.400	0.588
Alanine	197	230	176	218	1.20	14.5	0.049	0.198	0.718
Isoleucine	97.0	109	92.8	104	1.12	5.16	0.045	0.413	0.974
Hormones, ng/mL									
IGF-1	417	638	381	536	1.47	66.9	0.049	0.100	0.397

Table 6. Plasma metabolites and hormones affected by residual feed intake (RFI) in 12 growing-fattening bulls identified as extreme RFI animals during ad libitum feeding (0–70 days) and subsequently analysed during feed restriction (78–148 days) (Group A). *P values with a star are also FDR < 0.05. ¹Animals in the second test were the same as in the first test and kept the same RFI ranking (lowest vs highest) observed ad-libitum.

	Pearson correlation coefficient		P value ¹		
	Group B—Restricted (0–70 days) ¹	Group B—Ad-libitum (78–148 days)	RFI	FL	RFI × FL
5-Aminovaleric acid	0.72	0.74	0.005*	<0.001*	0.110
Serine	0.52	0.80	0.012	<0.001*	0.725
Betaine	0.58	0.57	0.042	<0.001*	0.272
Sarcosine	–0.03	0.73	0.166	<0.001*	0.012

Table 7. Plasma metabolites correlated with residual feed intake (RFI) in 12 growing-fattening bulls identified as extreme RFI animals during feed restriction (0–70 days) and subsequently analysed during ad libitum feeding (78–148 days) (Group B). *P values with a star are also FDR < 0.05. ¹Animals in the first RFI test were the same as those in the second RFI test and maintained the same RFI values (ranging from –0.98 to 0.61 kg/days) observed ad libitum.

observed approximately 8–10% variation in feed intake (% BW) among selected extreme RFI animals. In contrast, during feed restriction, we successfully minimized this variation to < 1%, which was statistically nonsignificant (Table 3) and much lower than the 12% difference observed during restriction in a previous study conducted in cattle²¹. Likewise, no differences in DM intake relative to body weight were reported when two chicken lines divergent in RFI were evaluated during feed restriction²⁰. Similarly, studies in pigs²² and mink²³ involving high and low RFI genetic lines or phenotypes, respectively, subjected to feed restriction were assessed at comparable feed intake levels. As argued by Metzler-Zebeli et al.²⁰, maintaining a similar intake level between both RFI lines is essential for understanding the contributions of feeding behaviour (intake) versus host metabolic mechanisms in plasma metabolite concentrations. Consequently, to distinguish between the contributions of intake and host metabolic factors to variation in systemic plasma metabolite concentrations, we adopted a similar rationale in the present study.

Conversely, our methodology for implementing feed restriction was responsible for the minimal variation observed in RFI values across individuals (coefficient of variation (CV) close to 0.6% during restriction vs. 4–5% during ad libitum feeding; Supplementary Table S2). Given that we ensured feed distribution was adjusted in accordance with BW during feed restriction, it is by design that nearly all the observed variation in DM intake was attributed to the metabolic BW predictor in the RFI model. Although we initially selected and sampled the most-extreme RFI animals during the first test period, extreme RFI bulls from Group B starting with feed restriction exhibited only a 1.7% CV in RFI, compared to an 11.3% CV in Group A during ad-libitum conditions (Supplementary Table S2). Consequently, and considering that RFI values were not correlated across feeding levels (Supplementary Table S2), we deemed the animals' RFI phenotype to be that observed during ad-libitum conditions (first test for Group A and second test for Group B), maintaining the same ranking when animals underwent the restriction phase. As a result, and considering our experimental design (Fig. 1), relationships between RFI phenotype and plasma parameters were analysed by repeated measurement ANOVA in Group A (comprising two contrasted RFI groups selected during the first period [*ad-libitum*]) and by mixed model linear regression analysis in Group B (comprising animals with continuous RFI values observed during the second test [*ad-libitum*]).

Plasma parameters associated with residual feed intake irrespective of the feeding level

Gaining a deeper understanding of the associations between blood plasma parameters and phenotypic traits is crucial before considering them as reliable biomarkers for aiding in genetic selection and precision feeding decisions. The findings of our study allow us, for the first time, to identify metabolites in beef cattle that show an association with the RFI phenotype even when animals are exposed to a uniform feeding level (i.e., eating the same amount of feed per kg of metabolic BW). This implies that these metabolites can be regarded as reliable RFI biomarkers, as they remain linked to this phenotype despite fluctuations in feed intake. The connection of these metabolites to RFI in a manner independent of intake would imply that they reflect host metabolic processes underlying individual variability in how animals convert feed and nutrients into growth²⁰. This unique characteristic of a biomarker, combined with a proven genetic correlation, may confer a notable benefit when assisting multi-objective breeding programs aimed at enhancing metabolic feed efficiency regardless of intake. In this context, when the energy use efficiency was compared between divergent genetic RFI lines subjected to a similar feed intake (based on BW), it was concluded that selection for RFI in cattle may not necessarily lead to an improved efficiency of energy use²⁴. They found that most of the metabolic differences observed across cattle RFI lines were attributed to variations in their feeding levels. Our results highlight the presence of some metabolic distinction between RFI groups that cannot be solely attributed to differences in feed intake levels.

It is important to note that the only metabolites associated with RFI independent of feed intake in our study (FDR < 0.05), namely, α -amino adipic acid and 5-aminovaleric acid (also known as 5-aminopentonic acid), for Groups A and B, respectively, both belong to the same metabolic pathway: lysine degradation. Previous studies in beef cattle also found lower plasma concentrations of α -amino adipic acid²⁵ and 5-aminovaleric acid¹⁷ in efficient (low-RFI) compared to inefficient (high-RFI) cattle and proposed them as candidate biomarkers of RFI. One of the first studies to propose plasma metabolomics to identify RFI biomarkers in beef cattle¹² also found that the lysine degradation pathway was linked to the RFI phenotype. The results from our team have similarly identified the lysine degradation pathway as a key metabolic pathway associated with differences in divergent RFI young bulls¹¹. Furthermore, we have previously demonstrated that lysine plasma concentration, along with that of




Figure 1. Experimental design of the study. Both Group A and Group B consisted of 30 growing-fattening beef cattle each (illustrated as circles within boxes, with boxes representing pens). During the first residual feed intake (RFI) test (0–70 days), Group A was fed ad libitum, while Group B underwent feed restriction. In both groups, 6 animals with the lowest and 6 with the highest RFI values were identified and sampled for blood (depicted as 24 black circles during the first RFI test). Following a 1-week transition period, the treatments were switched for the second RFI test (78–148 days): Group A underwent feed restriction, and Group B was fed ad libitum. Only the 24 cattle with the most extreme RFI values, sampled during the first RFI test, were resampled during the second RFI test (depicted as 24 black circles during the second RFI test). In Group A, the same RFI grouping established during ad libitum conditions (highest [H] and lowest [L] RFI) was maintained during feed restriction. In contrast, due to minimal RFI variation during feed restriction, the RFI ranking in group B (ranking number within black circles), as established during ad libitum conditions (second RFI test), was retained during feed restriction as well. For both groups and RFI tests, the retained RFI values were those obtained in ad-libitum conditions.

branched-chain amino acids, decreased in efficient compared to inefficient RFI heifers¹⁵. Likewise, studies in pigs have also linked plasma α -amino adipic acid levels and the lysine degradation pathway to the RFI phenotype²⁶.

Lysine degradation intermediates identified in this study may have both endogenous and microbial origins^{27–29}. Consequently, our study does not allow us to distinguish whether these differences between efficient and inefficient RFI cattle originate from host metabolism or host-related microbiota. Interestingly, a recent study conducted in beef heifers³⁰ identified lysine degradation as one of the unique pathways downregulated in the rumen of efficient RFI cattle. Similarly, in a study conducted in beef steers³¹, they found lower concentrations of several metabolites related to lysine metabolism, primarily N(6)-methylhistidine and cadaverine (the immediate precursor of 5-aminovaleric acid), in the rumen of more efficient RFI individuals. In a metatranscriptomic profiling study in beef cattle³², it was found that the lysine metabolism pathway, along with four other amino acids, was more active in the rumen microbiota of high-RFI individuals. Collectively, these results, combined with the findings of our study, may support the association between rumen microbiota composition or activity and the RFI phenotype^{30,33}, which is not explained by differences in feed intake. However, further studies are warranted to assess whether these two RFI biomarkers, which remain consistent regardless of the feed intake level, are related to endogenous host or microbial metabolism.

Other amino acids and related metabolites were found to be associated ($P < 0.05$) with RFI, regardless of the feeding level; however, they did not reach statistical significance after FDR correction ($FDR > 0.05$), likely due to the limited number of animals used in this experiment ($n = 6$ per classification). Among these, it is interesting to note that isoleucine, one of the three BCAAs, was 12% higher in less efficient animals from Group A. The same trend was also observed for the other two BCAAs, leucine and valine, which tended to differ between RFI phenotypes, regardless of the feeding level (raw P values < 0.10 ; data not shown). This confirms results reported by others in beef cattle^{11,14,16} and growing lambs³⁴, showing lower plasma concentrations of BCAAs in efficient RFI individuals. Our findings, even though they did not reach statistical significance after false positive correction, align with previous conclusions obtained in a meta-analysis of genome-wide association studies in beef cattle¹⁸, where BCAA degradation was identified as the only metabolic pathway associated with RFI. In contrast, the plasma concentration of none of the BCAAs was correlated with RFI values obtained in animals from Group B, those starting with feed restriction. Given the link between adiposity and BCAA plasma concentration in cattle³⁵, we cannot rule out the possibility that slight differences in adipogenesis in animals from Group A vs. B (Table 1) may mask the relationships between BCAA and RFI in those animals starting with feed restriction (Group B). Finally, it is interesting to note that the plasma concentration of IGF-1, although not statistically significant after FDR correction, exhibited higher values in efficient RFI cattle (raw P value = 0.046), whether they were fed ad libitum or restricted to a similar feeding level as their inefficient counterparts (RFI \times feeding level; $P > 0.10$).

The plasma concentration of IGF-1 has been shown to correlate with RFI in various cattle studies^{15,36}. However, diet-dependent relationships were highlighted in a review of the literature⁴, with a negative relationship, such as the one found in the present study, consistently observed only with low-concentrate and low-energy diets.

As opposed to lysine-degradation intermediates found in the present study, our results suggest that the plasma concentration of sarcosine, previously reported to differ across divergent RFI cattle fed diets rich in grass silage³⁷, could be associated with this phenotype only because it covaries with feed intake (RFI × FL interaction; $P = 0.012$ in Group B; Table 7). Similar results were found for glycine, its natural precursor, in genetically divergent chicken lines for RFI when fed ad libitum or under restrictive feeding²⁰. This finding is also consistent with previous studies, indicating that glycine is an amino acid whose plasma concentration in ruminants is inversely correlated with dietary protein intake³⁸.

The impact of feeding level on plasma parameters

Many plasma parameters evaluated here were significantly impacted by feeding level ($FDR < 0.05$), regardless of whether feed restriction was applied in the first or second test. The high correlation between feed intake and RFI, on the one hand, and feed intake and plasma metabolite concentration, on the other, may pose a challenge in determining whether plasma RFI biomarkers will remain consistent when animals are fed at similar feed intake levels. Among the plasma parameters which increased their values during feed restriction were some endogenous metabolites, such as methyl-histidines, which are known to increase during muscle degradation and mobilization in cattle³⁹, as well as other specific markers of (1) lipid mobilization, such as NEFA (highest-fold change in our study) or individual fatty acids⁴⁰ and (2) protein mobilization, such as $\delta^{15}N$ values^{41,42}. The values of these plasma markers of body mobilization agree with the effect of feeding level on the measured fat and muscle depth change, where feed restriction was associated with much lower tissue depth changes (close to 0 during the second test) compared to ad libitum feeding.

Furthermore, our results agree with previous findings showing that glycine and betaine plasma concentrations strongly increased when feed (protein) intake decreased in humans and mammals⁴³ and during feed restriction in female mink²³, respectively. In addition, we observed that serine and sarcosine, two metabolites related to glycine metabolism⁴³, also increased their plasma concentration during feed restriction, which suggests that the serine-glycine biosynthesis pathway is strongly regulated by feed intake. On the other hand, few plasma parameters decreased their values during feed restriction ($FDR < 0.05$). The lower plasma concentrations of β -hydroxybutyrate and hippuric acid during feed restriction likely indicated lower absorption of rumen butyrate (one of the three main volatile fatty acids) and benzoic acids (aromatic compounds found in forages), their main substrates, respectively^{44,45}. The third plasma metabolite that decreased in concentration when feed intake decreased was β -alanine. This result could be expected given that β -alanine is a breakdown product from the muscle-rich dipeptides anserine and carnosine, the latter also decreasing its concentration during feed restriction (raw P value = 0.018).

Our study revealed several plasma parameters that, although affected by feeding level, enabled the ranking of animals across time (between-feeding levels). The repeatability quantifies the proportion of total phenotypic variation that is attributed to differences between individuals and may suggest that genetic factors are likely to be important. The moderate-to-high repeatability observed for some plasma parameters in this study indicates that they are somehow inherent to animal metabolism beyond the effects of feeding level. To have confidence in their predictive ability, it is important that biomarkers consistently rank individual animals even when the environment changes. 5-Aminovaleric acid and α -amino adipic acid, the only two metabolites related to RFI in a feeding level-independent manner, showed moderate (0.59) to high repeatability (0.65), respectively. Among the plasma parameters showing the highest repeatability between-feeding levels, there were endogenous metabolites such as 1 and 3-methylhistidine, alkaline phosphatase and carnosine and others with a known microbial origin, such as indoleacetic acid (major product of tryptophan metabolism by microbial population in the rumen⁴⁶). Plasma metabolites such as methylhistidines and carnosine, with high repeatability across feeding levels, are endogenous metabolites that share a common feature: originating from skeletal muscle turnover and metabolism⁴⁷. Furthermore, alkaline phosphatase is an enzyme related to bone turnover and metabolism, and its blood levels have been shown to be genetically regulated in mice, with a heritability of 0.56, and related to bone mass density⁴⁸. Given that the proportion of carcass, comprising both muscle and bones, to total body weight are heritable traits in beef cattle⁴⁹ ($h^2 = 0.51$) it is not surprising that plasma metabolites reflecting muscle and bone turnover rates (mainly 3-methylhistidines and phosphatase alkaline) showed very high repeatability across time in the present study.

The plasma creatinine concentration showed moderate repeatability in our study (0.66), consistent with findings in dairy cows^{50,51}. Most highly conserved metabolites are also highly heritable⁵² and less influenced by environmental factors. Indeed, in a genome-wide association study conducted in crossbred beef cattle, creatinine ranked as the second highest plasma metabolite in terms of heritability⁵³ ($h^2 = 0.35$) and showed moderate heritability (0.45) when measured in cow's milk⁵⁰. Plasma creatinine is a product of muscle metabolism and has been proposed to reflect lean body mass in ruminants⁵⁴, an animal trait with moderate heritability and phenotypic and genetic correlations with RFI⁴⁹. Although several studies identified a relationship between plasma creatinine concentration and RFI^{11,12}, overall, we observed no correlation in our study. The study by Karisa et al.¹² highlighted that, in addition to explaining 26% of the phenotypic RFI variation observed in beef cattle, plasma creatinine also contributed to explaining 13% of DM intake variation. Further studies are needed to confirm whether the relationship between plasma creatinine concentration and RFI is confounded by covariations in DM intake.

Conclusions

In summary, our study revealed that metabolites belonging to the lysine degradation pathway, including α -amino adipic acid and 5-aminovaleric acid, emerged as consistent and repeatable biomarkers for RFI. These metabolites displayed a remarkable characteristic of remaining linked to RFI regardless of fluctuations in feed intake, likely reflecting underlying host metabolic processes associated with individual variability in feed efficiency. Our results support the notion that some of the phenotypic variance of RFI may not be solely attributed to differences in feed intake. Under the conditions of this study, we cannot confirm our hypothesis that the association between plasma branched-chain amino acids and RFI is independent of fluctuations in feeding level. Furthermore, our research has underscored changes in plasma parameter values during feed restriction, primarily associated with muscle and adipose tissue mobilization. Further investigations are needed to further validate our findings and to determine whether the identified RFI biomarkers have their origins in microbial or host metabolism. This step is essential before considering their utilization in assisting breeding programs and precision feeding strategies.

Methods

This study was conducted at the Teagasc Animal & Grassland, Research and Innovation Centre (Grange, Ireland) between 2021 and 2022. All experimental procedures involving animals were approved by the Teagasc Animal Ethics Committee (TAEC No. TAEC2021-304) and the Irish Health Products Regulatory Authority (HPRA, License number AE19132/P141) in accordance with the European Union (Protection of Animals used for Scientific Purposes) Regulations 2012 (S.I. No. 543/2012). Prior to commencing the study, all animals were treated for the control of internal parasites, skin lice, and vaccinated against clostridial and respiratory diseases. This experiment complied with the ARRIVE guidelines.

Animals, dietary treatments and experimental design

Sixty Charolais crossbred young bulls, with an average age of 13 ± 1.4 months and an initial body weight (BW) of 427 ± 20.4 kg, were utilized in a crossover design study, encompassing two 70-d FE test periods, as illustrated in Fig. 1. Bulls were allocated to one of two 'groups' (Group A, $n = 30$, and Group B, $n = 30$) in a randomized block design, with blocking based on initial live weight. Bulls were accommodated in a slatted floor building, with five pens of six animals per treatment. Pens within treatment were distributed throughout the building.

Prior to the commencement of the first RFI test, animals underwent a 21-day adaptation period to acclimatize to the diet and environment. Group A was provided ad libitum feeding during the initial 70-day RFI test, followed by a period of feed restriction in the subsequent 70-day RFI test. Conversely, Group B experienced the opposite sequence, commencing with feed restriction in the first test and then transitioning to ad libitum feeding for the second test. A one-week transition interval was implemented between the two RFI test periods.

Throughout the study, all animals were fed the same total mixed ration (TMR), which comprised a blend of grass silage and concentrate at a 50:50 ratio on a DM basis. The concentrate portion of the diet was formulated using rolled barley, soybean meal, and cane molasses plus minerals and vitamins. The TMR had a DM concentration of 528 g/kg, a crude protein (CP) content of 138 g/kg DM, and an estimated net energy (NE) concentration for maintenance and growth of 1.58 Mcal/kg DM². The chemical composition and in vitro digestibility of the grass silage and concentrate feed offered are detailed in Supplementary Table S1. Animals had free access to clean, fresh drinking water.

Bulls were individually offered their respective dietary feeding level through electronically controlled Calan doors (American Calan Inc., Northwood, NH, USA) once daily after 08:00. Ad libitum feed intake was based on a targeted refusal rate of approximately 5–10%; refused feed was weighed daily and discarded twice weekly. Feed restriction was set at an intermediate point between ad-libitum conditions and a feeding level that only met the energy requirements for maintenance (zero-growth). Considering the theoretical chemical composition and dietary values of the feed, along with the observed feeding level during the adaptation period (1.88% of body weight), the feed restriction level equated to 1.45% of body weight on a DM basis for each individual animal. Accordingly, the quantity of feed offered to individual animals in the restricted group(s) was adjusted every 2 weeks based on individual animal live weight. This adjustment resulted in a projected average daily live weight gain of 0.62 kg for a 400 kg young bull, as opposed to the 1.35 kg/day expected under ad-libitum conditions.

Representative samples of the grass silage and concentrates offered were collected three and two times weekly, respectively. Silage DM was determined by oven drying samples at 85 °C for 16 h; DM values were corrected for loss of volatiles using equation 1 proposed in⁵⁵. Concentrate samples were oven-dried at 98 °C for 16 h to determine DM. Concentrations of crude protein, neutral detergent fibre, acid detergent fibre, ash, starch and oil-B and in vitro measures of digestibility were determined in the feedstuffs (wet chemistry), as described previously⁵⁶.

Animal live weight was recorded on two consecutive days at the beginning and the end of each feed efficiency test period and weekly throughout using a calibrated scale. Weighing was carried out in the morning prior to feeding. The depth of subcutaneous fat (13th rib, third lumbar vertebra and the rump) and *M. longissimus* muscle (third lumbar vertebra) was measured at the beginning and end of each test period using an automatic real-time scanner (model—ECM ExaGo Veterinary scanner, with a 3.5 MHz linear transducer, IMV imaging, Meath, Ireland), as described previously⁵⁷.

Blood sampling and analysis

Blood was collected prior to the morning feeding at the end of the first FE test period from the 12 most divergent RFI (6 high-RFI and 6 low-RFI) animals from each group; the same animals were blood sampled again at the end of the second FE test period (Fig. 1). Blood samples were obtained using jugular venipuncture into two 9-mL evacuated vials containing lithium heparin (Vacurette; Cruinn Diagnostics, Dublin, Ireland) and two

8.5-mL evacuated vials containing silica clot activator (BD Vacutainer SST (II) Advance, Unitech). The lithium heparin blood tubes were placed in ice, and the serum separator tubes were left for approximately 24 h at 4 °C to permit clotting prior to centrifugation (1600 × g for 10 min). The resulting plasma and serum were divided into 1.5 mL plastic vials and stored at – 80 °C until subsequent analyses. The resulting supernatant was subjected to metabolomic profiling analysis at the core lab of Biocrates Life Science AG (Innsbruck, Austria).

All 48 plasma samples were stored upon receipt at – 80 °C and thawed on ice before analysis. The thawed samples were immediately pipetted to ensure that all samples were analyzed within 30 min, thereby avoiding any changes that could affect metabolite levels. To directly quantify known metabolites, we employed a commercially available kit (MxP Quant 500 Assay) utilizing liquid chromatography-tandem mass spectrometry, allowing for the quantification of up to 106 metabolites. This targeted metabolic approach included the analysis of one alkaloid, one amine oxide, 20 amino acids, 30 amino acid-related metabolites, 14 bile acids, nine biogenic amines, one carbohydrate and related, seven carboxylic acids, one cresol, 12 fatty acids, four hormones and related, four indoles and derivatives, two nucleobases and related and one vitamin. Only 71 out of the 106 targeted metabolites had values above the limit of detection for all observations and were used in further analysis. Among these metabolites, several were previously identified as candidate RFI biomarkers through untargeted and targeted approaches in beef cattle studies^{11,12,15}. Additionally, the plasma samples were subjected to spectrophotometric quantification of various metabolites, including glucose, urea, nonesterified fatty acids, β-hydroxybutyrate, as detailed in Jorge-Smeding et al.¹¹ and hepatic transaminases as previously described⁵⁸, some of which were identified as potential RFI biomarkers by Richardson et al.⁵⁹. The natural abundance of ¹⁵N, a promising biomarker of FE in beef cattle⁶⁰, was determined following the methodology outlined in Cantalapiedra-Hijar et al.⁶¹. Finally, using commercial ELISA kits, plasma insulin (bovine insulin ELISA kit Cat. no. EB0092, FineTest, China) and IGF-1 (bovine IGF-1 ELISA kit Cat. No. 221121, Mediadiagnost, GmbH, Tübingen, Germany) concentrations were also analysed from blood serum samples. Both hormones were also found to differ across extreme RFI cattle in previous studies^{11,15,62}.

Calculations and statistical analysis

The RFI values of each animal were computed as the difference between observed and predicted DM intake. DM intake was predicted from the following factors: block, pen within block, mid-test metabolic body weight (mBW^{0.75}), ADG and changes in backfat and muscle ultrasound measurements between the end and beginning of the test as commonly recommended⁵. This computation followed the equation provided below:

$$\text{DM intake} = \beta_0 + \text{Block} + \text{Pen} + \beta_1 \times \text{mBW}^{0.75} + \beta_2 \times \text{ADG} + \beta_3 \times \text{ultrasound depth change} + \varepsilon;$$

where β_0 represents the regression intercept, β_1 the partial regression coefficient of mean metabolic BW, β_2 the partial regression coefficient of ADG, β_3 the partial regression for the ultrasound depth change and ε is the residual error of the regression for each animal, or RFI.

The ADG used in the RFI model was determined as the regression of BW over time, except in the case of ADG measured in group B during the first test period, where nonlinear growth ($r^2 < 0.9$ for some animals) necessitated calculation as the difference between final BW and initial BW divided by 70 days.

Separate RFI models were established for each group and test period, totaling four conditions. Consequently, the models featured distinct significant variables depending on the specific condition, as outlined in Supplementary Table S2. All statistical analyses were performed using R software (version 2021.09.0; R Core Team, 2021). Variables that did not adhere to a normal distribution ($P < 0.05$ after the Shapiro–Wilk test) were log-transformed prior to analysis. We utilized a mixed-effects model analysis with the nlme package in R, consistently including “animal” as a random effect when repeated measures were analysed. Different fixed effects were examined in separate analyses. To assess the effect of feeding level, we included it as a fixed effect, along with the effect of the test period and their interaction. To evaluate the effect of RFI across the two feeding levels, both factors and their interaction were included as fixed effects in the model. However, as elaborated later, we treated the RFI effect differently according to the Group: as a qualitative variable in Group A (categorizing animals into high vs. low RFI extremes) and as a continuous variable in Group B (with values ranging from positive to negative, including values close to zero). The raw P values obtained from the analysis of plasma variables ($n = 83$) were adjusted using the Benjamini–Hochberg false discovery rate (FDR) correction. The significance threshold was set at $\text{FDR} \leq 0.05$. However, because some variables have already been identified as candidate plasma biomarkers in previous studies, we also examined raw P values. In this context, the likelihood of a false positive is lower when compared to new plasma biomarkers. The repeatability of plasma parameters across the two feeding levels was calculated as the intraclass correlation coefficient (i.e., animal variance divided by the sum of animal variance and residual variance) in a model that explained the variation in plasma parameters using feeding level, test period, and their interaction as fixed effects, along with the random animal effect.

Data availability

Data is provided within the manuscript or supplementary information files. Raw data generated and/or analysed during the current study are available from the corresponding author upon reasonable request.

Received: 18 December 2023; Accepted: 8 April 2024

Published online: 12 April 2024

References

1. OECD & Food and Agriculture Organization of the United Nations. *OECD-FAO Agricultural Outlook 2021–2030*. (OECD, 2021). <https://doi.org/10.1787/19428846-en>.

2. INRA. *INRA Feeding System for Ruminants* (Wageningen Academic Publishers, 2018). <https://doi.org/10.3920/978-90-8686-292-4>.
3. National Academies of Sciences *et al. Nutrient Requirements of Beef Cattle, 8th Revised Edition* 19014 (National Academies Press, 2015). <https://doi.org/10.17226/19014>.
4. Cantalapiedra-Hijar, G. *et al.* Review: Biological determinants of between-animal variation in feed efficiency of growing beef cattle. *Animal* **12**, s321–s335 (2018).
5. Berry, D. P. & Crowley, J. J. Cell biology symposium: Genetics of feed efficiency in dairy and beef cattle. *J. Anim. Sci.* **91**, 1594–1613 (2013).
6. Byrne, T. J. *et al.* New breeding objectives and selection indices for the Australian dairy industry. *J. Dairy Sci.* **99**, 8146–8167 (2016).
7. Tedeschi, L. O. *et al.* A glimpse of the future in animal nutrition science. 2. Current and future solutions. *R. Bras. Zootec.* **46**, 452–469 (2017).
8. Pomar, C., Andretta, I. & Remus, A. Feeding strategies to reduce nutrient losses and improve the sustainability of growing pigs. *Front. Vet. Sci.* **8**, 742220 (2021).
9. Taussat, S. *et al.* Gene networks for three feed efficiency criteria reveal shared and specific biological processes. *Genet. Sel. Evol.* **52**, 67 (2020).
10. Koch, R. M., Swiger, L. A., Chambers, D. & Gregory, K. E. Efficiency of feed use in beef cattle. *J. Anim. Sci.* **22**, 486–494 (1963).
11. Jorge-Smeding, E. *et al.* Common and diet-specific metabolic pathways underlying residual feed intake in fattening Charolais yearling bulls. *Sci. Rep.* **11**, 24346 (2021).
12. Karisa, B. K. *et al.* Plasma metabolites associated with residual feed intake and other productivity performance traits in beef cattle. *Livestock Sci.* **165**, 200–211 (2014).
13. Richardson, E. C. & Herd, R. M. Biological basis for variation in residual feed intake in beef cattle. 2. Synthesis of results following divergent selection. *Aust. J. Exp. Agric.* **44**, 431 (2004).
14. Foroutan, A., Fitzsimmons, C., Mandal, R., Berjanskii, M. V. & Wishart, D. S. Serum metabolite biomarkers for predicting residual feed intake (RFI) of Young Angus Bulls. *Metabolites* **10**, 491 (2020).
15. Jorge-Smeding, E. *et al.* Untargeted metabolomics confirms the association between plasma branched chain amino acids and residual feed intake in beef heifers. *Plos One* **17**, e0277458 (2022).
16. Li, J., Mukiibi, R., Wang, Y., Plastow, G. S. & Li, C. Identification of candidate genes and enriched biological functions for feed efficiency traits by integrating plasma metabolites and imputed whole genome sequence variants in beef cattle. *BMC Genom.* **22**, 823 (2021).
17. Taiwo, G. *et al.* Chemical group-based metabolome analysis identifies candidate plasma biomarkers associated with residual feed intake in beef steers. *Front. Anim. Sci.* **2**, 783314 (2022).
18. Duarte, D. A. S. *et al.* Genome-wide association studies pathway-based meta-analysis for residual feed intake in beef cattle. *Anim. Genet* **50**, 150–153 (2019).
19. Patton, R. A., Hristov, A. N., Parys, C. & Lapierre, H. Relationships between circulating plasma concentrations and duodenal flows of essential amino acids in lactating dairy cows. *J. Dairy Sci.* **98**, 4707–4734 (2015).
20. Metzler-Zebeli, B. *et al.* Feed restriction reveals distinct serum metabolome profiles in chickens divergent in feed efficiency traits. *Metabolites* **9**, 38 (2019).
21. Andreini, E. M., Augenstein, S. M., Fales, C. S., Sainz, R. D. & Oltjen, J. W. Effects of feeding level on efficiency of high- and low-residual feed intake beef steers. *J. Anim. Sci.* **98**, skaa286 (2020).
22. Boddicker, N., Gabler, N. K., Spurlock, M. E., Nettleton, D. & Dekkers, J. C. M. Effects of ad libitum and restricted feeding on early production performance and body composition of Yorkshire pigs selected for reduced residual feed intake. *Animal* **5**, 1344–1353 (2011).
23. Hedemann, M. S. & Damgaard, B. M. Metabolomic study of plasma from female mink (*Neovison vison*) with low and high residual feed intake during restrictive and ad libitum feeding. *Compar. Biochem. Physiol. Part D: Genom. Proteom.* **7**, 322–327 (2012).
24. Lines, D. S., Pitchford, W. S., Bottema, C. D. K., Herd, R. M. & Oddy, V. H. Selection for residual feed intake affects appetite and body composition rather than energetic efficiency. *Anim. Prod. Sci.* **58**, 175 (2018).
25. Goldansaz, S. A. *et al.* Candidate serum metabolite biomarkers of residual feed intake and carcass merit in sheep. *J. Anim. Sci.* **98**, skaa298 (2020).
26. Carmelo, V. A. O., Banerjee, P., da Silva Diniz, W. J. & Kadarmideen, H. N. Metabolomic networks and pathways associated with feed efficiency and related-traits in Duroc and Landrace pigs. *Sci. Rep.* **10**, 255 (2020).
27. Broquist, H. P. Lysine-pipecolic acid metabolic relationships in microbes and mammals. *Annu. Rev. Nutr.* **11**, 435–448 (1991).
28. Callery, P. S. & Geelhaar, L. A. Biosynthesis of 5-aminopentanoic acid and 2-piperidone from cadaverine and 1-piperidine in mouse. *J. Neurochem.* **43**, 1631–1634 (1984).
29. Tucker, H. A., Hanigan, M. D., Escobar, J., Doane, P. H. & Donkin, S. S. Hepatic expression of amino adipate semialdehyde synthase is unchanged by post-ruminal lysine supply in lactating dairy cows. *J. Dairy Sci.* **100**, 1009–1018 (2017).
30. Liu, Y. *et al.* Rumen microbiome and metabolome of high and low residual feed intake angus heifers. *Front. Vet. Sci.* **9**, 812861 (2022).
31. Sidney, T. *et al.* Rumen fluid amine/phenol-metabolome of beef steers with divergent residual feed intake phenotype. *Ruminants* **3**, 1–8 (2023).
32. Li, F. & Guan, L. L. Metatranscriptomic profiling reveals linkages between the active rumen microbiome and feed efficiency in beef cattle. *Appl. Environ. Microbiol.* **83**, e00061-e117 (2017).
33. Clemmons, B. A. *et al.* Ruminal bacterial communities and metabolome variation in beef heifers divergent in feed efficiency. *Ruminants* **2**, 282–296 (2022).
34. Touitou, F. *et al.* Evaluation of the links between lamb feed efficiency and rumen and plasma metabolomic data. *Metabolites* **12**, 304 (2022).
35. Webb, L. A. *et al.* Branched-chain amino acids: Abundance of their transporters and metabolizing enzymes in adipose tissue, skeletal muscle, and liver of dairy cows at high or normal body condition. *J. Dairy Sci.* **103**, 2847–2863 (2020).
36. Johnston, D. J. *et al.* Evidence of IGF1 as a genetic indicator of feed efficiency traits in beef cattle. In *7th World Cong. Genet. Appl. Livest. Prod., CD ROM Communication. No. 10–16* (2023).
37. Meale, S. J. *et al.* Exploration of biological markers of feed efficiency in young bulls. *J. Agric. Food Chem.* **65**, 9817–9827 (2017).
38. Broderick, G. A., Satter, L. D. & Harper, A. E. Use of plasma amino acid concentration to identify limiting amino acids for milk production. *J. Dairy Sci.* **57**, 1015–1023 (1974).
39. Houweling, M., van der Drift, S. G. A., Jorritsma, R. & Tielens, A. G. M. Technical note: Quantification of plasma 1- and 3-methylhistidine in dairy cows by high-performance liquid chromatography–tandem mass spectrometry. *J. Dairy Sci.* **95**, 3125–3130 (2012).
40. van der Drift, S. G. A., Houweling, M., Schonewille, J. T., Tielens, A. G. M. & Jorritsma, R. Protein and fat mobilization and associations with serum β -hydroxybutyrate concentrations in dairy cows. *J. Dairy Sci.* **95**, 4911–4920 (2012).
41. Chen, Y. T., McNamara, J. P., Ma, G. L., Harrison, J. H. & Block, E. Milk 13C and 15N discriminations as biomarkers of feed efficiency and energy status in early lactation cows. *Anim. Feed Sci. Technol.* **269**, 114638 (2020).
42. Correa-Luna, M. *et al.* Nitrogen isotopic discrimination as a biomarker of between-cow variation in the efficiency of nitrogen utilization for milk production: A meta-analysis. *J. Dairy Sci.* **105**, 5004–5023 (2022).
43. Wang, W. *et al.* Glycine metabolism in animals and humans: Implications for nutrition and health. *Amino Acids* **45**, 463–477 (2013).

44. Loncke, C. *et al.* Empirical prediction of net splanchnic release of ketogenic nutrients, acetate, butyrate and β -hydroxybutyrate in ruminants: A meta-analysis. *Animal* **9**, 449–463 (2015).
45. Martin, A. K. The origin of urinary aromatic compounds excreted by ruminants 2. The metabolism of phenolic cinnamic acids to benzoic acid. *Br. J. Nutr.* **47**, 155–164 (1982).
46. Allison, M. J., Robinson, I. M. & Baetz, A. L. Tryptophan biosynthesis from indole-3-acetic acid by anaerobic bacteria from the rumen. *J. Bacteriol.* **117**, 175–180 (1974).
47. Nishizawa, N. *et al.* NT-Methylhistidine content of organs and tissues of cattle and an attempt to estimate fractional catabolic and synthetic rates of myofibrillar proteins of skeletal muscle during growth by measuring urinary output of N T -methylhistidine. *Br. J. Nutr.* **42**, 247–252 (1979).
48. Srivastava, A. K., Masinde, G., Yu, H., Baylink, D. J. & Mohan, S. Mapping quantitative trait loci that influence blood levels of alkaline phosphatase in MRL/MpJ and SJL/J mice. *Bone* **35**, 1086–1094 (2004).
49. Taussat, S. *et al.* Relationship between feed efficiency and slaughter traits of French Charolais bulls. *J. Anim. Sci.* **97**, 2308–2319 (2019).
50. Buitenhuis, A. J. *et al.* Estimation of genetic parameters and detection of quantitative trait loci for metabolites in Danish Holstein milk. *J. Dairy Sci.* **96**, 3285–3295 (2013).
51. Peterson, R. G. & Waldern, D. E. Repeatabilities of serum constituents in holstein-friesians affected by feeding, age, lactation, and pregnancy. *J. Dairy Sci.* **64**, 822–831 (1981).
52. Yousri, N. A. *et al.* Long term conservation of human metabolic phenotypes and link to heritability. *Metabolomics* **10**, 1005–1017 (2014).
53. Li, J. *et al.* Genomic heritability and genome-wide association studies of plasma metabolites in crossbred beef cattle. *Front. Genet.* **11**, 538600 (2020).
54. Caldeira, R. M., Belo, A. T., Santos, C. C., Vazques, M. I. & Portugal, A. V. The effect of body condition score on blood metabolites and hormonal profiles in ewes. *Small Ruminant Res.* **68**, 233–241 (2007).
55. Porter, M. G. & Murray, R. S. The volatility of components of grass silage on oven drying and the inter-relationship between dry-matter content estimated by different analytical methods. *Grass Forage Sci.* **56**, 405–411 (2001).
56. Kennedy, R., Moloney, A. P., O’Riordan, E. G., Kelly, A. K. & McGee, M. Intake, growth and carcass traits of steers offered grass silage and concentrates based on contrasting cereal grain types supplemented with field beans, peas or maize by-products. *Animals* **13**, 1209 (2023).
57. Conroy, S. B. *et al.* Predicting beef carcass meat, fat and bone proportions from carcass conformation and fat scores or hindquarter dissection. *Animal* **4**, 234–241 (2010).
58. Delosière, M., Thomas, A., Terlouw, C. & Durand, D. Plasma indicators of bovine health: Impacts of diet supplementations and pre-slaughter stress. *Data Brief* **21**, 2496–2503 (2018).
59. Richardson, E. C., Herd, R. M., Archer, J. A. & Arthur, P. F. Metabolic differences in Angus steers divergently selected for residual feed intake. *Aust. J. Exp. Agric.* **44**, 441 (2004).
60. Cantalapiedra-Hijar, G. *et al.* Identifying cattle with superior growth feed efficiency through their natural ¹⁵N abundance and plasma urea concentration: A meta-analysis. *Peer Commun. J.* **2**, e31 (2022).
61. Cantalapiedra-Hijar, G. *et al.* Diet–animal fractionation of nitrogen stable isotopes reflects the efficiency of nitrogen assimilation in ruminants. *Br. J. Nutr.* **113**, 1158–1169 (2015).
62. Moore, K. L., Johnston, D. J., Graser, H.-U. & Herd, R. Genetic and phenotypic relationships between insulin-like growth factor-I (IGF-I) and net feed intake, fat, and growth traits in Angus beef cattle. *Aust. J. Agric. Res.* **56**, 211 (2005).

Acknowledgements

We acknowledge the technical assistance of M.J. Kelly, Teagasc Grange, the Grange farm staff for animal management, and Dr. A. Ikoyi and Grange Laboratories personnel for feed analyses and sample processing. We thank Céline Chantelauze and Marine Gauthier from INRAE UMR Herbivores for their technical assistance in laboratory analysis. This work received funding from SmartCow, a project funded by the European Union’s Horizon 2020 research and innovation program under Grant Agreement no. 730924. The hormone analysis was funded by a scientific project titled “Development of Research and Innovation at Trakia University in Service of Health and Sustainable Well-Being,” sponsored by the Bulgarian Ministry of Education and Science (Grant number: BG-RRP-2.004-0006).

Author contributions

G.C.-H., K.N., P.C., M.M. designed the experiment. P.C. and M.M. performed the experiment. GCH analysed and interpreted the experimental data. G.C.-H. wrote the draft manuscript. P.C. and M.M. reviewed and edited the manuscript. All the authors have read and approved the final version.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-024-59253-7>.

Correspondence and requests for materials should be addressed to G.C.-H.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher’s note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2024