



**HAL**  
open science

## Determination of digestive enzymes activities in human duodenal fluids

Olivia Ménard, Yann Le Gouar, Marlies Braeckmans, Patrick Augustijns,  
Didier Dupont

► **To cite this version:**

Olivia Ménard, Yann Le Gouar, Marlies Braeckmans, Patrick Augustijns, Didier Dupont. Determination of digestive enzymes activities in human duodenal fluids. <https://www.icfd2024.com/>. 8th International Conference on FOOD DIGESTION, Apr 2024, Porto, Portugal. , 2024. hal-04546976

**HAL Id: hal-04546976**

**<https://hal.inrae.fr/hal-04546976>**

Submitted on 15 Apr 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

# Determination of digestive enzymes activities in human duodenal fluids

Olivia Ménard<sup>1</sup>, Yann Le Gouar<sup>1</sup>, Marlies Braeckmans<sup>2</sup>, Patrick Augustijns<sup>2</sup>, Didier Dupont<sup>1</sup>

<sup>1</sup>INRAE, Institut Agro, STLO, 35042 Rennes, France

<sup>2</sup>Department of Pharmaceutical and Pharmacological Sciences, KU Leuven, Belgium


## Introduction

### Context

The improvement of *in vitro* digestion models is essential and depends, among others, on a better knowledge of the enzymatic activities in the human digestive tract. Within the framework of a collaboration between INFOGEST and UNGAP, a COST Action on oral drug absorption, access to intestinal fluids was possible.

### Objectives

Determination of enzymes activities in duodenum effluents : trypsin, chymotrypsin and lipase and pH measurements




New set of data for enzymes activities in human duodenum

### Analysis


## Materials and methods

- 13 healthy adult volunteers were used for the determination of digestive enzyme activities
- Meal : \* 240 ml of **water (fasted)**  
Or \* 400 ml of **Ensure Plus vanilla** +240 mL of water (**fed**)
- Kinetics of sampling : from -60 min before ingestion to 110 min after water or meal ingestion
- Samples storage: samples were mixed with glycerol (1:1) and inhibitors before storage at -20°C
- Statistical analysis: paired t-test


Duodenum fluids aspirates n=13



Enzyme activities Total samples n=88




pH duo n=88




Enzymatic assays : Brodkord et al;2019 Supplementary Material except that trypsin and chymotrypsin activities were determined at 37°C instead of 25°C, for lipase no modification (37°C).

## Results

### Stage effect: fasted vs fed-state



### Post-prandial time effect



- Stage effect on lipase activity
- Average trypsin: mean<sub>fasted</sub> = 63.2 ± 30.6 vs mean<sub>fed</sub> = 64.9 ± 17.2 U/ml
- Average chymotrypsin: mean<sub>fasted</sub> = 9.9 ± 3.9 vs mean<sub>fed</sub> = 6.7 ± 1.7 U/ml
- Average lipase: mean<sub>fasted</sub> = 457 ± 299 vs mean<sub>fed</sub> = 1093 ± 376 U/ml
- Average pH: pH<sub>fasted</sub> = 6.5 ± 1.3 vs pH<sub>fed</sub> = 6.0 ± 0.5
- Fasted: 1 time point / volunteer
- Fed 2 to 9 time points /volunteer

- No effect of post-prandial time on enzyme activities / effect on pH
- Average: mean<sub>trypsin activity</sub> = 62.1 ± 29.9 U/ml / mean<sub>chymotrypsin</sub> = 6.1 ± 2.6
- Average: mean<sub>lipase activity</sub> = 1129 ± 612 U/ml / pH<sub>average</sub> = 6.1 ± 2.6
- Ratio<sub>trypsin/chymotrypsin</sub> = 10.2
- 25 vs 37°C trypsin & chymotrypsin temperature assay : a 1.8 factor

	Trypsin (U/ml of intestinal content)	Lipase (U/ml of intestinal content)	assay temperature (°C)
Infogest digestion model	100	2000	25
In vivo (fed state)	34.5		25
In vivo (fed state)	62.1	1129	37

Care should be taken to use similar conditions for enzymatic assays to compare the data.

## Conclusions

There is currently little data in the literature on enzyme levels in the GI tract, so this new data set is of great importance. These results highlight the need for additional in vivo fluid characterisation and would help the scientific community to adapt and improve *in vitro* digestion models.