



**HAL**  
open science

## Socio-metabolism approaches and their relevance for analysing agricultural transitions

Claire Aubron, Charlotte Hemingway, Olivier Philippon, Laurent Ruiz,  
Mathieu Vigne

► **To cite this version:**

Claire Aubron, Charlotte Hemingway, Olivier Philippon, Laurent Ruiz, Mathieu Vigne. Socio-metabolism approaches and their relevance for analysing agricultural transitions. Sustainability Transitions in Agri-food Systems: Evidence and Options for Action in India and France, Marie Dervillé, Rajeswari Raina, CEFIPRA, Mar 2024, Roorkee, India. hal-04547323


**HAL Id: hal-04547323**

**<https://hal.inrae.fr/hal-04547323>**

Submitted on 15 Apr 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# SOCIO-METABOLISM APPROACHES AND THEIR RELEVANCE FOR ANALYSING AGRICULTURAL TRANSITIONS

Claire Aubron,  
Charlotte Hemingway,  
Olivier Philippon,  
Laurent Ruiz,  
Mathieu Vigne


# TWO SOCIO-METABOLISM CASE STUDIES

- Socio-metabolism research: a wide range of approaches with a common interest in **flows** of matter and energy
- Our approach in this presentation:



- Two case-studies in contrasted areas of India, not particularly involved in agroecological transitions but which we use as a basis for drawing lessons

# 1<sup>ST</sup> CASE-STUDY: NITROGEN METABOLISM AND GHG EMISSIONS IN A GUJARATI VILLAGE (2014)



**PLATEAU and HIGHLANDS**  
Hard rock aquifer

KARNATAKA

1. Channagiri (Davangere District)
2. Gundlupet (Chamarajanagar District)

ANDHRA PRADESH

3. Vinukonda (Guntur District)
4. Banagana Palli (Kurnool District)
5. Palamaner (Chittoor District)

GUJARAT

6. Gondal (Rajkot District)

WEST BENGAL

7. Hirbandh (Bankura District)

**ALLUVIAL PLAINS**

BIHAR

8. Bodhgaya (Gaya District)
9. Ekangarsarai (Nalanda District)

GUJARAT

10. Petlad (Anand District)
11. Dharampur (Valsad District)

**DELTA**

WEST BENGAL

12. Debra (Medinipur District)
13. Bangaon (North 24 Parganas District)


Petlad block, an alluvial plain with semi-arid climate

Densely populated by **people** (550 inhab./km<sup>2</sup>) and **animals** (230 bovines/km<sup>2</sup>) and intensively cultivated with **irrigation**




## A highly unbalanced nitrogen metabolism

- Water pollution: 11 of the 16 samples taken are over the NO<sub>3</sub> potability limit of 50 mg/L
- Nitrogen balance at village scale: 600 kgN/ha/yr for crops (22 times higher than the French average); total surplus (including livestock activities): 900 kg/ha/yr




## Map of the N surplus at field scale in the village (N balance for crops)



## High GHG emissions at village level coming from a diversity of sources

- 37 tCO<sub>2</sub>eq/ha at village level, 15 times higher than the Indian national average
- Bovine enteric fermentation is the main source of GHG emissions (25%) but synthetic fertilizers and imported feed account for 42% of the total:



## Re-integrating crops and livestock: a solution?

- Fostering the cultivation of fodder crops or crops that produce residues (≠ tobacco) to feed animals from local resources.
- Managing animal manure to limit the use of synthetic fertilizers.

Yes, but... it is not the way things are evolving!

Division of land through inheritance

Agrarian reforms

Green revolution

White revolution

1950

1990


2014


Mid-size and large farms

Small farms

Pastoral communities

Landless labourers keeping livestock






## Conclusion for the 1<sup>st</sup> case-study

- Farmers are not encouraged to use synthetic nitrogen fertilizers sparingly, due to subsidies to the fertilizer industry.
- Social categories **with sufficient access to land** (>1 ha) have no interest in developing livestock farming, as they obtain high incomes from irrigated tobacco growing.
- Social categories **with limited access to land** have an interest in developing livestock farming to generate additional income through the sale of milk, but lack the resources to feed their animals.

## 2<sup>ND</sup> CASE STUDY: GROUNDWATER AND FOSSIL ENERGY CONSUMPTION IN ANDHRA PRADESH (2021)

Rapthadu mandal, a semi-arid area on the Deccan plateau




An **rainfed groundnut** area with a late and incomplete development of irrigation (vegetables and **fruit trees**)




... also going in the wrong direction from an environmental perspective

# Declining use of animal energy, rising consumption of fossil energy



*All values are expressed in GJ of energy*


# Territorial water balance: still positive but fragile




**Tipping point?**

**+ 130 ha of fruit plantations**


**Net recharge 2 947 601 m<sup>3</sup>**



**Net recharge 2 374 689 m<sup>3</sup>**



**Net recharge = 0 m<sup>3</sup>**



**Hemingway, 2023**

## Managing water better in order to avoid reaching the tipping point


- Favouring the cultivation of irrigated crops during the monsoon/rabi seasons and avoiding cultivation during the dry season, when evapotranspiration is high
- Limiting the expansion of fruit plantations, which require water all year round

In theory possible ... but that's not what is currently happening!

# Evolution of irrigated land and social diversity (1950-2022)


% households belonging to each social category

**1950s :**  
**4% land irrigated**  
 (Wash depressions)




**2022**  
**15% land irrigated**  
 (Wash pediment)

**1950**





**2022**




Huge income gaps, with access to irrigation (especially fruit cultivation) being the game changer

### Agricultural income (Rs/family worker/yr)



# 2022: the poorer social categories recharge groundwater for the wealthier ones



## Conclusion of the 2<sup>nd</sup> case-study

- Farmers are not encouraged to use groundwater sparingly because thanks to subsidies electricity is free and available 6-hours a day
- Social categories coming from former large holder farmers who accumulated capital with groundnut cultivation were able to **invest earlier in borewells** and irrigated crops in the 1990s. Part of them became **fruit growers** in the 2000s, earning today the highest agricultural incomes and pumping the largest quantities of groundwater in the area
- Today **groundnut growers**, for most of them former landless agricultural labourers who benefited from the agrarian reform in the 1970s, have remained in the blind spot of irrigation. They get very low incomes from rainfed cultivation and depend on the farmers having access to irrigation for employment opportunities. They make a major contribution to water recharge.

# CONCLUSION OF THE TWO CASE-STUDIES


- There are good reasons to reintegrate crops and livestock in Petlad and to manage groundwater differently in Rapphadu.
- The technical terms of such evolutions are relatively well known.
- But the political, social and economic conditions are not conducive to the implementation of agroecological practices:
  - Subsidies to chemical fertilizers and electricity for pumping water, on the contrary, encourage the use of inputs
  - Social organization hinders the changes: the well-off farmers have no interest in implementing these agroecological practices, while others do not have the means to do so.

# RELEVANCE OF SUCH SOCIO-METABOLISM APPROACH FOR AGROECOLOGICAL TRANSITION

Territorial level (ex:  
*water balance between  
recharge and  
groundwater pumping on  
different areas*)

Farm level (ex:  
*crop-livestock  
integration*)


Plot/herd level (ex:  
*irrigation, animal  
feeding practices...*)






Environmental impacts differentiated amongst farm types, with possible interdependences (ex: *water recharge/withdrawal*)

Inclusion of all social categories, with their specific means and interests for changes (ex: *landless workers, pastoralists, farmers without irrigation...*)





Thanks for your attention!



credit photo : Charlotte Hemingway