



HAL
open science

Influence of horn status on bovine Semitendinosus muscle proteome: An analysis of the canonical pathways, upstream regulators and interactive networks

Sudipta Hazra, Ruth M. Hamill, Mohammed Gagaoua, Joe P. Kerry, Anne Maria Mullen, Anna Reiche, Paolo Silacci, Claudia Terlouw

► To cite this version:

Sudipta Hazra, Ruth M. Hamill, Mohammed Gagaoua, Joe P. Kerry, Anne Maria Mullen, et al.. Influence of horn status on bovine Semitendinosus muscle proteome: An analysis of the canonical pathways, upstream regulators and interactive networks. 51st Annual Food Science and Technology Conference, IFSTI, Dec 2023, Dublin, Ireland. hal-04552688

HAL Id: hal-04552688

<https://hal.inrae.fr/hal-04552688>

Submitted on 19 Apr 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



51st Annual Food Science and Technology Conference

Food Systems in a Changing World- the Science, Challenges and Opportunities

December 6-7th,
Teagasc Food Research Centre,
Ashtown, Dublin



IUFoST



Influence of horn status on bovine *Semitendinosus* muscle proteome: An analysis of the canonical pathways, upstream regulators and interactive networks

Author Information

Sudipta Hazra^{1,2}, Ruth M. Hamill¹, Mohammed Gagaoua³, Joe P. Kerry², Anne Maria Mullen¹, Anna Reiche⁴, Paolo Silacci⁴ and Claudia Terlouw⁵

¹ Food Quality and Sensory Science Department, Teagasc Food Research Centre, Ashtown, D15 DY05 Dublin, Ireland

² School of Food and Nutritional Sciences, College of Science, Engineering and Food Science, University College Cork, T12 K8AF Cork, Ireland

³ PEGASE, INRAE, Institut Agro, 35590 Saint-Gilles, France

⁴ Ruminant Nutrition and Emissions, Agroscope Posieux, Switzerland

⁵ Université Clermont Auvergne, INRAE, VetAgro Sup, UMR Herbivores, F-63122 Saint-Genès-Champagnelle, France

Abstract

Stress levels within bovine animals are influenced by several factors, one of which is the horn status of the animals in the herd. Previous research from our group identified 95 differentially abundant proteins (DAP) in *M. semitendinosus* in relation to horn status, when other factors were controlled [1]. Here, a bioinformatics analysis was applied to these DAP to identify canonical pathways, upstream regulators and interactive networks associated with differences in bovine horn status. The most activated canonical pathway in muscle from disbudded bovines was “Glycolysis I” with 10 of 27 proteins of this pathway observed, all more abundant in muscle of disbudded animals. Several pathways were inhibited in disbudded animals, including some involving proteasome proteins (“Regulation of Apoptosis” and “Metabolism of Polyamines”). The most significant upstream regulators were DMD (predicted activated, Z-score +2.6) and sesaminol (predicted inhibited, Z-score -3.2). The most activated and inhibited upstream transcription factors were MEF2C (Z-score +2.4) and NFYC (Z-score -2.8), respectively. The results demonstrate that energy metabolism was enhanced in disbudded animals, while protein turnover was inhibited. Previous studies also showed that the presence of horns has implications for the animal’s metabolic and physiological status. More specifically, horn growth is dependent upon the presence of various hormones associated with energy metabolism, including; growth and thyroid hormones, prolactin, and testosterone. Horn status appears to influence overall energy and physiological status of the animal, with consequences for specific metabolic muscle characteristics as evidenced in this study.

Reference:

Ben Mbarek, R., Terlouw, C., Hamill, R., Kerry, J., Picard, B., Mullen, A.M., Reiche, A-M., Silacci, P. & Gagaoua, M. (2022). Effect of rearing practices and pre-slaughter handling on the *Longissimus thoracis* and the *Semitendinosus* muscle proteomes of young bulls. *In*: Proceedings of the 68th International Congress of Meat Science and Technology, pp. 168. 22th to 25th August, Kobe, Japan.